



# Kent Academic Repository

Deave, Toity, Kendall, Sally, Lingam, Raghu, Day, Crispin, Trudy, Goodenough, Bailey, Elizabeth, Ginja, Sam, Nightingale, Sam and Coad, Jane (2018) *A study to evaluate the effectiveness of Best Beginnings' Baby Buddy phone app in England: a protocol paper*. Primary Health Care Research and Development, 20 . ISSN 1463-4236.

## Downloaded from

<https://kar.kent.ac.uk/68537/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1017/S1463423618000294>

## This document version

Author's Accepted Manuscript

## DOI for this version

## Licence for this version

UNSPECIFIED

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

## Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

# **A study to evaluate the effectiveness of Best Beginnings' Baby Buddy phone app in England: a protocol paper**

(Short title: Bumps And BaBies Longitudinal Study – BaBBLeS)

Authors: Deave T<sup>1</sup>, Kendal S<sup>2</sup>, Lingam R<sup>3</sup>, Day C<sup>4</sup>, Goodenough T<sup>1</sup>, Bailey E<sup>5</sup>, Ginja S<sup>6</sup>,  
Nightingale S<sup>5</sup>, Heeley M<sup>1</sup>, Coad J<sup>5</sup>.

Corresponding author: Toity Deave<sup>1</sup>

Address:

Centre for Child & Adolescent Health  
University of the West of England, Bristol  
Faculty of Health & Applied Sciences  
Oakfield House  
Oakfield Grove  
Bristol BS8 2BN, UK.  
Tel: 0117 3314085  
Email: [toity.deave@uwe.ac.uk](mailto:toity.deave@uwe.ac.uk)

Word count: 3964

2. Centre for Health Services Studies, University of Kent, Canterbury, UK.
3. School of Women's & Children's Health, University of New South Wales, Australia.
4. Child & Adolescent Mental Health Service Research Unit, Guy's Munro Centre, London, UK.
5. Centre for Technology Enabled Health Research, Coventry University, Coventry, UK.
6. Institute of Health & Society, Newcastle University, Newcastle upon Tyne, UK

# **A study to evaluate the effectiveness of Best Beginnings' Baby Buddy phone app in England: a protocol paper**

(Short title: Protocol for the Bumps And BaBies Longitudinal Study – BaBBLeS)

## **Abstract**

**Introduction:** Developments in information and communication technologies have enabled electronic health and seen a huge expansion over the last decade. This has increased the possibility of self-management of health issues.

**Purpose:** to assess the effectiveness of the Baby Buddy app on maternal self-efficacy and mental wellbeing three months post-birth in a sample of mothers recruited antenatally. In addition, to explore when, why and how mothers use the app and consider any benefits the app may offer them in relation to their parenting, health, relationships or communication with their child, friends, family members or health professionals.

**Methods:** We will use a mixed-methods approach, a cohort study, a qualitative element and analysis of in-app data. Participants will be first-time pregnant women, aged 16-years and over, between 12-16 weeks gestation and recruited from five English study sites.

**Evaluation plan:** we will compare maternal self-efficacy and mental health at 3 months post-delivery in mothers who have downloaded the Baby Buddy app compared to those that have not downloaded the app, controlling for confounding factors. Women will be recruited antenatally between 12-16 weeks gestation. Further follow-ups will take place at 35 weeks gestation and 3 months post-birth. Data from the cohort study will be supplemented by in-app data that will include, for example, patterns of usage. Qualitative data will assess the impact of the app on the lives of pregnant women and health professionals using both focus groups and interviews.

**Ethics:** approval from the West Midlands-South Birmingham Research Ethics Committee (NRES) (16/WM/0029) and the University of the West of England, Bristol, Research Ethics Committee (HAS.16.08.001).

**Dissemination:** findings of the study will be published in peer reviewed and professional journals, presented locally, nationally and at international conferences. Participants will receive a summary of the findings and the results will be published on Best Beginnings' website.

**Key words:** pregnancy, Best Beginnings, evaluation, parenting, app.

## Introduction

There has been a huge increase in the last 10 years of electronic health (e-Health) and mobile health (m-Health) solutions to increase self-management of health problems in high, medium and low income countries (Goetz *et al.*, 2017). This has been made possible by developments in information and communication technologies. Mobile phones have capabilities that were, previously, only available by using computers and portable devices, such as laptops. Access to the internet is now widespread and this has underpinned the development of ‘applications’ or apps; these have led to a fundamental change in the methods that people use to access information. Apps offer wider opportunities for the provision of healthcare and information than have ever been known before. This has also brought about changes in the demands that patients make of healthcare services (Hussain *et al.*, 2015).

The population of app users includes expectant and new mothers with whom internet-based information about pregnancy has been popular for nearly a decade (Bernhardt & Felter 2004, Declerq *et al.*, 2013). In 2013, around 1500 pregnancy apps were available on the Google Play app store and the Apple iStore (Tripp *et al.*, 2014) and there are likely to be many more if this search was repeated. Whilst e-Health and m-Health have the potential for enhancing traditionally delivered healthcare services (Tripp *et al.*, 2014, Lau *et al.*, 2015), little is known about the use and effectiveness of pregnancy- or perinatal-related smartphone apps (Rodger *et al.*, 2013). A systematic review of apps for women’s health concluded that more research was needed, especially in relation to the use and embedding of apps in healthcare settings (Derbyshire & Dancey, 2013). Most expectant women access the internet daily (Waring *et al.*, 2014) and pregnant women who were asked about using the internet for searching for information perceived the information that they accessed to be reliable but they had not discussed them with their midwife (Larsson, 2009). The most popular apps by consumer rating were those that were interactive (Tripp *et al.*, 2014).

The Baby Buddy app was launched in England in November 2014. It was developed by the national child health and wellbeing charity, ‘Best Beginnings’, who describes the app as ‘an electronically delivered health intervention’ that is intended to support and guide women through pregnancy (antenatal side of the app) and the first six months of their child’s life (postnatal side of the app). Best Beginnings designed the app to inform and empower women of all ages and from wide sociodemographic backgrounds so that all expectant parents could search for and benefit from high quality information and support during pregnancy. Many women may access information online that is unreliable (Rodger *et al.*, 2013) but Best Beginnings worked closely with health professionals to ensure that Baby Buddy provides evidence-based, trustworthy information (<https://www.bestbeginnings.org.uk/vision>). It provides interactive information to help parents look after their own physical and mental health during pregnancy and early parenthood and support them with caring for the physical and emotional health and wellbeing of their child. It was designed to complement maternity and postnatal services. The content of the app is written so that it can be understood by anyone with a reading age of 11 years or above with a ‘read aloud’ element available. It can be downloaded free of charge from the Apple iStore and the Google Play app store by anyone with a suitable smartphone, including fathers and other family members or friends.

The Baby Buddy app has been developed specifically to target expectant mothers who are under 25 years-old but also be attractive to wide range of mothers. Young (teenage) mothers

in the UK are less likely to both engage with formal maternity service provision early in their pregnancy and attend their appointments (Department of Health 2009; Bradshaw *et al.*, 2014); these are risk factors for maternal and infant mortality (Confidential Enquiry into Maternal and Child Health 2007).

The BaBBLeS study (Bumps And BaBies Longitudinal Study) aims to assess the effectiveness of the Baby Buddy app on improving maternal self-efficacy and mental wellbeing. It also aims to understand when, why and how mothers use the app and any benefits the app may offer them in relation to their parenting, health, relationships or communication with their child, friends and family members or health professionals.

## **Objectives**

1. To assess the effectiveness of the Baby Buddy app, at three months post-delivery on maternal self-efficacy and well-being in those mothers who used the Baby Buddy app compared to those who did not use the app;
2. To assess the effectiveness of the Baby Buddy app, at three months post-delivery, on maternal self-efficacy and well-being in those mothers who were shown how to use the app by a health professional compared to those who were not shown or did not download it;
3. To explore how the app has affected the day-to-day lives of participant mothers, specifically around their self-efficacy, parenting ability, health behaviour, interactions and communications with their friends, family and health professionals;
4. To obtain in-depth information from health professionals around their awareness of the Baby Buddy app and barriers and facilitators to integration of the app into routine health care.
5. To describe data on the uptake, patterns of usage and detailed analytics of key factors within the app.

## **Methodology**

This longitudinal study will be conducted, in parallel, in five geographical sites in England and will employ a mixed methods approach. These areas have been chosen since they are geographically, ethnically and socio-economically diverse and are areas where the app was reported to be well-embedded.

This study has three component parts that address the stated objectives: a cohort study, a qualitative element and analysis of in-app data. The cohort study will compare the self-reported self-efficacy and mental wellbeing of mothers three months post-delivery who downloaded the Baby Buddy app compared to those who did not download the app. The study will control for baseline characteristics, including demographic characteristics, use of technology, social support and baseline levels of self-efficacy and mental wellbeing. The qualitative element will include mothers and professionals taking part in one-one interviews or focus groups. The in-app data on the uptake, patterns of usage and detailed analytics of key factors within the app will be collected from downloads provided by Best Beginnings (see below).

## **Recruitment**

Recruitment will take place between June and October 2016. Women who aged 16 years and over, with no previous live child and between 12-16 weeks and six days gestation will be identified from the maternity units in each of the five study sites. Maternity unit administrative clerks, with support of the research midwives in each of the five geographical areas will be asked to undertake a database search for women who fulfil the inclusion criteria. Each woman who is identified will be sent or given a participant information booklet combining the study invitation letter and information booklet by the midwifery services staff and opportunities to discuss it by telephone, text or email. Women in three of the sites will also be invited to take part in further, nested qualitative studies. Mothers can choose to complete the baseline questionnaire online or on paper and will be offered a £5 thank you voucher. A two week reminder will be sent to each woman who has not responded.

Midwives and health visitors who work in three study sites will be invited to take part in a focus group between September 2016 and February 2017. If more than 10 health professionals agree to take part in each site purposive sampling will be employed to ensure that all geographical areas are covered.

Mothers and health professionals will be informed they can withdraw from the project at their own request. The mothers will be made aware that it would not affect their future care or support from the health professionals.

## **Data collection**

### **Quantitative data**

Baseline data will be collected at the same time as women agree to participate in the study through completion of a questionnaire. Baseline and follow-up data are being collected by self-completion questionnaire from the mothers. The baseline questionnaire asks questions about the mothers' socio-demographic details, validated tools to measure parenting self-efficacy, anticipated feeding practices, pregnancy dates (Expected Date of Delivery, EDD) use of the internet, apps and social media, sources of information about pregnancy and motherhood and social support.

Two follow-up questionnaires will be administered with reminders, using methods outlined above, at 35 weeks of pregnancy and at three months post-birth. The 35 week questionnaire will repeat the validated tools with a postal reminder. At three months post-birth, the questionnaire will repeat the validated tools and will also include questions relating to app usage, health service use, interaction with health services, childbirth experience; the initial reminder will be by telephone. Quantitative data will be entered onto one database at the University of the West of England, Bristol and stored on the university server.

### **Qualitative data:**

For those women who agree to take part in the qualitative study (as part of the recruitment process in the three selected evaluation sites), they will be invited to take part in in-depth semi-structured interviews and/or a focus group (up to 10 interviews per three geographical sites and to attend with their partner/chosen friend. The interviews will be offered both via the telephone or face-to-face in each geographical area as per individual preference. Their data will be recorded digitally and transcribed verbatim. Data from the one-to-one interviews

will be structured in a similar way as the focus group discussions to ensure congruence of data.

For the focus groups, an Appreciative Inquiry (AI) approach will be used with prompts, as developed by the research team and advisors. This element will aim to explore when, why and how mothers use the app and the perceived benefits the app gives them in relation to their parenting, health, relationships and communication with their child, friends and other family members. It will specifically explore how the app is used by mothers to enhance communication with healthcare providers to make “every contact count” (PHE, 2016) and the reasons behind differences in rates of usage, including what makes the app enjoyable to use and responses to embedding activities.

One-one interviews and focus groups will take place between three and six months post-delivery; focus groups will be 'baby friendly' to accommodate additional needs.

#### Health professionals

Health professionals' awareness of the Baby Buddy app and the barriers and facilitators to the integration of the app to usual service delivery will be collected.

Midwives and health visitors in the selected three study sites will be invited to take part in the study; the sample will be up to 10 professionals in each area (n = 30). Their views will also be sought to explore any 'added value' of the apps in supporting mothers, partners/supporters and their babies. Either face-to-face or telephone interviews and/or focus groups will be offered to professionals in each chosen geographical area between April 2017 and October 2017. Their data will be recorded digitally and transcribed verbatim. They will be encouraged to contribute to a shared, integrated perspective in evaluating the app of 'what works well' and 'what could work better' in integrating the use of the app into healthcare encounters. This is an important aspect of the study, in which the process of using the app will be explored. We will also determine preference/choice through an action learning set approach.

#### In-app data

At the 35 week gestation data collection, mothers will receive an information sheet and consent form to take part in this element of the study. The majority of the patterns of Baby Buddy app use are recorded and stored on secured databases hosted by Best Beginnings as part of a standard procedure necessary for managing and debugging the app. Best Beginnings will provide the research team with limited and secured download access to the database with the ability to obtain specific in-app data from the users of the app. These specific data will include information such as duration of sessions using the app, count of these sessions, flow of app use and general user information. The obtained data will be downloaded and stored in the encrypted format on highly secured university computers. Once all data collection is complete, relevant data will be merged onto one database prior to analysis but with an appropriate encoding of personal information (i.e. anonymised ID codes, not email addresses or postcodes) in order to preserve high level of user confidentiality and anonymity.

## **Outcome measures**

One main aim for the app is that it increases mothers' confidence and self-efficacy with regard to pregnancy, baby-care and early parenthood. Therefore, our primary outcome measure was chosen to reflect this, the Tool to measure Parenting Self-Efficacy (TOPSE) (Kendall and Bloomfield, 2005, Bloomfield and Kendall 2007). The theoretical underpinning of TOPSE is based on the self-efficacy theory developed by Albert Bandura (Bandura, 1982; 1986; 1989). TOPSE measures change in self-efficacy between time-points and is therefore ideally placed to measure outcomes in this study. The 0-6 month version of TOPSE will be adapted, with the author's help, to measure parenting self-efficacy expectations during pregnancy. TOPSE is a multi-dimensional instrument of 48 statements within eight scales, each scale having six statements and representing a distinct dimension of parenting: emotion and affection, play and enjoyment, empathy and understanding, control, discipline and boundaries, pressures, self-acceptance, learning and knowledge. The items are rated on an 11-point Likert scale where 0 represents completely disagree and 10 represents completely agree. The scale contains positive and negatively worded items and the responses are summed to create a total score; the lower the score, the lower the level of parenting self-efficacy. Internal reliability coefficients for the subscales ranged from 0.80 to 0.89 and the overall scale reliability was 0.94. External reliability coefficients ranged from  $r_s = 0.58$  ( $n=19$ ,  $p<0.01$ ) to  $r_s = 0.88$  ( $n=19$ ,  $p<0.01$ ).

The secondary outcomes are the Warwick-Edinburgh Mental Well-Being Scale (WEMWBS) and the in-app data. The WEMWBS is a 14 item scale of mental well-being covering subjective well-being and psychological functioning, in which all items are worded positively and address aspects of positive mental health (Tennant et al., 2007). The scale is scored by summing responses to each item answered on a 1 to 5 Likert scale. The minimum scale score is 14 and the maximum is 70. WEMWBS has been validated for use in the UK with those aged 16 and above. From the in-app data, the uptake, patterns of usage and detailed analytics of key factors within the app will be analysed (see below). This scale comprises of 14 statements describing feelings (eg., 'I have been feeling useful') and functional aspects (eg., 'I've been dealing with problems well') over the previous two weeks. Items were scored from 1 (none of the time) to 5 (all of the time) and summed to provide an overall score between 14 and 70, where higher scores corresponded to greater frequency. WEMWBS has demonstrated good content and criterion-related validity and high test-retest reliability (0.83) when used in different samples and public health contexts, including parenting programmes (Stewart-Brown et al., 2011).

Additional data to be collected and measures to be used include the mother's sociodemographic factors (age, ethnic group, highest level of education attained, relationship status and employment), intention and actual breastfeeding, the multidimensional scale of perceived social support (MCSDS)(Zimet *et al.*, 1988) and use of social media and technologies, adapted from the Media and Technology Usage and Attitudes Scale (Rosen *et al.* 2013). In both the second and final questionnaires, mothers will be asked whether they had downloaded and/or used the Baby Buddy app and how they had heard about it (source of information). In the final, three month post-delivery questionnaire, the Childbirth Experience Questionnaire (Walker *et al.*, 2015) is included.

## **Sample sizes**

Quantitative



The average percentage of app download across our five sites is 13.6%, based on data provided by Best Beginnings. Assuming that the percentage of mothers who downloaded the app is slightly below that rate, at 12.5% and assuming a 30% loss to follow-up, the study will need to recruit 559 women into the study to have 90% power to detect a 0.5 SD difference in the overall TOPSE (primary outcome) across domains. If the proportion of mothers in the final sample who downloaded the app changes then the required sample size will change accordingly.

#### Qualitative

The sample sizes for the qualitative study are based on an estimate to expect saturation of data on themes emerging from the study. A maximum of 30 in-depth interviews (n=30, 10 from each of the selected three sites) will be conducted with women who have used the app and who have consented to participate. There will be three focus groups (n=30 mothers), one group per three sites with 10 attendees in each group. A balanced group of recruits to interviews and focus groups will be spread across the three sites to allow for local differences in experience with the app use and professional encounters.

The same rationale is used for health professionals' views (n=30, 10 from each of three sites). The invitation to health professionals will be to attend a focus group with the option of telephone interview if unable to attend groups and to ensure the target sample size is achieved. This will facilitate reaching data saturation against the backdrop of 'real world' demands on professionals' time and availability.

#### Data analysis

##### Quantitative data:

Descriptive statistics will be used to describe the sample, including the mothers' age, socio-demographic, ethnicity, access to and use of technology and the overall scores for the outcome measures. Linear regression models will be used to compare primary and secondary outcomes in mothers who used the app compared to those who did not use the app, depending on outcome data distribution. If outcome data are not normally distributed, the possibility of logistic regression analysis will be explored, using binary outcomes. Key baseline demographic variables will be used to control for potential confounding variables, including maternal age, education, employment, relationship status, recruitment site, social support, general technology use and use of other pregnancy apps. Baseline levels of the outcome variables, e.g., maternal self-efficacy, will be controlled for in the final analysis. Analysis will be as per protocol and analysis plan. Regression methods will control for baseline demographic characteristics. All analyses will be carried out using Stata software. Missing data will not be accounted for using multiple imputation, unless 30% of the data or more is missing.

A second analysis will compare primary and secondary outcomes, as stated above, between those mothers who heard about the app from a health professional (embedded use) and those women who did not hear about it or who did not download the app by three months post-delivery. The above statistical methods will be used again to compare outcomes between 'download and heard about the app from healthcare professional' and 'downloaded but did not hear about the app from a healthcare professional, or did not download it'.

##### Qualitative data:

Divergence and concordance of opinion will be explored in the data from the interviews and focus groups.

#### **In-app data:**

Data orientation will be undertaken and then formatted for analysis. This will include an exploratory analysis of the basic socio-demographic information and profiling of app users (e.g. age, occupation, education, ethnic origin); a description of the patterns of app use: the creation of the avatar; uses of the goal setting function, 'ask me a question'; 'what does that mean; the media downloaded; a detailed analysis of interactions between various factors (e.g. age by frequency of use, location by number of downloads).

The findings from each section of the study will be discussed with the user group and their views of interpretation from their perspectives will be sought. A synthesis of all the information collected will be collated to form the final report.

#### **Anonymity and data protection**

All participants are being assigned a unique identity code (UIC) to facilitate the conduct and analysis of the study. These will be used on all paper and electronic versions of questionnaires, interview transcriptions and databases. Recruitment logs will serve as a separate confidential record of the participants' details. This will permit identification of all participants recruited to the study, in accordance with regulatory requirements and for follow-up, as required. Two logs will be created: one for women participants and one for the health professionals.

In addition, the following data will be collected and stored on a separate database: mother's UIC, name, address, telephone and other relevant contact details, e.g., date follow-up questionnaires due, preferred format for administration of follow up questionnaires, email address.

For the interview data, with the permission of the interviewees, interviews will be digitally audio-recorded. They will be anonymised, transcribed verbatim and stored with the other study data. The transcribed interviews will be treated as confidential documents and held securely with no identifiable information on the transcripts. These will be stored on a secure password protected University server.

All data collected will be held securely, in a locked room, or locked cupboard or cabinet. Access to the information will be limited to the study research team. Computer held data, including study databases, will be held on secure, password protected University or NHS servers and will be backed up as a minimum, every 24 hours. Personal laptops will not be used. Where University or NHS laptops are used, these will be password protected and data will be transferred to the University/NHS server as soon as possible. At the end of the study, data from all centres will be archived by the University of the West of England, Bristol (UWE) for a period of seven years or longer, if required, as set out in UWE's Code of Research Conduct.

#### **Adverse events**

The occurrence of adverse events, as a result of participation in this study, is not anticipated. We believe the risks for participants in this study are minimal. However, should an adverse

event occur this will be recorded and reported to the Chief Investigator. Any adverse events will also be reported in the final study report.

### **Ethical and regulatory aspects**

This study has received a favourable opinion from the NHS Research Ethics Committee (NRES) West Midlands-South Birmingham REC (16/WM/0029), the University of the West of England, Bristol Research Ethics Committee (HAS.16.08.001), and the respective study site's National Health Service (NHS) Research & Development (R&D) departments.

### **Strengths and limitations of the study**

#### **Strengths**

- Multi-centre longitudinal study
- Large sample size of over 500 participants to be recruited
- We are using mixed methods to help assess and explore the impact of the Baby Buddy app as well as how the app is being used by women and health professionals
- The primary outcome (TOPSE, parenting self-efficacy) is generic across all domains of parenting over time

#### **Limitations**

- The ante-natal version of the TOPSE has been adapted for this study.

### **Publication and Dissemination Policy**

The research team regard active, widespread dissemination of outputs from this research as essential and this will include peer reviewed and professional journals, conference presentations and national and local seminars. The research team members will contribute towards drafting the papers, reporting the findings of the study and will be named authors on those papers, providing they fulfil the Vancouver criteria for authorship. If a member of the research team wishes to analyse and report other findings from the study they can do so on the agreement of the other research team members.

The contribution of Best Beginnings and the funders will be acknowledged in any publication of dissemination of information about the study.

### **Authors' contributions**

TD led the process of writing the protocol. All authors had the opportunity to feedback prior to submission to the NRES. All authors have had input to the current paper.

### **Funding statement**

This work was supported by the Big Lottery via Best Beginnings as a competitive tender.

**Competing interests:** the authors have no competing interests.

## References

Bandura A. Self-efficacy mechanism in human agency. *American Psychologist* 1982;37:122–147.

Bandura A. Regulation of cognitive processes through perceived self-efficacy. *Developmental Psychology* 1989,25:729–735.

Bandura A. *Exercise of personal and collective efficacy in changing societies*. In *Self-Efficacy in Changing Societies* (Bandura A., ed), Cambridge University Press, Cambridge, 1995:1–45.

Bernhardt J, Felter E. Online Paediatric Information Seeking Among Mothers of Young Children: Results from a Qualitative Study Using Focus Groups. *Journal of Medical Internet Research* 2004;6(1):e7. Online. Available at <http://www.jmir.org/2004/1/e7>. Accessed 31/07/2017.

Bloomfield L, Kendall S. Testing a parenting programme evaluation tool as a pre-and post-course measure of parenting self-efficacy. *Journal of Advanced Nursing* 2007;60(5):487-493.

Bradshaw P, Schofield L, Maynard L. *The experiences of mothers aged under 20: Analysis of data from the Growing Up in Scotland Study*. Scottish Government, 2014. Online. Available at <http://www.gov.scot/Publications/2014/02/8789>. Accessed 31/07/2017.

Declercq E, Sakala C, Corry MP, et al. *Listening to mothers III: Report of the third national U.S. survey of women's childbearing experiences*. New York, NY: Childbirth Connection 2013. Available from:<http://transform.childbirthconnection.org/reports/listeningtomothers/>

Derbyshire E, Dancey D. Smartphone Medical Applications for Women's Health: What Is the Evidence-Base and Feedback? *International Journal of Telemedicine and Applications* 2013;Vol.2013, article ID 782074, 10pages. doi.org/10.1155/2013/782074. Accessed 31/07/2017.

Department of Health. *Getting maternity services right for pregnant teenagers and young fathers*. Revised edition. Department of Health, London, 2009. Online. Available at <http://www.education.gov.uk/publications/eOrderingDownload/DCSF-00673-2009.pdf>. Accessed 31/07/2017.

Goetz M, Müller M, Matthies LM, et al. Perceptions of Patient Engagement Applications During Pregnancy: A Qualitative Assessment of the Patient's Perspective. *JMIR Mhealth Uhealth* 2017,5(5):e73. DOI: 10.2196/mhealth.7040.

Hussain M, Al-Haiqi A, Zaidan A, et al. The landscape of research on smartphone medical apps: Coherent taxonomy, motivations, open challenges and recommendations. *Computer Methods Programs in Biomed* 2015; pii: S0169-2607(15)00225-4.

Kendall S, Bloomfield L. Developing and validating a tool to measure parenting self-efficacy. *Journal of Advanced Nursing*, 2005, 51(2): 174-181.

- Lau Y, Htun TP, Tam WSW, Klainin-Yobas P. Efficacy of e-technologies in improving breastfeeding outcomes among perinatal women: a meta-analysis. *Maternal & Child Nutrition* 2016;(12):381-401. doi: 10.1111/mcn.12202.
- Larsson M. A descriptive study of the use of the Internet by women seeking pregnancy-related information. *Midwifery* 2009;25(1):14–20.
- Rodger D, Skuse A, Wilmore M, *et al.* Pregnant women's use of information and communications technologies to access pregnancy-related health information in South Australia. *Australian Journal of Primary Care* 2013, <http://dx.doi.org/10.1071/PY13029>.
- Rosen LD, Whaling K, Carrier LM, *et al.* The media and technology usage and attitudes scale: An empirical investigation, *Computers in Human Behavior* 2013;29(6):2501-2511.
- Stewart-Brown SL, Platt S, Tennant A, *et al.* The Warwick-Edinburgh Mental Well-being Scale (WEMWBS): a valid and reliable tool for measuring mental well-being in diverse populations and projects. *Journal of Epidemiology & Community Health* 2011;65(Suppl 2):A38-A39.
- Tennant R, Hiller L, Fishwick R, *et al.* The Warwick-Edinburgh mental well-being scale (WEMWBS): development and UK validation. *Health and Quality of life Outcomes* 2007;5(1):63
- Tripp N, Hainey K, Liu A, Poulton A, *et al.* An emerging model of maternity care: smartphone, midwife, doctor? *Women Birth* 2014;27(1):64-67.
- Walker KF, Wilson P, Bugg GJ, *et al.* Childbirth experience questionnaire: validating its use in the United Kingdom. *BMC Pregnancy and Childbirth* 2015;15:86. doi:10.1186/s12884-015-0513-4.
- Waring ME, Moore Simas TA, Xiao RS, *et al.* Pregnant women's interest in a website or mobile application for healthy gestational weight gain. *Sexual Reproductive Healthcare* 2014;5(4):182–184. doi:10.1016/j.srhc.2014.05.002.
- Zimet GD, Dahlem NW, Zimet SG, Farley GK. The multidimensional scale of perceived social support. *Journal of Personality Assessment* 1988;52(1):30-41.