



Kent Academic Repository

Hilton, Kira, Tolley, Howard, Ortega-Roldan, Jose L., Thompson, Gary S., Sutton, Prof. Mark, Hind, Charlotte K. and Hiscock, Jennifer R. (2023) *Phospholipid headgroup composition modulates the molecular interactions and antimicrobial effects of sulfobetaine zwitterionic detergents against the "ESKAPE" pathogen Pseudomonas aeruginosa*. *Chemical Communications*, 59 (70). pp. 10504-10507. ISSN 1359-7345.

Downloaded from

<https://kar.kent.ac.uk/102336/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1039/D3CC02320F>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

For the purpose of open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

COMMUNICATION

Phospholipid headgroup composition modulates the molecular interactions and antimicrobial effects of sulfobetaine zwitterionic detergents against the “ESKAPE” pathogen *Pseudomonas aeruginosa*[†]

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Kira L. F. Hilton,^a Howard Tolley,^b Jose L. Ortega-Roldan,^a Gary S. Thompson,^a J. Mark Sutton,^{b,c} Charlotte K. Hind^{*b} and Jennifer R. Hiscock^{*a}

We determine the efficacy for three known structurally related, membrane active detergents against multidrug resistant and wild type strains of *Pseudomonas aeruginosa*. Accessible solution state NMR experiments are used to quantify phospholipid headgroup composition of the microbial membranes and to gain molecular level insight into antimicrobial mode of action.

Antimicrobial resistance (AMR) is a significant threat to world health,^{1,2} with bacteria now found to be resistant to all currently marketed antimicrobials,³ including the membrane-active peptide, colistin, one of the antibiotics of last resort.⁴ One mechanism of AMR which remains less well understood, involves alteration of the phospholipid composition of the bacterial membrane,^{5,6} which decreases effective drug delivery^{7–9} and/or alters the drug target.^{10–12} For example, daptomycin susceptible (S447) and resistant (R446) strains of *Enterococcus faecium* exhibit different phospholipid membrane profiles: PG:L-PG:CL:DAG 34:14:39:13 (S447) and PG:L-PG:CL:DAG 15:16:47:23 (R446).¹³ These differences in phospholipid content were paired with increased daptomycin resistance, which was thought to be due to increased membrane rigidity and changes in biophysical properties. As is common for studies of this type, the specific interactions between the antimicrobial agent and phospholipids within the cell membrane were not identified (see Figure 1 for a summary of phospholipid headgroup structures). Therefore, the development of methodologies, such as those described herein,

which enable the characterisation of molecular level interaction events with the cell membrane has become vital to inform the identification of antimicrobial therapeutic regimes.




Figure 1 a) General structure of a phospholipid and 1 – 3. Purple = hydrophilic phosphate headgroup, orange = glycerol linking group, pink = hydrophobic hydrocarbon residue, which can differ in chain length and degree of saturation. R groups: b) phosphatidylethanolamine (PE); c) phosphatidylglycerol (PG); d) phosphatidylserine (PS); e) phosphatidylinositol (PI); f) cardiolipin (CL).

Within this proof of principle study, we utilise a unique combination of solution state multinuclear NMR techniques, phospholipid nanodiscs produced from lipids obtained from target bacterial species, alongside standard antimicrobial efficacy studies and membrane fluidity experiments (Section 10[†]). This produces a body of data that enables us to understand the relationship between phospholipid membrane headgroup composition, molecular level membrane interaction events and antimicrobial activity against multiple microbial species taking

^a Division of Natural Sciences, University of Kent, Canterbury, CT2 7NH, UK. Email: J.R.Hiscock@Kent.ac.uk

^b UKHSA, Science Group, Manor Farm Road, Salisbury, SP4 0JG, UK. Email: Charlotte.Hind@UKHSA.gov.uk

^c Institute of Pharmaceutical Sciences, School of Cancer & Pharmaceutical Sciences King's College London, SE1 9NQ Address here.

[†]Electronic Supplementary Information (ESI) available: This includes experimental details, materials and methods, DLS data, microscopy data, multinuclear NMR spectroscopy data, nanodisc characterisation data, ¹H NMR titration data, MIC and MIC₅₀ data, membrane permeabilization and fluidity data. See DOI: 10.1039/x0xx00000x

advantage of widely accessible, automated, NMR based methodologies. This removes limitations such as the need for intrinsic molecular fluorescence and traditional bespoke experimental facilities. To demonstrate this, we target a series of three zwitterionic, sulfobetaine membrane disrupting agents **1** – **3** (Figure 1),¹⁴ which have been known as detergents for biological use. We chose **1** – **3** because of their stepwise molecular structure modification and lack of evidence describing molecular level mode of action.¹⁵

Initially, the minimum inhibitory concentration (MIC – defined as the lowest concentration of a compound required to inhibit visual growth) was determined for **1** – **3** against two strains of *Pseudomonas aeruginosa* (*P. aeruginosa*), Table 1. This pathogen was selected as it is a microbe identified by the WHO as urgently requiring new treatment options.¹⁶ NCTC 13437 and PAO1 are both derived from clinical isolates with differing antimicrobial resistance profiles. Whilst PAO1 is sensitive to the majority of clinically relevant antibiotic classes, NCTC 13437 is resistant to multiple classes including fluoroquinolones, aminoglycosides, carbapenems, cephalosporins and colistin.

Table 1 Top: MIC and MIC₅₀ values determined for **1** – **3** against two strains of *P. aeruginosa*. Values are presented as modal values (n = 3), or as a range due to the unusual concentration ranges needed to inhibit microbial growth (Figures S71 and S72, and Figure 2, line graphs). An MIC₅₀ value was calculated where an MIC could not be determined. Bottom: Total phospholipid headgroup composition data for lipids extracted from the cell membranes of PAO1 and NCTC 13437.

Bacterial Strain	MIC (mM)			MIC ₅₀ (mM)		
	1	2	3	1	2	3
PAO1	12.5	>100	100	3.12	1.56-50	0.4
NCTC 13437	50	>100	100	6.25	0.4	0.8
	Phospholipid headgroup composition (%)					
	PE	PG	PI	PS	CL	Other
PAO1	73.1	19.9	4.3	2.7	0	0
NCTC 13437	53.4	46.6	0	0	0	0

MIC values obtained against both strains of *P. aeruginosa* show this microorganism to be more susceptible to the effects of **1** over **2** and **3**. However, when comparing MIC₅₀ values, **3** is found to inhibit 50 % growth at much lower concentrations when compared to **1**, 0.8 mM and 6.25 mM respectively, against NCTC 13437. Interestingly, there is also some variation between the strains susceptibility for **1** – **3**. For example, the MIC of **1** against PAO1 was 12.5 mM, whilst for NCTC 13437, the MIC of **1** increased to 50 mM.

To confirm the membrane disruption mechanism of action for **1** – **3** against PAO1 and NCTC 13437, an outer membrane permeabilisation assay was performed (Figure 2, bar charts). Here, *N*-phenyl-naphthylamine (NPN) acts as a fluorescent reporter. When the outer membrane is intact, this hydrophobic reporter is excluded from the microbial membrane. However, should the microbial membrane become disrupted, the NPN reporter is able to enter the membranes hydrophobic environment resulting in an increased fluorescence signal. We observe a definitive inverse relationship between membrane

permeabilisation and percentage growth for **1**, as shown in Figure 2. This relationship was also observed for **2** and **3** however, the relationship is less obvious due to the unusual plateau effect associated with bacterial growth. Therefore, these data support the hypothesis that membrane disruption is the mechanism of action for **1** – **3**, as previously reported.¹⁴ These findings are further supported by the results of scanning electron microscopy (SEM), Section 12[†], where holes in the bacterial membranes of NCTC 13437 were identified in the presence of **1** – **3**.




Figure 2 The outer membrane permeabilization (bar chart, right hand y-axis) of a) PAO1 and b) NCTC 13437 when exposed to **1** (purple), **2** (orange) and **3** (pink), compared to the percentage endpoint growth for PAO1 and NCTC 13437 after treatment with **1** – **3** (line graph, left hand y-axis) compared to an untreated bacterial control.

In addition, **1** shows a correlation between increasing concentration and increasing percentage membrane permeabilization against both strains of *P. aeruginosa*. However, **2** and **3** show limited increases in percentage membrane permeabilization and growth at concentrations above 0.1 mM. We believe that this observation is due to the critical aggregation concentration (CAC) of **2** and **3**, 0.1 mM¹⁷ and 0.4 mM¹⁸ respectively. The CAC is defined as the concentration at which any additional compound added to a solution will result in the formation of higher order self-associated aggregates, e.g. micelles.¹⁹ Therefore, at concentrations above CAC, free **1** – **3** will become incorporated into aggregated species. In the case of **2** and **3**, we believe this limits the concentration of compound available to interact with the microbial membranes and thus limits the activity of these agents at concentrations above this value, explaining the differences in antimicrobial efficacy identified from MIC as opposed to MIC₅₀ values (Table 1). Compound **1** exhibits a much higher CAC at 6.4 mM²⁰ and thus the antimicrobial activity of this compound is less affected by increasing concentration.

To determine if there was any difference in the total phospholipid membrane headgroup composition of these two strains of *P. aeruginosa*, the membranes were extracted,²¹

analysed using 2D ^1H - ^{31}P HSQC NMR experiments, and compared to previously published control data.²² Here we observe clear differences in the phospholipid headgroup composition of the membranes extracted from wild type PAO1 and multidrug resistant NCTC 13437 (Table 1), meaning that changes in the phospholipid headgroup composition of microbial membranes could be a cause for, or a result of, the AMR observed for the NCTC 13437 strain of *P. aeruginosa*. For example, the presence of PS (2.7 %) and PI (4.3 %) could only be identified with PAO1. However, the greatest differences in total membrane phospholipid headgroup composition were identified when considering the percentage of PE (~20 % lower for NCTC 13437) and PG (>25 % higher for NCTC 13437) present within the membranes of these two bacterial strains. These substantial differences in phospholipid membrane headgroup composition support a hypothesis that differences in MIC values obtained for detergents **1** – **3** could be due to changes in phospholipid membrane headgroup composition.

To test the validity of this hypothesis, and confirm if alterations in phospholipid headgroup composition alter the molecular interaction events of **1** – **3**, we performed a series of ^1H NMR phospholipid nanodisc experiments. To date, phospholipid nanodiscs have been mainly used to enable the study of membrane bound proteins under native conditions.^{23–25} However, we have also recently expanded the use of these systems to enable the study of small molecule-phospholipid membrane interaction events, adapting traditional ^1H NMR titration based methodologies (Section 7 – 9).^{26,27,28} Absolute integration of ^1H NMR resonances, corresponding to the different ^1H environments of **1** – **3**, and subsequent fitting of these data to Hill plot kinetics using Origin 2018 software,^{28,29} enabled the effective concentration of nanodisc needed to coordinate 50 % of **1** – **3** (EC_{50}) at 100 μM to be determined (Figure 3). Put simply the greater the $1/\text{EC}_{50}$ the greater the affinity of **1** – **3** for the phospholipid nanodisc. When comparing EC_{50} values generated from different residues of the same molecules, we may gain some insight into the mode of action for a specific agent. For example, the sterol head group of **1** (Figure 3a, resonances d – f) was shown to exhibit a greater affinity towards both types of phospholipid nanodiscs, compared to the polar tail group (Figure 3a, resonances a – c). Thus, we hypothesise that the lipophilic sterol group is able to embed itself into the nanodisc, leaving the tail free to rotate within the solution. In addition, we observed increased $1/\text{EC}_{50}$ values for **1** against nanodiscs derived from PAO1 membranes, indicating a stronger interaction, which correlates with increased antimicrobial efficacy for **1** against this same strain of *P. aeruginosa* and changes in phospholipid headgroup composition (Table 1).

With the sterol group in **1** substituted for the alkyl group in **2** and **3**, the disparity in interaction between the two ends of the antimicrobial agent decreases. The step-wise modification between **2** and **3** shows that the amide linker of **2** is responsible for the specificity of this detergent molecule for nanodiscs formed from PAO1 over NCTC 13437 membranes, and therefore believe that introduction of this functionality results in decreased affinity for PG phospholipids. However, there is no

longer a correlation observed between increased antimicrobial interaction and increased antimicrobial efficacy, attributing this effect to **1** only, pointing to differing modes of action for this group of three antimicrobial agents.




Figure 3 Graphs summarizing $1/\text{EC}_{50}$ (μM^{-1}) values obtained from the fitting of a) **1**, b) **2** or c) **3** nanodisc titration data to Hill Plot kinetics using Origin 2018 software.²⁹ V_{max} was fixed to 100 %. Where co-ordination did not reach 50 %, data was not fitted. Purple = results from PAO1 nanodiscs ^1H NMR titration studies, orange = results from NCTC 13437 nanodisc ^1H NMR titration studies. The Hill coefficient (n) produced through the fitting of this data provides a measure of the cooperative/non-cooperative nature of binding events.³⁰ As shown in Table S2-S4 and Section 9,[†] the Hill coefficients generated from these data sets exhibit a value $n > 1$ in general, providing evidence that these phospholipid bilayer binding events are co-operative. Fitting these data to a simple binding isotherm was not possible due to the suspected complexity of interactions present between **1** – **3** and nanodiscs derived from PAO1 and NCTC 13437.

In conclusion, we establish the efficacy of membrane active antimicrobial agents, **1** – **3**, against different strains of the “ESCAPE” pathogen *P. aeruginosa*, adding molecular level insight to the mode of antimicrobial action for these agents for the first time. We show differences in the phospholipid membrane headgroup composition of wild type PAO1 and multidrug resistant clinical isolate NCTC 13437 and, through the incorporation of these phospholipids into nanodiscs, observe molecular level antimicrobial co-ordination events for **1** – **3**. This leads us to hypothesise that the sterol unit of **1** embeds into the nanodisc, leaving the hydrophilic tail substituent free to interact with the solution environment. Interestingly, the increased

strength of the sterol headgroup interaction events was also found to correlate with increased antimicrobial efficacy. Finally, we also suggest that decreased CAC may also limit the antimicrobial efficacy of an agent at concentrations above this value, further demonstrating the need to determine both MIC and MIC₅₀ values for membrane active amphiphilic agents such as **1–3**, dependent on CAC. These findings support the need for ongoing investigations in this area, initially identifying and then quantifying the role changes in phospholipid headgroup composition plays in AMR. This work was supported by JH's UKRI Future Leaders Fellowship (MR/T020415/1), the University of Kent and UKHSA.

Author Contributions

KLHF: Investigation; Validation; Writing - original draft, review & editing. **HT:** Investigation. **JLOR, GST and JMS:** Investigation; Validation; Writing - review & editing. **CKH and JRH:** Conceptualization; Funding acquisition; Project administration; Supervision; Writing – original draft, review & editing.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- C. J. Murray, K. S. Ikuta, F. Sharara, L. Swetschinski, G. Robles Aguilar, A. Gray, C. Han, C. Bisignano, P. Rao, E. Wool, S. C. Johnson, A. J. Browne, M. G. Chipeta, F. Fell, S. Hackett, G. Haines-Woodhouse, B. H. Kashef Hamadani, E. A. P. Kumaran, B. McMani-gal, R. Agarwal, S. Akech, S. Albertson, J. Amuasi, J. Andrews, A. Aravkin, E. Ashley, F. Bailey, S. Baker, B. Basnyat, A. Bekker, R. Bender, A. Bethou, J. Bielicki, S. Boonkasidecha, J. Bukosia, C. Car-valheiro, C. Castañeda-Orjuela, V. Chansamouth, S. Chaurasia, S. Chiurchiù, F. Chowdhury, A. J. Cook, B. Cooper, T. R. Cressey, E. Criollo-Mora, M. Cunningham, S. Darboe, N. P. J. Day, M. De Luca, K. Dokova, A. Dramowski, S. J. Dunachie, T. Eckmanns, D. Eibach, A. Emami, N. Feasey, N. Fisher-Pearson, K. Forrest, D. Garrett, P. Gastmeier, A. Z. Giref, R. C. Greer, V. Gupta, S. Haller, A. Haselbeck, S. I. Hay, M. Holm, S. Hopkins, K. C. Iregbu, J. Jacobs, D. Jarovsky, F. Javanmardi, M. Khorana, N. Kisoona, E. Kobeissi, T. Kostyanov, F. Krapp, R. Krumkamp, A. Kumar, H. H. Kyu, C. Lim, D. Limmathu-rotsakul, M. J. Loftus, M. Lunn, J. Ma, N. Mturi, T. Munera-Huertas, P. Musicha, M. M. Mussi-Pinhata, T. Nakamura, R. Nanavati, S. Nangia, P. Newton, C. Ngoun, A. Novotney, D. Nwakanma, C. W. Obiero, A. Olivas-Martinez, P. Olliaro, E. Ooko, E. Ortiz-Brizuela, A. Y. Peleg, C. Perrone, N. Plakkal, A. Ponce-de-Leon, M. Raad, T. Ramdin, A. Riddell, T. Roberts, J. V. Robotham, A. Roca, K. E. Rudd, N. Russell, J. Schnall, J. A. G. Scott, M. Shivamallappa, J. Sifuentes-Osornio, N. Steenkeste, A. J. Stewardson, T. Stoeva, N. Tasak, A. Thaiprakong, G. Thwaites, C. Turner, P. Turner, H. R. van Doorn, S. Velaphi, A. Vongpradith, H. Vu, T. Walsh, S. Waner, T. Wangrang-simakul, T. Wozniak, P. Zheng, B. Sartorius, A. D. Lopez, A. Stergachis, C. Moore, C. Dolecek and M. Naghavi, *The Lancet*, 2022, **399**, 629–655.
- S. Davies, J. Farrar, J. Rex, L. White, R. Murry and J. O'Neill, Antimicrobial Resistance: Tackling a crisis for the health and wealth of nations, 2014.
- E. D. Brown and G. D. Wright, *Nature*, 2016, **529**, 336–343.
- K. Talkington, C. Shore and P. Kothari, *Pew Charit. Trust Phila. PA USA*, 2016, **1**, 33–39.
- A. York, *Nat. Rev. Microbiol.*, 2020, **18**, 63–63.
- C. P. J. Vendrik, J. J. Bergers, W. H. De Jong and P. A. Steerenberg, *Cancer Chemother. Pharmacol.*, 1992, **29**, 413–429.
- K. L. F. Hilton, C. Manwani, J. E. Boles, L. J. White, S. Ozturk, M. D. Garrett and J. R. Hiscock, *Chem. Sci.*, 2021, **12**, 13273–13282.
- B. Chertok, M. J. Webber, M. D. Succi and R. Langer, *Mol. Pharm.*, 2013, **10**, 3531–3543.
- G. Tiwari, R. Tiwari, B. Sriwastawa, L. Bhati, S. Pandey, P. Pandey and S. K. Bannerjee, *Int. J. Pharm. Investig.*, 2012, **2**, 2–11.
- Y. Ma, K. Gao, H. Yu, W. Liu, Y. Qin, R. Xing, S. Liu and P. Li, *Carbohydr. Polym.*, 2021, **261**, 117821–117821.
- H. Cheng, R.-R. Zheng, G.-L. Fan, J.-H. Fan, L.-P. Zhao, X.-Y. Jiang, B. Yang, X.-Y. Yu, S.-Y. Li and X.-Z. Zhang, *Biomaterials*, 2019, **188**, 1–11.
- M. Rucevic, D. Hixson and D. Josic, *Electrophoresis*, 2011, **32**, 1549–1564.
- N. N. Mishra, A. S. Bayer, T. T. Tran, Y. Shamoo, E. Mi-leykovskaya, W. Dowhan, Z. Guan and C. A. Arias, *PLoS One*, 2012, **7**, e43958–e43958–e43958–e43958.
- M. Nazari, M. Kurdi and H. Heerklotz, *Biophysical Journal*, 2012, **102**, 498–506.
- D. Wiczorek, D. Gwiazdowska, K. Staszak, Y.-L. Chen and T.-L. Shen, *Journal of Surfactants and Detergents*, 2016, **19**, 813–822.
- WHO publishes list of bacteria for which new antibiotics are urgently needed, <https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed>, (accessed 27 January 2023).
- M. G. D'Andrea, C. C. Domingues, S. V. P. Malheiros, F. G. Neto, L. R. S. Barbosa, R. Itri, F. C. L. Almeida, E. de Paula and M. L. Bianconi, *Langmuir*, 2011, **27**, 8248–8256.
- P. Di Profio, R. Germani, A. Fontana and V. Canale, *J. Mol. Liq.*, 2019, **278**, 650–657.
- A. Piñeiro, X. Banquy, S. Pérez-Casas, E. Tovar, A. García, A. Villa, A. Amigo, A. E. Mark and M. Costas, *J. Phys. Chem. B*, 2007, **111**, 4383–4392.
- A. Chattopadhyay and K. g. Harikumar, *FEBS Lett.*, 1996, **391**, 199–202.
- J. Folch, M. Lees and G. H. Sloane Stan-ley, *J. Biol. Chem.*, 1957, **226**, 497–509.
- G. Townshend, G. S. Thompson, L. J. White, J. R. Hiscock and J. L. Ortega-Roldan, *Chem. Commun.*, 2020, **56**, 4015–4018.
- T. J. Knowles, R. Finka, C. Smith, Y.-P. Lin, T. Dafforn and M. Overduin, *J. Am. Chem. Soc.*, 2009, **131**, 7484–7485.
- S. C. Lee, T. J. Knowles, V. L. G. Postis, M. Jamshad, R. A. Parslow, Y. Lin, A. Goldman, P. Sridhar, M. Overduin, S. P. Muench and T. R. Dafforn, *Nat. Protoc.*, 2016, **11**, 1149–1162.
- M. Orwick-Rydmark, J. E. Lovett, A. Graziadei, L. Lind-holm, M. R. Hicks and A. Watts, *Nano Lett.*, 2012, **12**, 4687–4692.
- H. Y. Carr and E. M. Purcell, *Phys. Rev.*, 1954, **94**, 630–638.
- S. Meiboom and D. Gill, *Rev. Sci. Instrum.*, 1958, **29**, 688–691.
- J. E. Boles, C. Bennett, J. Baker, K. L. F. Hilton, H. A. Kotak, E. R. Clark, Y. Long, L. J. White, H. Yuk Lai, C. K. Hind, J. Mark Sutton, M. D. Garrett, A. Cheasty, J. L. Ortega-Roldan, M. Charles, C. J. E. Haynes and J. R. Hiscock, *Chem. Sci.*, 2022, **13**, 9761–9773.
- OriginPro (version 9.9.0.225 (Learning Edition)) *OriginLab Corporation, Northampton, MA, USA*.
- S. Goutelle, M. Maurin, F. Rougier, X. Barbaut, L. Bour-guignon, M. Ducher and P. Maire, *Fundam. Clin. Pharmacol.*, 2008, **22**, 633–648.