



# Kent Academic Repository

O'Driscoll, Jamie M., Edwards, Jamie J., Coleman, Damian A., Taylor, Katrina Ann, Sharma, Rajan and Wiles, Jonathan D. (2022) *One year of isometric exercise training for blood pressure management in men: a prospective randomized controlled study*. *Journal of Hypertension*, 40 (12). pp. 2406-2412. ISSN 1473-5598.

## Downloaded from

<https://kar.kent.ac.uk/102895/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1097/hjh.0000000000003269>

## This document version

Author's Accepted Manuscript

## DOI for this version

## Licence for this version

UNSPECIFIED

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

### Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

**One-year of isometric exercise training for blood pressure management in men: a prospective randomized controlled study.**

Jamie M. O'Driscoll<sup>1,2</sup>., Jamie J. Edwards<sup>1</sup>., Damian A. Coleman<sup>1</sup>., Katrina A. Taylor<sup>3</sup>.,  
Rajan Sharma<sup>2</sup>., Jonathan D. Wiles<sup>1</sup>

<sup>1</sup>School of Psychology and Life Sciences, Canterbury Christ Church University, Kent, CT1  
1QU

<sup>2</sup>Department of Cardiology, St George's University Hospitals NHS Foundation Trust,  
Blackshaw Road, Tooting, London, SW17 0QT.

School of Sport and Exercise Science, University of Kent, Canterbury, CT2 7NU

**Corresponding Author:** Correspondence to Dr Jamie O'Driscoll, School of Human and Life  
Sciences, Canterbury Christ Church University, Kent, CT1 1QU. Email:

[jamie.odriscoll@canterbury.ac.uk](mailto:jamie.odriscoll@canterbury.ac.uk); Telephone: 01227 782711

**Conflicts of Interest and Source of Funding:** None

**Running Title:** Longitudinal IET and blood pressure

**Keywords:** Blood pressure, hypertension, isometric exercise training

**Word Count:** 2427

## **Abbreviations**

BP – blood pressure

IET – isometric exercise training

CO – cardiac output

dBp – diastolic blood pressure

ECG – electrocardiogram

HR – heart rate

mBP – mean blood pressure

sBP – systolic blood pressure

SV – stroke volume

TPR – total peripheral resistance

**Abstract**

**Objective:** Isometric exercise training (IET) over 4-12 weeks is an effective anti-hypertensive intervention. However, blood pressure (BP) reductions are reversible if exercise is not maintained. No work to date has investigated the long-term effects of IET on resting blood pressure.

**Methods:** We randomised 24 unmedicated patients with high-normal BP to a 1-year wall squat IET intervention or non-intervention control group. Resting BP and various clinically important haemodynamic variables including heart rate (HR), stroke volume (SV), cardiac output (CO) and total peripheral resistance (TPR) were measured pre and post the 1-year study period.

**Results:** 1-year of IET produced statistically significant reductions in resting systolic ( $-8.5 \pm 5$  mmHg,  $p < 0.001$ ) and diastolic ( $-7.3 \pm 5.8$  mmHg,  $p < 0.001$ ) BP compared to the control group. There was also a significant reduction in resting HR ( $-4.2 \pm 3.7$  b $\cdot$ min $^{-1}$ ,  $p = 0.009$ ) and significant increase in SV ( $11.2 \pm 2.8$  ml,  $p = 0.012$ ), with no significant change in CO ( $0.12 \pm 2.8$  L $\cdot$ min $^{-1}$ ,  $p = 0.7$ ). TPR significantly decreased following IET ( $-246 \pm 88$  dyne $\cdot$ s $\cdot$ cm $^{-5}$ ,  $p = 0.011$ ). Adherence to the IET sessions was 77% across all participants (3x IET sessions per week), with no participant withdrawals.

**Conclusion:** This novel study supports IET as an effective long-term strategy for the management of resting BP, producing clinically important, chronic BP adaptations in patients at risk of hypertension. Importantly, this work also demonstrates impressive long-term adherence rates, further supporting the implementation of IET as a means of effective BP management in clinical populations.

## Introduction

Isometric exercise training (IET) is now well established as an efficacious anti-hypertensive intervention, capable of producing greater reductions in resting blood pressure (BP) when compared to the standard recommendations of exercise guidelines [1]. Specifically, recent meta-analysis evidence has shown statistically significant systolic and diastolic BP (sBP and dBP) reductions of 8.5 and 4.1 mmHg, respectively, following relatively short-term (4-12 weeks) IET interventions [2]. The magnitude of these reductions are comparable to standard anti-hypertensive pharmacotherapy [3], with the added benefit of no adverse side effects and reduced costs. Given that hypertension is the leading attributable risk factor for cardiovascular disease and all-cause mortality [4], with a prevalence of greater than 1.1 billion worldwide [5,6], IET may have substantial clinical implications for the future management of resting BP.

A number of trials have demonstrated clinically significant reductions in sBP and dBP following as little as 4-weeks of IET [7–10]; however, no work to date has investigated the efficacy of an IET intervention beyond 12-weeks in duration. Consequently, there is currently no evidence to support long-standing adaptations in resting BP or the long-term adherence to IET. Research from Millar et al. [11] demonstrated linear negative trends in resting sBP and dBP over an 8-week intervention with no plateau in reductions over this timeframe. In addition, a meta-analysis from Inder et al. [12], reported larger reductions in sBP in participants undertaking  $\geq 8$  weeks of IET compared to  $< 8$  weeks of IET. However, while this work may indicate greater reductions from a longer intervention duration, conclusive inferences cannot be made through the extrapolation of such limited data. As such, prospective randomised controlled trials are required to truly understand the long-term

effectiveness of IET. Furthermore, the potential future clinical implementation of IET is, at least somewhat, based on the presumption that adherence to IET is likely to be greater than that of other exercise modes due to its time-efficient and practical nature. However, without sufficient longitudinal investigation of IET, the validity of this presumption is not known.

This study aimed to perform the first longitudinal randomised controlled study, measuring the effects of a 12-month IET intervention on resting BP in patients with high normal BP. We also aimed to measure important mechanistic haemodynamic variables including cardiac output (CO) and total peripheral resistance (TPR).

## **Methods**

### ***Participant population and ethical approval***

Twenty-four male participants (median (IQR) age 45 (41.3, 49) years) were recruited to this single-centre prospective, randomised controlled trial. Participants were physically inactive and classified as high normal in accordance with the relevant guidelines [13]. None of the participants were under any acute or chronic pharmacotherapy and all presented with a normal clinical cardiovascular examination and 12-lead electrocardiogram (ECG) with no cardiac or metabolic disease. All participants were non-smokers. Written informed consent was obtained from all participants before testing and this study conformed to the Declaration of Helsinki principles with approval from the local ethics committee (Ref:12/SAS/122).

### ***Study Procedures***

Eligible participants were randomly allocated to a 12-month IET intervention or to the control group. Participants assigned to the IET group were required to attend the Canterbury Christ Church University laboratory on three separate occasions, while the control group attended twice. The IET group performed an extra initial visit which comprised an incremental isometric wall-squat test to determine the individualised knee joint angle for appropriate IET intensity prescription, as previously described [7]. The remaining sessions involved the acquisition of the relevant pre- and post-intervention cardiovascular parameters. All participants were required to maintain normal dietary and circadian routines throughout the study and each phase of testing, as well as refrain from alcohol and caffeine consumption for 24-hours and fast for at least 4 hours before testing.

### ***Resting clinic blood pressure***

Resting clinic brachial artery BP was acquired using a validated automated device (Dinamap Pro 200 Critikon; GE Medical Systems, Freiburg, Germany) and according to current guidelines [13]. All measures were recorded in a temperature-controlled room at the same time of day ( $\pm 1$ -hour) pre and post the 12-month IET intervention and control period.

### ***Haemodynamics***

All haemodynamic variables were acquired using the Task Force<sup>®</sup> Monitor, which is a validated non-invasive beat-to-beat monitoring system that provides automatic calculations of all outputs. Heart rate (HR) was acquired through a six-channel ECG and stroke volume (SV) was recorded via impedance cardiography through three electrode bands, two of which were in line with the xiphoid process and adjacent to the thorax and the other placed on the nape of the neck. Cardiac output (CO) was automatically derived from HR and SV. Total peripheral resistance (TPR) was separately calculated in accordance with Ohm's law. Both CO and TPR were indexed to body surface area following acquisition of participant height and mass using a SECA 213 stadiometer and SECA 700 mechanical column scales (SECA GmbH & Co., Hamburg, Germany) respectively. All measures were recorded following 15 minutes of supine rest with a continuous recording period of 5 minutes.

### ***Isometric exercise training intervention***

Participants allocated to the training group performed 3 weekly wall squat IET sessions at 95% HR peak for 12-months in an unsupervised home-based setting. As previously described [14–16], each participant performed a maximal incremental wall squat test with beat-to-beat HR recording in order to determine HR peak corresponding to the prescribed knee angle. Following this test, participants were issued a 'Bend and Squat' device, which is a simple device



developed in-house allowing for the alignment of a participant's feet and back placement into the correct position to elicit the appropriate HR response [9]. Using the Bend and Squat device, the home-based wall squat protocol involved participants resting their back against a fixed wall with their feet placed parallel and shoulder width apart, while their arms remained relaxed down by their side (see Supplement File for further details). Each session involved of 4 x 2-min bouts of isometric wall squat, separated with 2-min rest intervals, with at least 48-hours recovery between every session [7–9]. To allow for close monitoring of intensity and adherence, all participants recorded HR and rate of perceived exertion data following each IET bout and uploaded their data to a personal online database which was monitored by the investigators. The control group were required to maintain their usual routine and daily activities, abstaining from any form of exercise they did not habitually perform for the full 12-month period. The control group were routinely contacted by the investigators to confirm maintenance of circadian habits.

### *Sample size estimation*

Based on previous studies utilizing this protocol, we have previously reported extremely large changes in sBP [7] (reductions of >12mmHg) equivalent to an effect statistic of >2.2 with similar intervention. Using G\*Power software [17], a more conservative (albeit still large) effect statistic of 1.2 was selected, and with power of 80% and  $p < 0.05$ , the estimated sample size of 12 participants per group were recruited.

### *Statistical analysis*

Continuous variables are expressed as mean $\pm$ SD. Post-trial group differences for sBP, mBP, dBP and haemodynamic parameters were assessed using Analysis of Covariance (ANCOVA), with baseline scores for each parameter included as a covariate in the model. All data were

analysed using the statistical package for social sciences (SPSS 26 release version for Windows; SPSS Inc., Chicago, Illinois, USA).

## **Results**

All participants completed the intervention safely, with no adverse events following any IET session. There were no significant differences between the groups at baseline for any of the physiological variables or participant characteristics, including height (IET:  $178.6 \pm 8$  cm; Con:  $178.2 \pm 7.2$ ) and mass (IET:  $88.5 \pm 11.1$ ; Con:  $90.4 \pm 13.9$ ).

Adherence to the IET sessions was 77% across all participants (3x IET sessions per week). None of the participants withdrew from the intervention. One participant re-performed the IET incremental wall squat test after 7-months to re-calculate a sufficient knee joint angle, as routine monitoring of HR data recorded during training sessions demonstrated the training did not elicit the appropriate intensity.

### **Resting blood pressure and haemodynamics**

As seen in Table 1 and Figure 1, the IET group demonstrated significant reductions in resting clinic sBP ( $-8.5 \pm 5$  mmHg,  $p < 0.001$ ) and dBP ( $-7.3 \pm 5.8$  mmHg,  $p < 0.001$ ) compared to the control group. Figure 2 illustrates the individual changes in BP pre and post IET and control period. There was also a significant reduction in resting HR ( $-4.2 \pm 3.7$  b $\cdot$ min $^{-1}$ ,  $p = 0.009$ ) with adjacent significant increases in SV ( $11.2 \pm 2.8$  ml,  $p = 0.012$ ) and SV index ( $5.3 \pm 1.9$  ml $\cdot$ m $^2$ ,  $p = 0.013$ ), but no significant change in CO ( $0.12 \pm 2.8$  L $\cdot$ min $^{-1}$ ,  $p = 0.7$ ) or CO index ( $0.03 \pm 0.2$  L $\cdot$ min $^{-1}$  $\cdot$ m $^2$ ,  $p = 0.82$ ). Separately, there was a significant reduction in TPR ( $-246 \pm 88$  dyne $\cdot$ s $\cdot$ cm $^{-5}$ ).

<sup>5</sup>,  $p=0.011$ ) (Figure 3). Figure 4 illustrates the individual changes in HR, SV, CO and TPR pre and post IET and control period.

## **Discussion**

This is the first study to investigate the longitudinal (one-year) efficacy of IET as an anti-hypertensive intervention. This novel research supports IET as an effective long-term strategy in the management of resting BP, capable of both producing and maintaining clinically important BP adaptations in patients with high normal BP. Additionally, the findings of this study support the function of IET in sustaining clinically relevant haemodynamic adaptations, with significant reductions in resting HR and TPR and significant increase in SV. Importantly, this work also evidences substantially higher adherence to IET than that typically seen when following the current recommended exercise guidelines (44.6%) [18] or with clinical pharmacotherapy [19].

While the ability of IET to produce statistically significant reductions in resting BP is well-established [2], this study provides the first evidence that such adaptations can be sustained long-term. We observed reductions in sBP and dBP by 10.5 and 8 mmHg, respectively, following 1-year of IET, which is both greater than the average reduction seen from other modes of exercise [20,21] as well as that of standard dosage anti-hypertensive monotherapy [3]. In particular, Anderssen et al. [22] investigated the effects of a 1-year aerobic exercise intervention and reported reductions of 4.1 and 5.1 mmHg sBP and dBP, respectively. Despite having a lower baseline BP, the present study demonstrates more than twice the magnitude of reduction over the same intervention duration. Additionally, the average BP reduction across the five primary anti-hypertensive drug classes has been reported at 9.1 and 5.5 mmHg for sBP

and dBP, respectively [23], which is inferior to the reductions we report following IET. In terms of their clinical importance, previous large-scale data suggest these IET reductions to be associated with a reduced risk of ischaemic heart disease, stroke, cardiovascular disease events and cardiovascular mortality by 16.8%, 27.3%, 21% and 10.5% respectively [24].

Of interest, the dBP reductions from the present study are particularly substantial, being of a far greater magnitude than generally achievable with any other form of anti-hypertensive intervention [21,23] and greater than that observed following short term IET [9,25]. While the underpinnings of this finding are unknown, acute data shows that wall squat IET produces large post-exercise hypotensive effects on dBP, with previous studies reporting reductions close to 20 mmHg from baseline to the recovery phase in both normotensive [26], and hypertensive participants [16]. This response is certainly much larger than seen following other modes of exercise [27–29]. Indeed, the consistent repetition of such acute hypotensive responses elicited from each IET session may translate into the chronic reductions reported from this 1-year intervention, and may be an important mechanism for the cardiovascular adaptations previously reported [7,30].

This study echoes the wider mechanistic literature in that IET predominantly induces reductions in resting BP independent of changes in CO [1,7,8]. While significant reductions are seen in resting HR, CO remains unchanged as the consequence of a significant increase in SV, which is likely mediated through the significant left ventricular functional and mechanical adaptations recently reported following IET [30]. As such, IET induced BP reductions appear to be primarily modulated through TPR, such as altered autonomic and baroreflex vasomotor control [7,31], local endothelial-dependant mechanisms [7,32] and systematic structural

vascular remodelling [33]. However, the exact contribution of each of these potential processes is not yet known.

While the study sample size of this research is limited, the overall adherence to this 1-year intervention was impressive. Poor adherence rates have been deemed a primary contributor to the current ineffectiveness of both the present exercise guidelines and medical therapy [34,35]. Current physical inactivity rates in accordance with recommended exercise guidelines have been cited at 27.5% [35], or even as low as 5% via objective accelerometer measures [36], while the average adherence to anti-hypertensive medical therapy is typically reported at less than 50% 1-year following prescription [19]. As the first study to provide long-term adherence data in IET, these findings support the hypothesis that adherence to IET is likely to be greater than other anti-hypertensive means, especially given the well-cited report that 50% of people who start an exercise programme will abstain within 6 months [37]. At 77% adherence from a 1-year intervention, this work strengthens IET as a successful intervention in clinical practice. However, comparative large-scale randomised controlled data is needed to truly discern the differing adherence rates between IET and the current physical activity guidelines.

### ***Limitations***

This research was single centre with a limited sample size and only included males. Therefore, future multi-centre work in a large-scale population including female and different ethnic groups is warranted. In addition, this work was not statistically powered to detect haemodynamic changes, and thus only limited inferences can be made from this data. Finally, the exclusion of smokers from this work limits the relative applicability of these findings to such individuals who tend to derive an even greater benefit from effective anti-hypertensive interventions.

**Conclusion**

This is the first study to investigate the long-term efficacy of IET in the management of resting BP. 1-year of IET produces and maintains statistically and clinically significant reductions in resting BP, superior to that of the current exercise guidelines as well as anti-hypertensive pharmacotherapy. These reductions are likely mediated through peripheral vascular adaptations, while also producing significant improvements in resting HR and SV. Importantly, this work also demonstrates impressive long-term adherence rates, further supporting the clinical implementation of IET as a means of BP management.

## References

- 1 Millar PJ, McGowan CL, Cornelissen VA, Araujo CG, Swaine IL. Evidence for the role of isometric exercise training in reducing blood pressure: Potential mechanisms and future directions. *Sports Med.* 2014; 44:345–356.
- 2 Edwards J, Caux A De, Donaldson J, Wiles J, O’Driscoll J. Isometric exercise versus high-intensity interval training for the management of blood pressure: a systematic review and meta-analysis. *Br J Sports Med* 2021; :bjsports-2021-104642.
- 3 Law MR, Morris JK, Wald NJ. Use of blood pressure lowering drugs in the prevention of cardiovascular disease: Meta-analysis of 147 randomised trials in the context of expectations from prospective epidemiological studies. *BMJ* 2009; 338:1245.
- 4 Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, *et al.* A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380:2224–2260.
- 5 Forouzanfar MH, Liu P, Roth GA, Ng M, Biryukov S, Marczak L, *et al.* Global Burden of Hypertension and Systolic Blood Pressure of at Least 110 to 115 mm Hg, 1990-2015. *JAMA* 2017; 317:165.
- 6 WHO. Blood Pressure. Geneva 2018.
- 7 Taylor KA, Wiles JD, Coleman DA, Leeson P, Sharma R, O’Driscoll JM. Neurohumoral and ambulatory haemodynamic adaptations following isometric exercise training in unmedicated hypertensive patients. *J Hypertens* 2019; 37:827–836.
- 8 De Caux A, Edwards JJ, Swift H, Hurst P, Wiles JD, O’Driscoll J. Isometric exercise training for the management of blood pressure: A randomised sham-controlled study. *Physiol Rep* 2021. 10:e15112.
- 9 Wiles JD, Goldring N, Coleman D. Home-based isometric exercise training induced reductions resting blood pressure. *Eur J Appl Physiol* 2017; 117:83–93.
- 10 Baddeley-White DS, McGowan CL, Howden R, Gordon BD, Kyberd P, Swaine IL. Blood pressure lowering effects of a novel isometric exercise device following a 4-week isometric handgrip intervention. *Open access J Sport Med* 2019; 10:89–98.

- 11 Millar PJ, Bray SR, McGowan CL, MacDonald MJ, McCartney N. Effects of isometric handgrip training among people medicated for hypertension: a multilevel analysis. *Blood Press Monit* 2007; 12:307–314.
- 12 Inder JD, Carlson DJ, Dieberg G, Mcfarlane JR, Hess NCL, Smart NA. Isometric exercise training for blood pressure management: A systematic review and meta-analysis to optimize benefit. *Hypertens Res* 2016; 39:89–94.
- 13 Williams B, Mancia G, Spiering W, Rosei EA, Azizi M, Burnier M, *et al.* 2018 ESC/ESH Guidelines for the management of arterial hypertension. *Eur Heart J.* 2018; 39:3021–3104.
- 14 Wiles J, Goldring N, O’Driscoll J, Taylor K, Coleman D. An alternative approach to isometric exercise training prescription for cardiovascular health. *Transl J Am Coll Sport Med* 2018; 3:10–18.
- 15 Goldring N, Wiles JD, Coleman D. The effects of isometric wall squat exercise on heart rate and blood pressure in a normotensive population. *J Sports Sci* 2014; 32:129–136.
- 16 Taylor KA, Wiles JD, Coleman DD, Sharma R, O’Driscoll JM. Continuous cardiac autonomic and hemodynamic responses to isometric exercise. *Med Sci Sports Exerc* 2017; 49:1511–1519.
- 17 Faul F, Erdfelder E, Lang AG, Buchner A. G\*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007; 39:175–191.
- 18 Ussery EN, Fulton JE, Galuska DA, Katzmarzyk PT, Carlson SA. Joint Prevalence of Sitting Time and Leisure-Time Physical Activity Among US Adults, 2015-2016. *JAMA* 2018; 320:2036–2038.
- 19 Vrijens B, Vincze G, Kristanto P, Urquhart J, Burnier M. Adherence to prescribed antihypertensive drug treatments: Longitudinal study of electronically compiled dosing histories. *BMJ* 2008; 336:1114–1117.
- 20 Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: A meta-analysis of randomized, controlled trials. *Ann Intern Med* 2002; 136:493–503.
- 21 Cornelissen VA, Smart NA. Exercise training for blood pressure: a systematic review



- and meta-analysis. *J. Am. Heart Assoc.* 2013; 2. doi:10.1161/JAHA.112.004473
- 22 Anderssen S, Holme I, Urdal P, Hjermmann I. Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Press* 1995; 4:343–349.
- 23 Law MR, Wald NJ, Morris JK, Jordan RE. Value of low dose combination treatment with blood pressure lowering drugs: analysis of 354 randomised trials. *BMJ Br Med J* 2003; 326:1427.
- 24 Adler A, Agodoa L, Algra A, Asselbergs FW, Beckett NS, Berge E, *et al.* Pharmacological blood pressure lowering for primary and secondary prevention of cardiovascular disease across different levels of blood pressure: an individual participant-level data meta-analysis. *Lancet* 2021; 397:1625–1636.
- 25 Carlson DJ, Inder J, Palanisamy SKA, McFarlane JR, Dieberg G, Smart NA. The efficacy of isometric resistance training utilizing handgrip exercise for blood pressure management: A randomized trial. *Medicine (Baltimore)* 2016; 95.
- 26 O’Driscoll JM, Boucher C, Vilda M, Taylor KA, Wiles JD. Continuous cardiac autonomic and haemodynamic responses to isometric exercise in females. *Eur J Appl Physiol* 2021; 121:319–329.
- 27 Chen CY, Bonham AC. Postexercise Hypotension: Central Mechanisms. *Exerc Sport Sci Rev* 2010; 38:122.
- 28 Pescatello LS, Guidry MA, Blanchard BE, Kerr A, Taylor AL, Johnson AN, *et al.* Exercise intensity alters postexercise hypotension. *J Hypertens* 2004; 22:1881–1888.
- 29 Edwards JJ, Wiles JD, Vadaszy N, Taylor KA, O’Driscoll JM. Left ventricular mechanical, cardiac autonomic and metabolic responses to a single session of high intensity interval training. *Eur J Appl Physiol* 2021 1222 2021; 122:383–394.
- 30 O’Driscoll JM, Edwards JJ, Wiles JD, Taylor KA, Leeson P, Sharma R. Myocardial work and left ventricular mechanical adaptations following isometric exercise training in hypertensive patients. *Eur J Appl Physiol* 2022; 122:727-734.
- 31 Taylor AC, McCartney N, Kamath M V., Wiley RL. Isometric training lowers resting blood pressure and modulates autonomic control. *Med Sci Sports Exerc* 2003; 35:251–256.

- 32 Correia MA, Oliveira PL, Farah BQ, Vianna LC, Wolosker N, Puech-Leao P, *et al.* Effects of Isometric Handgrip Training in Patients With Peripheral Artery Disease: A Randomized Controlled Trial. *J Am Heart Assoc* 2020; 18:e013596.
- 33 Cahu Rodrigues SL, Farah BQ, Silva G, Correia M, Pedrosa R, Vianna L, *et al.* Vascular effects of isometric handgrip training in hypertensives. *Clin Exp Hypertens* 2020; 42:24–30.
- 34 Williamson W, Foster C, Reid H, Kelly P, Lewandowski AJ, Boardman H, *et al.* Will Exercise Advice Be Sufficient for Treatment of Young Adults With Prehypertension and Hypertension? A Systematic Review and Meta-Analysis. *Hypertension*. 2016; 68:78–87.
- 35 Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1·9 million participants. *Lancet Glob Heal* 2018; 6:e1077–e1086.
- 36 Troiano RP, Berrigan D, Dodd KW, Mâsse LC, Tilert T, Mcdowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc* 2008; 40:181–188.
- 37 Dishman RK. *Exercise adherence: its impact on public health*. Human Kinetics Books; 1988.

Table 1: Resting office blood pressure and haemodynamic adaptations pre and post isometric exercise training and control period.

	Control (n=12)		Isometric Exercise Training (n=12)	
	Pre	Post	Pre	Post
<b>Resting Office BP</b>				
sBP (mmHg)	132.4 ± 5.7	130.4 ± 6.5	132.3 ± 5.8	121.8 ± 3.9**
mBP (mmHg)	98.5 ± 6.1	96.5 ± 5.8	98.6 ± 5.4	88.7 ± 3.2**
dBP (mmHg)	81.5 ± 6.7	81.3 ± 6.4	81.7 ± 6.9	73.7 ± 5.1**
<b>Resting Haemodynamic Variables</b>				
Heart Rate (b·min <sup>-1</sup> )	58.5 ± 6.6	59.5 ± 6.8	61.8 ± 10.0	57.9 ± 8.8*
Stroke Volume (ml)	85.8 ± 13.2	85.2 ± 11.3	82.5 ± 12	94.3 ± 13.7*
Stroke Index (ml·m <sup>-2</sup> )	40.8 ± 5.7	40.6 ± 6.3	39.7 ± 6.7	45.2 ± 6.6*
Cardiac Output (L·min <sup>-1</sup> )	4.8 ± 1	5.1 ± 0.9	5.3 ± 0.8	5.5 ± 1.3
Cardiac Index (L·min <sup>-1</sup> ·m <sup>-2</sup> )	2.3 ± 0.4	2.4 ± 0.5	2.5 ± 0.4	2.6 ± 0.6
TPR (dyne·s·cm <sup>-5</sup> )	1622 ± 328	1565 ± 312	1650 ± 288	1355 ± 314*

Note: Values are mean ± SD. \* p<0.05, \*\* p<0.001 differences between change scores from control and IET condition. BP = blood pressure; sBP = systolic blood pressure; mBP = mean blood pressure; dBP = diastolic blood pressure; TPR = total peripheral resistance; TPRI = total peripheral resistance index.

**Figure Legend:**

Figure 1: Mean sBP (A) and dBP (B) change values following control (closed circles) and IET (open circles) conditions. Note: \*\*Significant ( $P < 0.001$ ) difference in the control and IET change value.

Figure 2: Individual changes in sBP and dBP following control and IET conditions. Grey lines represent individual participants and black lines represent the mean changes at baseline and following 1-year for IET and control groups.

Figure 3: Mean HR (A), SV (B), CO (C) and TPR (D) change values following control (closed circles) and IET (open circles) conditions. Note: \*Significant ( $P < 0.05$ ) difference in the control and IET change value.

Figure 4: Individual changes in HR, SV, CO and TPR following control and IET conditions. Grey lines represent individual participants and black lines represent the mean changes at baseline and following 1-year for IET or control groups.