



Kent Academic Repository

Abbot-Smith, Kirsten, Matthews, Danielle, Bannard, Colin, Nice, Joshua, Malkin, Louise, Williams, David M. and Hobson, William (2024) *Conversational topic maintenance and related cognitive abilities in autistic versus neurotypical children.* Autism . ISSN 1362-3613. (In press)

Downloaded from

<https://kar.kent.ac.uk/107156/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1177/13623613241286610>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal** , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Conversational topic maintenance and related cognitive abilities in autistic and neuro-typical children

Kirsten Abbot-Smith¹

Danielle Matthews²

Colin Bannard³

Joshua Nice¹

Louise Malkin¹

David Williams¹

William Hobson²

¹School of Psychology, University of Kent, UK

²School of Psychology, University of Sheffield, UK

³Department of Linguistics and English Language, University of Manchester, UK

Contact: Kirsten Abbot-Smith

Dr Kirsten Abbot-Smith
Kent Child Development Unit, School of Psychology,
University of Kent,
Keynes College, Canterbury CT2 7NP, UK
Email: K.Abbot-Smith@kent.ac.uk
Tel: +44 (0)1227 823016

Acknowledgments

These studies were funded by Kent Health, a School of Psychology Seed Fund and a PhD scholarship to Louise Malkin. A special 'thank you' to the children, to the parents who brought their children into the Kent Child Development Unit and to the staff and caregivers of local primary schools in Kent, UK. Additional thanks to the British National Health Service for ethical approval for Study 3. Thanks go to John Allen for providing some assistance with R. Many thanks to Sophie Darroch, Chantel Smith, Dr Lauren Jenner, Laetitia Rater and Rebecca Steele (SLT) for testing in Studies 2 and 3. Thanks to Hannah Meurice for coding in Studies 2 and 3 and to Caitlin Moriarty for coding reliabilities for Study 1. Thanks also to Yemi Bank-Anthony, Hannah Browes, Dr Rebecca Dole, Hannah Doolan, Rhiannon Howells-Harris, Dr Lauren Jenner, Caitlin Moriarty, Corrina Penn, Helen Petterson, Amanda Norman, Gail O'Connor, Caitlin Tuttle, and Mia van der Klashorst for transcription. Thanks to Bejana Rai for theory of mind test transcription and coding.

Abstract

Keeping a conversation going is the social glue of friendships. The DSM criteria for autism list difficulties with back-and-forth conversation but does not necessitate that all autistic children will be equally impacted. We carried out three studies (two pre-registered) with verbally-fluent school children (age 5-9 years) to investigate how autistic and neurotypical children maintain a conversation topic. We also investigated within-group relationships between conversational ability and cognitive and socio-cognitive predictors. Study 1 found autistic children were more likely than neurotypical controls to give off-topic and generic minimal responses (e.g. 'mm', 'oh') and were less likely to give non-verbal responses (e.g. nodding or use of facial affect to respond). Nonetheless, the autistic group provided topic-supporting responses 62% of the time, indicating some aspects of conversation topic maintenance are a relative strength. Studies 2 and 3 found large individual differences in topic-supporting conversational responding amongst both neurotypical and autistic children. These were positively related to theory of mind ability and age in both groups. Conversational skills lie on a continuum for the general population and differences by diagnostic group are a matter of degree. Given the importance for peer relationships, we suggest a whole-classroom approach to supporting conversation skills in all children.

Lay abstract

Children who struggle to maintain conversation with peers often have fewer friends and lower popularity ratings, which can affect wellbeing. Verbal social communication more broadly is linked to both behavioural difficulties and emotional problems. We carried out three studies to examine children's ability to provide responses which keep a back and forth conversation going. The first study found that while autistic children had on average greater difficulties than their neurotypical peers with certain aspects of conversation topic

maintenance, for other aspects the autistic group showed considerable strengths. Both studies 2 (neurotypical children) and 3 (autistic children) found relationships between, on the one hand, conversational ability, and on the other, the ability to consider another's viewpoint and the ability to maintain and update information in short term memory. We suggest support for social conversation skills should be part of mainstream classroom curricula for autistic and neurotypical children alike.

Introduction

One of the two symptom domains of the diagnostic criteria for autism entails ‘persistent deficits in social communication and social interaction’ manifested by difficulties with social-emotional reciprocity, nonverbal communication and with developing and maintaining relationships (American Psychiatric Association, 2013). Particular hurdles within the first category - social-emotional reciprocity - are specified in the DSM-5 to include difficulties with social approach, with back-and-forth conversation, reduced sharing of interests, emotions or affect and failure to initiate or respond to social interactions (American Psychiatric Association, 2013). Thus, the ability to maintain a conversation topic is not only highlighted in the diagnostic criteria but is also critical for most of the other points listed in the domain of social interaction and social communication since conversation skills are central for establishing and maintaining relationships (Friedman et al., 2019). The ‘gold standard’ assessment tools for autism (ADOS-2, ADI-R) involve direct and indirect assessment of conversational ability. Indeed, many have noted that the most common communication deficits seen in intellectually-able autistic youth pertain to conversation (Paul et al., 2009).

Yet, to date, there has been a lack of clarity around a) the distribution of conversational skills in neurotypical school-aged children (i.e., those with no diagnoses or suspected disorders) and how this overlaps with that of autistic children and b) the cognitive and social-cognitive skills that are correlated with conversation skill for both groups of children. This is because there are few quantitative studies that have used direct measurements to examine individual differences in how neurotypical children fare with maintaining conversation (see Abbot-Smith et al., 2023 for a review).

To keep a conversation topic going, speakers need to be able to do two key things. First, a speaker needs to be able to provide ‘topic-supporting’ responses. These include

‘contingent responses’, that is, those which provide information that is both relevant to the immediately preceding conversational turn and add information (Bloom et al., 1976; Pagmar et al., 2022). Another way to support a conversation topic is to provide ‘specific’ minimal responses (e.g. ‘Really?’) or even in certain contexts non-verbal responses, such as laughing, gestures or affective reactions (e.g. an expression of shock) (Bavelas et al., 2000). All of these responses indicate that one has processed and is interested in what the speaker is saying. In contrast, non-contingent (off-topic) responses can provide a stumbling block for conversation topics (Hale & Tager-Flusberg, 2005; Nadig et al., 2010). Children in upper primary school and adolescents – both autistic and neurotypical – give significantly lower ratings of intentions to interact socially with children they have heard responding non-contingently (McGuinness et al., in press; Place & Becker, 1991). An example of a non-contingent response from our dataset is as follows:

RESEARCHER: “I like cooking biscuits with chocolate chips in them. They’re my favourite.”

P2: “There’s twins in my class. They’re identical.”

The second key conversational ability required to maintain a conversation topic involves restricting the length of each turn to avoid ‘monologuing’ (i.e., talking excessively without allowing the partner a turn. While non-contingency and excessive verbosity have often been discussed in relation to autism (Cola et al., 2022; Nadig et al., 2010; Ying Sng et al., 2018), it is unclear to what degree autistic children as a group show conversational impairments, particularly now that far more verbally-fluent autistic children are receiving autism diagnoses (e.g. McConkey, 2020).

Moreover, there are large individual differences amongst neurotypical children in the ability to maintain a conversation topic (see e.g. De Rosnay et al., 2014 for five- to eight-

year-olds). Conversational contingency relates to sociometric ratings by peers in four-year-olds (Hazen & Black, 1989) and training conversation skills leads to higher rates of playground social interaction in neurotypical 10- and 11-year-olds (Bierman & Furman, 1984). Even amongst neurotypical adults, conversational ability relates to ratings (by others) of attraction (Wheless et al., 1992) and interpersonal relationship satisfaction (Miczo et al., 2001). Indeed, as it has been demonstrated experimentally that adults rate other adults less positively on scales of likeability when the latter are less conversationally responsive (Mein et al., 2016). Given the implications for lifelong social relations, we need a better understanding of the degree of heterogeneity in conversational ability within both populations (autistic and neurotypical) and we also need to know the degree to which conversation skills show distinct relationships with key cognitive abilities in each group.

The ‘key suspects’ in the literature regarding the cognitive underpinnings of conversational ability are theory of mind and / or executive functioning abilities such as working memory or inhibitory control (see Matthews et al., 2018 for a review). There are some studies which have examined relationship between either theory of mind or executive functioning and parent questionnaire measures of pragmatics or conversation (Baixauli-Fortea et al., 2019; Hutchison et al., 2020). However, there are very few studies that have examined either theory of mind or executive functioning in relation to *directly* elicited child conversation. Nonetheless, of those that have, there is evidence of links between contingent responding and theory of mind in both autistic primary-school-aged children (Hale & Tager-Flusberg, 2005) and neurotypical children (Slomkowski & Dunn, 1996). However, the degree to which theory of mind uniquely accounts for contingent responding is not clear given that individual differences in theory of mind itself are correlated with both working memory and inhibitory control (Fizke et al., 2014; Lecce et al., 2017). Furthermore, the one prior study that examined relationships between direct measures of child conversation and executive

functioning found that working memory (Backwards Digit Span) correlated positively with contingent responding and ‘fluidity’ whereas inhibitory control correlated negatively with both child talkativeness and number of words per utterance (Blain-Brière et al., 2014). There is therefore a question as to whether any given candidate cognitive underpinning explains unique variance when other candidates are controlled for.

Blain-Brière’s (2014) finding of a relationship between inhibitory control and talkativeness speaks to the second key ability required to maintain a reciprocal conversation, namely that it should involve both conversation partners (as opposed to monologuing). However, talkativeness per se is not an inherently negative trait (Coplan et al., 2011). Talkativeness is considered ‘excessive’ or ‘verbose’ – and leads to negative conversation satisfaction ratings - when speakers also tend to veer off-topic during the turn (Pushkar et al., 2000; Ruscher & Hurley, 2000). The only prior study which has directly examined ‘problematic verbosity’ in children is that of Nadig et al. (2010), who termed this ‘self-contingent elaboration’ and found certain group differences between autistic and matched neurotypical 10- and 11-year-olds. However, while a small number of verbally-fluent, cognitively-able autistic children are ‘excessively verbose’, this is not necessarily a general characteristic of this group (Adams et al., 2002).

In this article we report on studies with children of primary school age, all of whom had non-verbal reasoning and vocabularies within the neurotypical normal range for their age. In Study 1, we investigate differences and similarities between autistic and neurotypical (NT) primary school children in their ability to maintain a conversation. In Studies 2 and 3, we investigate individual differences within a group of neurotypical primary-school children and within a group of autistic primary school children. Here we also ask whether theory of mind and working memory (and – for one study – inhibitory control) account of individual

differences conversation topic maintenance where all predictors are included in the same regression model.

We carried out three studies, all with children tested in southern England. (There was no overlap in participants between any of the studies). Each child was assessed individually in verbal interaction with a psychology researcher or clinician. In the first study, we compared autistic five- to seven-year-olds with neurotypical children matched for age, non-verbal reasoning, core language and sex. In the second study, we elicited extended naturalistic conversations from neurotypical six-year-olds and also assessed their theory of mind, working memory, inhibitory control, vocabulary and non-verbal reasoning abilities. In the third study, conversation was elicited by a clinical psychologist (a co-author on this article) from five- to nine-year-olds who went on to receive an autism diagnosis from a multi-disciplinary assessment in the British National Health Service (NHS). The same children were also assessed on theory of mind, working memory, vocabulary and non-verbal reasoning tasks. The second and third studies were pre-registered prior to the completion of transcription and prior to start of data analysis: <https://osf.io/q7wa4/registrations>

Ethical and sampling statement

Ethical approval was obtained from the first author's institution for all three studies. In addition, for the third study, NHS ethical approval was also obtained. All parents provided full written informed consent. Socioeconomic circumstances¹ and ethnicity were not recorded. The adult autism community was not involved in this study.

¹ This is a limitation. However, the vast majority of the neuro-typical sample in Study 1 were recruited from and tested in low-mid to low decile schools and thus greater conversational proficiency in the NT group is highly unlikely to be related to higher SES.

Study 1

Method

Participants

We included 30 autistic (7 female) and 30 neurotypical (7 female), monolingual British-English-speaking 5- to 7-year-olds (see Table 1 for demographics). The sample size was agreed between the lead, second and fifth authors prior to data collection as the maximum sample size manageable since the data were collection alongside the fifth author's final PhD project between January 2019 and March 2020, when the UK COVID-19 lockdown commenced. No participant had hearing difficulties or ADHD. All autistic children had a diagnosis requiring multi-disciplinary consensus within the British National Health Service. All scored above the threshold for autism on the parent-completed Social Responsiveness Scale 2 (SRS) (Costantino & Gruber, 2005). Of the autistic sample, 37% were recruited via an autism charity. The rest were recruited via their school. Of the NT sample, 87% were recruited via schools and only one scored above threshold (T-score of 62 - mild range) on the SRS-2.

Groups were matched on chronological age, sex, non-verbal reasoning (assessed by the 'Matrices' sub-test of the British Ability Scales; Eliot & Smith, 2011), vocabulary as assessed by both the 'Expressive Vocabulary' sub-test of the *Clinical Evaluation of Language Fundamentals 4* (Semel et al., 2006) and sentence comprehension ability as assessed by the 'Sentence Structures' sub-test of the CELF-5 (Wiig et al., 2013).

Table 1. Means (SD in brackets) for participant characteristics

	Autistic (<i>n</i> =30)	NT (<i>n</i> =30)		
	Mean (SD)	Mean (SD)	<i>p</i>	<i>d</i>
Gender split	23 M; 7 F	23 M; 7 F	-	-
Chronological Age (Months)	77.37 (10.58)	76.94 (9.08)	.87	0.04
Sentence Comprehension CELF-5 Scaled Score	9.97 (2.46)	10.2 (1.94)	.68	0.10
Expressive Vocabulary CELF-4 Scaled Score	8.87 (2.70)	9.13 (1.78)	.65	0.11
Non-Verbal Reasoning: BAS T-Score	41.53 (7.82)	38.77 (7.06)	.16	0.37
Social Responsiveness Scale T-score	84.57 (7.72)	44.37 (6.16)	<.001	5.76

Procedure

In this study, continuous conversation was not elicited. Instead, the experimenter (the fifth author) uttered pre-planned statements (probes) such as “*I like to eat fruit*” interspersed between standardised tests or games. There were 16 such statements, organised by topic into groups of four (Appendix A). The first statement of a block related to a task or game the child had just completed. The immediate child responses to these 16 conversation probes were video- and audio-recorded and analysed.

Data exclusion

If the child did not respond verbally but the experimenter did not leave at least 2000 msec between the offset of her pre-planned statement and the onset of her following conversational statement, then this item was excluded from analyses. Furthermore, if the experimenter deviated from the syntax of the planned conversational probe, this item was excluded from analyses.

Coding conversational responses

For all conversation probes which met the inclusion criteria (see above), each child's response was coded as either 'verbal response' or 'no language used'. An explanation of the coding scheme, together with some example participant responses to the probe "I have a pet", is outlined in Table 2 below.

Table 2: Conversation coding scheme

	Coding category	Definition	Example response from Study 1 data
Sentential response	Contingent Responses	Multi-word utterances which are relevant and which elaborate on the content of the conversation partner's turn (Bloom et al., 1976)	P12: 'I've got 2 pets' P18: 'Is it a tiger?'
	Noncontingent Responses	Either (i) were not on the topic of (or only very tangentially related to) the immediately preceding turn, ii) had a very unclear meaning or iii) were 'bizarre' or iv) switching to talk about something in the environment (Nadig et al., 2010)	P51: 'Yeah there's a new boy and he punched him like some chocolate and he went [NOISE] in the face
Minimal response	Specific minimal responses	Were either (i) one word utterances which were clearly specific to the content of the preceding turn (e.g. E: 'I have a pet' C: '(A) Doggy?') or ii) short phrases which encouraged the conversation partner to keep talking (e.g. 'Did you?', 'Was it?') but which did not elaborate on the topic (Bavelas et al., 2000)	P15: A doggy! P31: What? P28: Have you? (Other possible responses: "Really?", "Cool!")
	Generic minimal responses	Were single words or short phrases which might be acknowledgements of the conversation partner's turn but which could potentially be uttered without the speaker listening to or processing the conversation partner's turn (Bavelas et al., 2000; Jefferson, 1984)	P13: 'Mm-hmm' (Other possible responses: 'Yeah' 'Oh' 'Uh-huh' 'Right')
No language used	Nonverbal Responses	Non-verbal behaviour which acknowledge or respond to the conversation partner's turn	Nods Raises eyebrows

			Smiles in response
	Null response	The child either looked at something in the room or else looked at the experimenter's face with no discernible reaction	

We also included repetitions (partial or full) in with ‘generic minimal’ – although these can serve a communicative function (Luyster et al., 2022) – since there were only four repetitions in our whole dataset.

Inter-rater reliability (IRR)

The verbal responses for 23.3% of the data were second coded by a native English-speaking Psychology student, who was ‘blind’ to both the ratings of the main coder as well as to the diagnostic status of the child. There was excellent agreement between the coders ($k = .92$). For non-verbal behaviours, 30% of the dataset was second coded in the same way and reliability was good (Cohen's $k = .83$).

Results

The full anonymised dataset is available on the Open Science Framework (OSF) here <https://osf.io/q7wa4/> The mean proportion of each type of conversational response is shown in Table 3.

Table 3. Mean (SD in brackets) proportion of responses to conversational probes (see Table 2 for definitions)

		Autistic	NT
		Mean (SD)	Mean (SD)
Sentential response	Contingent	.484 (.5)	.565 (.496)
	Noncontingent	.135 (.342)	.042 (.202)
Minimal response	Specific	.120 (.33)	.116 (.32)
	Generic	.097 (.30)	.038 (.19)
No language used	Non-verbal Responses	.015 (.121)	.078 (.268)
	Null response	.145 (.352)	.15 (.356)

We fitted a mixed-effects multinomial logistic regression model using STAN via *brms* (Bürkner, 2017) with the default priors. The six response types (contingent, non-contingent, specific minimal, generic minimal, non-verbal response, null response) were the outcome variables, with null response being the reference response. Diagnostic group (Autistic vs. Neurotypical) was included as an effect-coded fixed effect and participant and item were included as random effects on the intercept.




Figure 1: Log odds of each response (relative to the odds of producing no response) for the two participant groups with 95% credible intervals.

Figure 1 shows the estimated log odds ratios for each of the responses for each group, relative to the odds of producing no reaction. The dot for each response/group represents the best estimate for the log odds of that response in that group. The bars represent the uncertainty around this estimate — the range in which the value falls with 95% probability. The vertical dashed line represents the log odds of producing no response. Only the estimates for contingent responding are greater than the estimates for not responding. Critically for each response type, where the best estimate for one group falls outside the interval for the other, we can say that there is < 0.05 probability (two-tailed) that that response has the same likelihood of occurrence for that group as for the other. This is the case for three of the six response categories, with neurotypical children showing lower rates of non-contingent (off-

topic) responses ($p_{\text{mcmc}} < 0.005$) and generic minimal responses (i.e. where it is difficult to know for sure that the child processed the content) ($p < 0.05$) than autistic children, but higher rates of non-verbal responses such as nodding and / or smiling in response ($p_{\text{mcmc}} < 0.0005$).

Thus, overall the findings are mixed. The three aforementioned significant differences between groups fit with the diagnostic criteria and with previous literature (e.g., Nadig et al., 2010). However, Table 3 and Figure 1 reveal striking similarities between groups in their ability to use contingent responses and specific minimal responses to keep a conversation topic going.

To further explore this skill amongst the autistic group, we then followed the method in Pagmar et al. (2022) in conflating over the three response categories which could be termed '*topic-supporting*' (which Pagmar et al., 2022, refer to as 'appropriate') in a conversation, namely contingent responses, specific minimal responses (see Table 2 for definitions) and non-verbal responses. For this conflated 'overall topic-supporting responses' category, we built a binary logistic regression model and found that there was a significant between-groups difference (Est = 0.85, SE = 0.34, $z = 2.516$, $p = .012$), largely driven by lower rates in the autistic group of non-verbal responding (i.e. gestures or use of facial affect to respond). However, it is important to highlight that our sample of young primary-school-aged autistic children gave topic-supporting responses to 62% of the conversation probes, indicating that even at this young age, verbally fluent autistic children are able to respond in a way that maintains a conversation topic a large proportion of the time.

Interim Discussion

Although for many of conversational response types, the above findings are consistent with claims (and diagnostic criteria) that highlight conversational weaknesses in autistic children,

for other response types there is notable overlap between the diagnostic groups. Moreover, we noticed high variability in both populations. In Study 2 we explored these individual differences in neurotypical children further by investigating the cognitive correlates of conversation ability. We also elicited full conversations as opposed to responses to conversation probes. We then explored individual difference in autistic children in Study 3.

Study 2

Method

Participants

The study included 48 neurotypical, monolingual British-English-speaking children (24 female), aged between 5;11 and 6;11 years ($M = 77.56$ months/6;5 years, $SD = 3.77$), recruited via and tested in the Kent Child Development Unit between July 2017 and February 2018.

All parents reported that their children had no known or suspected hearing, learning or language difficulties and none had been referred for any neuro-developmental disorders. The sample size was originally planned to be 75 but had to be reduced to 48 due to competing constraints on participant availability (i.e. external funding was obtained by another researcher in the department).

Elicitation of conversation

Conversation was elicited by one of two research assistants (RAs), first in relation to one or more of the child's interests and then on one or more of the specific topics of siblings, pets and families. The RAs (female Psychology graduates in their early twenties) were trained to avoid using a questioning style. The RAs prepared two or three declarative statements per conversation to use at topic-relevant moments. The majority of conversation was unplanned and free-flowing (see Appendix 2 for details).

Additional Measures and procedure

First, *Expressive Vocabulary* was assessed using a sub-test of the CELF-4 (Semel et al., 2006). Then, *Theory of Mind (ToM)* was assessed via the following commonly administered vignettes: ‘Birthday Surprise’ (Sullivan et al., 1994), ‘Robots’ (Coull et al., 2006), two of Happé’s (1994) strange stories - lying to persuade (‘Kittens’) and Malkin et al.’s (2018) version of ‘double-bluff’. For the ToM verbal responses 14.6% of the data were coded by a second coder, who was blind to scores of the main coder (Cohen’s $k = .96$). For analyses, we summed across all ToM measures.

After this first bank of tests, the first conversation was elicited following the procedure above. This was followed by a second bank of tests. *Inhibitory Control (IC)* was assessed via a child-friendly version of the Flanker task (Blakey et al., 2020), where the measure is the difference between mean reaction times in the incongruent and congruent conditions. *Working Memory* was assessed using the Backwards Digit Span sub-scale of the Wechsler Intelligence Scale for Children (WISC) (Wechsler, 2003). Then, a second conversation was elicited. Finally, *Non-verbal Reasoning* was assessed using the ‘Matrices’ sub-test BAS-3 (Eliot & Smith, 2011). Of the predictors, only vocabulary was significantly correlated with other predictors, namely age ($r(45) = .34, p = .021$), and Theory of Mind ($r(45) = .37, p = .013$).

All except one parent completed four sub-scales of the Children’s Communication Checklist 2 (Bishop, 2003), namely D (Coherence), E (Inappropriate Initiation), F (Stereotyped Language) and G (Context) in the waiting room during the testing session².

² Following a reviewer’s suggestion, we examined relationships between the composite of these CCC2 subscales and contingent responding and found there was indeed a weak but significant relationship ($r = .34, p = .02$).

Conversation coding and Inter-rater reliability (IRR)

Child immediate responses to experimenter turns were coded by the first author using the categories used for Study 1. Since Studies 2 and 3 elicited conversation (as opposed to merely a response to one statement), here we also coded (following Nadig et al., 2010) child utterances that followed on from their own prior utterance as ‘elaborations’ (see details below). Five children (i.e., 390 child utterances) were second coded by a Linguistics Masters student blind to the original coding (Cohen’s $k = .88$).

Results

We followed a pre-registered analysis plan (<https://osf.io/q7wa4/registrations>) to first explore children’s responses to experimenter declarative statements and, second, to explore children’s elaborations on their own prior turn.

Child direct responses to experimenter turns

Distribution of conversational response types

Following our pre-registered analysis plan, we focused on the analyses of the following contrasts:

- a) overall *topic-supporting* responses (the sum of contingent responses, specific minimal responses and non-verbal responses) as opposed to responses which are either a hurdle for the topic or do not particularly support it (noncontingent, generic minimal responses, null responses);
- b) *contingent* responses as opposed to all *other* response types (i.e., non-contingent, all minimal and all non-verbal and null responses);
- c) *non-contingent* responses as opposed to all *other* response types (i.e., contingent, all minimal and all non-verbal and null responses).

For comparability with Study 1, mean proportions and mean raw utterances for all fine-grained coded categories of direct responses to an experimenter declarative statement (as well as the summed measure of overall topic-supporting responses) are provided in Table 4.

Table 4: Study 2 (neurotypical) child direct response to experimenter turn (mean proportion, mean raw by child)

Superordinate	Utterance type	Mean prop (SD)	Mean raw (SD)
Topic-supporting	Contingent	0.51 (0.18)	23.75 (10.19)
	Specific minimal	0.09 (0.07)	4.54 (3.86)
	Non-verbal	0.09 (0.09)	4.17 (3.89)
Overall topic-supporting responses to experimenter turns		0.69	32.46
	Non-contingent	0.06 (0.05)	2.94 (2.35)
	Generic minimal	0.14 (0.11)	7.38 (6.76)
	Null response	0.10 (0.11)	4.46 (5.11)

Investigating the role of cognitive and socio-cognitive predictors

For each of the three outcome variables (overall Topic-supporting responses, Contingent and Non-contingent), we built mixed effects logistic regression models with Age, Expressive Vocabulary, Theory of Mind, Inhibitory Control, Working Memory and Nonverbal Reasoning as predictor variables (converted to z scores) and participant as a random effect on the intercept. The p value for each predictor was calculated by using likelihood ratio tests to compare the full model to a model without that particular predictor. For these analyses below, due to missing data for some predictor variables, the sample size

was 39 because non-verbal reasoning scores were missing for five participants, vocabulary scores were missing for an additional three participants and the incorrect Flanker task version was used for an additional participant. (However, zero order correlations included the full sample, where possible, see Appendix 4).

For overall topic-supporting responses, the only predictors to explain unique variance were age ($b = 0.28$, $r^2 = 6.11$, $p = .01$) and the theory of mind composite ($b = 0.27$, $r^2 = 4.43$, $p = .04$), which were both positively related³. For contingent responses, the same pattern was found, where both age ($b = 0.48$, $r^2 = 11.11$, $p < 0.001$) and theory of mind ($b = 0.33$, $r^2 = 4.52$, $p = .03$) were positive predictors⁴.

For non-contingent responses the only predictors to approach significance were the theory of mind composite ($b = -0.30$, $r^2 = 3.29$, $p = .07$) and working memory ($b = -0.26$, $r^2 = 2.92$, $p = .087$), such that as these skills increased, non-contingent responses became less likely⁵.

Although non-contingent responses were a relatively infrequent utterance category (see Table 3 above), the individual differences in their usage deserve highlighting. On the one hand, 17% of these children never gave a non-contingent response. Conversely, 30% of the children gave non-contingent responses to an experimenter turn in more than 10% of their responses. Children's non-contingent responses tended to either involve thinking of a more

³ The same pattern was found when each Theory of Mind measure was entered as a separate predictor, whereby age was then marginally significant ($b = 0.23$, $r^2 = 3.82$, $p = .051$), one ToM measure – Strange Stories was a significant positive predictor ($b = 0.29$, $r^2 = 5.78$, $p = .016$) but the other ToM measure (Knowledge-Belief) was not significant ($p = .55$).

⁴ The same pattern of results was found when each Theory of Mind measure was entered separately, namely age was a significant positive predictor ($b = 0.43$, $r^2 = 8.37$, $p = 0.004$) and the Strange Stories Theory of Mind measure was also a significant positive predictor ($b = 0.31$, $r^2 = 4.4$, $p = 0.036$), whereas the Knowledge-Belief predictor was not significant ($p = .38$).

⁵ The same pattern of results was seen when the two Theory of Mind measures were entered as separate predictors whereby here one of the Theory of Mind measures (Knowledge-Belief) then reached significance ($b = -0.55$, $r^2 = 10.28$, $p = .0014$) and Strange Stories did not ($p = .35$). Working Memory remained as a marginally significant negative predictor ($b = -0.26$, $r^2 = 3.52$, $p = .061$).

interesting topic (see P2 in the introduction), becoming distracted by the environment (P13 below) or returning to a previous topic (P4 below).

RESEARCHER (R): “He’s like a fish.”

P13: [REFERRING TO HER SHOES] “I’m trying to do with these over a bit so they’re a bit tighter.”

RESEARCHER: “So all his powers were gone and he couldn’t do anything about it.”

[TALKING ABOUT BATMAN]

P4: “I might do that with toothpaste and sugar.” [RETURNS TO PREVIOUS TOPIC ABOUT PLAYING PRANKS ON SIBLINGS]

Child elaborations on their own preceding turn

In addition to responding to researcher turns, children frequently elaborated on their own prior turn. Following Nadig et al. (2010: 2735), we categorised elaborations into ‘contingent’ elaborations and ‘self-contingent elaborations’. Contingent elaborations involved extended talk which stayed on the sub-topic established by the experimenter. The other two categories were self-contingent and non-contingent elaborations. Self-contingent elaborations were utterances where the child started (during his / her extended response) to veer off from the topic of the experimenter’s last utterance. Non-contingent elaborations involved the child switching from (her own) topic to another. Some illustrations from the dataset are provided in the coding manual (on OSF).

F k u v t k d w v k q p " q h " e j k n f " ÷ e q p x g t u c v k q p " g n c d q t c v

Table 5 below indicates the mean proportion and mean raw frequency of each sub-type of child elaboration (follow in) on his / her own response.

Table 5: Study 2 (neurotypical) child elaboration type (mean proportion, mean raw by child)

Superordinate	Utterance type	Mean prop. (SD)	Mean raw (SD)
Topic-supporting	Contingent Elaboration	0.72 (0.21)	25.69 (15.50)
Non-reciprocal	Self-contingent Elaboration	0.24 (0.20)	11.29 (11.46)
Non-reciprocal	Non-contingent Elaboration	0.04 (0.07)	1.31 (2.51)
	Elaboration - indeterminate	0.00 (0.00)	0.02 (0.14)

Investigating the role of cognitive and socio-cognitive predictors

Following the pre-registered analysis plan, we fitted regression models for each type of elaboration. Here we deviated slightly from the pre-registration by carrying out mixed effects logistic regression (and not linear regression) with the target type of elaboration coded as ‘1’ and the other types of child elaboration coded as zero)⁶. For self-contingent elaborations, the only predictor to explain unique variance was working memory ($b = -0.50$, $SE = 0.23$, $\chi^2 = 5.1$, $p = 0.024$), which was negatively related. For contingent elaborations, there was a marginally significant positive relationship with working memory ($b = 0.33$, $SE = 0.17$, $\chi^2 = 3.81$, $p = 0.051$).

⁶ A logistic regression was employed as more appropriate to proportions which are bounded by 0 and 1, than linear regression which assumes a continuous scale. However, in fact, we obtain the same pattern of results if we follow the pre-registration exactly and treat self-contingent elaborations as proportion scores (using all elaborations as the denominator) and running multiple linear regression analyses. The same goes for contingent elaborations, although in this case the trend in the direction of a positive significant relationship for working memory is less marked ($p = .07$).

It is important to highlight the wide individual differences; 19% of children never produced self-contingent or non-contingent elaborations. Nonetheless, indicators of off-topic verbosity was common in these neurotypical six-year-olds; 25% of children produced self-contingent elaborations in 20% or more of their responses.

Discussion

There is a great deal of variability amongst neurotypical, monolingual six-year-olds regarding if and how they manage to maintain reciprocal conversation. Around a third demonstrate quite a high degree of either non-contingent responding and / or off-topic verbosity. In line with the (relatively scant) literature to date, Study 2 indicates that these individual differences in conversational topic maintenance align with individual differences in theory of mind (De Rosnay et al., 2014; Slomkowski & Dunn, 1996) and working memory (Blain-Brière et al., 2014). Our next study addressed the question of whether a similar pattern of interrelationships is seen for individual differences amongst autistic children regarding their conversational topic maintenance.

Study 3

Method

Participants

Participants were scheduled for an Autism Diagnostic Observation Schedule (ADOS) 2 (Lord et al., 2012) module 3 assessment with a clinical psychologist (the 4th author) as part of an NHS multi-disciplinary autism assessment. 50 children were tested (13 female) between January 2019 and March 2020. Of these, 41 received an autism diagnosis and all scored above threshold on the ADOS-2 total algorithm scores (mean score = 12.7). In accordance

with the requirements of module 3, part of the eligibility criteria included speaking English fluently. All children indeed spoke fluently in sentences and with southern English accents.

One autistic child did not provide any conversation data during the ADOS-2 and was excluded from analyses. Thus, the final sample comprised 40 children (10 female) aged between 5;4 and 9;10. We had originally aimed to include 60 autistic children (sample size determined by budget) but direct assessment was curtailed due to the outbreak of COVID-19 and first UK national lockdown.

Conversation

Conversation was elicited as part of the ADOS-2 module 3 administration which assesses the child's ability to "build on [psychologist's] statements ... and take a full role in back-and-forth conversation.". We excluded all verbal interaction that was part of the elicited narratives, make-believe play, interactive play, construction task, interview about friendships and other ADOS-2-related tasks. The administrator always included the majority of the pre-planned statements listed in the supplementary materials. The transcribed conversations were all coded by the same Linguistics Masters student who coded Study 2 using the same criteria (see <https://osf.io/q7wa4/> for the coding manual). The first author coded five children (i.e., 755 child utterances) blind to the original coding with good inter-rater reliability (Cohen's $k = .872$).

Additional Measures and procedure

After the child had completed the ADOS-2, an RA assessed the child on the following measures in the following order.

Non-verbal reasoning was assessed using raw scores on the Matrices sub-test in the Wechsler Preschool and Primary Scales of Intelligence (Wechsler, 2012). Then we carried out the

Backwards Word Span measure of verbal working memory (Blakey et al., 2020). After that, *Expressive Vocabulary* was assessed using raw scores on the CELF-4 (as for Study 2).

Following this, theory of mind was assessed in ‘stepwise’ manner. First, all children were administered two first-order false belief measures (Perner et al., 1987; Wimmer, 1983). If a child passed both, then the RA administered the advanced test items ‘Birthday Surprise’ (Sullivan et al., 1994) and ‘Robots’ (Coull et al., 2006). If, however, the child failed one or both first-order ToM tests, then it was assumed that the latter advanced measures would be failed and instead three measures of ‘precursor’ theory of mind were administered (Pillow, 1989; Wellman & Liu, 2004). In this way, without administering too many tests, it was possible to allocate a child a score on a theory of mind composite, which was then used in analyses.

Finally, the RA administered the Backwards Digit Span from the WISC (Wechsler, 2003). In analyses, for ‘*Working Memory*’ we used a composite (mean z scores) of the Backwards Digit raw scores and Backwards Word Span raw scores, which were highly positively correlated ($r(40) = .56, p < .001$).

All the above predictor variables were significantly positively correlated (see Table 9 Supplemental), with the exception of age, which was only correlated with the vocabulary raw score ($r(df = 38) = .41, p = .01$).

Results

This study was pre-registered (see <https://osf.io/q7wa4/registrations>) and the same analysis plan employed as for Study 2 with one exception — we did not test inhibitory control. (See Supplemental Table 10 for correlations).

Child direct responses to experimenter turns

Distribution of conversational response types

Table 6 below illustrates mean proportions and mean raw frequencies for all coded categories of direct responses to a conversational turn by the ADOS-2 administrator. Due to the NHS ethics required for this study, we did not have access to video-recordings and thus could not code non-verbal responses.

Table 6: Study 3 (autistic) child direct response to ADOS-2 administrator turn (mean proportion, mean raw frequency)

Superordinate	Utterance type	Mean prop (SD)	Mean raw (SD)
Topic supporting	Contingent	0.33 (0.16)	26.98 (14.31)
	Specific Minimal	0.21 (0.08)	16.90 (7.32)
	Laughing response	0.03 (0.03)	2.13 (2.41)
<i>Overall proportion topic-supporting responses to experimenter turns</i>		0.57	46.00
	Non-contingent	0.10 (0.10)	7.53 (6.48)
	Generic minimal	0.19 (0.11)	15.88 (11.48)
	No audible response	0.14 (0.14)	10.88 (10.25)

Investigating the role of cognitive and socio-cognitive predictors

No predictors explained unique variance in contingent responding. However, as for Study 2, unique variance in all topic-supporting responding was explained by theory of mind ($b = 0.38$, $SE = 0.18$, $r^2 = 4.39$, $p = .036$). For the non-contingent responses, age was the only predictor to explain unique variance ($b = -0.64$, $SE = 0.27$, $r^2 = 5.64$, $p = .018$). This was

negatively related such that (verbally-fluent) autistic children become less non-contingent with age.

Child elaborations on their own preceding turn

Table 7 indicates the mean proportion and mean raw frequencies of each sub-type of child elaboration. No predictor explained unique variance for any of the elaboration sub-types.

Table 7: Study 3 (autistic) child elaboration type (mean proportion, mean raw by child)

Superordinate	Utterance type	Mean prop (SD)	Mean raw (SD)
Topic-supporting	Contingent Elaboration	0.60 (0.24)	20.58 (14.50)
Non-reciprocal	Self-contingent Elaboration	0.33 (0.21)	14.65 (16.03)
Non-reciprocal	Non-contingent Elaboration	0.07 (0.09)	2.50 (3.15)
	Elaboration - indeterminate	0.00 (0.00)	0.03 (0.16)

Individual differences

Around a third of our autistic sample here showed either non-contingent responding to the experimenter or showed self-contingent elaborations (off-topic verbosity) in 20% or more of their utterances. While a very small handful of autistic participants showed fairly extreme tendencies in this regard, at the other extreme there was also a handful of autistic children who never responded non-contingently nor exhibited any off-topic verbosity.

Discussion

Compared to study 2, the relationship between the cognitive and socio-cognitive predictors and measures of conversation was less clear. Nonetheless, topic-supporting responding overall was explained by theory of mind as for study 2. Autistic children also became less

likely to respond non-contingently with age. Finally, as for Study 2 (with neurotypical children), the autistic children in Study 3 showed great heterogeneity in their conversational behaviours.

General Discussion

We carried out three studies in which we examined conversational skill by categorising children's turns during live interaction. Study 1 examined conversational differences and similarities between verbally-fluent autistic 5- to 7-year-olds and their neurotypical peers. Study 2 investigated relationships between conversation and theory of mind, working memory and inhibitory control in neurotypical 6-year-olds. Study 3 investigated relationships between conversation, theory of mind and working memory within autistic 5- to 9-year-olds.

Our main analyses in all three studies examined direct responses to statements uttered by the conversation partner. We first conflated all responses which supported the topic either by being relevant or by acknowledging the specific content of the immediately preceding turn. However, not all 'topic-supporting' responses contribute equally to maintaining a conversation topic; topic-supporting responses include specific minimal responses (such as 'Wow!') and non-verbal gestures and facial affect responses. While the latter acknowledge the conversation partner's contribution and encourage him or her to continue, they do not extend a conversation. Therefore, we also investigated 'contingent' responses, which by definition keep a conversation going since they elaborate on the topic of the conversation partner's turn. In terms of responses, which do not support the conversation topic, we focused, first, on 'non-contingent' responses, since these are highlighted in the literature as being particularly problematic for conversational back-and-forth (Bloom et al., 1976; Hale & Tager-Flusberg, 2005; Nadig et al., 2010). Second, in Studies 2 and 3 (for which we elicited extended

conversation), we also examined ‘self-contingent elaboration’ (Nadig et al., 2010), which is akin to the ‘off-topic verbosity’ (Pushkar et al., 2000; Ruscher & Hurley, 2000).

In Study 1, we found that autistic children were less likely to give non-verbal responses (gestures or facial affect responses) and more likely to respond non-contingently and / or to give generic minimal responses (e.g. ‘mm’ or ‘oh’). This findings is in keeping with the diagnostic criteria for autism (American Psychiatric Association, 2013) and highlights an aspect of difficulty that contributes to wider social-communication differences in autism. However, it is important to stress that the between-group differences in Study 1 suggest a *relative* difficulty with conversation among (verbally-fluent) autistic children, rather than an inability to support a conversation topic, with a number of autistic children showing highly skilled conversational turn taking. Indeed, across conversational response types, the evidence from Study 1 points to important similarities, as well as differences, in terms of conversation skills across neurotypical and (verbally-fluent) autistic children.

Moreover, although we cannot compare Study 2 and Study 3 directly, it is interesting that in both studies both working memory and theory of mind emerge as correlates of conversational topic maintenance. In Study 2, working memory related negatively to those types of conversational behaviour, which are a hurdle to reciprocal conversation; working memory was related to self-contingent elaboration (off-topic verbosity) and was marginally related to non-contingent responding to statements in regression analyses. Similarly, there were indications in Study 3 that working memory limits may play a role in difficulties in maintaining a conversation topic; although working memory was not a significant predictor of any conversation variables in the main regression analyses (responses to clinician declarative statements), it was correlated (see Appendix 10) and it was a significant positive predictor of contingent responses to the clinician, if we included responses all clinician turns including questions (Appendix 13). The findings for working memory extend those found by

Blain-Brière et al. (2014). Furthermore, in Study 2, working memory related negatively to types of utterances that are lengthy but tend to elicit negative social impression ratings (self-contingent elaboration or off-topic verbosity) whereas working memory related positively to those which are lengthy but which do not veer off the original topic (contingent elaboration or ‘general chattiness’). We also found a marginal relationship between working memory and noncontingent responses to the experimenter in Study 2. One interpretation of these findings could be that working memory is required to retain the current conversation topic at the forefront of one’s mind.

Regarding theory of mind, over Studies 2 and 3, this emerged as a significant predictor variable for various conversational response types (when controlling for all the other key variables: age, vocabulary, non-verbal reasoning, working memory and (for Study 2) inhibitory control). While previous studies have shown a relationship between contingent responses in conversation and theory of mind in both neurotypical (Slomkowski & Dunn, 1996) and autistic children (Hale & Tager-Flusberg, 2005), none have shown this relation to be independent of working memory ability. This is the first study to show this and provides support for proposals that an in-depth consideration of the conversation partner’s perspectives is intertwined with conversational proficiency (De Rosnay et al., 2014; Tomasello, 2018). That the pattern of relationships with conversation topic maintenance appear to be similar among autistic and neurotypical children suggests that individual differences in theory of mind and working memory influence conversational ability regardless of diagnostic status.

Limitations

While the current studies benefitted from eliciting relatively natural, live conversation and a rich, linguistically-informed coding scheme, there are a number of limitations. Sample sizes were small for regression analyses. Moreover, some of the cognitive and social-cognitive

measures relied on a single test. In addition, stipulations from NHS ethics meant that we were not allowed to store video-recordings of the ADOS and were thus unable to score non-verbal responses for Study 3. Ideally, we would liked to have worked with autistic adolescents to discuss the conversation coding scheme.

Important steps for future studies

To gain a more precise insight into the degree to which autistic and neurotypical child conversation shows similar patterns of relationships to theory of mind and working memory, an important future step for a future study would be to carry out (large scale) individual differences studies of both autistic and neurotypical children in which conversations are elicited in the same manner and in the same settings. Ideally such future studies should also include individual differences measures of both language processing speed and attention (or tendency for distraction). There is evidence both from populations with no diagnoses (Barnett et al., 2022) and autistic children (Howard et al., 2023) that the ability to sustain attention is related to the amount of expressive language produced. Regarding language-processing speed, a number of studies have found a relationship between autism severity and prolonged turn-taking gaps in conversation for both autistic children (McKernan et al., 2022; Parish-Morris et al., 2016) and adults (Ochi et al., 2019). Therefore, the contributions of attention and language processing speed to social conversational ability need exploration.

Implications

Perhaps the most important finding of the current study is that the conversation skills that are characteristically difficult to deploy for autistic children are also difficult for very many neurotypical children, apparently for similar reasons. Given that conversation skills are

part of the primary school curriculum in many countries, this might be taken to suggest that there should be an increased emphasis on empowering mainstream teachers to scaffold and allow practice time for social conversation in classroom settings (Abbot-Smith et al., 2023). The evidence base regarding how to do this effectively is currently incredibly thin. Future research should test the value of different approaches to supporting conversation skill for a diverse range of children. This might include, for example, following up on the finding that theory of mind is related to the ability to maintain a conversation by testing whether actively encouraging consideration of conversation partner's current mental states (and how to determine / read these) is helpful (Adams et al., 2012; Winner, 2007). Likewise testing ways to mitigate the effects of limited working memory would be worthwhile. All such research should be conducted with teachers and children and would also open the opportunity to talk meta-linguistically about differences in what each of us needs and prefers when having a conversation.

Conclusions

Although Study 1 found group-level differences between verbally-fluent autistic primary-school children and their neurotypical peers in their ability to support and maintain a conversation topic, there were also striking similarities. Moreover, across both Studies 2 and 3 we found relationships between topic-supporting responding and theory of mind, with some evidence for the cost of limited working memory. Continuity across diagnostic groups regarding responses which are not topic-supporting suggests a need for universal support for social conversation skills within schools, particularly since social communication predicts peer relations, emotional stability and behavioural problems (Conti-Ramsden et al., 2019; Law et al., 2014; Roy & Chiat, 2014; Saul et al., 2022).

References

- Abbot-Smith, K., Dockrell, J., Sturrock, A., Matthews, D., & Wilson, C. (2023). Topic maintenance in social conversation: What children need to learn and evidence this can be taught. *First Language* 43(6), 614–642. <https://doi.org/10.1177/01427237231172652>
- Adams, C., Green, J., Gilchrist, A., & Cox, A. (2002). Conversational behaviour of children with Asperger syndrome and conduct disorder. *Journal of Child Psychology and Psychiatry* 43(5), 679–690. <https://doi.org/10.1111/1469-7610.00056>
- Adams, C., Lockton, E., Freed, J., Gaile, J., Earl, G., McBean, K., Nash, M., Green, J., Vail, A., & Law, J. (2012). The Social Communication Intervention Project: A randomized controlled trial of the effectiveness of speech and language therapy for school-age children who have pragmatic and social communication problems with or without autism spectrum disorder: RCT of speech/language therapy for children with pragmatic language impairment. *International Journal of Language & Communication Disorders* 47(3), 233–244. <https://doi.org/10.1111/j.1460-6984.2011.00146.x>
- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders* (Fifth Edition). American Psychiatric Association. <https://doi.org/10.1176/appi.books.9780890425596>
- Baixauli-Fortea, I., Miranda Casas, A., Berenguer-Forner, C., Colomer-Diago, C., & Roselló-Miranda, B. (2019). Pragmatic competence of children with autism spectrum disorder. Impact of theory of mind, verbal working memory, ADHD symptoms, and structural language. *Applied Neuropsychology: Child* 10(2), 101–112. <https://doi.org/10.1080/21622965.2017.1392861>
- Barnett, M., Hephrey, J., & Childers, L. (2022). Attention problems and off-topic verbosity among young adult and older adult age cohorts. *Applied Neuropsychology: Adult* 41(8). <https://doi.org/10.1080/23279095.2022.2147009>
- Bavelas, J. B., Coates, L., & Johnson, T. (2000). Listeners as co-narrators. *Journal of Personality and Social Psychology* 79(6), 941–952. <https://doi.org/10.1037/0022-3514.79.6.941>

Bierman, K. L., & Furman, W. (1984). The Effects of Social Skills Training and Peer Involvement on the Social Adjustment of Preadolescents. *Child Development*, 55(1), 151.

<https://doi.org/10.2307/1129841>

Bishop, D. V. M. (2003). (Vol. 2). Harcourt Assessment.

Blain-Brière, B., Bouchard, C., & Bigras, N. (2014). The role of executive functions in the pragmatic skills of children age 4 to 5. *Frontiers in Psychology*

<https://doi.org/10.3389/fpsyg.2014.00240>

Blakey, E., Matthews, D., Cragg, L., Buck, J., Cameron, D., Higgins, B., Pepper, L., Ridley, E., Sullivan, E., & Carroll, D. J. (2020). The Role of Executive Functions in Socioeconomic Attainment Gaps: Results From a Randomized Controlled Trial. *Child Development*, 91(5), 1594–1614.

<https://doi.org/10.1111/cdev.13358>

Bloom, L., Rocissano, L., & Hood, L. (1976). Adult-child discourse: Developmental interaction between information processing and linguistic knowledge. *Cognitive Psychology*, 8(4), 521–552. [https://doi.org/10.1016/0010-0285\(76\)90017-7](https://doi.org/10.1016/0010-0285(76)90017-7)

Bürkner, P.-C. (2017). **brms**: An RPackage for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, 80(1). <https://doi.org/10.18637/jss.v080.i01>

Cola, M., Zampella, C. J., Yankowitz, L. D., Plate, S., Petrulla, V., Tena, K., Russell, A., Pandey, J.,

Schultz, R. T., & Parish-Morris, J. (2022). C

with autism: Differences in talkativeness across contexts. *Autism Research*, 15(6), 1090–

1108. <https://doi.org/10.1002/aur.2693>

Conti-Ramsden, G., Mok, P., Durkin, K., Pickles, A., Toseeb, U., & Botting, N. (2019). Do emotional difficulties and peer problems occur together from childhood to adolescence? The case of children with a history of developmental language disorder (DLD). *European Child & Adolescent Psychiatry*, 28(7), 993–1004. <https://doi.org/10.1007/s00787-018-1261-6>

Coplan, R. J., Hughes, K., Bosacki, S., & Rose-Krasnor, L. (2011). Is silence golden? Elementary school

teachers' strategies and beliefs regarding h y

- children. *Journal of Educational Psychology*, 100(4), 939–951.
<https://doi.org/10.1037/a0024551>
- Costantino, J., & Gruber, C. (2005). *Social Responsiveness Scale*. Western Psychological Services.
- Coull, G. J., Leekam, S. R., & Bennett, M. (2006). Simplifying Second-order Belief Attribution: What Facilitates Children's Performance on Social Measurement Developmental (52), 260–275. <https://doi.org/10.1111/j.1467-9507.2006.00340.x>
- De Rosnay, M., Fink, E., Begeer, S., Slaughter, V., & Peterson, C. (2014). Talking theory of mind talk: Young school-aged children's everyday conversation and emotion. *Journal of Child Language*, 41(5), 1179–1193.
<https://doi.org/10.1017/S0305000913000433>
- Eliot, C., & Smith, P. (2011). *British Ability Scales, BAS-2*. Assessment.
- Fizke, E., Barthel, D., Peters, T., & Rakoczy, H. (2014). Executive function plays a role in coordinating different perspectives, particularly when one's Cognition, 13(3), 315–334. <https://doi.org/10.1016/j.cognition.2013.11.017>
- Friedman, L., Sterling, A., DaWalt, L. S., & Mailick, M. R. (2019). Conversational Language Is a Predictor of Vocational Independence and Friendships in Adults with ASD. *Journal of Autism and Developmental Disorders*, 49(10), 4294–4305. <https://doi.org/10.1007/s10803-019-04147-1>
- Hale, C. M., & Tager-Flusberg, H. (2005). Social communication in children with autism: The relationship between theory of mind and discourse development. *Autism*, 9(2), 157–178.
<https://doi.org/10.1177/1362361305051395>
- Happé, F. G. E. (1994). An advanced test of theory of thoughts and feelings by able autistic, mentally handicapped, and normal children and adults. *Journal of Autism and Developmental Disorders*, 24(1), 129–154.
<https://doi.org/10.1007/BF02172093>

Hazen, N. L., & Black, B. (1989). Preschool Peer Communication Skills: The Role of Social Status and Interaction Context. *Child Development*, 60(4), 867. <https://doi.org/10.2307/1131028>

Howard, J., Herold, B., Major, S., Leahy, C., Ramseur, K., Franz, L., Deaver, M., Vermeer, S., Carpenter, K. L., Murias, M., Huang, W. A., & Dawson, G. (2023). Associations between executive function and attention abilities and language and social communication skills in young autistic children. *Autism*, 27(7), 2135–2144. <https://doi.org/10.1177/13623613231154310>

Hutchison, S. M., Müller, U., & Iarocci, G. (2020). Parent Reports of Executive Function Associated with Functional Communication and Conversational Skills Among School Age Children With and Without Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 50(6), 2019–2029. <https://doi.org/10.1007/s10803-019-03958-6>

Jefferson, G. (1984). Notes on a systematic deep
“Minimalist Paper in Linguistics”, 197–216. <https://doi.org/10.1080/08351818409389201>

Law, J., Rush, R., & McBean, K. (2014). The relative roles played by structural and pragmatic language skills in relation to behaviour in a population of primary school children from socially disadvantaged backgrounds. *Emotional and Behavioural Difficulties*, 19(1), 28–40. <https://doi.org/10.1080/13632752.2013.854960>

Lecce, S., Bianco, F., Devine, R. T., & Hughes, C. (2017). Relations between theory of mind and executive function in middle childhood: A short-term longitudinal study. *Journal of Experimental Child Psychology*, 169, 69–86. <https://doi.org/10.1016/j.jecp.2017.06.011>

Lord, C., Rutter, M., DiLavore, P., Risi, S., Gotham, K., Bishop, S., Luyster, R., & Guthrie, W. (2012). *Autism diagnostic observation schedule, Second Edition (ADOS-2)*. Western Psychological Services,.

Luyster, R. J., Zane, E., & Wisman Weil, L. (2022). Conventions for unconventional language:

Revisiting a framework for spoken language features in autism. *Autism & Developmental Language Impairment*, 7, 239694152211054. <https://doi.org/10.1177/23969415221105472>

Malkin, L., Abbot-Smith, K., Williams, D., & Ayling, J. (2018). When do Children with Autism Spectrum

Disorder Take Common Ground into Account During Communication?: Malkin et al./ASD common ground. *Autism Research*, 11(10), 1366–1375. <https://doi.org/10.1002/aur.2007>

Matthews, D., Biney, H., & Abbot-Smith, K. (2018). Individual Differences

Ability: A Review of Associations with Formal Language, Social Cognition, and Executive Functions. *Language Learning and Development*, 14(3), 186–223.

<https://doi.org/10.1080/15475441.2018.1455584>

McConkey, R. (2020). The rise in the numbers of pupils identified by schools with autism spectrum

disorder (ASD): A comparison of the four countries in the United Kingdom. *Support for Learning*, 35(2), 132–143. <https://doi.org/10.1111/1467-9604.12296>

McGuinness, L., Abbot-Smith, K., & Gambi, C. (in press). Seeing it in others versus doing it yourself:

Social desirability judgements and conversation production data from autistic and non-autistic children. *Autism*

McKernan, E. P., Kumar, M., Di Martino, A., Shulman, L., Kolevzon, A., Lord, C., Narayanan, S., & Kim,

S. H. (2022). Intra-topic latency as an automated behavioral marker of treatment response in autism spectrum disorder. *Scientific Reports*, 12(1), 3255. <https://doi.org/10.1038/s41598-022-07299-w>

Mein, C., Fay, N., & Page, A. C. (2016). Deficits in joint action explain why socially anxious individuals

are less well liked. *Journal of Behavior Therapy and Experimental Psychology*, 50(1), 147–151.

<https://doi.org/10.1016/j.jbtep.2015.07.001>

Miczko, N., Segrin, C., & Allspach, L. E. (2001). Relationship between nonverbal sensitivity, encoding,

and relational satisfaction. *Communication Reports*, 14(1), 39–48.

<https://doi.org/10.1080/08934210109367735>

- Nadig, A., Lee, I., Singh, L., Bosshart, K., & Ozonoff, S. (2010). How does the topic of conversation affect verbal exchange and eye gaze? A comparison between typical development and high-functioning autism. *Neuropsychologia*, *48*(8), 2730–2739. APA PsycInfo.
<https://doi.org/10.1016/j.neuropsychologia.2010.05.020>
- Ochi, K., Ono, N., Owada, K., Kojima, M., Kuroda, M., Sagayama, S., & Yamasue, H. (2019). Quantification of speech and synchrony in the conversation of adults with autism spectrum disorder. *PLOS ONE*, *14*(12), e0225377. <https://doi.org/10.1371/journal.pone.0225377>
- Pagmar, D., Abbot-Smith, K., & Matthews, D. (2022). *contingency* [PDF]. <https://doi.org/10.34842/2022-511>
- Parish-Morris, J., Liberman, M., Ryant, N., Cieri, C., Bateman, L., Ferguson, E., & Schultz, R. (2016). Exploring Autism Spectrum Disorders Using HLT. *Proceedings of the Third Workshop on Computational Linguistics and Clinical Psychology*.
<https://doi.org/10.18653/v1/W16-0308>
- Paul, R., Orlovski, S. M., Marcinko, H. C., & Volkmar, F. (2009). Conversational Behaviors in Youth with High-functioning ASD and Asperger Syndrome. *Journal of Autism and Developmental Disorders*, *39*(1), 115–125. <https://doi.org/10.1007/s10803-008-0607-1>
- Perner, J., Leekam, S. R., & Wimmer, H. (1987). Three-year-olds' difficulty with false-beliefs for a conceptual deficit. *British Journal of Developmental Psychology*, *5*(2), 125–137.
<https://doi.org/10.1111/j.2044-835X.1987.tb01048.x>
- Pillow, B. H. (1989). Early understanding of perception as a source of knowledge. *Journal of Experimental Child Psychology*, *47*(1), 116–129. [https://doi.org/10.1016/0022-0965\(89\)90066-0](https://doi.org/10.1016/0022-0965(89)90066-0)
- Place, K. S., & Becker, J. A. (1991). The influence of context on the reading comprehension of school children. *Discourse Processes*, *14*(2), 227–241.
<https://doi.org/10.1080/01638539109544783>

- Pushkar, D., Basevitz, P., Arbuckle, T., Nohara-LeClair, M., Lapidus, S., & Peled, M. (2000). Social behavior and off-target verbosity in elderly people. *Psychology and Aging*, *15*(2), 361–374. <https://doi.org/10.1037/0882-7974.15.2.361>
- Roy, P., & Chiat, S. (2014). Developmental pathways of language and social communication problems in 9–11 year olds: Unpicking the heterogeneity. *Research in Developmental Disabilities*, *35*(10), 2534–2546. <https://doi.org/10.1016/j.ridd.2014.06.014>
- Ruscher, J. B., & Hurley, M. M. (2000). Off-Target Verbosity Evokes Negative Stereotypes of Older Adults. *Journal of Language and Social Psychology*, *19*(1), 141–149. <https://doi.org/10.1177/0261927X00019001007>
- Saul, J., Griffiths, S., & Norbury, C. F. (2022). Prevalence and functional impact of social (pragmatic) communication disorders. *Journal of Child Psychology and Psychiatry*, *63*(13), 1370–1375. <https://doi.org/10.1111/jcpp.13705>
- Semel, E., Wiig, E., & Secord, W. (2006). *Clinical evaluation of language fundamentals, fourth edition (CELF4)*. Pearson: Psychological Corporation.
- S l o m k o w s k i , C . , & D u n n , J . (1 9 9 6) . Y o u n g c h i l d r e n ' s feelings and their connected communication with friends. *Developmental Psychology*, *32*(3), 442–447. <https://doi.org/10.1037/0012-1649.32.3.442>
- Sullivan, K., Zaitchik, D., & Tager-Flusberg, H. (1994). Preschoolers can attribute second-order beliefs. *Developmental Psychology*, *30*(3), 395–402. <https://doi.org/10.1037/0012-1649.30.3.395>
- Tomasello, M. (2018). How children come to understand false beliefs: A shared intentionality account. *Proceedings of the National Academy of Sciences*, *115*(14), 8491–8498. <https://doi.org/10.1073/pnas.1804761115>
- Wechsler, D. (2003). *The Wechsler intelligence scale for children (The Psychological Corporation)*. Psychological Corporation.
- Wechsler, D. (2012). *Wechsler Preschool and Primary Scale of Intelligence, Fourth Edition*. Psychological Corporation.

Wellman, H. M., & Liu, D. (2004). Scaling of Theory-of-Mind Tasks. *Child Development*, 75(2), 523–541. <https://doi.org/10.1111/j.1467-8624.2004.00691.x>

Wheeless, L. R., Frymier, A. B., & Thompson, C. A. (1992). A comparison of verbal output and receptivity in relation to attraction and communication satisfaction in interpersonal relationships. *Communication Quarterly*, 4(2), 102–115. <https://doi.org/10.1080/01463379209369826>

Wiig, E., Semel, E., & Secord, W. (2013). *Clinical Evaluation of Language Fundamentals*. Pearson / PsychCorp.

Wimmer, H. (1983). Beliefs about beliefs: Representation and constraining function of wrong beliefs in young children's *Cognition*, 13(1), 103–128. [https://doi.org/10.1016/0010-0277\(83\)90004-5](https://doi.org/10.1016/0010-0277(83)90004-5)

Winner, M. G. (2007). *Thinking about you, thinking about them*. Social Publishing.

Ying Sng, C., Carter, M., & Stephenson, J. (2018). A systematic review of the comparative pragmatic differences in conversational skills of individuals with autism. *Autism & Developmental Language Impairment*, 3, 239694151880380. <https://doi.org/10.1177/2396941518803806>

SUPPLEMENTAL MATERIALS / APPENDICES

Appendix 1: Study 1 Pre-planned conversation probes

ANIMALS SET

- a) I really like animals (*This always followed playing a pet shop game*)
- b) I have got a pet
- c) My pet has a funny name
- d) My pet does something really silly

FRUIT SET (*This always followed playing a fruit sorting game ó see pic*)



- e) I like to eat fruit
- f) I think apples are really tasty
- g) Some people cook apples
- h) I can bake something nice with apples

HOLIDAYS SET (*This always followed the administration of the CELF Expressive Vocabulary ó the experimenter pointed to the picture of an island*)

- i) Oh look! A beach!
- j) I went to the beach in the summer
- k) It was a beach somewhere really special
- l) I went on such an amazing trip



GAMES SET (*This always followed a fishing game ó see*

- m) I really like that game
- n) But it's not my favourite game
- o) There's another game I really like playing
- p) The game has a surprise in it

pic)

Appendix 2: More detailed description of instructions given to RAs for Study 2

Each child was pre-assigned to one of the two RAs. The children were told that they would carry out some word naming tests, some story tests, some puzzles and a ‘fish’ game but that because some of these were a “bit boring”, they would have some chats in between – and also a ‘biscuit break’ whenever they liked.

During the first ‘chat’, the RA showed the child a picture or object which related to a topic which the child’s parent had named during the recruitment phase as being one of the child’s particular interests. Once conversation was underway, the picture was removed. During the second ‘chat’ break, the picture was of the RA and her brother (as both had brothers) and the elicited conversation revolved around one or more of the following generic topics: siblings, pets and families. The RAs were trained to avoid using an overly questioning style. To aid this, they prepared two or three declarative statements per conversation to use at a topic-appropriate moment. For example, for the ‘swimming’ topic, which was a frequently cited favourite hobby, the RA might prepare a statement about once getting out of her depth before she could swim properly and playing games in the pool. For the sibling topic, the RA usually mentioned her brother once annoyed her in a zoo by pretending to be a monkey and also (at another point in the conversation) mentioned that her brother liked to play tricks on her, such as putting salt in her tea or jumping out and saying ‘Boo!’.

For each of these conversations, the RA was told to aim for around five minutes but not to cut the child off if he or she was (for example) part-way through talking about a topic. The RAs were also told that if it was painfully awkward to continue a conversation, then they could cut it short after three minutes. (This occurred for a few children). Since we used logistic regression with random intercepts for participants, differences between children in conversation length were taken into account.

Example excerpt from conversation (from very proficient child):

RA Statement: Once I went to the hospital and they gave me free lollies. [LAUGHS]. They were tasty.

P11: Once I went to a doctor and then they gave me a sweet

RA Minimal: Ah! You too!

P11: Well I think it was because of my ears or because of my cough because sometimes I have a cough

RA Question: Okay. And you went for your ear because // because it was hurting?

P11: hmm because // because it – no because it – because of the problem with my ears which it doesn’t really hear very well

RA Statement: Okay well that’s what doctors are for. They make things better. They help out when // something isn’t working quite well//

P11: [Sneezes]

RA Minimal: [LAUGHS] Yeah. Right

P11: Sometimes we – we live inside the [ANONYMISED]

RA Minimal: Wow!

P11: And sometimes the police come with their cars – not their motorbikes, though.

RA Statement: Hmm – yeah, I saw a policeman on a motorbike once and he fell because he was turning and then he turned too fast and he fell over. Yeah.

P11: And did he hurt himself?

RA Statement: Yeah a little bit but I think he got back on the bike quite quickly. I think if you if you get back on it you’re not as scared after.

P11: Well probably not because he wants to finish off his job, doesn’t he?

RA Statement: Exactly! He has some people to save so he couldn’t -

P11: - well probably not saved but arrest. [LAUGHS]

Appendix 3: Study 2 (Neurotypical 6-year-olds) Correlation matrix - predictor variable relationships with one another ($N = 48$ except where stated)

	Age in months	Expressive Vocabulary (raw)	NV reasoning (raw)	WM Backwards Digit Span (raw)	Flanker RT Diff Score (IC) (raw)	Knowledge & Belief ToM (raw)
Vocabulary (raw) (N = 45)	.34*					
NVIQ (raw) (N = 42)	.24	-.08				
WM Digit (raw)	.05	.27	.09			
Flanker RT Diff Score (IC) (N = 45)	.11	-.13	.07	-.08		
Knowledge & Belief ToM	-.04	.34*	.18	.27 [∇] ($p = .06$)	-.35*	
Strange Stories ToM	.25	.17	.05	.09	-.01	.06

[∇] = $p < .08$

* = $p < .05$

** = $p < .01$

*** = $p < .001$

Appendix 4: Study 2 (Neurotypical 6-year-olds) Correlation matrix – proportion of direct responses to all experimenter declarative utterances

	Contingent direct responses	Non-contingent direct responses	All topic-supporting direct responses
Age in months (N = 48)	.28 [∇] (<i>p</i> = .058)	-.12 (<i>p</i> = .4)	.16 (<i>p</i> = .27)
CELF-4 Expressive Vocabulary raw score (N = 45)	-.031 (<i>p</i> = .84)	.01 (<i>p</i> = .97)	-.11 (<i>p</i> = .46)
BAS Matrices (NVIQ) raw score (N = 42)	.06 (<i>p</i> = .7)	-.33* (<i>p</i> = .04)	-.01 (<i>p</i> = .3)
WISC Backwards Digit Span raw score (N = 48)	-.18 (<i>p</i> = .23)	-.15 (<i>p</i> = .3)	-.14 (<i>p</i> = .33)
Flanker Incongruent – Congruent RT Difference Score (N = 45)	.05 (<i>p</i> = .7)	.006 (<i>p</i> = .97)	.04 (<i>p</i> = .79)
Knowledge-Belief (‘Puppy’ + ‘Robots’) ToM raw score (N = 48)	.06 (<i>p</i> = .70)	-.29* (<i>p</i> = .048)	-.007 (<i>p</i> = .96)
Strange Stories (‘Kittens’ + ‘Biscuits’) ToM raw score (N = 48)	.21 (<i>p</i> = .15)	-.06 (<i>p</i> = .69)	.21 (<i>p</i> = .15)
Total ToM raw score (N = 48)	.17 (<i>p</i> = .24)	-.26 [∇] (<i>p</i> = .077)	.12 (<i>p</i> = .41)

[∇] = *p* < .08

* = *p* < .05

** = *p* < .01

*** = *p* < .001

Appendix 5: Study 2 (Neurotypical 6-year-olds) Correlation matrix – proportion of child elaborations (i.e. child turns which followed on from the child's own turn)

Elaborations (NB: One child produced no elaborations at all and is excluded from the following analyses)

	Contingent elaborations	Self-contingent elaborations
Age in months (N = 47)	.13 (<i>p</i> = .39)	.04 (<i>p</i> = .8)
CELF-4 Expressive Vocabulary raw score (N = 44)	.21 (<i>p</i> = .16)	-.23 (<i>p</i> = .14)
BAS Matrices (NVIQ) raw score (N = 42)	.22 (<i>p</i> = .16)	.11 (<i>p</i> = .47)
WISC Backwards Digit Span raw score (N = 47)	.33* (<i>p</i> = .024)	-.36** (<i>p</i> = .01)
Flanker Incongruent – Congruent RT Difference Score (N = 45)	.09 (<i>p</i> = .58)	-.15 (<i>p</i> = .34)
Knowledge-Belief ('Puppy' + 'Robots') ToM raw score (N = 47)	.30* (<i>p</i> = .04)	-.31* (<i>p</i> = .03)
Strange Stories ('Kittens' + 'Biscuits') ToM raw score (N = 47)	.06 (<i>p</i> = .67)	.01 (<i>p</i> = .96)
Total ToM raw score (N = 47)	.27 [∇] (<i>p</i> = .06)	-.24 (<i>p</i> = .11)

[∇] = *p* < .08

* = *p* < .05

** = *p* < .01

*** = *p* < .001

Appendix 6: Study 3 Means (SDs) for age, cognitive and socio-cognitive raw scores

	Mean	SE	Minimum	Maximum
Age in months	87.12	13.62	64	118
WPPSI Matrices (raw)	15.10	5.54	4	24
CELF-4 Expressive Vocabulary (raw)	26.80	11.34	6	52
WISC Backwards Digit Span (raw)	5.05	1.75	0	9
Backwards Word Span (raw)	4.78	2.22	0	11
Theory of Mind Composite (raw)	7.92	3.87	0	13

Appendix 7: Study 3 (Autistic 5 to 9-year-olds) Correlation matrix – predictor variable

relationships with one another

	Age in months	WPPSI Matrices	Express Vocab	Backwards Digit Span	Backwards Words
WPPSI Matrices (raw)	.15	-			
CELF-4 Expressive Vocabulary (raw)	.41**	.45***	-		
WISC Backwards Digit Span (raw)	.12	.33*	.43**	-	
Backwards Word Span (raw)	.29 [∇]	.29 [∇]	.56***	.56***	-
Theory of Mind Composite (raw)	.13	.34*	.69***	.36*	.57***

[∇] = p < .08

* = p < .05

** = p < .01

*** = p < .001

Appendix 8: Study 3 (Autistic 5 to 9-year-olds) Correlation matrix – proportion of direct responses to all clinician declarative utterances

	Contingent responses	Non-contingent responses	All Topic-Supporting Responses
Age in months	-.06 (<i>p</i> = .73)	-.3 [∇] (<i>p</i> = .062)	-.01 (<i>p</i> = .94)
CELF 4 Expressive Vocabulary raw score	.05 (<i>p</i> = .76)	-.16 (<i>p</i> = .33)	.23 (<i>p</i> = .15)
WPPSI Matrices (NVIQ) raw score	-.03 (<i>p</i> = .84)	.08 (<i>p</i> = .63)	.16 (<i>p</i> = .32)
WISC Backwards Digit Span raw score	.19 (<i>p</i> = .23)	-.1 (<i>p</i> = .56)	.12 (<i>p</i> = .46)
Backwards Words Span raw score	.33* (<i>p</i> = .04)	-.04 (<i>p</i> = .83)	.31 [∇] (<i>p</i> = .056)
Working Memory composite	.29 [∇] (<i>p</i> = .068)	-.08 (<i>p</i> = .65)	.23 (<i>p</i> = .15)
Theory of Mind (stepwise) composite	.24 (<i>p</i> = .15)	-.20 (<i>p</i> = .21)	.41** (<i>p</i> = .008)

[∇] = *p* < .08


* = *p* < .05

** = *p* < .01

*** = *p* < .001

Appendix 9: Relationship between the ADOS A8 ‘Conversation’ score and our conversation assessment measure

There was a significant negative correlation between the ADOD ‘A8’ (Conversation) score and our direct measure of ‘all topic-supporting’ responding ($r(36) = -.51, p = 0.001$) – see figure below. The same relationship was found for contingent responding but the effect size was less ($r = .34, p < .05$).



Appendix 10: Study 3 (Autistic 5 to 9-year-olds) Correlation matrix – proportion of child elaborations (i.e. child turns which followed on from the child's own turn)

	Contingent elaborations	Self-contingent elaborations
Age in months	.17 (<i>p</i> = .3)	-.01 (<i>p</i> = .96)
CELF 4 Expressive Vocabulary raw score	.22 (<i>p</i> = .18)	.08 (<i>p</i> = .64)
WPPSI Matrices (NVIQ) raw score	.11 (<i>p</i> = .51)	.03 (<i>p</i> = .87)
WISC Backwards Digit Span raw score	.32* (<i>p</i> = .045)	.01 (<i>p</i> = .93)
Backwards Words Span raw score	.26 (<i>p</i> = .11)	.13 (<i>p</i> = .44)
Working Memory composite	.33* (<i>p</i> = .04)	.08 (<i>p</i> = .64)
Theory of Mind (stepwise) composite	.19 (<i>p</i> = .23)	-.02 (<i>p</i> = .89)

∇ = *p* < .08

* = *p* < .05

** = *p* < .01

*** = *p* < .001

Appendix 11: Study 3 (Autistic 5 to 9-year-olds) Logistic regression - direct responses to all clinical conversational turns (including questions)

Contingent responses

Random effects:

Groups	Name	Variance	Std.Dev.
Participant	(Intercept)	0.4498	0.6707

Number of obs: 2612, groups: Participant, 40

Fixed effects:

	B	SE	Chi-square	p-value
(Intercept)	-0.50797	0.11624		
Age in months	0.04918	0.13374	0.1355	0.7128
CELF	-0.19453	0.20099	0.9254	0.3361
NVIQ	-0.11931	0.13226	0.8064	0.3692
Working Memory Composite	0.34111	0.16471	4.0815	0.04335 *
ToM_scale_composite	0.12833	0.17062	0.5617	0.4536

Non-contingent responses

Random effects:

Groups	Name	Variance	Std.Dev.
Participant	(Intercept)	1.199	1.095

Number of obs: 2612, groups: Participant, 40

Fixed effects:

	B	SE	z value	Pr(> z)	Chi	P-val	comp
(Intercept)	-2.58	0.20	-12.60	< 2e-16	***		
Age in months	-0.64	0.23	-2.77	0.006	**	7.35	0.007 **
CELF	0.17	0.34	0.49	0.62		0.2367	1 0.626
NVIQ	0.23	0.22	1.02	0.31		1.0454	1 0.3066
Working Memory Composite	0.01	0.27	0.04	0.97		0.0016	1 0.9679
ToM_scale_composite	-0.30	0.29	-1.02	0.31		1.0023	1 0.3168

All topic-supporting responses

Random effects:

Groups	Name	Variance	Std.Dev.
Participant	(Intercept)	0.1792	0.4233

Number of obs: 2612, groups: Participant, 40

Fixed effects:

	B	SE	z value	Pr(> z)	Chi	P-val	comp
(Intercept)	0.59	0.08	7.31	2.66e-13	***		
Age in months	0.04	0.09	0.41	0.69		0.1639	0.69
CELF	-0.07	0.14	-0.52	0.60		0.2689	0.60
NVIQ	0.06	0.09	0.62	0.54		0.382	0.54
Working Memory Composite	0.07	0.11	0.61	0.54		0.3747	0.54
ToM_scale_composite	0.20	0.12	1.67	0.095	.	2.7193	0.099 ^v .