



Cui, Jingjing, He, Yiyi, Souza, Cauê P., Weiser, Jonas, Dietz, Maximilian, Krummenacher, Ivo, Dewhurst, Rian D., Braunschweig, Holger and Fantuzzi, Felipe (2024) *Single electron oxidation, chloride abstraction, and hydride-induced decomposition of a dichloro-bis(germylene)*. European Journal of Inorganic Chemistry . ISSN 1434-1948.

Downloaded from

<https://kar.kent.ac.uk/107276/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1002/ejic.202400422>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

EurJIC

European Journal of Inorganic Chemistry

 **Chemistry
Europe**

European Chemical
Societies Publishing

Accepted Article

Title: Single-Electron Oxidation, Chloride Abstraction, and Hydride-Induced Decomposition of a Dichloro-Bis(Germylene)

Authors: Jingjing Cui, Yiyi He, Cauê P. Souza, Jonas Weiser, Maximilian Dietz, Ivo Krummenacher, Rian D. Dewhurst, Holger Braunschweig, and Felipe Fantuzzi

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Eur. J. Inorg. Chem.* **2024**, e202400422

Link to VoR: <https://doi.org/10.1002/ejic.202400422>

WILEY-VCH

RESEARCH ARTICLE

Single-Electron Oxidation, Chloride Abstraction, and Hydride-Induced Decomposition of a Dichloro-Bis(Germylene)

Yiyi He,^[a] Cauê P. Souza,^[b] Jonas Weiser,^[c] Maximilian Dietz,^[d,e] Ivo Krummenacher,^[d,e] Rian D. Dewhurst,^[d,e] Holger Braunschweig,^{*[d,e]} Felipe Fantuzzi,^{*[b]} and Jingjing Cui^{*[a]}

Abstract: In this work we explore the reactivity of a dichloro-bis(germylene) compound that features a naphthyridine diimine (NDI) ligand, denoted NDI-Ge₂Cl₂. Upon reaction with the oxidant [Cp₂Fe][BAR^F₄] (Cp = C₅H₅, BAR^F₄ = [{3,5-(CF₃)₂C₆H₃}_4B]), we observed the formation of the radical species [NDI-Ge₂Cl₂]^{•+}[BAR^F₄]⁻. Furthermore, the introduction of Na[BAR^F₄] induced a single chloride anion abstraction process, resulting in the formation of [NDI-Ge₂Cl][BAR^F₄]. The addition of Li[HB(^tBu)₃] to NDI-Ge₂Cl₂ led to the generation of NDI-Ge, presumably via a Cl/H exchange pathway followed by a decomposition process.

Introduction

Germynes, heavier analogs of carbenes, exhibit diverse reactivities, including oxidation, reduction, coordination to Lewis acids or bases, and activation of small molecules.^[1] In the case of germynes featuring a Ge–X (X = halogen) bond, further derivations, such as nucleophilic substitution, halogen extraction, and halogen/H exchange, become conceivable.^[1a,1f,2] However, the exploration of dihalo-digermynes (Figure 1, I–III) and their reactivity is rather limited, with a predominant focus on their reduction chemistry.^[3] The contribution from the Rivard group^[4] represents a rare example of a detailed investigation into the derivation of dichloro-digermylene, providing invaluable insights into this specific class of compounds. This situation is in sharp contrast to the extensive reactivity studies on halomonogermynes.

Recently, utilizing a redox-active naphthyridine–diimine ligand (NDI) (Figure 2, IV), we successfully synthesized dichloro-digermylene **1** (NDI-Ge₂Cl₂) and subsequently obtained a six-electron donating digermylene NDI-Ge₂ (**V**) via the reduction of **1**. Additionally, we achieved the salt elimination of **1** by employing a transition-metal-based nucleophile.^[5] In this work, we demonstrate single-electron oxidation, chloride anion abstraction, halogen/H exchange, and subsequent disproportionation of compound **1**.




Figure 1. Structurally characterized dihalo-digermynes (Ar = 2,6-diisopropylphenyl, Mes = 2,4,6-trimethylphenyl).





Figure 2. Structures of IV, V and **1** (Dipp = 2,6-diisopropylphenyl).

Results and Discussion

Upon the addition of equimolar amounts of oxidant [Cp₂Fe][BAR^F₄] (Cp = C₅H₅, BAR^F₄ = [{3,5-(CF₃)₂C₆H₃}_4B]) to a C₆D₆ solution of **1** at ambient temperature, the appearance of the reaction mixture remained unchanged. However, the ¹H NMR spectrum exclusively exhibited a signal for Cp₂Fe, suggesting the formation of radical species **2** (Scheme 1).



Scheme 1. Synthesis of compound **2** (Dipp = 2,6-diisopropylphenyl).

After removing Cp₂Fe by hexane washing in an ultrasound bath, single crystals suitable for X-ray diffraction analysis were obtained. The structure of the cationic part of **2** (Figure 3) strongly resembles that of **1**.^[6] The mean planes of the two C₂N₄Ge₂ moieties display a slightly smaller dihedral angle of 11.70°

- [a] Y. He, Dr. J. Cui
School of Chemistry and Environmental Engineering
Wuhan Institute of Technology
Wuhan 430205 (P.R. China)
E-mail: 17061701@wit.edu.cn
- [b] C. P. Souza, Dr. F. Fantuzzi
School of Chemistry and Forensic Science
University of Kent
Park Wood Rd, Canterbury CT2 7NH (UK)
E-mail: f.fantuzzi@kent.ac.uk
- [c] J. Weiser
Institute for Physical and Theoretical Chemistry
Julius-Maximilians-Universität Würzburg
Emil-Fischer-Str. 42, 97074 Würzburg (Germany)
- [d] Dr. M. Dietz, Dr. I. Krummenacher, Dr. R. D. Dewhurst, Prof. Dr. H. Braunschweig
Institute for Inorganic Chemistry
Julius-Maximilians-Universität Würzburg
Am Hubland, 97074 Würzburg (Germany)
E-mail: h.braunschweig@uni-wuerzburg.de
- [e] Dr. M. Dietz, Dr. I. Krummenacher, Dr. R. D. Dewhurst, Prof. Dr. H. Braunschweig
Institute for Sustainable Chemistry & Catalysis with Boron
Julius-Maximilians-Universität Würzburg
Am Hubland, 97074 Würzburg (Germany)

Supporting information for this article is given via a link at the end of the document.

RESEARCH ARTICLE

compared to that observed in **1** (15.86°),^[5] indicating a distorted arrangement of the naphthyridine-diimine-Ge₂ unit. The distance between the two Ge atoms increased from 2.8944(9) Å in **1** to 3.0021(15) Å in **2**. The Ge–Cl bonds [2.2218(18) and 2.2424(15) Å], the Ge–N bonds [2.015(3)–2.049(4) Å] and the sum of the bond angles (268.37° – 270.34°) around the Ge atoms are all comparable to those of **1** [Ge–Cl bond 2.2916(7) Å, Ge–N bonds 1.9663(15) and 2.0011(16) Å, and the bond angle sum of Ge $273.11(15)^\circ$]. Adding two equivalents of [Fc][BAR^F₄] to **1** led to the same result, even with heating to 90 °C.




Figure 3. Solid-state structure of the cation of **2** (hydrogen atoms and ellipsoids of the Dipp groups are omitted for clarity). Thermal ellipsoids are set at the 50% probability level. Selected bond lengths [Å] and angles [°]: Ge1...Ge2, 3.001(2); Ge1–Cl1, 2.2424(15); Ge1–N1, 2.049(4); Ge1–N13, 2.015(3); Ge2–Cl2, 2.2218(18); Ge2–N8, 2.044(4); Ge2–N16, 2.020(3); C2–C11, 1.429(6); C7–C14, 1.442(6); Cl1–Ge1–N1, 95.91(11); Cl1–Ge1–N13, 92.55(11); N1–Ge1–N13, 79.92(13); Cl2–Ge2–N8, 94.78(11); Cl2–Ge2–N16, 95.41(11); N8–Ge2–N16, 80.17(13).

The electron paramagnetic resonance (EPR) spectrum of a toluene solution of compound **2** at room temperature exhibited a single broad peak ($g = 2.0029$) with a peak-to-peak linewidth of 1.4 mT (Figure 4). Simulating the spectrum is unfeasible due to complicated overlapping couplings that are difficult to resolve. For a better understanding of the radical cation [NDI-Ge₂Cl₂]^{•+}, we performed DFT calculations at the M06-D3/def2-SVP level of theory.^[7] We selected this level based on our preliminary benchmark investigation, as it yielded the most accurate structural agreement with the available experimental X-ray data.^[5,8] The canonical Kohn–Sham singly-occupied molecular orbital (SOMO, Figure 5A) indicates a delocalization of the radical electron via the NDI ligand instead of the Ge atoms, in agreement with the experimental findings. Additional analysis of the spin density (Figure 5B) corroborated this conclusion. The positive charge is evenly distributed and is localized on the Ge atoms, each possessing a Mulliken charge of 0.64.

The combined information from the EPR study and the DFT calculations suggested that the radical predominantly localizes on the ligand rather than on the Ge atoms, as observed previously for [(dpp-BIAN)GeCl]^{•+}[GeCl₃][−] (dpp-BIAN = 1,2-Bis{(2,6-diisopropyl-phenyl)imino}acenaphthene) (Figure 6, **VI**),^[9] 2,6-diiminophenyl-supported germylidendiide dianion radical **VII**,^[10] β-diketiminato germanium radical complex •LGe (•L = •[PhC(PhCN-Dipp)₂][−]) (**VIII**),^[11] the dipyrromethene-based neutral radical **IX**,^[12]

and the open-shell germylene stabilized by a phenalenyl-based bidentate ligand **X**.^[13]




Figure 4. Experimental continuous-wave (CW) X-band EPR spectrum of **2** in toluene at room temperature. Experimental parameters: microwave frequency = 9.86 GHz; microwave power = 2 mW; modulation amplitude = 0.2 G; conversion time = 40 ms; modulation frequency = 100 kHz.




Figure 5. (A) Kohn–Sham SOMO (isovalue: 0.03) and (B) total spin density (isovalue: 0.015) of compound **2**. Hydrogen atoms are omitted for clarity. Level of theory: M06-D3/def2-SVP.





Figure 6. Structurally characterized delocalized germanium radicals (Ar = 2,6-diisopropylphenyl, Mes = 2,4,6-trimethylphenyl, Trip = 2,4,6-triisopropylphenyl, Dipp = 2,6-diisopropylphenyl),

RESEARCH ARTICLE

The reaction between one equivalent of $\text{Na}[\text{BAR}^{\text{F}}_4]$ and **1** in benzene proceeded smoothly at ambient temperature, resulting in a red-violet solution. Despite several attempts to optimize the recrystallization conditions or to employ different counteranions, all efforts to obtain crystallographic-grade single crystals proved unsuccessful. However, high-resolution mass spectrometry (HRMS) at $m/z = 713.1662$ confirmed the successful removal of one chlorine anion (Scheme 2). The attempt to remove the second chlorine atom resulted in decomposition at 130°C .

In the ^1H NMR spectrum of **3**, we observed signals corresponding to the naphthyridine protons at 6.81 and 6.28 ppm, exhibiting a downfield shift compared to those of **1** (6.16 and 5.38 ppm),^[5] in line with the electron-deficient nature of the cation. Notably, the difference between the two signals in **3** decreased to 0.53 ppm, compared to 0.78 ppm in **1**. Additionally, a single peak was observed at 2.33 ppm with an integration of four, corresponding to the CHMe_2 group (Figure S1). This prompted us to explore the possibility of a structure featuring a bridging chloride, as such a motif has been reported in the literature, specifically in the bis-germylene $[(\text{MeCNDipp})_2\text{C}=\text{CHGe}]_2(\mu\text{-Cl})$.^[4]



Scheme 2. Synthesis of compound **3** (Dipp = 2,6-diisopropylphenyl).

In the absence of an experimental crystal structure, we used DFT calculations to investigate the potential structures of compound **3** (Figure 7) and determine their NMR data, the latter computed at the $\omega\text{B97X-D3}/\text{pcSseg-2}$ level of theory.^[14] Our theoretical investigation revealed a structure (**3'**, Figure 7B) featuring a bridging chloride positioned in a bent arrangement relative to the naphthyridine ligand. However, this structure exhibited an imaginary frequency of $98i\text{ cm}^{-1}$, indicating that it is not a viable minimum energy structure but rather a transition state (TS).

Additionally, a second structure (**3''**, Figure 7B) featuring a bridging chloride in a quasi-planar orientation was obtained through constrained optimization. This geometry was significantly less energetically favorable, being located at an energy level $61.4\text{ kcal mol}^{-1}$ higher than that of **3'**. Further examination with various functionals consistently indicated the absence of any minimum energy structure with a bridging chloride.

Following the direction of the imaginary mode of **3'**, we identified a plausible structure for **3** (Figure 7A), wherein the chloride is bound to only one of the germanium centers. The computational analysis of the chemical shifts for this structure revealed that the average deviation of the chemical shifts for the four CHMe_2 signals was merely 0.34 ppm, which could explain the appearance of a single peak for these protons in the experimental spectrum.




Figure 7. Optimized structures of **3**, **3'**, and **3''**. Energies (ΔE) are relative to that of **3**. Structure **3'** has an imaginary frequency and is a transition state connecting two enantiomers of **3**. Structure **3''** is located $61.4\text{ kcal mol}^{-1}$ above **3'**. Hydrogen atoms are omitted for clarity. Level of theory: M06-D3/def2-SVP.

To gain deeper insight into the electronic structure of **3**, we conducted a computational investigation of its electronic absorption features. For this purpose, time-dependent DFT (TD-DFT) calculations were performed at the $\omega\text{B97X-D3(BJ)}/\text{def2-SVPD}$ level of theory,^[15] with the results shown in Figure 8. Our calculations reveal that, unlike systems **1** and **2**, where the $\text{S}_0 \rightarrow \text{S}_1$ transitions occur in the near-infrared region (815 nm and 893 nm, respectively, see Figures S11–S12 in the SI), the corresponding transition in **3** is observed at 578 nm (Figure 8A).




Figure 8. (A) TD-DFT vertical excitations of **3** at the $\omega\text{B97X-D3(BJ)}/\text{def2-SVPD}$ level of theory. The inset figure shows the CDD plot of the $\text{S}_0 \rightarrow \text{S}_1$ excitation, where the charge flows from red to blue (isovalue: 0.0015). Hydrogen atoms are omitted for clarity. (B) NTOs of the $\text{S}_0 \rightarrow \text{S}_1$ excitation of **3**.

RESEARCH ARTICLE


Analysis of the natural transition orbitals (NTOs,^[16] Figure 8B) of **3** indicates that this transition is characterized as a $\pi \rightarrow \pi^*$ excitation. The hole NTO spans the entire π framework of the molecule, but is predominantly concentrated in the GeC_2N_2 ring where Ge is not bound to Cl. In contrast, the electron NTO shows a greater degree of localization in the $\text{Ge}(\text{Cl})\text{C}_2\text{N}_2$ ring, including a significant contribution from the Cl atom. These findings suggest that the transition has an appreciable charge transfer (CT) character, which is blueshifted relative to compounds **1** and **2** due to the asymmetry of the GeC_2N_2 rings.

The CT character is further supported by the charge density difference (CDD) plot (Figure 8A) and the interfragment charge transfer (IFCT) method, as implemented in Multiwfn 3.8.^[17] The CDD plot clearly shows electron flow from the GeC_2N_2 ring to the $\text{Ge}(\text{Cl})\text{C}_2\text{N}_2$ ring. The IFCT analysis quantifies this charge transfer, showing 72.5% CT character, with the GeC_2N_2 ring losing 38.8% of its electron density during excitation, while the $\text{Ge}(\text{Cl})\text{C}_2\text{N}_2$ ring gains 32.6% and the remainder of the molecule gains 6.2%.

These combined results not only explain the red-violet color of compound **3**, consistent with its absorption maximum at 578 nm, but also support the placement of the Cl atom as a terminal ligand rather than a bridging one.

Our calculations confirmed that **3** exhibits only one Ge–Cl single bond, with a characteristic bond length of 2.245 Å. A distance of 3.393 Å between the chloride and the uncoordinated Ge atom ruled out the possibility of chloride bridging. Furthermore, bonding analyses based on the intrinsic bond orbital (IBO)^[18] and natural bond orbital (NBO)^[19] methods provided no evidence of chloride bonding to the uncoordinated Ge atom. Second-order perturbation energies from NBO calculations indicated the presence of a weak hyperconjugative effect of 6.68 kcal mol⁻¹ between the chloride and the uncoordinated Ge atom. However, given the relatively flat potential energy surface concerning the position of the chloride, as evidenced by **3'** being merely 5.1 kcal mol⁻¹ above **3**, we cannot rule out the possibility of a fluxional structure. Further investigation is warranted to explore this aspect.

For the reaction between **1** and two equivalents of $\text{Li}[\text{HB}(\text{sBu})_3]$ in C_6D_6 , the in situ ¹H NMR spectrum indicates the generation of **V** and an unsymmetrical compound species which features four doublets at 5.11, 5.42, 5.94, and 6.13 ppm (Figure S5). Moreover, the singlet at 8.01 ppm is comparable to that of $[\{\text{HC}-(\text{CMeAr})_2\}\text{GeH}]$ (Ar = 2,6-*i*-Pr₂C₆H₃) (8.08 ppm),^[20] implying that hydrido-germylene **4** is a possible intermediate. A prolonged reaction time or removal of the solvent led to the generation of **5** as the major product (Scheme 3).



Scheme 3. Synthesis of compound **5** (Dipp = 2,6-diisopropylphenyl).




Figure 9. Solid-state structure of **5** (hydrogen atoms and ellipsoids of the Dipp groups are omitted for clarity). Thermal ellipsoids are set at the 50% probability level. Selected bond lengths [Å] and angles [°]: Ge1–N1, 1.8925(17); Ge1–N13, 1.8777(17); N1–C2, 1.409(3); C2–C11, 1.371(3); C11–N13, 1.387(3); C3–C4, 1.347(3); N8–C7, 1.346(3); C7–C14, 1.500(3); C14–N16, 1.277(3); N1–Ge1–N13, 81.91(7); N1–C2–C11, 113.42(18); C2–C11–N13, 112.93(18); C2–N1–C9, 120.09(17); C2–C3–C4, 121.14(19); N8–C7–C14, 117.00(19); C7–C14–N16 115.6(2); C7–N8–C9, 117.58(18); C5–C6–C7, 118.6(2).

X-ray diffraction analysis (Figure 9) revealed that the three annulated rings within compound **5** adopt a coplanar arrangement. The largest deviation from the mean plane defined by these rings is only 0.062 Å, attributed to the Ge atom. The coordination-free half of **5**, denoted **5**(free), resembles the free ligand **IV**. In contrast, the germanium-containing half of **5**, denoted as **5**(Ge), is similar to compound **V**. The bond distances of compounds **1**, **2**, **5**, **IV**, and **V** suggest that increasing the degree of ligand reduction leads to a shortening of the $\text{C}_{\text{imine}}-\text{C}_{\text{Np}}$ bonds, along with a lengthening of the $\text{C}_{\text{Np}}-\text{N}_{\text{Np}}$ and $\text{N}_{\text{imine}}-\text{C}_{\text{imine}}$ bonds (Figure 10). This trend correlates with observations from Uyeda's study involving dinuclear nickel complexes of NDI.^[21]




Figure 10. Average bond distances (Å) of the $\text{N}_{\text{imine}}-\text{C}_{\text{imine}}-\text{C}_{\text{Np}}-\text{N}_{\text{Np}}$ moieties of compounds **IV**, **5**, **2**, **1**, and **V**. The coordination-free and germanium-containing parts of **5** are denoted as **5**(free) and **5**(Ge), respectively. The estimated standard deviations are ≤ 0.006 Å for all structures. Data are derived from Table S2.

Conclusion

In conclusion, we have investigated the reactivity of **1**, a dichloro-bis(germylene) compound incorporating a naphthyridine-diimine (NDI) ligand. The reaction with oxidant $[\text{Cp}_2\text{Fe}][\text{BAR}^{\text{F}}_4]$ led to the formation of radical species **2**. Moreover, addition of $\text{Na}[\text{BAR}^{\text{F}}_4]$ to **1** resulted in the abstraction of a single chloride anion,

RESEARCH ARTICLE

forming compound **3**. When $\text{Li}[\text{HB}(\text{sBu})_3]$ was used as a hydride source, in-situ ^1H NMR spectroscopy provided evidence for the generation of a mixed hydrido/chlorido bis(germylene), which decomposed to the thermally stable monogermanium species **5**.

Supporting Information

The authors have cited additional references within the Supporting Information.^[22–38]

Acknowledgements


Financial support from the Julius-Maximilians-University of Würzburg is gratefully acknowledged. Y.H. and J.C. thank the Postgraduate Innovation Foundation of the Wuhan Institute of Technology (No. CX2024340), the Open/Innovation Project of the Engineering Research Center of Phosphorus Resources Development and Utilization of the Ministry of Education (LCX202202), and the Hubei Three Gorges Laboratory (SC240003) for financial support. This work was supported by the Engineering and Physical Sciences Research Council (grant number EP/W52461X/1). This article is also based upon work from the COST Action CA20129 MultiChem, supported by COST (European Cooperation in Science and Technology).

Keywords: Single-electron oxidation • Chloride abstraction • Bis-germylene • Radical cation

- 1) a) M. Asay, C. Jones, M. Driess, *Chem. Rev.* **2011**, *111*, 354–396; b) M. Walewska, J. Baumgartner, C. Marschner, L. Albers, T. Müller, *Chem. Eur. J.* **2016**, *22*, 18512–18521; c) M. Walewska, J. Hlina, J. Baumgartner, T. Müller, C. Marschner, *Organometallics* **2016**, *35*, 2728–2737; d) H. Wang, Z. Xie, *Eur. J. Inorg. Chem.* **2017**, *2017*, 4430–4435; e) N. Del Rio, M. Lopez-Reyes, A. Baceiredo, N. Saffon-Merceron, D. Lutters, T. Müller, T. Kato, *Angew. Chem. Int. Ed.* **2017**, *56*, 1365–1370; f) R. Dasgupta, S. Khan, in *Advances in Organometallic Chemistry*, Vol. 74 (Ed.: P. J. Pérez), Academic Press, **2020**, pp. 105–152; g) J. A. Cabeza, P. García-Álvarez, C. J. Laglera-Gándara, *Eur. J. Inorg. Chem.* **2020**, *2020*, 784–795; h) R. J. Somerville, J. Campos, *Eur. J. Inorg. Chem.* **2021**, *2021*, 3488–3498; i) M. Ghosh, N. Sen, S. Khan, *ACS Omega* **2022**, *7*, 6449–6454; j) V. Y. Lee, *Eur. J. Inorg. Chem.* **2022**, *2022*, e202200175; k) Y. Qin, Y. Kang, L. Zhang, J. Sun, Z. Zhang, J. Xu, F. Zeng, A. Li, W. Wang, W. Shi, *Chin. J. Chem.* **2023**, 108691–108694; l) K. V. Arsenyeva, A. V. Piskunov, *J. Struct. Chem.* **2023**, *64*, 1–45.
- [2] a) Y. N. Lebedev, U. Das, G. Schnakenburg, A. C. Filippou, *Organometallics* **2017**, *36*, 1530–1540; b) A. Schulz, T. L. Kalkuhl, P. M. Keil, T. J. Hadlington, *Angew. Chem. Int. Ed.*, e202305996; c) A. F. Richards, A. D. Phillips, M. M. Olmstead, P. P. Power, *J. Am. Chem. Soc.* **2003**, *125*, 3204–3205.
- [3] a) Y. Li, K. C. Mondal, P. Stollberg, H. Zhu, H. W. Roesky, R. Herbst-Irmer, D. Stalke, H. Fiegl, *Chem. Commun.* **2014**, *50*, 3356–3358; b) Y. Su, Y. Li, R. Ganguly, R. Kinjo, *Eur. J. Inorg. Chem.* **2018**, *2018*, 2228–2231; c) X. Gao, Y. He, J. Cui, *J. Organomet. Chem.* **2024**, *1012*, 123146–123162.
- [4] E. Hupf, F. Kaiser, P. A. Lummis, M. M. D. Roy, R. McDonald, M. J. 2228Ferguson, F. E. Kühn, E. Rivard, *Inorg. Chem.* **2020**, *59*, 1592–1601.
- [5] J. Cui, J. Weiser, F. Fantuzzi, M. Dietz, Y. Yatsenko, A. Hafner, S. Nees, I. Krummenacher, M. Zhang, K. Hammond, P. Roth, W. Lu, R. D. Dewhurst, B. Engels, H. Braunschweig, *Chem. Commun.* **2022**, *58*, 13357–13360.
- [6] Deposition Number(s) <url href="https://www.ccdc.cam.ac.uk/services/structures?id=doi:10.1002/ejic.202400422">2278647 (for **2**) and 2312125 (for **5**)</url> contain(s) the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe <url href="http://www.ccdc.cam.ac.uk/structures">Access Structures service</url>.
- [7] a) Y. Zhao, D. G. Truhlar, *Theor. Chem. Acc.* **2008**, *120*, 215–241; b) S. Grimme, J. Antony, S. Ehrlich, H. Krieg, *J. Chem. Phys.* **2010**, *132*, 154104; c) F. Weigend, R. Ahlrichs, *Phys. Chem. Chem. Phys.* **2005**, *7*, 3297–3305.
- [8] J. Weiser, J. Cui, R. D. Dewhurst, H. Braunschweig, B. Engels, F. Fantuzzi, *J. Comput. Chem.* **2023**, *44*, 456–467.
- [9] I. L. Fedushkin, N. M. Khvoynova, A. Y. Baurin, G. K. Fukin, V. K. Cherkasov, M. P. Bubnov, *Inorg. Chem.* **2004**, *43*, 7807–7815.
- [10] S.-P. Chia, E. Carter, H.-W. Xi, Y. Li, C.-W. So, *Angew. Chem. Int. Ed.* **2014**, *53*, 8455–8458.
- [11] X. Lu, H. Cheng, Y. Meng, X. Wang, L. Hou, Z. Wang, S. Chen, Y. Wang, G. Tan, A. Li, W. Wang, *Organometallics* **2017**, *36*, 2706–2709.
- [12] Y. Dai, M. Bao, W. Wang, Z. Xie, C. Liu, Y. Su, *Chinese J. Chem.* **2022**, *40*, 2387–2392.
- [13] T. Kodama, K. Uchida, C. Nakasuji, R. Kishi, Y. Kitagawa, M. Tobisu, *Inorg. Chem.* **2023**, *62*, 7861–7867.
- [14] a) Y.-S. Lin, G.-D. Li, S.-P. Mao, J.-D. Chai, *J. Chem. Theory Comput.* **2013**, *9*, 263–272; b) F. Jensen, *J. Chem. Theory Comput.* **2015**, *11*, 132–138.
- [15] A. Najibi, L. Goerigk, *J. Chem. Theory Comput.* **2018**, *14*, 5725–5738.
- [16] R. L. Martin, *J. Chem. Phys.* **2003**, *118*, 4775–4777.
- [17] T. Lu, F. Chen, *J. Comput. Chem.* **2012**, *33*, 580–592.
- [18] G. Knizia, *J. Chem. Theory Comput.* **2013**, *9*, 4834–4843.
- [19] F. Weinhold, C. R. Landis, E. D. Glendening, *Int. Rev. Phys. Chem.* **2016**, *35*, 399–440.
- [20] a) L. W. Pineda, V. Jancik, K. Starke, R. B. Oswald, H. W. Roesky, *Angew. Chem. Int. Ed.* **2006**, *45*, 2602–2605; b) Y. Ding, H. Hao, H. W. Roesky, M. Noltemeyer, H.-G. Schmidt, *Organometallics* **2001**, *20*, 4806–4811.
- [21] Y. Y. Zhou, D. R. Hartline, T. J. Steiman, P. E. Fanwick, C. Uyeda, *Inorg. Chem.* **2014**, *53*, 11770–11777.
- [22] R. Ditchfield, *Mol. Phys.* **1974**, *27*, 789–807.
- [23] K. Wolinski, J. F. Hinton, P. Pulay, *J. Am. Chem. Soc.* **1990**, *112*, 8251–8260.
- [24] F. Calderazzo, G. Pampaloni, L. Rocchi, U. Englert, *Organometallics* **1994**, *13*, 2592–2601.
- [25] J. R. Cheeseman, G. W. Trucks, T. A. Keith, M. J. Frisch, *J. Chem. Phys.* **1996**, *104*, 5497–5509.
- [26] S. Hirata, M. Head-Gordon, *Chem. Phys. Lett.* **1999**, *314*, 291–299.
- [27] J. Le Bras, H. Jiao, W. E. Meyer, F. Hampel, J. A. Gladysz, *J. Organomet. Chem.* **2000**, *616*, 54–66.
- [28] D. L. Reger, T. D. Wright, C. A. Little, J. J. Lamba, M. D. Smith, *Inorg. Chem.* **2001**, *40*, 3810–3814.
- [29] M. Cossi, N. Rega, G. Scalmani, V. Barone, *J. Comput. Chem.* **2003**, *24*, 669–681.
- [30] F. Neese, *J. Comput. Chem.* **2003**, *24*, 1740–1747.
- [31] G. M. Sheldrick, *Acta Cryst.* **2008**, *A64*, 112–122.
- [32] F. Neese, F. Wennmohs, A. Hansen, U. Becker, *Chem. Phys.* **2009**, *356*, 98–109.
- [33] F. Neese, *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **2012**, *2*, 73–78.
- [34] G. M. Sheldrick, *Acta Cryst.* **2015**, *A71*, 3–8.
- [35] A. L. Spek, *Acta Crystallogr. C Struct. Chem.* **2015**, *71*, 9–18.
- [36] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, Williams, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, D. J. Fox, *Gaussian 16*, Revision C.01. Wallingford, CT, **2016**.
- [37] W. J. Kerr, R. J. Mudd, J. A. Brown, *Chem. Eur. J.* **2016**, *22*, 4738–4742.
- [38] F. Neese, *WIREs Comput. Mol. Sci.* **2022**, *12*, e1606.

RESEARCH ARTICLE

Entry for the Table of Contents



A dichloro-bis(germylene) ligated with a naphthyridine diimine (NDI)-Ge₂Cl₂ undergoes single-electron oxidation, chloride abstraction, and germanium elimination when it reacts with [Cp₂Fe][BARF₄] (Cp = C₅H₅, BARF₄ = [{3,5-(CF₃)₂C₆H₃}]₄B), Na[BARF₄], and Li[HB(sBu)₃], respectively. The NDI-Ge₂ClH intermediate, which is formed via single Cl/H exchange, is proposed to lead to the formation of the thermally stable product NDI-Ge.

Institute and/or researcher Twitter usernames:

@uni_WUE

@ICBWuerzburg

@felipefantuzzi

@LabBraunschweig