



Kent Academic Repository

Lo, Michaela, Morgans, Courtney L., Santika, Truly, Mumbunan, Sonny, Winarni, Nurul, Supriatna, Jatna, Voigt, Maria, Davies, Zoe G. and Struebig, Matthew J. (2024) *Nickel mining reduced forest cover in Indonesia but had mixed outcomes for well-being*. One Earth . ISSN 2590-3322.

Downloaded from

<https://kar.kent.ac.uk/107575/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1016/j.oneear.2024.10.010>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts


If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal** , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Nickel mining reduced forest cover in Indonesia but had mixed outcomes for well-being

Graphical abstract



Authors

Michaela G.Y. Lo,
Courtney L. Morgans, Truly Santika, ...,
Maria Voigt, Zoe G. Davies,
Matthew J. Struebig

Correspondence

m.lo@kent.ac.uk (M.G.Y.L.),
m.j.struebig@kent.ac.uk (M.J.S.)

In brief

As global demand for nickel rises, it is vital to understand the socio-environmental impacts of nickel mining in Indonesia, a country key to global biodiversity and nickel supply. However, the extent of these impacts has not been thoroughly examined. Using a comprehensive analysis of 7,721 villages between 2011 and 2018, we found that nickel mining contributed to deforestation and environmental pollution in local communities. Stakeholders must adopt more sustainable mining practices to protect ecosystems while safeguarding the well-being of local communities.

Highlights

- Nickel mining amplified deforestation in Sulawesi, Indonesia
- Mining led to mixed outcomes for the well-being of local communities
- Improved living standards were outweighed by declines in other well-being dimensions
- Biophysical and socio-demographic factors moderated well-being outcomes



Article

Nickel mining reduced forest cover in Indonesia but had mixed outcomes for well-being

Michaela G.Y. Lo,^{1,7,*} Courtney L. Morgans,¹ Truly Santika,² Sonny Mumbunan,^{3,4} Nurul Winarni,⁵ Jatna Supriatna,^{5,6} Maria Voigt,¹ Zoe G. Davies,¹ and Matthew J. Struebig^{1,*}

¹Durrell Institute of Conservation and Ecology (DICE), University of Kent, Canterbury CT2 7NR, UK

²Natural Resources Institute (NRI), University of Greenwich, Chatham Maritime ME4 4TB, UK

³Center for Climate and Sustainable Finance, Faculty of Mathematics and Natural Sciences, University of Indonesia (UI), Depok 16424, Indonesia

⁴Master of Public Policy in Climate Change, Faculty of Social Sciences, Indonesian International Islamic University (UIII), Depok 16416, Indonesia

⁵Research Center for Climate Change, Faculty of Mathematics and Natural Sciences, University of Indonesia (UI), Depok 16424, Indonesia

⁶Department of Biology, Faculty of Mathematics and Natural Sciences, University of Indonesia (UI), Depok 16424, Indonesia

⁷Lead contact

*Correspondence: m.lo@kent.ac.uk (M.G.Y.L.), m.j.struebig@kent.ac.uk (M.J.S.)

<https://doi.org/10.1016/j.oneear.2024.10.010>

SCIENCE FOR SOCIETY Indonesia, a global hotspot for forests and biodiversity, is also the world's largest producer of nickel. The rising demand for nickel to facilitate the low-carbon transition creates a dilemma: mining can boost economies and support climate goals but might also harm ecosystems and local communities. Accounting for the impacts of nickel production is urgently needed. Our examination of 7,721 villages in Sulawesi, the primary nickel-producing region in Indonesia, showed that deforestation nearly doubled between 2011 and 2018 in nickel-mining villages. The impacts on local community well-being—encompassing living standards, environment, infrastructure, health, social cohesion, and education—varied at different mining stages. Over the whole period, improvements to infrastructure and living standards were outweighed by worsening environmental well-being. We urge policymakers and nickel-mining companies to implement regulations and practices that mitigate environmental damage.

SUMMARY

Soaring demand for nickel to support the low-carbon transition is driving extensive mining in mineral-rich countries, but the environmental and social impacts of nickel mining remain underexplored. Here, we use a counterfactual approach to examine nickel-mining outcomes on forests and the well-being of nearby communities in Sulawesi, Indonesia—a region renowned for its biodiverse tropical forests and now a global center of nickel production. By examining changes across 7,721 villages between 2011 and 2018, we show that deforestation in nickel-mining villages nearly doubled. During the early stages of mining, environmental well-being, living standards, and education outcomes declined, but improvements were observed in health, infrastructure, and social relations. Environmental well-being continued to substantially deteriorate in the later stages of mining production, especially in villages with already high poverty. These findings highlight the environmental and social consequences of nickel mining, underscoring the need for greater accountability of local outcomes if the sector is to support a just and sustainable low-carbon transition.

INTRODUCTION

The development and adoption of low-carbon technologies is crucial for reducing the effects of climate change and meeting the Paris Climate Agreement targets.¹ As it stands, less than one-fifth of energy production comes from renewables,² so dramatic expansion of the sector is expected in the coming decades.³ Yet renewable energy production is highly mineral inten-

sive, and an additional 3 billion metric tons of metal is expected to be required to realize the Paris Agreement goals.¹

The global transition to low-carbon energy will not be possible without nickel extraction.¹ Nickel is a critical component in rechargeable batteries and is widely used in stainless-steel production. Between 2011 and 2018, demand for nickel increased by 43% globally,⁴ and it is increasingly recognized as a critical mineral for energy (Table S1). Over one-third of all nickel is mined






Figure 1. Mining concessions across Sulawesi and Indonesia

(A) Nickel mining (orange) concessions and other mining (purple) concessions in production phase in Sulawesi before (2000–2010, left) and after (2011–2020, right) mining policy changes were introduced. Forest cover in 2011 is shown in green.

(B) Total area of mining concessions for the top five mineral commodities produced in Sulawesi (black), before and after the mining policy changes, in relation to the rest of Indonesia (gray), according to the Indonesian Ministry of Energy and Mineral Resources.

in Indonesia, making it the largest producer in the world,⁵ employing 1.3 million people.⁶ In 2021, >1 million metric tons was produced in the country, a near 4-fold increase of nickel produced in a decade, with most of this coming from Sulawesi (Figure 1).⁴ The role of extractive industries in alleviating poverty and creating job opportunities has been used to justify policies that ease business and foreign direct investment into the nickel-mining industry.⁷ However, this justification has also been criticized for overlooking the detrimental environmental and social impacts on communities directly affected by mining operations. Environmental damage and communal violence associated with nickel mining have already been reported in the region.⁸

It is widely acknowledged that mining is one of the major drivers of land-cover change, having significant implications on

biodiversity and ecosystem functioning.⁹ Similarly, the expansion of mining has the potential to radically transform local societies and economies.¹⁰ Yet few studies address the observed impacts of nickel mining specifically. Previous studies evaluating mining impacts have typically aggregated multiple mineral commodities together^{11,12} rather than examining variation among them.¹¹ This is in stark comparison to individual agricultural commodities,^{13,14} for which environmental and social effects of producing specific crops are thoroughly evaluated (e.g., Santika et al.¹⁴). Indeed, the impacts of mining for different mineral commodities will vary depending on, for example, the amount of land required, the extraction technique, water used, and pollution generated. In Sulawesi, nickel is mostly derived from nickel laterite ores, close to the earth's surface.¹⁵ The extraction

process typically involves digging shallow open-cut mines to access the nickel ore, which can lead to large areas of land being cleared.¹⁶ Moreover, nickel mining relies heavily on machinery, requiring a workforce with specialized skills and higher levels of education compared to more labor-intensive forms of mineral extraction.^{17,18} Nickel mining may have limited local employment opportunities if communities do not match the required human capital.¹⁷ It is crucial to account for these variations to robustly assess land-use change patterns and understand how the benefits and costs of producing different mineral commodities impact local communities.¹¹

While there is some evidence evaluating the impact of nickel mining, the highly localized context of these studies makes it challenging to derive general insights. Some case studies have found that nickel mining has contributed to increasing local income,¹⁹ while in others waste from nickel-mining extraction damaged local fishing and farming production, resulting in greater economic losses overall.⁸ Such case studies offer valuable insights into the effects of nickel mining on local communities and ecosystems, but few are widely generalizable due to methodological inconsistencies.²⁰ In contrast, aggregated measures, such as the percentage of national GDP attributed to the nickel-mining sector, risk overlooking the spatial characteristics that can reveal patterns resulting from nickel-mining activities.²¹ A robust impact evaluation of nickel mining that accounts for spatial variation over broad areas could yield crucial insights for improving policy design and implementation, ultimately supporting better environmental and social outcomes.²²

Evaluating the causal effects of mining interventions is complex due to confounding factors influencing both the outcomes of interest and the underlying pattern of where production occurs (e.g., biophysical characteristics; land governance). Impact-evaluation methodologies can help address this problem by comparing outcomes to a counterfactual condition where no intervention has occurred.²² Such evaluations can be enhanced by tracking outcomes over time, both before and after the mining intervention, further strengthening estimates of causal effects.

To date, large-scale mining-impact studies have highlighted the potential environmental and social risks involved,^{23,24} especially those that overlap protected areas or indigenous lands.²⁵ Yet impact-evaluation studies addressing ongoing mining impacts tend to focus exclusively on either environmental outcomes^{9,11,26,27} or well-being^{28,29} but rarely both. Furthermore, well-being is often limited to a single economic or institutional indicator to signal societal development. Indeed, the impacts of land-use change may be experienced differently across various types of communities and the type of livelihood activities in which they engage.³⁰ Using a combination of indicators can simultaneously capture the benefits and costs for both the environment and local communities³¹ and help to identify the contexts in which synergies or trade-offs between environmental and well-being outcomes exist.³²

Here, we use rigorous impact-evaluation methods to examine and quantify the environmental and social impacts of nickel-mining extraction, focusing on forest-cover change and the well-being of villages across Sulawesi, Indonesia. By collating data at the village level, we capture local effects over a large spatial scale and reveal general patterns. We specifically investigate the effects of nickel extraction compared to

other types of mining to understand the relative implications on forests and well-being. We found that deforestation was higher in villages with nickel mining compared to those with no mining. For villages overlapping nickel-mining areas, we found slower improvements to overall well-being and mixed outcomes across different well-being dimensions. We also identify the biophysical and socio-economic factors moderating the intensity of these impacts. Our findings highlight the extent to which nickel mining affects forest ecosystems and community livelihoods. These insights can inform policymakers and mining companies aiming to implement sustainable practices, ensuring that nickel-mining production aligns with environmental and development objectives.

RESULTS

Methods summary

Nickel-mining data were derived from Indonesia's mining concession map released by the Ministry of Energy and Mineral Resources. Mining concessions were overlaid with high-resolution estimates of forest cover as well as socio-economic data from Indonesia's *Potensi Desa* (or "village potential," PODES) census to estimate the environmental and social changes attributed to nickel-mining production and how this compared with other mineral commodities. To establish a causal estimate of nickel-mining impacts on environmental and social outcomes, we apply statistical matching within a before-after-control-impact (BACI) analytical design to measure counterfactual outcomes in the absence of mining.³³ Such an approach avoids misleading findings by (1) accounting for the contextual dynamics through which forest cover and well-being might have occurred while (2) isolating impacts of nickel-mining extraction from other biophysical, political, and social factors that influence environmental and social outcomes.

To measure the impact of nickel mining (and other mineral commodities) on deforestation, we quantify the change in forest cover across individual villages in Sulawesi throughout the 8-year evaluation period. If the expansion of nickel mining caused deforestation, we would expect to observe a greater reduction in forest cover within nickel-mining villages compared to that experienced in non-mining (control) villages. Forest cover was estimated in the years 2011, 2014, and 2018 to match the time points of the PODES census. Well-being outcomes were tracked using 18 indicators derived from PODES, which we evenly grouped into six dimensions³⁴ (Table 1): education (access to education facilities and support), environment (the occurrence of natural disasters and pollution), health (accessibility of health facilities and cases of disease), infrastructure (access to markets and financial support), living standards (basic living conditions), and social relations (cooperation and incidence of conflict). The overall well-being score is the total combination of all six dimensions, each given equal weighting. If nickel mining improved the well-being of local communities, we would expect a positive change in well-being relative to non-mining villages. Again, well-being was measured in the years 2011, 2014, and 2018 to follow the years the PODES census was implemented. All references to the datasets cited can be found in Table S2.

Table 1. Construction of a well-being index for Indonesia

Dimension ^a	PODES indicator	Description	Score of 0 (low well-being) or 1 (high well-being). A score of 1 is given if:
Living standards	WATER	main source of drinking water	water is obtained from bottled water, refill water, plumbing with meter, or plumbing without meter
	TOILET	type of toilet facility	majority of households have a private facility
	FUEL	type of cooking fuel	fuel source is LPG or gas
Health	FACILITIES	availability of healthcare facilities	there are healthcare facilities, and the nearest polyclinic is <19 km away
	MALNUTRITION	cases of malnutrition	where less than two cases of malnutrition are reported per 1,000 of the village population in the last year
	DISEASE	mortality due to malaria or vomiting	no mortality has occurred due to malaria nor diarrhea reported in the last year
Education	PRIMARY	presence of primary school	at least one primary school is present
	JUNIOR	presence of junior high school	junior high school is <3 km away
	LITERACY	at least one literacy support program	at least one literacy support program is available
Environment	WATER	occurrence of water pollution	no water pollution has occurred in the last 3 years
	AIR	occurrence of air pollution	no air pollution has occurred in the last 3 years
	DISASTERS	occurrence of natural disaster	no natural disaster has occurred in the last 3 years
Infrastructure	SKTM	number of households with poverty (SKTM) letters	the number of families with SKTM letters is no more than 10% of village household population
	MARKET	presence of permanent or semi-permanent market in village	where nearest permanent or semi-permanent market is <10 km away
	CREDIT	availability of food credit, small business credit, people business credit, and housing credit	village with any sort of access to credit support
Social	COOPERATION	mutual cooperation activities	mutual cooperation activities are present
	CRIME	any serious crimes occur in the last 3 years	village reporting three crimes or less in the past year
	CONFLICT	a report of mass conflict in the past year	no report of mass conflict in the last year

^aEighteen PODES³⁵ indicators were grouped to represent six dimensions of well-being (living standards, environment, infrastructure, health, social, and education).³⁴ To calculate overall well-being, each indicator was given an equal weighting within a dimension (1/3). Each individual indicator was given the binary score of 0 or 1, where 0 denotes a village falling below the acceptable threshold specific to that indicator. References and visualization of the pathways between nickel mining and village well-being outcomes are shown in [Figure S3](#). [Table S6](#) provides details on the directionality of expected well-being outcomes and the possible causal pathway mechanisms.

Background changes in forest cover and well-being

After excluding villages with no forest within their boundaries, the average village in Sulawesi had around 36% forest cover in 2011 ($n = 4,458$, [Figure 2A](#)), which dropped to 34% by 2018 (a decline of 2 percentage points, equating to 15 ha for a median village area of 763 ha). Over the same period, village well-being increased across Sulawesi by 4 percentage points ([Figure 2B](#)). Living standards improved by nearly 40 percentage points in both mining and non-mining areas compared to the 2011 baseline ([Figure 2C](#)). Health also improved by 5 percentage points, while education, social, and environmental well-being declined by 3, 4, and 6 percentage points, respectively.

The impact of nickel mining and other mineral types

We estimated the relative impact of nickel mining on forest cover compared to non-mining (control) villages. The analysis indicates that nickel mining exacerbated deforestation over the 8-year period. Deforestation was 2 percentage points greater in nickel-mining villages relative to non-mining villages ($n = 132$, 95% confidence interval [CI]: -3.1 to -0.9 ; [Figure 3A](#)). Villages associated with other mineral types ($n = 115$) also experienced greater deforestation compared to controls; however, this difference was marginal (-0.4 percentage points, 95% CI: -1.5 to 0.7). Since the 2011 baseline year, forest cover declined by 4.4 percentage points in nickel-mining villages, while the decline was 2.4 percentage points where there was no mining




Figure 2. Background (i.e., unmatched) changes in forest cover and well-being between 2011 and 2018

(A) Changes in forest cover between 2011 and 2018, with *n* indicating the number of villages in each forest cover category.

(B and C) Change (B) in overall well-being and (C) across the six well-being dimensions (the indicators used within each dimension can be found in Table 1) by subgroups. Well-being changes in nickel-mining villages are shown in orange, other mining villages (all other mineral commodities excluding nickel) are shown in purple, and non-mining villages (no mines present from 2000 to 2018) are shown in white.

comes between mining of other commodities and non-mining villages.

Factors influencing the impacts of nickel mining

To further explore the heterogeneity of nickel-mining impacts, we selected potential characteristics that could be expected to moderate the effect size. Results in Table S5 and Figures S1 and S2 imply that the effects of nickel mining on deforestation and well-being are indeed highly heterogeneous. Nickel mining was associated with greater deforestation in villages

that were more accessible, as shown by the positive values for the interaction terms (Figure S1 and Table S5). Accessibility was evaluated using a travel-cost surface model that accounts for the topography, land cover, and road density (Table 2). Nickel mining was also associated with greater deforestation at higher elevations and steeper slopes.

We further explored whether other factors, such as poverty baseline conditions (low versus high well-being scores in 2011), livelihood type, and accessibility in villages, moderated the impacts of nickel mining on well-being. Smaller well-being improvements in nickel-mining villages were found where poverty conditions were already high (Figure S2A). Well-being impacts were worse in villages where capture fisheries were the dominant livelihood (Figure S2B). Accessibility had no observable influence on the effect size (Figure S2C). We further examined the specific pathways in which nickel-mining impacts could be moderated by poverty baseline conditions by running an additional matching and regression analyses by low- versus high-poverty villages. Within the first 3 years of production, deforestation was greatest in nickel-mining villages where poverty conditions were initially low (−1.4 percentage points, 95% CI: −2.4 to −0.4) rather than high (−1.1 percentage points, 95% CI: −2.1 to −0.1; Figure 5A). In the later stages of production, deforestation was similar between the two poverty groups. In villages where poverty levels were initially high, well-being declined by 1.8 percentage points (95% CI: −7 to 3.5) (Figure 5B). This decline was mostly driven by a 13.7-percentage-point decrease in environmental well-being (95% CI: −26.7 to −0.7) and a 13-percentage-point reduction in health outcomes (95%

(Figure 3B), representing a near 2-fold increase in deforestation associated with nickel mining compared to controls (Table S3). Villages overlapping nickel mining also experienced slower improvements in overall well-being, which is reflected in the negative coefficient in Figure 3C (−2.1 percentage points, 95% CI: −4.5 to 0.2; Table S4).

We further partitioned our analyses to examine mining outcomes at an early (1–3 years) and later (4–7 years) stage of production to reflect the accumulation of impacts over time. The negative coefficient in Figure 4A shows that nickel-mining villages lost 1.2 percentage points more forest than control areas in the first 3 years of production (95% CI: −2.3 to −0.1), and a further 0.4 percentage points (−1.6 percentage points, 95% CI: −2.5 to −0.7) thereafter. For villages associated with other mines, the impacts took longer to accrue (−1.8 percentage points at 4–7 years, 95% CI: −4.3 to 0.7; Figure 4A). Nickel-mining villages experienced a 5-fold decrease in their overall well-being between the early (−0.5 percentage points, 95% CI: −3.6 to 2.6) and later (−2.5 percentage points, 95% CI: −6.6 to 1.7) stages of production (Figure 4B). This was mostly driven by large reductions in environmental well-being (−11.3 percentage points, 95% CI: −21.8 to −0.8; Figure 4C). Education, health, and social well-being also declined over time, while infrastructure and living standards improved. However, the effect sizes were highly variable, as reflected by wide confidence intervals. For villages near other mines, overall well-being marginally improved compared to controls (0.2 percentage points 4–7 years after production, 95% CI: −3.7 to 4.2; Figure 4B). Across the six dimensions, there were marginal differences in well-being out-




Figure 3. Relative impact of mining nickel and other commodities on changes to forest cover and well-being in Sulawesi

Coefficient plot (A) compares forest-cover change in villages with nickel mines ($n = 132$, orange dots) or other mines ($n = 115$, purple triangles) relative to their respective non-mining control villages. The error bars represent 95% confidence intervals: if these cross the zero line there is no significant difference between mining interventions and non-mining control villages. Bar plots (B) depict the change in forest cover since the 2011 baseline year in nickel-mining villages and matched controls and in other mining villages versus their matched controls. Here, the confidence intervals show whether forest cover in 2018 differed from that in the baseline year (i.e., by not crossing the zero line). Plots (C) and (D) should be interpreted in the same way but for overall village well-being. Further details on the interpretation of the figure can be found in [Tables S3 and S4](#). [Table 1](#) provides an overview of the well-being dimensions and indicators, which together comprise the overall well-being score.

largest nickel-producing regions of the world. Combining rich spatial data across 7,721 villages, we highlight the differential impacts of mining nickel versus other mineral commodities, variations of outcomes over time, and the factors that minimized or exacerbated environmental and social outcomes.

Nickel mining impacts deforestation and well-being

Nickel mining contributed to a significant increase in deforestation since the expansion of the sector in Indonesia around 2011. Studies using similar methods elsewhere have found limited evidence of deforestation attributed to mining, as other drivers of forest-cover change would be present even in the absence of mining.²⁶ Conversely, others reported mining to be a key contributor to deforestation across Sulawesi.^{43,44} We find that the extent of deforestation associated

CI: -24.7 to -1.2) but was countered by improvements to living standards and infrastructure ([Figure 5C](#)).

In contrast, villages with initial low levels of poverty experienced short-term improvements to health (6.1 percentage points, 95% CI: 0.1 to 12.1; [Figure 5C](#)). At the same time, living standards improved substantially 4–7 years post production (10.6 percentage points, 95% CI: -3.9 to 25.2).

DISCUSSION

The demand for low-carbon energy is a key driver of nickel production.⁴² Examining empirical evidence on the ways in which landscapes and people are being transformed by nickel mining is crucial to addressing sustainability challenges. Our analysis identified the multiple impacts of nickel mining in one of the

with nickel mines was substantial—near double that observed in matched control areas. The environmental sustainability narrative frequently used to justify the increasing supply of nickel therefore risks overlooking other environmental consequences resulting from nickel mining. This is particularly relevant in Sulawesi, which is globally recognized for its unique ecosystems and biodiversity.⁴⁵ Concerted resources to mitigate against deforestation are therefore needed.

The marginal decline in overall well-being associated with nickel mining was largely driven by a decrease in environmental indicators. Waste materials and pollution resulting from nickel extraction and processing is a long-standing issue.¹⁵ Our analysis supports the notion that safeguards against pollution and mining-related disasters should be strengthened to minimize negative environmental impacts that affect people's well-being.




Figure 4. Relative mining impacts across villages in Sulawesi over time

Impact of nickel mining (orange dots) or other mining (purple triangles) relative to changes occurring in non-mining control villages 1–3 years ($n = 132$ for nickel; $n = 115$ for other mines) and 4–7 years ($n = 44$ for nickel; $n = 47$ for other mines) following production. The intervals of 1–3 and 4–7 years post mining are determined by the availability of census data. (A) and (B) show the impacts for forest cover and overall well-being, respectively. (C) shows the well-being outcomes by dimension. Error bars represent 95% confidence intervals between mineral commodities and control villages (the zero line). [Table 1](#) provides an overview of the six dimensions and 18 indicators, which together comprise the overall well-being score.

Incorporating environmental and social standards into existing international and national governance mechanisms, such as the Extractive Industries Transparency Initiative (EITI), could be a positive step toward such endeavors. However, such standards are currently lacking due to alleged inconsistencies between governing bodies of extractive industries at the local level and the EITI at the Indonesian national level.⁴⁶ Greater coordination and alignment between local, regional, and national governments in Indonesia are crucial to ensuring consistency in implementing environmental and social safeguards.^{46,47} Other actors, including local and international non-governmental organizations, along with industry groups, play crucial roles in holding mining companies accountable to adherence with international standards, such as the OECD's Due Diligence Guidance.⁴⁸ Due diligence protocols can help companies to better integrate human rights into environmental and social assessments and further avoid negative social and environmental impacts of nickel mining.

Reductions in social well-being were also typical of areas overlapping nickel mines. Conflict driven by environmental damage and land acquisition from nickel-mining activities has been reported in Indonesia.⁸ However, we also found the social well-being was highly variable between individual villages, making it difficult to draw conclusions. It is important to also acknowledge the positive contribution of nickel mining to local living conditions and infrastructural development. These improvements are likely to be attributed to the construction of transportation and water-based infrastructure.¹⁷ Revenue from nickel mining may have facilitated local government investment within communities.¹⁷ Greater variability across villages was also evident for these indicators, implying that both positive and negative outcomes are experienced collectively across Sulawesi, and more information

research progresses with new data and advancements in impact-evaluation methods, further work will play a key role in identifying these other factors that influence the impact of nickel-mining extraction.

Certain effects of nickel extraction, such as the decline in environmental well-being, were only detected 4–7 years after the issuance of mining leases. One explanation for this is that some indicators, such as environmental pollution and flooding, take time to accrue, making them detectable only after several years.⁵⁰ Indeed, given the relatively short- to mid-term trends we uncover, the longer-term effects of nickel mining should also be assessed, including after mine closure. Evidence from historical tin mining in Indonesia demonstrated negative effects on local employment once the mines were closed.⁵¹ When data become available in the nickel sector, such studies will be important in addressing the lasting effects of mining as well as opportunities for restoring and rehabilitating post-mining landscapes.

Comparisons between nickel and other mines

By disaggregating mining concessions by mineral commodity, we determined the impacts of nickel mining compared with other types of mines. Our results reveal that the onset of deforestation in nickel-mining villages occurs faster than it does in villages where other types of mines are present. This implies that the land-cover changes associated with mining depend on the mineral commodity extracted. We also found differences in well-being between mining nickel versus other minerals in terms of effect size and directionality. All mining was associated with worsening environmental well-being, but this was stronger in nickel-mining areas. Conversely, impacts on social well-being were divergent—nickel mining worsened social

Table 2. Covariates that influence the assignment of mining villages and trends in forest cover and well-being

Covariate type ^a	Covariate details ^b	Rationale ^b	Dataset ^b
Biophysical	average elevation (m.a.s.l.) (log(continuous))	correlated with deforestation and livelihood decisions ³⁶	SRTM 90 m Digital Elevation Database v.4.1
	average slope (degrees) (log(continuous))	correlated with deforestation and livelihood decisions ³⁶	SRTM 90 m Digital Elevation Database v.4.1
	average rainfall in dry season (continuous)	affects livelihood decisions ³⁶	WorldClim
	average rainfall in wet season (continuous)	affects livelihood decisions ³⁶	WorldClim
	area of limestone rock (%) (continuous)	geological characteristics shape mineral selection ³⁷ (only included in “other mineral commodity” impact evaluation)	RePPPProT
	area of volcanic rock (%) (continuous)	geological characteristics shape mineral selection ³⁸ (only included in “other mineral commodity” impact evaluation)	RePPPProT
	area of forest cover in 2011 (%)	previous area of forest cover is correlated with deforestation activities already existing	forest cover and forest-cover change layer
Land governance	area of village under production forest (<i>hutan produksi</i>) (continuous)	mining concessions allowed in production forest sites ⁶	Forest Zone Map
	area of village under protected forest (<i>hutan lindung</i>) (continuous)	underground mining activities can take place in protected forests. Open pit mining is not permitted ⁶	Forest Zone Map
	area of village that is non-forest estate (areal <i>penggunaan lain</i>) (continuous)	mining concessions are allowed on non-forest estate ³⁹	Forest Zone Map
	administrative provincial boundary where villages are located (categorical)	government decision-making at the province level influences distribution of resources ⁴⁰	Potensi Desa (PODES)
	area of village (km ²) (log(continuous))	village area linked to size of mining area	PODES
Socio-demographic	population density (capita/km ²) (log(continuous))	higher resource extraction is associated with high population levels ⁴¹	PODES
	slope, road, and land cover to measure level of accessibility to settlements (travel time, hours) (log(continuous))	greater access to markets and infrastructure can influence livelihood decisions and forest cover ⁴¹	accessibility layer
	village poverty score in 2011 (categorical [low, high])	poverty baseline level may moderate well-being outcomes	PODES
	primary village livelihood at baseline year (2011) (categorical [capture fisheries; commercial fisheries; market-orientated; subsistence; other])	livelihood type shapes well-being outcomes ³⁰	PODES

^aCovariates are grouped into three types: (1) biophysical, (2) land governance, and (3) socio-demographic.

^bDetails, rationale, and source of each covariate included in the analysis. The source of each dataset cited is listed in Table S2.

well-being, while slight improvements were associated with other mines. Incorporating mineral-commodity-level data into impact-evaluation assessments therefore helps inform specific mineral-commodity chains.^{11,32} Nonetheless, we also found large variations across individual villages, which could be due to other factors that have not been included in the analyses. The source dataset on mine locations classified concessions by mineral commodity but missed information on other mining characteristics that may have also influenced the type of deforestation and well-being patterns detected. For example, commercially sensitive information on the amount of production, extraction methods, and the type of legal ownership (such as being domestic or foreign owned) is not accessible. More detailed information on such mining characteristics will better capture how these specific attributes can shape well-being

and deforestation,⁵² but this requires these data to be made available.³²

Other factors moderating nickel-mining impacts

Identifying the underlying conditions that improve or hinder efforts toward reducing deforestation and improving the livelihoods of local communities is important for guiding the management of mining activities and land-use planning. As expected, more accessible villages lost more forest compared to less accessible areas, implying that accessibility of transport, facilities, and infrastructure to clear forests incentivizes companies to clear more land. That said, we also found that nickel mining had led to greater deforestation at higher elevations. While we may assume that upland sites might have more difficulty establishing mining operations and thus disincentivize forest




Figure 5. The role of poverty baseline conditions in determining forest-cover change and well-being outcomes in nickel-mining villages

Impact of nickel mining where 2011 poverty baseline conditions were low ($n = 61$ [1–3 years], $n = 19$ [4–7 years], yellow dots) or high ($n = 70$ [1–3 years], $n = 34$ [4–7 years], green dots) compared to non-mining (control) villages with the similar poverty baseline conditions. (A) and (B) show the impacts for forest cover and overall well-being, respectively. (C) shows the well-being outcomes by dimension. The intervals of 1–3 and 4–7 years post mining are determined by the availability of census data. Error bars represent 95% confidence intervals between nickel mining and control villages (the zero line). Table 1 provides an overview of the six dimensions and 18 indicators, which together comprise the overall well-being score.

clearance, such outcomes could be due to the relatively low rates of deforestation experienced in these areas more broadly. These results imply that nickel mining may be one of the key drivers of deforestation in upland regions, especially compared to other activities such as logging and agriculture, which tend toward greater deforestation in the lowlands.⁴⁴ Our analysis points to specific biophysical regions that require greater attention should policies focus on targets to avoid deforestation.

We also found that the well-being impacts of nickel mining differed according to livelihood type and poverty baseline conditions. Villages where the primary source of income was from capture fisheries experienced greater reductions in well-being compared to those where other livelihoods (e.g., commercial and market-orientated fisheries) were dominant. This is consistent with our findings of declines in environmental well-being attributed to nickel mining as well as observations in the broader literature. For example, water and environmental pollution from the largest nickel mine in Indonesia has led to reduced fish stocks in water bodies that are close to extraction sites.⁵³

We focus our analyses on land-based concessions and thus provide conservative estimates of the damage to water-based livelihoods, such as fisheries, from nickel-mining production. Nickel mining also extends to offshore locations employing large trawls carried across the ocean floor to extract nickel from the seabed.⁵⁴ Under Indonesia's Omnibus Law (*Undang-Undang Cipta Kerja*, Law 11/2023), offshore mineral mining is no longer limited within 12 km from the coastline and can now take place in all maritime (including deep-water) areas within the country's jurisdiction. While we were only able to account for mines that overlapped with onshore villages, the disturbance from offshore mining activities and the waste produced may have even greater

adverse effects on marine ecosystems and the livelihoods of fishers in the future.⁵⁵ Poorer villages were more likely to experience the negative effects of nickel mining on environmental well-being and health. Villages with high levels of deprivation may have limited resources and capacity to cope with environmental pollution associated with mining activities, leading to adverse health effects.⁵⁶ While initial poverty baseline conditions influenced the impacts of nickel mining, we are not suggesting that this should be the sole consideration when establishing concessions. On the contrary, engaging local communities in decision-making, development, and implementation processes can strengthen mining governance and planning⁵⁷ and further empower marginalized social groups.^{10,58} Indonesia has made significant developments in engaging communities in Environmental Impact Assessment regulations by making public participation mandatory.⁵⁹ To be effective, engagement efforts should be paired with greater resources and action to strengthen the capacity of local communities, thereby reducing their vulnerability to potential negative impacts of land-use activities.⁶⁰

Informing the low-carbon transition

There are multiple potential pathways to a low-carbon transition, including policy action that lowers nickel demand by improving the recyclability of renewable technologies and reducing overall consumption of energy.^{61,62} Yet despite such actions, there remain questions about whether this will be enough to curb projected mineral demand trends in coming decades.⁴² With more countries looking to electrify transport to achieve 2050 net-zero emission targets, heightened demand for nickel is anticipated, with some projections predicting a more than doubling of current levels.^{1,42,63} Around 75% of this supply is expected to come from Indonesia,⁴² which is seeking to attract more foreign investment and increase the ease of business, as evidenced in the Omnibus Law of 2023.⁶⁴ Nickel is expected to be a key resource at the center of Indonesia's business and investment ventures.⁶⁵

There are rising concerns about potential weakening of environmental and social regulations under the new Omnibus Law.⁶⁴ Yet since its initial announcement in 2020 the law has faced several implementation challenges, with few regulations fully realized.⁶⁴ Our study is therefore well timed to inform the design of implementing regulations under the new law, ensuring that the nickel-mining sector operates sustainably by balancing the economic gains with environmental and social responsibility. One potential approach is to harmonize policies between land-use sectors, including forestry and mining, which often remain largely separated. In the case of Indonesia, these divisions exist in licensing, development planning, and environmental impact assessments and have led to challenges in overlapping resource concessions and unclear tenure rights.⁶⁶ While permission may be granted to undertake mining operations within concessions, this does not necessarily cover indirect impacts in surrounding areas.¹¹ Mining extraction rarely occurs in isolation, with disturbances also occurring off-site via road development, energy infrastructure, and settlements. While distinguishing the direct and indirect impacts of mining was outside the scope of our work, previous studies have identified the need to include the cumulative impacts of deforestation outside of concessions.^{9,11} Governance mechanisms such as the One Map Policy (*Kebijakan Satu Peta*, KSP) in Indonesia can potentially overcome these challenges. The initiative aims to harmonize spatial data from government departments and incorporate them into a single database. Promoting initiatives that actively encourage the coordination and integration across sectors can lead toward a more cohesive multi-sectoral approach to natural-resource regulation in an attempt to reduce both deforestation and poverty.⁶⁷

As the mining of critical minerals continues to expand in Indonesia and other countries, it will be important to deepen our understanding of environmental and social impacts so that improvements to sustainability can be targeted. Our work presents a crucial synthesis of the environmental and social outcomes of nickel mining, providing a broad overview across the sector in Sulawesi. However, nickel mining on other Indonesian islands also warrants examination as relevant datasets become available. A caveat to our evaluation is that the outcomes are aggregated to the village administrative unit, reflecting the spatial scale of the census data. This village-level analysis can mask important variations between households or individuals within communities, where some may benefit more or less than the average village outcome. In-depth case studies on how different individuals and groups are impacted by mining will remain valuable for identifying and supporting those most vulnerable to negative impacts while also highlighting the conditions that foster greater benefits. The spatial nature of our analysis enables targeted case studies in areas where mining outcomes have been particularly positive or negative. Furthermore, studying mining outcomes in relation to subjective well-being, which considers how people evaluate their own well-being related to the aspects of life they consider as important,⁶⁸ could deepen our understanding of the diverse ways mining interventions impact local communities. Integrating qualitative methods into impact evaluations can offer a richer spectrum of possible outcomes, ensuring that the voices and perspectives of affected communities are included.⁶⁹

Nickel extraction is expected to boom in the coming years through a global low-carbon transition. We show how this can

conflict with other environmental and development objectives. Pinpointing where these environmental and social divergences occur is a crucial step toward addressing these challenges, minimizing trade-offs, and further promoting mining extraction, thus contributing to a sustainable future for both people and planet.

EXPERIMENTAL PROCEDURES

The study area

Approximately 20 million people live in Sulawesi across its six provinces.³⁵ The complex geological history of central Indonesia has also resulted in highly unique ecosystems, making Sulawesi a globally important region for biodiversity and endemism.⁴⁵ Compared to western Indonesian islands, small-scale agriculture and other subsistence-based livelihoods are more dominant in Sulawesi, and commodities such as coffee, cacao, and coconut are more commonly grown than industrial-scale products such as palm oil.⁴⁵ Consequently, deforestation rates have been much lower in Sulawesi compared to neighboring Borneo and Sumatra, where deforestation is primarily driven by the expansion of large-scale oil palm and paper-pulp plantations.^{43,70} However, a recent deforestation surge in Sulawesi has been linked, in part, to mining, a sector that has experienced rapid growth during the last decade.⁴⁴

The 2009 mining law in Indonesia decentralized the power of issuing mining permits, granting greater authority to local and regional officials.⁷¹ This shift, implemented in 2010,⁴⁵ resulted in a sharp increase in the issuance of mining licenses throughout the country, with most of the permits for establishing nickel-mining operations granted in Sulawesi (Figure 1). By 2020, over 65% (424,270 ha) of Indonesia's active nickel-mining concessions were on Sulawesi, with a further 672,100 ha under exploration. The rapid proliferation of mining permits reportedly led to several licensing issues, including overlapping boundaries of land-use activities and difficulties in monitoring and ensuring that mining companies adhered to national procedures.⁷²

Forest-cover data

To track annual forest-cover change between 2011 and 2018, we used data from the Global Forest Change (GFC) repository v.1.6 (Table S2). The GFC dataset provides consistent and accurate estimates of forest loss when applying national definitions of forest cover and was therefore appropriate for measurement of forest-cover change. Here, forest cover is defined as at least five hectares of >70% tree-canopy cover of natural composition and structure in a 30-m-resolution Landsat pixel,^{14,43} including mangrove forests (Table S2). This definition corresponds with those used for primary and secondary forest by Indonesia's Ministry of Environment and Forestry.⁷³ Using conservative measures of forest minimized the inclusion of deforestation that may have been temporary rather than permanent, and we excluded tree-cover changes in other land-cover types (plantations, agroforests, mixed gardens regrowth, and scrublands).⁴³ All forest maps were converted to the Asia South Albers Equal Area Conic projection to reduce distortions in area and distance. Spatial data were then aggregated to 180 × 180-m pixel size to ease computational processing. Using the conservative definition of forest cover, we restricted tree-cover pixels from the GFC dataset to form a baseline forest-cover map for 2000. We then applied the forest-loss data to the forest-cover map and calculated the proportion of village areas forested for the years 2011, 2014, and 2018 to match the years for which well-being data were available.

Well-being data

Data on well-being were extracted from Indonesia's village-level census, PODES, which provides a rich and extensive source of socio-economic and demographic information across Indonesia. PODES data are collected from village authorities around three times per decade, with the results aggregated at the village (*desa*) administration level (thus, inference should not be linked to the individual or household level). As with all census surveys, the accuracy of responses may vary depending on the capacities and resources available within each village. Rigorous protocols have been developed by the Indonesian Bureau of Statistics (BPS) to ensure that the collected data are consistent and accurate across villages.⁷⁴ Previous studies have used PODES to assess environmental and social effects of land-use policies and other types of interventions, including oil palm certification^{14,75} and protected areas.^{34,76} PODES

therefore remains the richest and most extensive source of well-being data covering the whole of the Indonesian archipelago. Well-being was characterized across three consecutive censuses in 2011, 2014, and 2018 (Table S2), and, due to changes in village boundaries, observations were harmonized to those in 2014. We chose 2011 as the baseline following the announcement of the new mining regulatory regime in 2009, which was officially implemented in 2010.⁷¹

We define well-being as a multi-dimensional concept that recognizes the multiple assets, abilities, and attributes that are needed to support and achieve a better life.⁷⁷ To capture these multiple facets, our overall well-being index comprises six dimensions: living standards, environment, infrastructure, health, social, and education (Table 1).³⁴ Each dimension was assigned three equally weighted indicators derived from the PODES questionnaire. To calculate overall well-being, each indicator was given an equal weighting within a dimension (1/3). Each individual indicator is given the binary score of 0 or 1, where 0 denotes a village falling below the acceptable threshold specific to that indicator. The overall well-being index was calculated as the average score across the six dimensions. The indicator dimensions, thresholds, and directionality of measures were informed by established well-being and poverty frameworks, including the global multi-dimensional poverty index³¹ and Indonesia's Village Development Index.⁷⁸ The possible mechanisms and directionality of well-being outcomes for each indicator are presented in Figure S3 and Table S6.

Mining data

We used national mining concession data from Indonesia's Ministry of Energy and Mineral Resources, containing detailed information on the type of commodity and stage of mining activity (Table S2). Compared to other mining databases available, such as the S&P Global Market Intelligence database⁷⁹ and the mining data from Maus et al.,⁸⁰ the national concession database provides greater coverage of mining areas in Indonesia,⁸¹ with more specific information on production status and mining characteristics.

Exploration and production require separate mining permits. We focus our analysis on the local impacts of mining concessions at the stage of production only. While the specific type of activity is not detailed in the mining concession data, construction, extraction, processing, refining, and transportation are all grouped into the stage of production.⁸² The unit of analysis was the village boundary level, matching the same scale as the PODES data. After narrowing our sample to Sulawesi, a total of 417 mining polygons were included in our analysis, covering approximately 540,000 ha (Table S7).

Villages exposed to mining were identified as those with concessions covering at least 15% of the village area (the median value across the Sulawesi dataset). Mining villages were then divided into two groups: nickel-mining villages, where nickel was the primary mineral commodity extracted; and other mining villages, where all other minerals were produced. Non-mining villages were considered as those where no mining activity occurred during the same period for any of the commodities. We excluded villages where mining production operations occurred 10 years before the baseline year, as well as villages where mining concessions covered more than 0% but less than 15% of the village area. The mining database provides a comprehensive list of concessions that have formally received a permit but does not include unlicensed operations. Therefore, to avoid the possible inclusion of informal mining activities, we excluded villages containing concession areas where any exploration activities had taken place prior to and during the study period (2000–2018). This assumes that expected informal mining may occur where exploration or scoping studies have been conducted, which also provide estimates of where nickel resources are located. Mining concessions were merged by mineral-commodity groups (nickel or other mineral) and year of production to avoid issues of overlapping boundaries. Villages with overlapping mining concessions producing a mixture of nickel and other mineral commodities were excluded from our sample.

Mining villages also include mining concessions that were issued mining production licenses in 2011 because (1) the 2011 PODES survey was carried out in April 2011, so information is likely to reflect the status of villages in the previous year,⁷⁴ and (2) the steps from the decision to the construction and startup of mines take time, and we would therefore expect a delay in observing impacts.⁸³ After applying our inclusion/exclusion criteria and removing villages with missing data, 461 villages were excluded, resulting in a sample of 7,721

villages across Sulawesi. Of this total, 7,474 villages experienced no mining activities between 2000 and 2018, 132 villages contained nickel-mining concessions that were in 1–7 years of production, and 115 villages overlapped with other mineral commodities excluding nickel (Table S8). Figure S4 maps mining polygons by nickel and other mineral commodities from 2011 to 2018.

Analytical framework and methodology

For the matching analysis, we carried out two assessments: the first comprised nickel-mining villages matched to non-mining villages; and the second matched other mining villages (excluding nickel) with non-mining villages. For the regression analysis we estimated and compared the difference in forest cover and well-being between the mining and non-mining villages within each set. To increase the robustness of our estimates, we included covariates at both stages, which served to reduce any leftover bias resulting from the matching process.⁸⁴

Covariates of forest-cover change and well-being

There are multiple factors other than mining that could affect how interventions are spatially distributed and further interact with the outcome of interest. Therefore, to reliably measure the impact of mining, other factors that might influence the assignment of mining concessions (e.g., proximity to roads) or forest-cover and well-being outcomes (e.g., poverty baseline conditions) should be controlled for.³³ We identified 16 covariates that may influence the selection process of mining and non-mining villages and influence forest-cover and well-being outcomes (Table 2). These reflected (1) biophysical conditions, (2) land governance, and (3) socio-demographic features of villages. This selection identifies factors that were known to affect the allocation of mining sites as well as other factors that influence forest-cover and well-being outcomes. Prior to the matching process, we log-transformed covariates that were highly skewed.

Statistical matching

For each set of mining interventions (nickel-mining villages and other mining villages), we performed a 1:1 paired matching analysis to balance observed covariates between the mining villages and non-mining villages and make them comparable. We also matched across two periods to account for possible time lags in the impacts from mining.⁵⁰ Throughout, we used genetic matching with replacement, a method that specifically uses a matching algorithm to iteratively search for the best balance.⁸⁵ Analyses were undertaken using the Matching and MatchIt packages in R.^{86,87} As villages were either assigned a binary value of being exposed to a mining intervention or not, using a logistic regression to estimate the propensity scores of villages from the covariates listed in Table 2 was the most appropriate statistical model. For both sets of interventions, all mining villages were matched to non-mining villages (Table S8). After comparing the covariate balance before and after matching, we observed a significant improvement in the overall distribution between mining and non-mining groups in both sets. The normalized differences for all covariates were below 0.2 (Tables S9 and S10), implying that a strong balance was found across our mining and non-mining villages.

Regression analysis

With the matched datasets, we implemented a BACI approach to infer the effects of mining. This approach first uses longitudinal data to compare changes in forest cover and well-being before and after a mining intervention takes place. This change is then compared with cross-sectional differences in forest cover and well-being between mining and non-mining villages²² (see visual diagram of the analysis in Figure S5). Changes in forest cover and well-being were measured over 1–7 years (i.e., up to two census intervals available in the data) after a mining intervention had been introduced, with another set of analyses to further observe changes 1–3 years and 4–7 years after the mining intervention (the intervals matching the census years). In the analysis, standard errors were clustered (1) by subclass, which represents pairs between the paired mining and non-mining village, (2) by village, which accounts for non-mining villages that were included more than once in our matched sample, and (3) at the regency (*kabupaten*) level, as most permits within our sample were issued at this level, therefore accounting for other unobserved political factors within regencies.⁸⁸

Interaction terms were used to examine the factors that might influence the intensity of nickel-mining impacts on deforestation and village well-being. We

hypothesize that villages with greater accessibility may increase the intensity of deforestation caused by nickel mining. As nickel mining is highly capital intensive, the additional costs of transporting resources and labor with limited infrastructure, as well as low access to markets for outputs, may result in a greater disincentive to clearing larger areas of forest lands.⁶⁷ Similarly, establishing nickel-mining concessions on steep slopes at high elevations may be more restricting in the expansion of mining operations, resulting in the greater retention of forest cover.

Another group of interactions with nickel mining was carried out to assess whether livelihood type, poverty baseline conditions, and accessibility moderated the outcomes of interest. Based on other literature, livelihoods are known to influence well-being outcomes.³⁰ Therefore, we hypothesized that livelihoods with a greater dependency on natural resources are more likely to be affected by the negative environmental externalities derived from mining operations. Furthermore, we may also expect poverty baseline conditions to moderate mining outcomes. According to the “natural resource curse” theory, natural-resource extraction exacerbates poverty¹⁷; therefore, we would expect poorer areas to experience a worsening in well-being. In contrast, following the “natural resource blessing” theory, we would expect to see the opposite trend.¹⁷ The poverty baseline conditions of villages were determined as the inverse of the well-being index (the negative directionality of each well-being indicator). Before matching, we grouped each village into two classes of equal intervals—low and high poverty—based on their poverty status in 2011. We also assessed whether the accessibility of villages moderated the impact of nickel-mining production on the well-being of local communities. We might also expect nickel-mining extraction in more accessible areas to be more profitable due to fewer additional costs in transport-related facilities as well as easier access to resources and markets. These profits may lead to greater investment in the local economy, with results showing a greater improvement in well-being over time.

To further assess the pathways through which baseline factors moderated the outcomes of deforestation and across various dimensions of well-being, we ran another matching analysis (Table S11) of the initial poverty status of villages. We reran the same statistical matching analysis across all poverty groups between nickel-mining and non-mining villages to assess deforestation trends and well-being outcomes across each dimension over time.

RESOURCE AVAILABILITY

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Michaela G.Y. Lo (m.lo@kent.ac.uk).

Materials availability

This study did not generate any new unique materials.

Data and code availability

Mining concession data were derived from the Ministry of Energy and Mineral Resources. Visualizations of the mining concessions maps are available at Nusantara Atlas of Deforestation and Industrial Plantations in Indonesia (nusantara-atlas.org) and ESDM One Map—Exploring Energy and Mineral Resources of Indonesia (<https://geoportal.esdm.go.id>). Data used to measure forest-cover change were derived from the 30-m-resolution GFC database: <https://glad.earthengine.app/view/global-forest-change>. Well-being and other socio-demographic data were sourced from the PODES census village survey led by the Indonesian Bureau of Statistics. These data can be visualized using the WebGIS PODES portal (<https://sig.bps.go.id/webmap/podes/>). The code for replication of the statistical analysis can be found at Zenodo: <https://doi.org/10.5281/zenodo.14032310> and GitHub: github.com/michaelaglyo/Nickel-mining-repository.

ACKNOWLEDGMENTS

This study was funded by the Newton Fund Wallacea Programme via the UK Natural Environment Research Council (NERC, NE/S007067/1) and the Indonesian Ministry for Higher Education, Research & Technology (Ristekdikti, NKB-2892/UN2.RST/HKP.05.00/2020 and 1/E1/KP.PTNBH/2019). M.G.Y.L. was supported by the University of Kent Global Challenges Doctoral Centre,

and M.J.S. was supported by a Leverhulme Trust Research Leadership Award. Research in Indonesia was authorized by Ristekdikti under permit 7/TKPIPA/E5/Dit.K1/VI/2019 and subsequently BRIN via permit 2/TU.B5.4/SIP/VIII/2021. We would like to thank Nicolas Deere for feedback on the data and code repository and the anonymous reviewers for their constructive feedback.

AUTHOR CONTRIBUTIONS

Conceptualization, M.G.Y.L. and M.J.S.; data curation, M.G.Y.L., C.L.M., and T.S.; methodology, M.G.Y.L., C.L.M., T.S., M.V., and M.J.S.; resources, T.S. and S.M.; formal analysis, M.G.Y.L., T.S., and M.V.; investigation, M.G.Y.L.; writing – original draft, M.G.Y.L., M.J.S., and Z.G.D.; writing – review & editing, all authors; visualization, M.G.Y.L. and M.J.S.; supervision, M.J.S. and Z.G.D.; project administration, M.J.S. and Z.G.D.; funding acquisition, M.J.S. and Z.G.D.

DECLARATION OF INTERESTS

The authors declare no competing interests.

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.oneear.2024.10.010>.

Received: April 3, 2023

Revised: May 5, 2023

Accepted: October 21, 2024

Published: November 15, 2024

REFERENCES

- Hund, K., Porta, D.L., Fabregas, T.P., Laing, T., and Drexhage, J. (2020). Minerals for Climate Action: The Mineral Intensity of the Clean Energy Transition (The World Bank Group). <https://documents1.worldbank.org/curated/en/099052423172525564/pdf/P16627806f5aa400508f8c0bdcbca0878a3e.pdf>.
- World Bank Group (2017). The Growing Role of Minerals and Metals for a Low Carbon Future (English) (The World Bank Group). <http://documents.worldbank.org/curated/en/207371500386458722/The-Growing-Role-of-Minerals-and-Metals-for-a-Low-Carbon-Future>.
- IEA (2015). Energy Technology Perspectives 2015: Mobilising Innovation to Accelerate Climate Action (International Energy Agency). <https://www.iea.org/reports/energy-technology-perspectives-2015>.
- Statista (2024). Mine production of nickel worldwide from 2010 to 2023. <https://www.statista.com/statistics/260748/mine-production-of-nickel-since-2006/>.
- USGS (2022). Nickel - U.S. Geological Survey, Mineral commodity summaries, January 2022. <https://pubs.usgs.gov/periodicals/mcs2022/mcs2022-nickel.pdf>.
- PwC (2019). Mining in Indonesia: Investment and taxation guide. <https://www.pwc.com/id/en/energy-utilities-mining/assets/mining/mining-guide-2019.pdf>.
- PwC (2023). Mining in Indonesia: Investment, taxation and regulatory guide. <https://www.pwc.com/id/en/energy-utilities-mining/assets/mining/mining-guide-2023.pdf>.
- Hudayana, B., Suharko, and Widyanta, A. (2020). Communal violence as a strategy for negotiation: community responses to nickel mining industry in Central Sulawesi, Indonesia. *Extr. Ind. Soc.* 7, 1547–1556. <https://doi.org/10.1016/j.exis.2020.08.012>.
- Sonter, L.J., Herrera, D., Barrett, D.J., Galford, G.L., Moran, C.J., and Soares-Filho, B.S. (2017). Mining drives extensive deforestation in the Brazilian Amazon. *Nat. Commun.* 8, 1013. <https://doi.org/10.1038/s41467-017-00557-w>.
- Bebbington, A.J., Humphreys Bebbington, D., Sauls, L.A., Rogan, J., Agrawal, S., Gamboa, C., Imhof, A., Johnson, K., Rosa, H., Royo, A.,

- et al. (2018). Resource extraction and infrastructure threaten forest cover and community rights. *Proc. Natl. Acad. Sci. USA* 115, 13164–13173. <https://doi.org/10.1073/pnas.1812505115>.
11. Giljum, S., Maus, V., Kuschner, N., Luckeneder, S., Tost, M., Sonter, L.J., and Bebbington, A.J. (2022). A pantropical assessment of deforestation caused by industrial mining. *Proc. Natl. Acad. Sci. USA* 119, e2118273119. <https://doi.org/10.1073/pnas.2118273119>.
12. Bhattacharyya, S., and Resosudarmo, B.P. (2015). Growth, growth accelerations, and the poor: Lessons from Indonesia. *World Dev.* 66, 154–165. <https://doi.org/10.1016/j.worlddev.2014.08.009>.
13. Pendrill, F., Gardner, T.A., Meyfroidt, P., Persson, U.M., Adams, J., Azevedo, T., Bastos Lima, M.G., Baumann, M., Curtis, P.G., De Sy, V., et al. (2022). Disentangling the numbers behind agriculture-driven tropical deforestation. *Science* 377, eabm9267. <https://doi.org/10.1126/science.abm9267>.
14. Santika, T., Wilson, K.A., Law, E.A., St John, F.A.V., Carlson, K.M., Gibbs, H., Morgans, C.L., Ancrenaz, M., Meijaard, E., and Struebig, M.J. (2021). Impact of palm oil sustainability certification on village well-being and poverty in Indonesia. *Nat. Sustain.* 4, 109–119. <https://doi.org/10.1038/s41893-020-00630-1>.
15. Mudd, G.M. (2010). Global trends and environmental issues in nickel mining: Sulfides versus laterites. *Ore Geol. Rev.* 38, 9–26. <https://doi.org/10.1016/j.oregeorev.2010.05.003>.
16. Jaffré, T., Munzinger, J., and Lowry, P.P. (2010). Threats to the conifer species found on New Caledonia's ultramafic massifs and proposals for urgently needed measures to improve their protection. *Biodivers. Conserv.* 19, 1485–1502. <https://doi.org/10.1007/s10531-010-9780-6>.
17. Gamu, J., Le Billon, P., and Spiegel, S. (2015). Extractive industries and poverty: a review of recent findings and linkage mechanisms. *Extr. Ind. Soc.* 2, 162–176. <https://doi.org/10.1016/j.exis.2014.11.001>.
18. Mudd, G.M., and Jowitt, S.M. (2022). The new century for nickel resources, reserves, and mining: reassessing the sustainability of the devil's metal. *Econ. Geol.* 117, 1961–1983. <https://doi.org/10.5382/econ-geo.4950>.
19. Karsadi, K., and Aso, L. (2023). Multidimensional impacts of nickel mining exploitation towards the lives of the local community. *JISH* 12, 222–227. <https://doi.org/10.23887/jish.v12i2.58881>.
20. Boldy, R., Santini, T., Annandale, M., Erskine, P.D., and Sonter, L.J. (2021). Understanding the impacts of mining on ecosystem services through a systematic review. *Extr. Ind. Soc.* 8, 457–466. <https://doi.org/10.1016/j.exis.2020.12.005>.
21. Naidoo, R., Gerkey, D., Hole, D., Pfaff, A., Ellis, A.M., Golden, C.D., Herrera, D., Johnson, K., Mulligan, M., Ricketts, T.H., and Fisher, B. (2019). Evaluating the impacts of protected areas on human well-being across the developing world. *Sci. Adv.* 5, eaav3006.
22. Blackman, A. (2013). Evaluating forest conservation policies in developing countries using remote sensing data: An introduction and practical guide. *For. Policy Econ.* 34, 1–16. <https://doi.org/10.1016/j.forpol.2013.04.006>.
23. Luckeneder, S., Giljum, S., Schaffartzik, A., Maus, V., and Tost, M. (2021). Surge in global metal mining threatens vulnerable ecosystems. *Global Environ. Change* 69, 102303.
24. Lèbre, É., Stringer, M., Svobodova, K., Owen, J.R., Kemp, D., Côte, C., Arratia-Solar, A., and Valenta, R.K. (2020). The social and environmental complexities of extracting energy transition metals. *Nat. Commun.* 11, 4823.
25. Siqueira-Gay, J., Soares-Filho, B., Sanchez, L.E., Oviedo, A., and Sonter, L.J. (2020). Proposed legislation to mine Brazil's indigenous lands will threaten amazon forests and their valuable ecosystem services. *One Earth* 3, 356–362. <https://doi.org/10.1016/j.oneear.2020.08.008>.
26. Morley, J., Buchanan, G., Mitchard, E.T.A., and Keane, A. (2022). Quasi-experimental analysis of new mining developments as a driver of deforestation in Zambia. *Sci. Rep.* 12, 18252. <https://doi.org/10.1038/s41598-022-22762-4>.
27. Devenish, K., Desbureaux, S., Willcock, S., and Jones, J.P.G. (2022). On track to achieve no net loss of forest at Madagascar's biggest mine. *Nat. Sustain.* 5, 498–508. <https://doi.org/10.1038/s41893-022-00850-7>.
28. Edwards, R.B. (2016). Mining away the Preston curve. *World Dev.* 78, 22–36. <https://doi.org/10.1016/j.worlddev.2015.10.013>.
29. Zabre, H.R., Farnham, A., Diabougba, S.P., Fink, G., Divall, M.J., Winkler, M.S., and Knoblauch, A.M. (2021). Changes in household wealth in communities living in proximity to a large-scale copper mine in Zambia. *Resour. Pol.* 74, 102395. <https://doi.org/10.1016/j.resourpol.2021.102395>.
30. Santika, T., Wilson, K.A., Budiharta, S., Law, E.A., Poh, T.M., Ancrenaz, M., Struebig, M.J., and Meijaard, E. (2019). Does oil palm agriculture help alleviate poverty? A multidimensional counterfactual assessment of oil palm development in Indonesia. *World Dev.* 120, 105–117. <https://doi.org/10.1016/j.worlddev.2019.04.012>.
31. Alkire, S., Roche, J.M., Ballon, P., Foster, J., Santos, M.E., and Seth, S. (2015). *Multidimensional Poverty Measurement and Analysis* (Oxford University Press).
32. Werner, T.T., Bebbington, A., and Gregory, G. (2019). Assessing impacts of mining: Recent contributions from GIS and remote sensing. *Extr. Ind. Soc.* 6, 993–1012. <https://doi.org/10.1016/j.exis.2019.06.011>.
33. Schleicher, J., Eklund, J., D Barnes, M., Geldmann, J., Oldekop, J.A., and Jones, J.P.G. (2020). Statistical matching for conservation science. *Conserv. Biol.* 34, 538–549. <https://doi.org/10.1111/cobi.13448>.
34. Morgans, C.L., Jago, S., Andayani, N., Linkie, M., Lo, M.G.Y., Mumbunan, S., St John, F.A.V., Supriatna, J., Voigt, M., Winarni, N.L., et al. (2024). Improving well-being and reducing deforestation in Indonesia's protected areas. *Conserv. Lett.* 17, e13010. <https://doi.org/10.1111/cons.13010>.
35. BPS (2018). Village Potential Statistics (PODES) 2011, 2014, and 2018. Bureau of Statistic Indonesia. silastik.bps.go.id/v3/index.php/site/login/.
36. Chomitz, K.M. (2007). At Loggerheads? Agricultural Expansion, Poverty Reduction, and Environment in the Tropical Forests (World Bank). <http://hdl.handle.net/10986/7190>.
37. Sarwanto, D., and Prayitno, C.H. (2015). The diversity and productivity of indigenous forage in former limestone mining quarry in karst mountain of Southern Gombong, Central Java Indonesia. *Anim. Prod.* 17, 69–75.
38. Blundy, J., Mavrogenes, J., Tattitch, B., Sparks, S., and Gilmer, A. (2015). Generation of porphyry copper deposits by gas–brine reaction in volcanic arcs. *Nat. Geosci.* 8, 235–240.
39. Hayati, T. (2019). Problematika perizinan di sektor pertambangan dan kehutanan (University of Indonesia). www.academia.edu/52935711/Problematika_Perizinan_di_Sektor_Pertambangan_dan_Kehutanan.
40. Camba, A. (2021). The unintended consequences of national regulations: Large-scale-small-scale relations in Philippine and Indonesian nickel mining. *Resour. Pol.* 74, 102213. <https://doi.org/10.1016/j.resourpol.2021.102213>.
41. Geist, H.J., and Lambin, E.F. (2002). Proximate causes and underlying driving forces of tropical deforestation. *Bioscience* 52, 143–150.
42. IEA (2024). Global Critical Minerals Outlook 2024 (IEA). <https://www.iea.org/reports/global-critical-minerals-outlook-2024>.
43. Voigt, M., Supriatna, J., Deere, N.J., Kastanya, A., Mitchell, S.L., Rosa, I.M.D., Santika, T., Siregar, R., Tasirin, J.S., Widianto, A., et al. (2021). Emerging threats from deforestation and forest fragmentation in the Wallacea centre of endemism. *Environ. Res. Lett.* 16, 094048. <https://doi.org/10.1088/1748-9326/ac15cd>.
44. Supriatna, J., Shekelle, M., Fuad, H.A., Winarni, N.L., Dwiyanreni, A.A., Farid, M., Mariati, S., Margules, C., Prakoso, B., and Zakaria, Z. (2020). Deforestation on the Indonesian island of Sulawesi and the loss of primate habitat. *Glob. Ecol. Conserv.* 24, e01205. <https://doi.org/10.1016/j.gecco.2020.e01205>.
45. Struebig, M.J., Aninta, S.G., Beger, M., Bani, A., Barus, H., Brace, S., Davies, Z.G., Brauwer, M.D., Diele, K., Djakiman, C., et al. (2022). Safeguarding imperiled biodiversity and evolutionary processes in the

- Wallacea center of endemism. *Bioscience* 72, 1118–1130. <https://doi.org/10.1093/biosci/biac085>.
46. Yanuardi, Y., Vijge, M.J., and Biermann, F. (2021). Improving governance quality through global standard setting? Experiences from the Extractive Industries Transparency Initiative in Indonesia. *Extr. Ind. Soc.* 8, 100905. <https://doi.org/10.1016/j.exis.2021.100905>.
47. Ardiansyah, F., and Jotzo, F. (2013). Decentralization and avoiding deforestation: The case of Indonesia. In *Federal Reform Strategies: Lessons from Asia and Australia*, S. Howes and M. Govinda Rao, eds. (Oxford University Press), pp. 274–302. <https://doi.org/10.1093/acprof:oso/9780198092001.003.0009>.
48. OECD (2016). *OECD Due Diligence Guidance for Responsible Supply Chains of Minerals from Conflict-Affected and High-Risk Areas: Third Edition* (OECD Publishing).
49. Asare, B.K., and Darkoh, M. (2001). Socio-economic and environmental impacts of mining in Botswana: a case Study of the Selebi-Phikwe Copper-Nickel Mine. *EASSRR* 17, 1–42.
50. Liu, W., and Agusdinata, D.B. (2021). Dynamics of local impacts in low-carbon transition: Agent-based modeling of lithium mining-community-aquifer interactions in Salar de Atacama, Chile. *Extr. Ind. Soc.* 8, 100927. <https://doi.org/10.1016/j.exis.2021.100927>.
51. Syahrir, R., Wall, F., and Diallo, P. (2020). Socio-economic impacts and sustainability of mining, a case study of the historical tin mining in Singkep Island-Indonesia. *Extr. Ind. Soc.* 7, 1525–1533. <https://doi.org/10.1016/j.exis.2020.07.023>.
52. Carr-Wilson, S., Pattanayak, S.K., and Weinthal, E. (2024). Critical mineral mining in the energy transition: A systematic review of environmental, social, and governance risks and opportunities. *Energy Res. Soc. Sci.* 116, 103672.
53. Murdifi, I., Pelu, M.F.A.R., Putra, A.H.P.K., Arumbarkah, A.M., and Rahmah, A. (2019). Environmental disclosure as corporate social responsibility: Evidence from the biggest nickel mining in Indonesia. *IJEPP* 9, 115–122. <https://doi.org/10.32479/ijepp.7048>.
54. Miller, K.A., Thompson, K.F., Johnston, P., and Santillo, D. (2018). An overview of seabed mining including the current state of development, environmental impacts, and knowledge gaps. *Front. Mar. Sci.* 4, 312755.
55. Orcutt, B.N., Bradley, J.A., Brazelton, W.J., Estes, E.R., Goordial, J.M., Huber, J.A., Jones, R.M., Mahmoudi, N., Marlow, J.J., Murdock, S., and Pachiadaki, M. (2020). Impacts of deep-sea mining on microbial ecosystem services. *Limnol. Oceanogr.* 65, 1489–1510. <https://doi.org/10.1002/lno.11403>.
56. Nieminen, P., Panychev, D., Lyalyushkin, S., Komarov, G., Nikanov, A., Borisenko, M., Kinnula, V.L., and Toljamo, T. (2013). Environmental exposure as an independent risk factor of chronic bronchitis in northwest Russia. *Int. J. Circumpolar Health* 72, 19742. <https://doi.org/10.3402/ijch.v72i0.19742>.
57. Kurniawan, N.I., Lujala, P., Rye, S.A., and Vela-Almeida, D. (2022). The role of local participation in the governance of natural resource extraction. *Extr. Ind. Soc.* 9, 101029. <https://doi.org/10.1016/j.exis.2021.101029>.
58. Gustafsson, M.T. (2017). The struggles surrounding ecological and economic zoning in Peru. *Third World Q.* 38, 1146–1163. <https://doi.org/10.1080/01436597.2016.1255141>.
59. Government of the Republic of Indonesia (2009). Law number 32 of 2009 on environmental protection and management (Ministry of Environment and Forestry (KLHK)). https://pslb3.menlhk.go.id/portal/uploads/laporan/1548333565_UU_NO_32_2009.pdf.
60. Zainuddin Rela, I., Awang, A.H., Ramli, Z., Taufik, Y., Md Sum, S., and Muhammad, M. (2020). Effect of corporate social responsibility on community resilience: Empirical evidence in the nickel mining industry in Southeast Sulawesi, Indonesia. *Sustainability* 12, 1395. <https://doi.org/10.3390/su12041395>.
61. Hickel, J., Kallis, G., Jackson, T., O'Neill, D.W., Schor, J.B., Steinberger, J.K., Victor, P.A., and Ürge-Vorsatz, D. (2022). Degrowth can work—here's how science can help. *Nature* 672, 400–403.
62. Castelvocchi, D. (2021). Electric cars and batteries: how will the world produce enough? *Nature* 596, 336–339.
63. Guohua, Y., Eishkaki, A., and Xiao, X. (2021). Dynamic analysis of future nickel demand, supply, and associated materials, energy, water, and carbon emissions in China. *Resour. Pol.* 74, 102432.
64. Sanders, A., Khatarina, J., Assegaf, R., Toubmourou, T., Kurniasih, H., and Suwarso, R. (2024). The Omnibus Law on Job Creation and its potential implications for rural youth and future farming in Indonesia. *Asia Pac. Viewp.* 65, 248–262. <https://doi.org/10.1111/apv.12408>.
65. Amatullah, N., Setyadani, N.A., and Ramadhanty, S. (2020). The Extension of the Special Business Mining License (IUPK) under The Law No. 3 of 2020 of the Coal and Mineral Mining: Pro or Cons? *Legal Brief* 10, 39–49.
66. Samadhi, T.N., and Mumbunan, S. (2015). *Tambang, hutan, dan kebun: Tata kelola perizinan dan penerimaan negara di sektor berbasis lahan*, 2nd Edition (IPB Press).
67. Agrawal, S., Bebbington, A.J., Imhof, A., Jebing, M., Royo, N., Sauls, L.A., Sulaiman, R., Toubmourou, T., and Wicaksono, A. (2018). Impacts of extractive industry and infrastructure on forests. <http://www.climateandlandusealliance.org/wp-content/uploads/2018/12/Indonesia-Impacts-of-EII-on-Forests-1.pdf>.
68. Camfield, L. (2006). The why and how of understanding 'subjective' well-being: Exploratory work by the WeD group in four developing countries. eprints.uea.ac.uk/id/eprint/24946/1/Camfield_2006_%28the_why_and_how_of_understanding_SWB%29.pdf.
69. Leuenberger, A., Winkler, M.S., Cambaco, O., Cossa, H., Kihwele, F., Lyatuu, I., Zabré, H.R., Farnham, A., Macete, E., and Mungambe, K. (2021). Health impacts of industrial mining on surrounding communities: Local perspectives from three sub-Saharan African countries. *PLoS One* 16, e0252433.
70. Austin, K.G., Schwantes, A., Gu, Y., and Kasibhatla, P.S. (2019). What causes deforestation in Indonesia? *Environ. Res. Lett.* 14, 024007. <https://doi.org/10.1088/1748-9326/aaf6db>.
71. Devi, B., and Prayogo, D. (2013). Mining and Development in Indonesia: An Overview of the Regulatory Framework and Policies (International Mining for Development Centre). <https://delvedatabase.org/uploads/resources/Mining-and-Development-in-Indonesia.pdf>.
72. Resosudarmo, I.A.P., Oka, N.P., Mardiah, S., and Utomo, N.A. (2014). Governing Fragile Ecologies: A Perspective on Forest and Land-Based Development in the Regions. In *Regional Dynamics in a Decentralised Indonesia*, H. Hill, ed. (Institute of Southeast Asian Studies–Yusof Ishak Institute), pp. 260–284. <https://www.cambridge.org/core/books/abs/regional-dynamics-in-a-decentralized-indonesia/governing-fragile-ecologies-a-perspective-on-forest-and-landbased-development-in-the-regions/1B3BA64D56EF24E9DEB337E35C2BE3A8>.
73. KLHK (2018). The state of Indonesia's forests 2018 (Ministry of Environment and Forestry, Republic of Indonesia). <https://ppid.menlhk.go.id/infografis/2985/buku-the-state-of-indonesias-forests-2018-dapat-download-disini>.
74. BPS (2011). *Statistik Potensi Desa Indonesia 2011* (Badan Pusat Statistik). <https://www.bps.go.id/id/publication/2011/11/22/f0f9b7483c34df5e5ae96be2/statistik-potensi-desa-indonesia-2011.html>.
75. Lee, J.S.H., Miteva, D.A., Carlson, K.M., Heilmayr, R., and Saif, O. (2020). Does oil palm certification create trade-offs between environment and development in Indonesia? *Environ. Res. Lett.* 15, 124064. <https://doi.org/10.1088/1748-9326/abc279>.
76. Ferraro, P.J., Hanauer, M.M., Miteva, D.A., Nelson, J.L., Pattanayak, S.K., Nolte, C., and Sims, K.R.E. (2015). Estimating the impacts of conservation on ecosystem services and poverty by integrating modeling and evaluation. *Proc. Natl. Acad. Sci. USA* 112, 7420–7425. <https://doi.org/10.1073/pnas.1406487112>.

77. Sen, A. (1993). Capability and well-being. In *The Quality of Life*, M. Nussbaum and A. Sen, eds. (Clarendon Press), pp. 30–53.
78. BPS (2018). Indeks Pembangunan Desa 2018 (Badan Pusat Statistik). <https://www.bps.go.id/publication/2019/05/09/4edae4bd6c18d24b1b4273fe/indeks-pembangunan-desa-2018.html>.
79. S&P Global Market Intelligence (2020). SNL Metals and Mining S&P Global. [www.marketplace.spglobal.com/en/datasets/snl-metals-mining-\(19\)#sample-data](http://www.marketplace.spglobal.com/en/datasets/snl-metals-mining-(19)#sample-data).
80. Maus, V., Giljum, S., Gutschhofer, J., da Silva, D.M., Probst, M., Gass, S.L.B., Luckeneder, S., Lieber, M., and McCallum, I. (2020). A global-scale data set of mining areas. *Sci. Data* 7, 289.
81. Maus, V., and Werner, T.T. (2024). Impacts for half of the world's mining areas are undocumented. *Nature* 625, 26–29.
82. Hamidi, J. (2015). Management of mining in Indonesia: Decentralization and corruption eradication. *JL Pol'y & Globalization* 44, 80–101. www.iiste.org/Journals/index.php/JLPG/article/viewFile/27744/28468.
83. Manalo, P. (2023). Discovery to production averages 15.7 years for 127 mines. <https://www.spglobal.com/marketintelligence/en/news-insights/research/discovery-to-production-averages-15-7-years-for-127-mines>.
84. Nguyen, T.L., Collins, G.S., Spence, J., Daurès, J.P., Devereaux, P.J., Landais, P., and Le Manach, Y. (2017). Double-adjustment in propensity score matching analysis: choosing a threshold for considering residual imbalance. *BMC Med. Res. Methodol.* 17, 78. <https://doi.org/10.1186/s12874-017-0338-0>.
85. Ribas, L.G.S., Pressey, R.L., and Bini, L.M. (2021). Estimating counterfactuals for evaluation of ecological and conservation impact: an introduction to matching methods. *Biol. Rev. Camb. Phil. Soc.* 96, 1186–1204. <https://doi.org/10.1111/brv.12697>.
86. Diamond, A., and Sekhon, J.S. (2013). Genetic matching for estimating causal effects: A general multivariate matching method for achieving balance in observational studies. *Rev. Econ. Stat.* 95, 932–945. https://doi.org/10.1162/REST_a_00318.
87. Sekhon, J.S. (2011). Multivariate and propensity score matching software with automated balance optimization: The matching package for R. *J. Stat. Software* 42, 1–52. <https://doi.org/10.18637/jss.v042.i07>.
88. Abadie, A., Athey, S., Imbens, G.W., and Wooldridge, J.M. (2023). When should you adjust standard errors for clustering? *Q. J. Econ.* 138, 1–35. <https://doi.org/10.1093/qje/qjac038>.