



Kent Academic Repository

Schrijvers, Karen, Cosma, Alina, Potrebny, Thomas, Thorsteinsson, Einar, Catunda, Carolina, Reiss, Franziska, Hulbert, Sabina, Kostičová, Michaela, Melkumova, Marina and Bersia, Michela (2024) *Three decades of adolescent health: Unveiling global trends across 41 countries in psychological and somatic complaints (1994–2022)*. *International Journal of Public Health*, 69 . ISSN 1661-8556.

Downloaded from

<https://kar.kent.ac.uk/108012/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.3389/ijph.2024.1607774>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).



Three Decades of Adolescent Health: Unveiling Global Trends Across 41 Countries in Psychological and Somatic Complaints (1994–2022)

Karen Schrijvers^{1*}, Alina Cosma^{2,3}, Thomas Potrebný⁴, Einar Thorsteinsson⁵, Carolina Catunda⁶, Franziska Reiss⁷, Sabina Hulbert⁸, Michaela Kostičová⁹, Marina Melkumova¹⁰, Michela Bersia¹¹, Helena Jeriček Klanšček¹², Tania Gaspar¹³ and Maxim Dierckens¹

¹Department of Public Health and Primary Care, Ghent University, Ghent, Belgium, ²School of Psychology, Trinity College Dublin, Dublin, Ireland, ³Olomouc University Social Health Institute, Palacky University Olomouc, Olomouc, Czechia, ⁴Department of Health and Functioning, Western Norway University of Applied Sciences, Bergen, Norway, ⁵School of Psychology, University of New England, Armidale, NSW, Australia, ⁶Department of Social Sciences, University of Luxembourg, Esch-sur-Alzette, Luxembourg, ⁷Department of Child and Adolescent Psychiatry, Psychotherapy, and Psychosomatics, University Medical Center Hamburg-Eppendorf, Hamburg, Germany, ⁸Centre for Health Services Studies (CHSS), University of Kent, Canterbury, United Kingdom, ⁹Institute of Social Medicine and Medical Ethics, Faculty of Medicine, Comenius University in Bratislava, Bratislava, Slovakia, ¹⁰Arabkir Medical Centre, Institute of Child and Adolescent Health, Yerevan, Armenia, ¹¹Department of Public Health and Pediatrics, University of Torino, Torino, Italy, ¹²National Institute of Public Health, Ljubljana, Slovenia, ¹³School of Psychology and Life Sciences, SPIC/Hei-Lab/Lusófona University, Lisbon, Portugal

OPEN ACCESS

Edited by:

Caroline Residori,
University of Luxembourg,
Luxembourg

Reviewed by:

Aleksandar Višnjić,
University of Niš, Serbia
Alessio Rebecchi,
University of Luxembourg,
Luxembourg

*Correspondence

Karen Schrijvers,
✉ karen.schrijvers@ugent.be

This Original Article is part of the IJPH Special Issue "Trends in Adolescent Health and Wellbeing - 40 Years of the HBSC Study"

Received: 15 July 2024

Accepted: 15 November 2024

Published: 02 December 2024

Citation:

Schrijvers K, Cosma A, Potrebný T, Thorsteinsson E, Catunda C, Reiss F, Hulbert S, Kostičová M, Melkumova M, Bersia M, Klanšček HJ, Gaspar T and Dierckens M (2024) Three Decades of Adolescent Health: Unveiling Global Trends Across 41 Countries in Psychological and Somatic Complaints (1994–2022). *Int J Public Health* 69:1607774. doi: 10.3389/ijph.2024.1607774

Objectives: This study examined (non-)monotonic time trends in psychological and somatic complaints among adolescents, along with gender differences.

Methods: Repeated cross-sectional Health Behaviour in School-aged Children (HBSC) data from 1994 to 2022 covering 15-year-old adolescents from 41 countries (N = 470,797) were analysed. Three polynomial logistic regression models (linear, quadratic, cubic) were tested for best fit, including separate analyses by gender and health complaints dimension.

Results: Time trend patterns varied by gender and health complaints dimension. Increases were found in 82.3% of cases (linear 25%, quadratic U-shaped 28.7%, cubic 28.7%), while 14% showed no clear trend, and 3.7% decreased. Boys typically showed linear increases or no clear trend over time, whereas girls generally showed cubic or U-shaped trends. Psychological complaints often displayed U-shaped or cubic patterns, whereas somatic complaints mostly showed linear increases.

Conclusion: Psychological and somatic complaints demonstrated diverse time trend patterns across countries, with non-monotonic patterns (U-shaped and cubic) frequently observed alongside linear increases. These findings highlight the complexity of changes within countries over three decades, suggesting that linear modelling may not effectively capture this heterogeneity.

Keywords: adolescence, mental health, gender differences, cross-national, HBSC

INTRODUCTION

Good health and wellbeing underpin crucial developmental tasks associated with adolescent years [1], of which mental health is a fundamental component. However, more than one-third of adolescents often experience a variety of psychological and somatic complaints [2, 3]. These complaints serve as important indicators of difficulties in mental health and wellbeing and can be stress reactions to psychosocial tensions in adolescents' lives without a clear organic cause [4, 5]. Moreover, psychological and somatic complaints can negatively affect adolescents' general health, functioning, and wellbeing, potentially leading to mental health issues later in life [6] that significantly contribute to the global disease burden [7]. Recent evidence has indicated an increase in psychological and somatic complaints over time [8], possibly influenced by national-level events and cross-national crises like the 2008 economic crisis and the COVID-19 pandemic as well as increased mental health awareness and availability of mental health services. Long-term time trend analyses are essential to fully understand these changes and guide efforts to monitor and improve the overall wellbeing of adolescents.

The four-yearly cross-national Health Behaviour in School-aged Children (HBSC) study provides a unique opportunity to monitor time trends in adolescent health and wellbeing using the HBSC Multiple Health Complaints scale. Initially introduced in 1986, the scale was refined in 1994 to measure eight items, encompassing somatic (headache, backache, stomachache, and dizziness) and psychological (feeling nervous, feeling low, feeling irritable, and difficulties in getting to sleep) complaints. Since then, the scale has remained unchanged [9]. Somatic and psychological complaints tend to be comorbid [10–13] with some researchers treating them as one factor called psychosomatic complaints [13, 14] and others as two different factors (i.e., psychological and somatic complaints) [11, 12, 15, 16]. Both approaches have solid validation evidence and have been used previously in adolescent health research.

Studies utilizing the one-factor approach reported similar trend patterns over time across different national contexts. For example, changes over time in psychosomatic complaints appeared to be relatively stable in the 1990s and early 2000s in countries such as Scotland, England, and the Netherlands [17–19]. However, there has been an increase in the proportion of adolescents reporting two or more psychosomatic complaints from 2010 onwards [17–20]. These results have been further confirmed through pooling of cross-national data [8].

Studies employing the two-factor approach revealed diverse changes over time in psychological and somatic complaints across different national contexts. These studies showed a relatively stable trend for *psychological complaints* during the 1990s and early 2000s, followed by an increase with some differences between countries following the early 2000s. For instance, Switzerland noted a stable trend from 1994 to 2006 [15], while in Norway, an upward trend during the same period was identified, followed by a decrease in 2014 [16]. Recent studies in Czechia (2002–2018), Italy (2010–2018) and Canada

(2022–2018, particularly among girls) reported an increase in psychological complaints [20–22]. A recent cross-national study showed a stable trend in psychological complaints between 2002 and 2010 followed by an increase towards 2018 [23], another study using pooled data (2002–2018) reported an increase in psychological complaints over time across the majority of countries [24]. In terms of *somatic complaints*, an increasing trend from the 1990s onwards was noted until the early 2000s in Switzerland, Norway, and Czechia [15, 16, 21], followed by a decrease in Norway (2010–2014) and Czechia (2010–2018) [16, 21]. Conversely, Italy and Canada showed an increase over this same period [20, 22]. An increasing trend in somatic complaints between 2002 and 2018 has also been confirmed in one cross-national study thus far [23]. Given these diverging trends for both psychological and somatic complaints and the lack of studies spanning a long time frame, it is essential to consolidate results in one study and explore both psychological and somatic complaints (i.e., two-factor approach) separately over an extended timeframe and across countries to further elucidate differences in time trends.

Previous studies have mostly employed a monotonic approach and examined linear changes over time in psychosomatic complaints. However, this may not be sufficient to capture the full complexity of these trends [25]. The presence of non-monotonic time trends, such as quadratic (U-shaped or inverted U-shaped) or cubic trends has rarely been examined, except for one study [25]. These non-monotonic time trend patterns are often not considered, although they are important, especially when examining trends over a longer period of time where multiple fluctuations may have occurred. Our study aims to improve the understanding of these developments over a longer period of time and update them with recent representative cross-national data by employing the two-factor approach.

Although divergent trends over time have been reported, some results were also consistent across studies in terms of prevalence rate and gender. Trend studies employing a two-factor approach have generally reported a lower prevalence of somatic compared to psychological complaints [15, 16, 20, 21, 26]. Moreover, a higher prevalence of health complaints, using either the one or two-factor approach, were noted in girls compared to boys [15–21, 26, 27]. Further investigation of the differences in prevalence rates according to gender and health complaints dimension within trend studies considering different trend patterns could provide a more nuanced understanding of these changes over time.

This Study

The aforementioned studies highlighted the necessity for further investigation into diverse patterns of change over time, beyond the commonly employed linear approach. Therefore, to address these gaps the current study aims to inform policy and signpost key target groups for tailored interventions such as addressing the specific needs of boys and girls in coping with different dimensions of health complaints over time. Moreover, using representative data from adolescents across 41 countries, this study aims to provide insights into potential cross-national

disparities, emphasizing the importance of considering diverse cultural and contextual factors alongside gender differences and differences by dimension of health complaints. To gain a deeper understanding of these long-term time trends, the present study spans a period of almost three decades in 41 countries. This research tests which pattern of change over time (i.e., linear, quadratic, cubic) most accurately describes the fluctuations observed over the past three decades in psychological and somatic complaints among adolescents for each country separately. Additionally, it explores whether these different patterns of change over time differ between boys and girls.

METHODS

Study Design and Sample

Data from the cross-national HBSC study on health and wellbeing in 11-, 13- and 15-year-old adolescents is collected every 4 years. A representative sample is surveyed within the school context in each region/country following the international research protocol [9]. The present study employed data from eight survey cycles, (i.e., 1994, 1998, 2002, 2006, 2010, 2014, 2018, and 2022) spanning three decades. A total of 41 countries were included in the sample, representing 79% of the 52 countries in the dataset. Countries with at least four data points, even if non-consecutive, were included to meet the minimum requirement for testing cubic trends. Systematic previous findings documented that the 15-year-olds report the highest prevalence, the largest gender differences and also the widest cross-national variations [8, 20, 22, 24], therefore this study included only participants from this age group. Our final sample included 470,797 15-year-old adolescents. Sample sizes by country are presented in **Supplementary Table S1**.

Measures

Adolescents' psychological and somatic complaints were measured using the HBSC Multiple Health Complaints scale [9]. Adolescents were asked to indicate how often in the past 6 months they had each of the following eight symptoms: feeling low, irritability or bad temper, feeling nervous, difficulties in getting to sleep, headache, backache, stomachache, and feeling dizzy. Response options ranged from 1 "about every day" to 5 "rarely or never." A two-dimensional approach was followed with the first four items as a measure for psychological complaints and the last four items indicating somatic complaints [11, 15, 16]. Both variables were dichotomized based on the cut-off of experiencing two or more complaints more than once a week [11, 15, 16, 20]. This cut-off point was used as it was suitable for both psychological and somatic complaints, where a more conservative approach of 3 or more complaints would result in low prevalence rates for somatic complaints and a less conservative approach of 1 or more complaints would result in high prevalence rates for psychological complaints (**Supplementary Figure S1**). The measure has shown good internal reliability, construct validity [12] and configural and metric cross-national invariance [13]. Gender was measured

by asking adolescents whether they are a boy or a girl. Age was determined by asking for the month and year of birth.

Statistical Analyses


Prevalence rates of psychological and somatic complaints were calculated by country, survey cycle, and gender. To determine the most suitable trend pattern over time within each country, logistic regression analyses were conducted for boys and girls, and for each health complaints dimension, resulting in 164 separate cases ($k = 164$; 41 countries \times 2 genders \times 2 dimensions of health complaints). The survey cycle variable (year of data collection) was centred, and orthogonal polynomials were used in the quadratic and cubic models. Three models were tested – including cubic, quadratic, and linear functions – and polynomial regression models were fitted using "glmnet" package in R v4.1-1 (1) [28]. The optimal trend function was determined based on the degree of change and the model significance, employing a conservative p-value criterion ($p < .001$) because of the large sample size [29]. Subsequently, two sensitivity analyses were conducted. First, one on the pooled dataset to evaluate the influence of the selected cut-off point (i.e., at least one, two or three complaints more than once a week) on the results of the trend patterns. Besides, also a sensitivity analysis to assess the COVID-19 impact on the overall trend patterns in adolescents' psychological and somatic symptoms. This involved rerunning the respective models without the 2022 data and comparing them to the full models with the 2022 data. Sensitivity testing could not be performed for Russia, Ukraine, and the United States due to the unavailability of 2022 data.

RESULTS

Descriptives

Figure 1 presents the observed prevalence over time by country, while **Supplementary Table S2** provides detailed information per survey cycle. Additionally, **Figure 2** illustrates the observed gap between the highest and lowest prevalence observed over the study period by country, highlighting a large range of differences over time between countries. Overall, the highest prevalence rates in somatic and psychological complaints were observed in 2022 across most countries for both boys (32 countries) and girls (36 countries). Conversely, there were wide cross-national variations in relation to the lowest prevalence for these indicators over time. More specifically, for psychological complaints, the lowest prevalence within each country could be observed at various points between 1994 and 2014. For somatic complaints within countries the lowest prevalence was found between 1994 and 2018. However, in 1994, data was collected from 20 countries with the prevalence for somatic complaints being the lowest for girls in 17 countries and boys in 13 countries as compared with data collected after 1994.

Across countries, the prevalence rates of psychological complaints were generally higher, ranging from 4.9% in boys from Austria in 1998 to 80.2% in girls from Italy in 2022. In contrast, somatic complaints ranged from 1.7% in boys from Latvia in 1994 to 16.1% in girls from Italy in 2022. Overall, for




both psychological and somatic complaints in Italy and Israel high prevalence rates were observed. Specifically for psychological complaints, Greece showed particularly higher prevalence rates. Furthermore, across countries the observed gap over time was generally larger in psychological complaints for both boys and girls, suggesting greater fluctuations over time in psychological complaints compared to somatic complaints.

Additionally, girls reported higher levels of psychological and somatic complaints compared to boys across all countries.

Trend Patterns

Notable cross-national differences in trend patterns for both dimensions of health complaints and genders were found (see **Tables 1, 2**). Within most countries and for both dimensions of



health complaints and genders, 82.3% ($k = 135$) of the observed trend patterns indicated an increase in complaints over the survey period. More specifically, three increasing trend patterns emerged: linear (25%; $k = 41$; i.e., steady increase since the start of the survey), quadratic U-shaped (28.7%; $k = 47$; i.e., an initial decrease followed by an increase in more recent years), and cubic (28.7%; $k = 47$; i.e., multiple fluctuations of increases and decreases over the years with an increase towards the latest datapoint). For a small number of cases (3.7%, $k = 6$) a decrease in psychological and somatic symptoms for boys and girls was found. More specifically, two trend patterns emerged. A linear decrease was observed in 1.8% of cases ($k = 3$), suggesting an overall steady decrease over the years. Additionally, in 1.8% of

cases ($k = 3$), an inverted U-shaped trend was identified, indicating an initial increase followed by a subsequent decrease. Finally, in 14% of the cases ($k = 23$) no clear trend pattern was observed, suggesting stability in complaints over the period studied (see also **Table 3** for an overview).

The findings also indicated several differences in the pattern of change for boys and girls. In general, linear increases (32.9%, $k = 27$) over time or no clear trend patterns (23.2%, $k = 19$) were more common among boys, whereas cubic (41.5%, $k = 34$) or U-shaped (35.4%, $k = 29$) trends were more common among girls. Therefore, there were greater fluctuations in trends over time among girls compared to boys within the countries included in this study. Furthermore, the results revealed notable differences




FIGURE 2 | Dumbbell plot showing highest and lowest prevalence rate of two or more psychological and somatic complaints by gender and country across all countries (Health Behaviour in School-aged Children study, 1994–2022 for 41 countries). Notes. The difference between the highest and lowest prevalence is displayed in grey between the dumbbells.

in trend patterns by *health complaints dimensions*. The most common patterns observed for psychological complaints were U-shaped (41.5%, $k = 34$) and cubic (34.1%, $k = 28$) time trends, while linear increases (35.4%, $k = 29$) were the predominant pattern for somatic complaints, followed by both cubic (23.2%, $k = 19$) and stable (20.7%, $k = 17$) time trends. Specifically, for *boys* non-monotonic trends were observed in *psychological symptoms* with U-shaped trends seen in 15 countries and cubic trends in 10 countries. In 8 countries a linear increase was found. Conversely, *somatic complaints* among *boys* predominantly exhibited linear increases in 19 countries, with U-shaped and cubic trends being less prevalent, observed in 2 and 3 countries respectively. In the case of *girls*, U-shaped trends were most common in *psychological complaints* across 18 countries, while cubic trends were prominent in 16 countries, with linear increases noted in 4 countries. Similarly, for *somatic complaints among girls*, cubic trends prevailed in 16 countries, followed by

U-shaped trends in 11 countries, with linear increases observed in 10 countries.

Sensitivity Testing

Sensitivity checks on the cut-off point of at least one, two or three complaints more than once a week on the pooled dataset showed similar conclusions for each selected cut-off for both boys and girls and both health complaints dimensions (**Supplementary Table S3**).

As COVID-19 may have had a considerable impact on the latest data [30], sensitivity testing was performed without the 2022 data (**Supplementary Tables S4, S5**). **Table 3** illustrates an overview of the findings from the trend pattern testing, including 2022 data, and the sensitivity testing. After excluding the 2022 data, the initially observed trend patterns changed in several countries, likely showing the impact of COVID-19 on the overall time trends.

TABLE 1 | Time trend pattern for psychological complaints by country and gender (Health Behaviour in School-aged Children study, 1994–2022 for 41 countries).

	Boys psychological complaints			Girls psychological complaints			Trend pattern	
	Linear trend	Quadratic trend	Cubic trend	Linear trend	Quadratic trend	Cubic trend	Boys	Girls
Armenia	0.002 (0.003)	0.471 (0.451)	-0.195 (0.451)	0.010 (0.002)***	2.311 (0.494)***	-0.751 (0.494)	—	QU
Austria	0.005 (0.000)***	2.242 (0.319)***	0.529 (0.319)	0.013 (0.001)***	5.032 (0.409)***	1.306 (0.409)**	QU	QU
Belgium - Flanders	0.003 (0.000)***	0.803 (0.372)*	2.120.371)***	0.011 (0.001)***	4.273 (0.437)***	2.453 (0.436)***	C	C
Belgium - Wallonia	0.002 (0.001)***	2.990 (0.436)***	0.006 (0.436)	0.008 (0.001)***	3.860 (0.486)***	1.711 (0.486)***	QU	C
Bulgaria	0.007 (0.001)***	-1.231 (0.460)**	-2.541 (0.458)***	0.009 (0.002)***	0.513 (0.497)	0.254 (0.498)	C	LI
Canada	0.002 (0.000)***	2.950 (0.399)***	0.311 (0.399)	0.011 (0.000)***	5.240 (0.471)***	0.692 (0.471)	QU	QU
Croatia	-0.000 (0.001)	0.998 (0.386)*	0.440 (0.386)	0.006 (0.001)***	3.646 (0.479)***	0.437 (0.479)	—	QU
Czechia	0.004 (0.001)***	0.370 (0.419)	0.770 (0.419)	0.012 (0.001)***	2.855 (0.481)***	1.566 (0.481)**	LI	QU
Denmark	0.003 (0.000)***	0.326 (0.313)	0.273 (0.313)	0.008 (0.001)***	2.504 (0.419)***	1.470 (0.419)***	LI	C
England	0.006 (0.001)***	2.318 (0.410)***	0.916 (0.410)*	0.015 (0.001)***	3.503 (0.464)***	1.970 (0.463)***	QU	C
Estonia	0.005 (0.001)***	2.099 (0.411)***	1.398 (0.410)***	0.0090.001)***	4.390 (0.477)***	3.140 (0.475)***	C	C
Finland	0.002 (0.001)***	0.863 (0.357)**	-0.095 (0.357)	0.010 (0.001)***	2.830 (0.439)***	0.660 (0.439)	LI	QU
France	0.003 (0.001)***	2.106 (0.419)***	-0.635 (0.418)	0.007 (0.001)***	4.284 (0.486)***	1.022 (0.486)*	QU	QU
Germany	0.004 (0.000)***	2.199 (0.320)***	1.104 (0.319)***	0.012 (0.001)***	6.287 (0.412)***	2.738 (0.411)***	C	C
Greece	0.001 (0.001)	3.299 (0.473)***	2.001 (0.473)***	0.007 (0.001)***	4.636 (0.488)***	2.966 (0.486)***	C	C
Greenland	0.004 (0.001)***	0.350 (0.365)	-0.213 (0.366)	0.010 (0.001)***	0.236 (0.457)	0.631 (0.457)	LI	LI
Hungary	0.003 (0.001)***	2.337 (0.433)***	0.929 (0.432)*	0.008 (0.001)***	6.123 (0.477)***	2.486 (0.476)***	QU	C
Iceland	0.002 (0.001)*	1.593 (0.400)***	-0.464 (0.400)	0.014 (0.001)***	3.206 (0.479)***	0.510 (0.479)	QU	QU
Ireland	0.003 (0.001)***	1.997 (0.405)***	0.212 (0.405)	0.013 (0.001)***	2.655 (0.467)***	1.072 (0.467)*	QU	QU
Israel	-0.001 (0.001)	1.161 (0.485)**	1.161 (0.485)*	0.005 (0.001)***	3.078 (0.497)***	3.994 (0.495)***	—	C
Italy	0.006 (0.001)***	1.256 (0.469)**	0.302 (0.469)	0.014 (0.001)***	2.857 (0.481)***	0.766 (0.481)	LI	QU
Latvia	0.005 (0.001)***	1.675 (0.403)***	0.938 (0.403)*	0.010 (0.001)***	3.837 (0.482)***	2.360 (0.481)***	QU	C
Lithuania	0.002 (0.001)***	1.485 (0.410)***	1.550 (0.410)***	0.005 (0.001)***	2.998 (0.492)***	4.635 (0.489)***	C	C
Luxembourg	0.005 (0.001)***	0.065 (0.419)	0.156 (0.419)	0.010 (0.001)***	2.436 (0.488)***	0.884 (0.487)	LI	QU
North-Macedonia	0.005 (0.001)***	1.274 (0.421)**	-0.597 (0.421)	0.008 (0.001)***	2.505 (0.490)***	-0.976 (0.490)*	LI	QU
Malta	0.004 (0.002)**	-0.433 (0.460)	1.284 (0.459)**	0.013 (0.002)***	0.768 (0.490)	1.514 (0.489)**	—	LI
Netherlands	0.005 (0.001)***	1.402 (0.326)***	0.181 (0.326)	0.011 (0.001)***	2.221 (0.433)***	0.930 (0.432)*	QU	QU
Norway	0.004 (0.001)***	1.005 (0.336)**	1.253 (0.336)***	0.009 (0.001)***	0.743 (0.431)	1.260 (0.430)**	C	LI
Poland	0.007 (0.001)***	0.859 (0.437)*	1.655 (0.437)***	0.010 (0.001)***	3.432 (0.485)***	2.613 (0.485)***	C	C
Portugal	0.004 (0.001)***	2.708 (0.364)***	0.001 (0.364)	0.011 (0.001)***	5.750 (0.462)***	0.406 (0.462)	QU	QU
Romania	0.006 (0.001)***	2.698 (0.449)***	0.457 (0.449)	0.005 (0.001)***	3.586 (0.497)***	-0.317 (0.497)	QU	QU
Russia	0.000 (0.001)	-0.014 (0.394)	0.289 (0.394)	0.000 (0.001)	1.455 (0.470)**	-0.862 (0.470)	—	—
Scotland	0.006 (0.001)***	2.487 (0.392)***	-0.238 (0.392)	0.012 (0.001)***	4.142 (0.460)***	-0.323 (0.460)	QU	QU
Slovakia	0.004 (0.001)***	0.187 (0.432)	1.815 (0.431)***	0.008 (0.001)***	3.555 (0.480)***	2.722 (0.479)***	C	C
Slovenia	0.005 (0.001)***	1.328 (0.357)***	-0.719 (0.356)*	0.018 (0.001)***	2.746 (0.460)***	-3.616 (0.457)***	QU	C
Spain	-0.003 (0.001)***	0.759 (0.377)*	0.233 (0.377)	0.001 (0.001)	3.419 (0.475)***	0.392 (0.475)	LD	QU
Sweden	0.005 (0.001)***	1.982 (0.411)***	1.562 (0.410)***	0.011 (0.001)***	2.329 (0.485)***	2.273 (0.484)***	C	C
Switzerland	0.003 (0.001)***	0.814 (0.348)*	0.791 (0.348)*	0.009 (0.001)***	4.959 (0.455)***	1.543 (0.455)***	LI	C
Ukraine	-0.001 (0.001)	1.726 (0.408)***	1.235 (0.408)**	-0.003 (0.001)*	2.033 (0.494)***	1.937 (0.493)***	QU	C
USA	-0.008 (0.002)***	0.019 (0.407)	-1.045 (0.407)*	-0.003 (0.002)	-0.150 (0.472)	-2.103 (0.471)***	LD	C
Wales	0.007 (0.000)***	3.682 (0.432)***	1.192 (0.432)**	0.014 (0.001)***	4.918 (0.479)***	0.632 (0.480)	QU	QU

Notes. Estimates (Standard error); $P < 0.001 = ***$; $P < 0.01 = **$; $P < 0.05 = *$; LI, linear increase; LD, linear decrease; QU, quadratic U-shaped; QIU, quadratic inverted U-shaped; C, cubic.

For *psychological complaints* among the 38 countries included in the sensitivity testing, the trend pattern changed in 26 countries for boys and in 18 countries for girls. For *boys*, the most common impact of 2022 data was that several countries showed no clear trend pattern until 2018, whereas by including the 2022 data this changed to a linear increase ($k = 6$), a U-shaped trend ($k = 6$), or a cubic trend ($k = 5$). For *girls*, the most common impact of the 2022 data encountered, was that the existing linear increase until 2018, became a more complex U-shaped ($k = 5$) or cubic trend ($k = 5$).

Similarly, concerning *somatic complaints*, sensitivity analyses showed that the trend pattern changed in 20 countries for boys and in 28 countries among girls when the 2022 data were included, as opposed to when it was excluded. In the case of somatic complaints in *boys*, the

most notable impact of including the 2022 data was that linear increases emerged ($k = 9$), whereas without the 2022 data, often no clear trend was observed. Among *girls*, similarly to psychological complaints, the most common change contains a linear increase observed until 2018, whereas when including the 2022 data, a cubic trend ($k = 9$) was found. In addition, five countries that showed no clear trend pattern until 2018, shifted to a cubic trend when 2022 data were included.

DISCUSSION

Our study – spanning almost three decades (from 1994 to 2022) and covering 41 countries – aimed to identify the most accurate

TABLE 2 | Time trend pattern for somatic complaints by country and gender (Health Behaviour in School-aged Children study, 1994–2022 for 41 countries).

	Boys somatic complaints			Girls somatic complaints			Trend pattern	
	Linear trend	Quadratic trend	Cubic trend	Linear trend	Quadratic trend	Cubic trend	Boys	Girls
Armenia	0.000 (0.002)	0.224 (0.305)	0.464 (0.305)	0.001 (0.002)	1.671 (0.363)***	0.393 (0.363)	—	QU
Austria	0.001 (0.000)***	0.494 (0.241)*	-0.214 (0.241)	0.006 (0.000)***	2.488 (0.375)***	1.098 (0.375)**	LI	QU
Belgium - Flanders	0.002 (0.000)***	-0.362 (0.271)	0.328 (0.271)	0.007 (0.000)***	1.354 (0.386)***	1.140 (0.386)**	LI	QU
Belgium - Wallonia	0.002 (0.000)**	-0.517 (0.311)	0.800 (0.311)**	0.005 (0.001)***	0.802 (0.430)	1.205 (0.430)**	—	LI
Bulgaria	0.005 (0.001)***	0.634 (0.349)	-0.993 (0.349)**	0.007 (0.001)***	1.247 (0.419)**	1.407 (0.419)***	LI	C
Canada	0.001 (0.000)	0.646 (0.304)*	0.386 (0.304)	0.005 (0.000)***	3.376 (0.432)***	0.847 (0.432)	—	QU
Croatia	0.000 (0.001)	0.116 (0.234)	0.003 (0.234)	0.003 (0.001)***	1.339 (0.381)***	0.364 (0.381)	—	QU
Czechia	0.000 (0.000)	0.224 (0.243)	0.640 (0.243)**	0.004 (0.001)***	0.544 (0.370)	1.453 (0.370)***	—	C
Denmark	0.002 (0.000)***	0.413 (0.225)	0.210 (0.225)	0.006 (0.000)***	1.443 (0.333)***	1.362 (0.332)***	LI	C
England	0.002 (0.001)***	1.027 (0.293)***	1.075 (0.293)***	0.009 (0.001)***	1.458(0.406)***	1.915 (0.405)***	C	C
Estonia	0.003 (0.000)***	0.991(0.300)**	1.039 (0.299)***	0.007 (0.001)***	2.279 (0.408)***	3.144 (0.406)***	C	C
Finland	0.001 (0.000)**	-0.188 (0.250)	-0.115 (0.250)	0.005 (0.001)***	1.090 (0.378)**	1.471 (0.378)***	—	C
France	0.002 (0.000)***	0.063 (0.281)	-0.240 (0.281)	0.005 (0.001)***	0.420 (0.403)	1.527 (0.403)***	LI	C
Germany	0.002 (0.000)***	0.391 (0.238)	0.342 (0.238)	0.006 (0.001)***	2.359 (0.380)***	0.614 (0.380)	LI	QU
Greece	0.002 (0.000)***	0.989 (0.287)***	0.352 (0.287)	0.008 (0.001)***	3.825 (0.416)***	2.375 (0.415)***	QU	C
Greenland	0.004 (0.001)***	-0.051 (0.297)	-0.508 (0.296)	0.007 (0.001)***	-0.312 (0.375)	-0.260 (0.375)	LI	LI
Hungary	0.003 (0.000)***	0.073 (0.294)	0.091 (0.294)	0.008 (0.001)***	1.751 (0.422)***	2.104 (0.422)***	LI	C
Iceland	0.000 (0.001)	0.826 (0.328)*	-0.215 (0.328)	0.008 (0.001)***	3.267 (0.446)***	0.346 (0.446)	—	QU
Ireland	0.003 (0.001)***	0.630 (0.295)*	0.537 (0.295)	0.010 (0.001)***	0.787 (0.398)*	1.016 (0.398)*	LI	LI
Israel	3.835 (0.000)	-1.453 (0.384)***	-0.131 (0.384)	0.004 (0.001)***	1.582 (0.469)***	2.291 (0.468)***	QIU	C
Italy	0.002 (0.001)*	0.216 (0.315)	0.387 (0.315)	0.009 (0.001)***	1.067 (0.465)*	1.469 (0.464)**	—	LI
Latvia	0.003 (0.000)***	-0.325 (0.264)	0.063 (0.264)	0.008 (0.001)***	2.419 (0.407)***	1.030 (0.407)*	LI	QU
Lithuania	0.001 (0.000)**	-1.001 (0.298)***	0.753 (0.298)*	0.003 (0.001)***	0.680 (0.422)	2.530 (0.421)***	QIU	C
Luxembourg	0.001 (0.001)	-0.355 (0.304)	0.848 (0.304)**	0.004 (0.001)**	0.944 (0.433)*	0.768 (0.433)	—	—
North-Macedonia	0.003 (0.001)***	0.486 (0.269)	0.054 (0.269)	0.005 (0.001)***	1.666 (0.372)***	-0.277 (0.372)	LI	LI
Malta	0.005 (0.001)***	-0.845 (0.369)*	0.761 (0.368)*	0.011 (0.001)***	-0.482 (0.447)	0.660 (0.446)	LI	LI
Netherlands	0.002 (0.001)***	0.644 (0.249)*	-0.044 (0.249)	0.006 (0.001)***	0.567 (0.390)	0.348 (0.390)	LI	LI
Norway	0.001 (0.000)**	0.261 (0.248)	-0.287 (0.248)	0.003 (0.001)***	0.398 (0.364)	0.478 (0.364)	—	LI
Poland	0.003 (0.000)***	-0.441 (0.282)	0.202 (0.282)	0.008 (0.001)***	1.520 (0.404)***	1.654 (0.404)***	LI	C
Portugal	0.001 (0.000)*	0.548 (0.241)*	-0.212 (0.241)	0.005 (0.001)***	2.420 (0.389)***	0.531 (0.389)	—	QU
Romania	0.001 (0.001)	0.016 (0.321)	0.662 (0.321)*	0.004 (0.001)***	1.691 (0.442)***	1.542 (0.441)***	—	C
Russia	0.003 (0.001)***	-1.171 (0.303)***	-0.193 (0.303)	0.003 (0.001)***	-1.039 (0.384)**	-0.461 (0.384)	QIU	LI
Scotland	0.002 (0.000)***	0.473 (0.275)	-0.103 (0.275)	0.006 (0.001)***	3.095 (0.403)***	0.701 (0.403)	LI	QU
Slovakia	0.003 (0.000)***	-0.828 (0.288)**	-0.119 (0.288)	0.007 (0.001)***	1.656 (0.405)***	1.061 (0.405)**	LI	QU
Slovenia	0.001 (0.001)*	0.252 (0.237)	0.410 (0.237)	0.007 (0.001)***	0.912 (0.355)**	-0.843 (0.354)**	LI	LI
Spain	-0.002 (0.001)**	0.136 (0.278)	-0.063 (0.278)	-0.002 (0.001)**	1.299 (0.406)**	-0.178 (0.406)	—	LI
Sweden	0.002 (0.000)***	0.436 (0.299)	0.904 (0.299)**	0.006 (0.001)***	1.442 (0.428)***	2.007 (0.427)***	LI	C
Switzerland	0.002 (0.000)***	0.492 (0.248)*	0.624 (0.248)**	0.007 (0.001)***	2.913 (0.389)***	2.231 (0.388)***	LI	C
Ukraine	0.000 (0.001)	-0.477 (0.272)	1.025 (0.272)***	0.000 (0.001)	0.344 (0.399)	1.170 (0.398)**	C	—
USA	-0.004 (0.001)**	0.385 (0.331)	0.218 (0.331)	-0.002 (0.002)	-0.901 (0.441)*	-1.140 (0.440)**	—	—
Wales	0.002 (0.000)***	1.622 (0.309)***	0.649 (0.309)*	0.009 (0.000)***	3.797 (0.444)***	2.401 (0.443)***	QU	C

Notes. Estimates (Standard error); $P < 0.001 = ***$; $P < 0.01 = **$; $P < 0.05 = *$; LI, linear increase; LD, linear decrease; QU, quadratic U-shaped; QIU, quadratic inverted U-shaped; C = cubic.

patterns of change over time (i.e., linear, cubic, and quadratic) in psychological and somatic complaints, as well as explore gender differences therein among representative samples of 15-year-old adolescents.

Within most countries an increase in both psychological and somatic complaints was observed since 1994 with either a steady increasing trend (i.e., linear trend), a decrease followed by an increase (i.e., U-shaped trend), or a fluctuating trend with an increase towards 2022 (i.e., cubic trend). Only within a small number of countries stability or a decrease in psychological and somatic complaints over time was observed. Furthermore, our analyses revealed that the trend patterns among girls exhibited greater complexity, characterised by a higher prevalence of U-shaped and cubic

patterns, in contrast to boys, where many patterns remained linear, particularly in somatic complaints. These more intricate patterns suggest heightened fluctuations over time. This might be linked to school stress, body image issues, obesity, and bullying which have been shown to disproportionately affect girls [24, 31].

In comparison to previous research [25], which used data from 1994 to 2010, a shift in trend patterns was observed. While Ottová-Jordan et al. [25] indicated a relatively even distribution between stable, increasing (linear or U-shaped), or decreasing trend patterns (linear, inverted-U-shaped), with only four countries exhibiting an unsteady trend pattern (i.e., cubic), the current study revealed a notable shift in this distribution, with increases becoming the predominant trend pattern by early

TABLE 3 | Overview of time trend patterns for psychological and somatic complaints by gender and country for 1994–2018 and 1994–2022 (Health Behaviour in School-aged Children study, 41 countries).

Country/region	Trend until 2018							Trend until 2022	
	Psychological complaints			Somatic complaints				Psychological complaints	Somatic complaints
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls	
Armenia ^e	–	–	–	Linear decrease	–	Quadratic—U-shaped*	–	Quadratic—U-shaped*	
Austria ^a	Linear increase	Quadratic—U-shaped	–	Linear increase	Quadratic—U-shaped*	Quadratic—U-shaped	Linear increase*	Quadratic—U-shaped*	
Belgium—Flanders ^a	–	Linear increase	Quadratic—inverted U-shaped	Cubic	Cubic*	Cubic*	Linear increase*	Quadratic—U-shaped*	
Belgium—Wallonia ^a	Quadratic—U-shaped	Quadratic—U-shaped	Quadratic—inverted U-shaped	Linear increase	Quadratic—U-shaped	Cubic*	—*	Linear increase	
Bulgaria ^d	Quadratic—U-shaped	Linear increase	Quadratic—U-shaped	–	Cubic*	Linear increase	Linear increase*	Cubic*	
Canada ^a	–	Quadratic—U-shaped	–	Linear increase	Quadratic—U-shaped*	Quadratic—U-shaped	–	Quadratic—U-shaped*	
Croatia ^c	–	Quadratic—U-shaped	–	–	–	Quadratic—U-shaped	–	Quadratic—U-shaped*	
Czechia	Linear increase	Linear increase	–	Quadratic—inverted U-shaped	Linear increase	Quadratic—U-shaped*	–	Cubic*	
Denmark ^a	Linear increase	Linear increase	Linear increase	Linear increase	Linear increase	Cubic*	Linear increase	Cubic*	
England ^b	Cubic	Linear increase	–	Linear increase	Quadratic—U-shaped*	Cubic*	Cubic*	Cubic*	
Estonia ^a	–	Cubic	–	Cubic	Cubic*	Cubic	Cubic*	Cubic	
Finland ^a	–	Quadratic—U-shaped	Linear increase	Linear increase	Linear increase*	Quadratic—U-shaped	—*	Cubic*	
France ^a	Quadratic—U-shaped	Quadratic—U-shaped	Cubic	Quadratic—inverted U-shaped	Quadratic—U-shaped	Quadratic—U-shaped	Linear increase*	Cubic*	
Germany ^a	–	Quadratic—U-shaped	–	Linear increase	Cubic*	Cubic*	Linear increase*	Quadratic—U-shaped*	
Greece ^b	Linear decrease	Cubic	–	–	Cubic*	Cubic	Quadratic—U-shaped*	Cubic*	
Greenland ^a	–	Linear increase	Linear increase	Linear increase	Linear increase*	Linear increase	Linear increase	Linear increase	
Hungary ^a	–	Cubic	Linear increase	Linear increase	Quadratic—U-shaped*	Cubic	Linear increase	Cubic*	
Iceland ^d	–	Linear increase	–	Cubic	Quadratic—U-shaped*	Quadratic—U-shaped*	/	Quadratic—U-shaped*	
Ireland ^b	–	Linear increase	–	Linear increase	Quadratic—U-shaped*	Quadratic—U-shaped*	Linear increase*	Linear increase	
Israel ^a	Linear decrease	–	–	–	—*	Cubic*	Quadratic—inverted U-shaped*	Cubic*	
Italy ^c	–	Linear increase	–	Linear increase	Linear increase*	Quadratic—U-shaped*	/	Linear increase	
Latvia ^a	Linear increase	Linear increase	Linear increase	Linear increase	Quadratic—U-shaped*	Cubic*	Linear increase	Quadratic—U-shaped*	
Lithuania ^a	–	Cubic	Quadratic - inverted U-shaped	–	Cubic*	Cubic	Quadratic-inverted U-shaped	Cubic*	
Luxembourg ^d	–	Cubic	–	–	Linear increase*	Quadratic—U-shaped*	–	–	

(Continued on following page)

TABLE 3 | (Continued) Overview of time trend patterns for psychological and somatic complaints by gender and country for 1994–2018 and 1994–2022 (Health Behaviour in School-aged Children study, 41 countries).

Country/region	Trend until 2018						Trend until 2022	
	Psychological complaints			Somatic complaints			Psychological complaints	Somatic complaints
	Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls
North-Macedonia ^c	—	Quadratic—U-shaped	—	—	Linear increase*	Quadratic—U-shaped	Linear increase*	Linear increase*
Malta ^c	—	Linear increase	Linear increase	Linear increase	—	Linear increase	Linear increase	Linear increase
Netherlands ^c	—	Linear increase	—	Cubic	Quadratic—U-shaped*	Quadratic—U-shaped*	Linear increase*	Linear increase*
Norway ^a	—	Linear increase	—	—	Cubic*	Linear increase	—	Linear increase*
Poland ^a	Cubic	Cubic	Quadratic—inverted U-shaped	Linear increase	Cubic	Cubic	Linear increase*	Cubic*
Portugal ^b	Quadratic—U-shaped	Quadratic—U-shaped	—	—	Quadratic—U-shaped	Quadratic—U-shaped	/	Quadratic—U-shaped*
Romania ^d	—/	Quadratic - U-shaped	—	—	Quadratic—U-shaped*	Quadratic—U-shaped	/	Cubic*
Russia ^a	—	—	Quadratic—inverted U-shaped	Linear increase	—	—	—	—
Scotland ^a	Quadratic—U-shaped	Quadratic—U-shaped	—	Quadratic—U-shaped	Quadratic—U-shaped	Quadratic—U-shaped	Linear increase*	Quadratic—U-shaped
Slovakia ^a	Linear increase	—	Linear increase	Linear increase	Cubic*	Cubic*	Linear increase	Quadratic—U-shaped*
Slovenia ^c	Quadratic—U-shaped	Cubic	—	Linear increase	Quadratic—U-shaped	Cubic	Linear increase*	Linear increase
Spain ^c	Linear decrease	Linear decrease	—	Linear decrease	Linear decrease	Quadratic—U-shaped*	—	Linear decrease
Sweden ^a	Linear increase	Linear increase	—	Linear increase	Cubic*	Cubic*	Linear increase*	Cubic*
Switzerland ^b	—	Quadratic—U-shaped	—	Linear increase	Linear increase*	Cubic*	Linear increase*	Cubic*
Ukraine ^c	Quadratic—U-shaped	Cubic	Cubic	—	—	—	—	—
USA ^b	Linear decrease	Cubic	—	—	—	—	—	—
Wales ^a	Cubic	Quadratic—U-shaped	—	Linear increase	Quadratic—U-shaped*	Quadratic—U-shaped	Quadratic—U-shaped*	Cubic*
<i>Total number of countries with</i>								
Linear decrease	4	1	0	2	1	0	0	1
Linear increase	6	14	7	20	8	4	19	9
Quadratic—inverted U-shaped	0	0	5	2	0	0	2	0
Quadratic—U-shaped	7	13	1	1	15	18	2	11
Cubic	3	9	2	4	10	16	2	16
None (—)	21	4	26	12	4	0	13	1

Notes *. Different trend pattern for 1994–2018 and 1994–2022.

^aData since 1994.

^bData since 1998.

^cData since 2002.

^dData since 2006.

^eData since 2010.

2020s. In general, the inclusion of 2022 data demonstrated the acceleration of this upward trend in recent years with non-monotonic time trends becoming more prevalent, which has also been confirmed by previous studies. This may be attributed to the impact of the COVID-19 pandemic, however, the limited number of decreasing time trends in 2018 and a higher number of increasing time trends compared to Ottová-Jordan [25] suggests that this concerning upwards trend was already established prior to the pandemic. Therefore, it is imperative to closely monitor future trends to provide insight into whether COVID-19 has temporarily increased the prevalence or whether the alarming time trend has continued.

In addition, considerable variations in trend patterns for psychological and somatic complaints were observed, with psychological complaints often characterised by complex fluctuations over time, in comparison to the more stable trajectory of somatic complaints. Moreover, numerous cross-national variations in the observed trend patterns by gender and health complaints dimensions were identified. However, no discernible clusters of countries with similar trend patterns were observed. These results indicate that a linear approach in many countries may not be sufficient to capture the complexities of trends over three decades, particularly in girls and in psychological complaints.

Previous research has also well-established a higher prevalence of psychological and somatic complaints among girls compared to boys, along with more pronounced increases over time in girls, which may be reflected in the more complex trend patterns observed in girls compared to boys [32, 33]. Moreover, some countries showed particularly high prevalence rates, like Israel and Italy for both Dimensions of health complaints, with Greece showing higher psychological complaints. Structural factors and macro-level changes may partially explain these findings. For instance, in Israel, political and economic instability and violence exposure contribute to adolescents' mental health issues [34, 35]. In Greece, the refugee crisis and economic hardship worsen health inequities and mental health problems among adolescents [36, 37]. In Italy, the severe COVID-19 impact and prolonged school closures, along with increased school pressure and social media influence, have led to high psychosomatic complaints in adolescents [38, 39].

The sensitivity test confirmed that for both boys and girls the patterns became more complex when 2022 data (i.e., post-COVID period) were added. For boys in many countries there was stability until 2018, which changed to an upward trend pattern when 2022 data were included. For girls most of the time, existing linear increases became more complex trend patterns with 2022 data included. This sensitivity test illustrated an acceleration of a pre-existing upward trend for girls, whereas for boys it illustrated the beginning of an increase.

In addition to the observed rise in psychological and somatic complaints, potentially attributable to COVID-19, there have been multiple notable fluctuations over time. While this study did not address these fluctuations in detail, it is plausible that various cross-national crises and societal developments have contributed to the observed rise in adolescent psychological and somatic complaints. For example, past crises such as the 2008 economic recession have

been associated with adverse effects on adolescent mental health [40]. Additionally, factors such as climate change [41, 42], and armed conflicts [35, 43], may have contributed to the increase in psychosomatic complaints in recent years.

Beyond crises, societal developments have played an important role, with both positive and negative effects on adolescent mental health. Developments such as the widespread availability of the internet since the 1990s and the emergence of social media in the mid-2000s are likely to have negatively influenced adolescent mental health [24, 44], although some show no impact or a positive one [45, 46]. Furthermore, increased awareness of mental health issues, coupled with greater availability of mental health services and increased prevention efforts, may have positively influenced mental health outcomes [47–51]. Also, individual-level factors and specific events at the national level may have influenced these time trends in psychological and somatic complaints as well. All these factors, and probably many more, are likely to have contributed to the observed variations in trend patterns. However, this was not the primary objective of this study, and it is therefore uncertain whether the observed fluctuations in adolescent psychological and somatic complaints can be attributed to these factors.

Strengths, Limitations and Future Research

The strength of the present study was the use of repeated cross-sectional data from the HBSC study in a large sample of countries, which has adhered to a standardised protocol over time. This provided a unique opportunity to examine 30-year trend patterns across multiple countries, enriching our understanding of the dynamics of adolescent health and wellbeing. However, some limitations need to be considered. The rapid expansion of the HBSC network over the past two decades, led to inconsistent data availability across countries, with some countries lacking data starting from 1994, thereby limiting the examination of trend patterns. Additionally, reliance on self-report measures introduces potential bias, as increased awareness of mental health issues may lead to more reported symptoms, aligning with the prevalence inflation hypothesis [50]. Also, although psychological and somatic complaints are more pronounced in 15-year-olds, studying a wider age range in future studies might provide a better overview of these time trends and their development. Furthermore, the different time trend patterns observed for psychological and somatic complaints in most countries raised questions about the relationship between these two dimensions. Future research is needed to investigate whether there might be a different relationship between the two dimensions over time. Future studies could also focus clusters of countries and include macro-level indicators such as cultural factors or characteristics of welfare and healthcare systems to examine these matters in greater depth. Lastly, the nature of the Multiple Health Complaints Scale, which primarily measures internalizing complaints, could bias the results regarding gender differences. From a developmental perspective, girls are generally more prone to internalizing complaints, whereas boys are more prone to externalizing complaints [5]. Therefore, internalising instruments might not fully capture psychological distress in boys and results should be interpreted with caution.

Conclusion

In conclusion, an upward trend in psychological and somatic complaints has been observed cross-nationally in recent years. Furthermore, non-monotonical trend patterns have been observed more frequently in girls compared to boys and in psychological compared to somatic complaints. The most prevalent patterns were U-shaped and cubic patterns, in addition to linear increases. However, the identified trend patterns included many cross-national variations. The diverse trend patterns highlight the complexity of the time trends and suggest that a linear approach might be insufficient for capturing the full scope of these trends over three decades.

ETHICS STATEMENT

The studies involving humans were approved by the Committee of the University Hospital of Ghent (ONZ20230093). The studies were conducted in accordance with the local legislation and institutional requirements. Participants provided informed consent for participation in this study.

AUTHOR CONTRIBUTIONS

KS, MD, and AC conceptualised the study. KS analyzed the data with the support of TP. KS drafted the first version of the manuscript. All

authors reviewed and edited several versions of the manuscript until an agreement was reached on its final version. All authors contributed to the article and approved the submitted version.

FUNDING

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. This work was supported by the government of Flanders (Belgium) and published with the support of the University Foundation of Belgium. The Health Behaviour in School-aged Children survey is funded by public sources in each member country. AC was supported by the project “Research of Excellence on Digital Technologies and Wellbeing CZ.02.01.01/00/22_008/0004583” which is co-financed by the European Union.

CONFLICT OF INTEREST

The authors declare that they do not have any conflicts of interest.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.ssph-journal.org/articles/10.3389/ijph.2024.1607774/full#supplementary-material>

REFERENCES

- Patton GC, Sawyer SM, Santelli JS, Ross DA, Afifi R, Allen NB, et al. Our Future: A Lancet Commission on Adolescent Health and Wellbeing. *Lancet* (2016) 387(10036):2423–78. doi:10.1016/s0140-6736(16)00579-1
- Cosma A, Abdrakhmanova S, Taut D, Schrijvers K, Catunda C, Schnohr C. A Focus on Adolescent Mental Health and Well-Being in Europe, Central Asia and Canada. In: *Health Behaviour in School-Aged Children International Report from the 2021/2022 Survey*. Copenhagen: WHO Regional Office for Europe (2023).
- Inchley J, Currie D, Budisavljevic S, Torsheim T, Jåstad A, Cosma A. Spotlight on Adolescent Health and Well-Being. Findings from the 2017/2018 Health Behaviour in School-Aged Children (HBSC) Survey in Europe and Canada. In: *International Report*. Copenhagen: WHO Regional Office for Europe (2020).
- Ravens-Sieberer U, Torsheim T, Hetland J, Vollebergh W, Cavallo F, Jericek H, et al. Subjective Health, Symptom Load and Quality of Life of Children and Adolescents in Europe. *Int J Public Health* (2009) 54(2):151–9. doi:10.1007/s00038-009-5406-8
- Beck JE. A Developmental Perspective on Functional Somatic Symptoms. *J Pediatr Psychol* (2008) 33(5):547–62. doi:10.1093/jpepsy/jsm113
- Grigorian K, Östberg V, Raninen J, Åhlén J, Brolin Läftman S. Prospective Associations between Psychosomatic Complaints in Adolescence and Depression and Anxiety Symptoms in Young Adulthood: A Swedish National Cohort Study. *SSM Popul Health* (2023) 24:101509. doi:10.1016/j.ssmph.2023.101509
- Wu Y, Wang L, Tao M, Cao H, Yuan H, Ye M, et al. Changing Trends in the Global Burden of Mental Disorders From 1990 to 2019 and Predicted Levels in 25 Years. *Epidemiol Psychiatr Sci* (2023) 32(e63):e63–9. doi:10.1017/s2045796023000756
- Cosma A, Stevens G, Martin G, Duinhof EL, Walsh SD, Garcia-Moya I, et al. Cross-National Time Trends in Adolescent Mental Well-Being From 2002 to 2018 and the Explanatory Role of Schoolwork Pressure. *J Adolesc Health* (2020) 66(6s):S50–S58–s8. doi:10.1016/j.jadohealth.2020.02.010
- Inchley J, Currie D, Piper A, Jåstad A, Cosma A, Nic Gabhainn S, et al. Health Behaviour in School-Aged Children (HBSC) Study Protocol: Background, Methodology, Mandatory Questions and Optional Packages for the 2021/22 Survey. In: *MRC/CSO Social and Public Health Sciences Unit*. Glasgow: University of Glasgow (2023).
- Potrebný T, Wium N, Lundegård MM-I. Temporal Trends in Adolescents’ Self-Reported Psychosomatic Health Complaints from 1980–2016: A Systematic Review and Meta-Analysis. *PLoS One* (2017) 12(11):e0188374. doi:10.1371/journal.pone.0188374
- Hetland J, Torsheim T, Aarø LE. Subjective Health Complaints in Adolescence: Dimensional Structure and Variation across Gender and Age. *Scand J Public Health* (2002) 30(3):223–30. doi:10.1080/140349402320290953
- Haugland S, Wold B, Stevenson J, Aarø LE, Woynarowska B. Subjective Health Complaints in Adolescence. A Cross-National Comparison of Prevalence and Dimensionality. *Eur J Public Health* (2001) 11(1):4–10. doi:10.1093/eurpub/11.1.4
- Heinz A, Sischka PE, Catunda C, Cosma A, Garcia-Moya I, Lyyra N, et al. Item Response Theory and Differential Test Functioning Analysis of the HBSC-Symptom-Checklist Across 46 Countries. *BMC Med Res Methodol* (2022) 22(1):253. doi:10.1186/s12874-022-01698-3
- Ravens-Sieberer U, Erhart M, Torsheim T, Hetland J, Freeman J, Danielson M, et al. An International Scoring System for Self-Reported Health Complaints in Adolescents. *Eur J Public Health* (2008) 18(3):294–9. doi:10.1093/eurpub/ckn001
- Dey M, Jorm AF, Mackinnon AJ. Cross-sectional Time Trends in Psychological and Somatic Health Complaints Among Adolescents: A

- Structural Equation Modelling Analysis of 'Health Behaviour in School-Aged Children' Data From Switzerland. *Soc Psychiatry Psychiatr Epidemiol* (2015) 50(8):1189–98. doi:10.1007/s00127-015-1040-3
16. Potrebny T, Wiium N, Haugstvedt A, Sollesnes R, Torsheim T, Wold B, et al. Health Complaints Among Adolescents in Norway: A Twenty-Year Perspective on Trends. *PLoS One* (2019) 14(1):e0210509. doi:10.1371/journal.pone.0210509
 17. Boer M, Van Dorsselaer S, De Looze M, De Roos S, Brons H, Van den Eijnden R, et al. *Gezondheid en welzijn van jongeren in Nederland*. Utrecht: Universiteit Utrecht (2022).
 18. Inchley J, Mabelis J, Brown J, Willis M, Currie D. *Health Behaviour in School-Aged Children (HBSC) 2022 Survey in Scotland: National Report*. Glasgow: MRC/CSO Social and Public Health Sciences Unit, University of Glasgow (2023).
 19. Hulbert S, Eida T, Ferris E, Hrytsenko V, Kendall S. *HBSC England National Report: Findings from the 2021–2022 HBSC Study for England*. Kent: University of Kent (2023).
 20. Bersia M, Berchiappa P, Charrier L, Lemma P, Borraccino A, Nardone P, et al. Mental Well-Being: 2010–2018 Trends Among Italian Adolescents. *Int J Environ Res Public Health* (2022) 19(2):863. doi:10.3390/ijerph19020863
 21. Cosma A, Kólto A, Badura P, Winkler P, Kalman M. Time Trends in Adolescent Mental Wellbeing in the Czech Republic Between 2002 and 2018: Gender, Age and Socioeconomic Differences. *Cent Eur J Public Health* (2021) 29(4):271–8. doi:10.21101/cejph.a6717
 22. Hammami N, Da Silva MA, Elgar FJ. Trends in Gender and Socioeconomic Inequalities in Adolescent Health Over 16 Years (2002–2018): Findings From the Canadian Health Behaviour in School-Aged Children Study. *Health Promotion Chronic Dis Prev Can* (2022) 42(2):68–78. doi:10.24095/hpcdp.42.2.03
 23. Mazur J, Klanšček HJ, Augustine L, Porwit K, Sigmund E, Šmigelskas K. Trends in Multiple Health Complaints in Polish Adolescents in Light of Data From 30 European Countries and Canada (2002–2018). *J Mother Child* (2021) 25(1):25–34. doi:10.34763/jmotherandchild.20212501.d-21-00011
 24. Boer M, Cosma A, Twenge J, Inchley J, Klanšček H, Stevens G. National-Level Schoolwork Pressure, Family Structure, Internet Use, and Obesity as Drivers of Time Trends in Adolescent Psychological Complaints Between 2002 and 2018. *J Youth Adolesc* (2023) 52:2061–77. doi:10.1007/s10964-023-01800-y
 25. Ottová-Jordan V, Smith ORF, Gobina I, Mazur J, Augustine L, Cavallo F, et al. Trends in Multiple Recurrent Health Complaints in 15-Year-Olds in 35 Countries in Europe, North America and Israel From 1994 to 2010. *Eur J Public Health* (2015) 25(Suppl. 1.2):24–7. doi:10.1093/eurpub/ckv015
 26. Hammami N, Da Silva MA, Elgar FJ. Evolution of Health Inequalities Among Adolescents Based on Gender and Socioeconomic Status Over 16 Years (2002 to 2018): Results of the Study on Health Behaviors of School-Aged Children. *Health Promotion Chronic Dis Prev Can* (2022) 42(2):77–88. doi:10.24095/hpcdp.42.2.03f
 27. Buli BG, Lehtinen-Jacks S, Larm P, Nilsson KW, Hellström-Olsson C, Giannotta F. Trends in Psychosomatic Symptoms Among Adolescents and the Role of Lifestyle Factors. *BMC Public Health* (2024) 24(1):878. doi:10.1186/s12889-024-18327-x
 28. Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via Coordinate Descent. *J Stat Softw* (2010) 33:1–22. doi:10.18637/jss.v033.i01
 29. Ioannidis JPA. What Have We (Not) Learnt from Millions of Scientific Papers With P-Values? *Am Statistician* (2019) 73:20–5. doi:10.1080/00031305.2018.1447512
 30. Cosma A, Bersia M, Abdrakhmanova S, Badura P, Gobina I. Coping through Crisis: COVID-19 Pandemic Experiences and Adolescent Mental Health and Well-Being in the WHO European Region: Impact of the COVID-19 Pandemic on Young People's Health and Well-Being from the Findings of the HBSC Survey Round 2021/2022. In: *Regional Office for Europe*. Copenhagen: World Health Organization (2023).
 31. Whitehead RD, Cosma A, Cecil J, Currie C, Currie D, Neville F, et al. Trends in the Perceived Body Size of Adolescent Males and Females in Scotland, 1990–2014: Changing Associations with Mental Well-Being. *Int J Public Health* (2018) 63(1):69–80. doi:10.1007/s00038-017-0997-y
 32. Keyes KM, Platt JM. Annual Research Review: Sex, Gender, and Internalizing Conditions Among Adolescents in the 21st Century - Trends, Causes, Consequences. *J Child Psychol Psychiatry* (2023) 65(4):384–407. doi:10.1111/jcpp.13864
 33. Bor W, Dean AJ, Najman J, Hayatbakhsh R. Are Child and Adolescent Mental Health Problems Increasing in the 21st Century? A Systematic Review. *Aust N Z J Psychiatry* (2014) 48(7):606–16. doi:10.1177/0004867414533834
 34. Mana A, Super S, Sardu C, Juvinya Canal D, Moran N, Sagy S. Individual, Social and National Coping Resources and Their Relationships with Mental Health and Anxiety: A Comparative Study in Israel, Italy, Spain, and the Netherlands during the Coronavirus Pandemic. *Glob Health Promotion* (2021) 28(2):17–26. doi:10.1177/1757975921992957
 35. Wagner G, Glick P, Khammash U, Shaheen M, Brown R, Goutam P, et al. Exposure to Violence and its Relationship to Mental Health Among Young People in Palestine. *East Mediterr Health J* (2020) 26(2):189–97. doi:10.26719/2020.26.2.189
 36. Diamantis DV, Stavropoulou I, Katsas K, Mugford L, Linos A, Kouvari M. Assessing Quality of Life in First- and Second-Generation Immigrant Children and Adolescents; Highlights From the DIATROFI Food Aid and Healthy Nutrition Promotion Program. *Int J Environ Res Public Health* (2023) 20(3):2471. doi:10.3390/ijerph20032471
 37. Karagianni P. Youth Mental Health Context in Greece. *The Eur Health Psychol* (2016) 18(3):119–22.
 38. Barbieri V, Wiedermann CJ, Kaman A, Erhart M, Piccoliori G, Plagg B, et al. Quality of Life and Mental Health in Children and Adolescents after the First Year of the COVID-19 Pandemic: A Large Population-Based Survey in South Tyrol, Italy. *Int J Environ Res Public Health* (2022) 19(9):5220. doi:10.3390/ijerph19095220
 39. Bersia M, Charrier L, Berchiappa P, Cosma A, Comoretto RI, Dalmasso P. The Mental Well-Being of Italian Adolescents in the Last Decade through the Lens of the Dual Factor Model. *Children-Basel* (2022) 9(12):1981. doi:10.3390/children9121981
 40. Hiilamo A, Hiilamo H, Ristikari T, Virtanen P. Impact of the Great Recession on Mental Health, Substance Use and Violence in Families with Children: A Systematic Review of the Evidence. *Child Youth Serv Rev* (2021) 121:105772. doi:10.1016/j.chilgyouth.2020.105772
 41. van Nieuwenhuizen A, Hudson K, Chen X, Hwong AR. The Effects of Climate Change on Child and Adolescent Mental Health: Clinical Considerations. *Curr Psychiatry Rep* (2021) 23(12):88. doi:10.1007/s11920-021-01296-y
 42. Martin G, Reilly K, Everitt H, Gilliland JA. Review: The Impact of Climate Change Awareness on Children's Mental Well-Being and Negative Emotions - A Scoping Review. *Child Adolesc Ment Health* (2022) 27(1):59–72. doi:10.1111/camh.12525
 43. Bendavid E, Boerma T, Akseer N, Langer A, Malembaka EB, Okiro EA, et al. The Effects of Armed Conflict on the Health of Women and Children. *Lancet* (2021) 397(10273):522–32. doi:10.1016/s0140-6736(21)00131-8
 44. Sohn SY, Rees P, Wildridge B, Kalk NJ, Carter B. Prevalence of Problematic Smartphone Usage and Associated Mental Health Outcomes Amongst Children and Young People: A Systematic Review, Meta-Analysis and Grade of the Evidence. *BMC Psychiatry* (2019) 19(1):356. doi:10.1186/s12888-019-2350-x
 45. Ferguson CJ, Kaye LK, Branley-Bell D, Markey P, Ivory JD, Klisanin D, et al. Like This Meta-Analysis: Screen Media and Mental Health. *Prof Psychol Res Pract* (2022) 53(2):205–14. doi:10.1037/pro0000426
 46. Radtke T, Apel T, Schenkel K, Keller J, von Lindern E. Digital Detox: An Effective Solution in the Smartphone Era? A Systematic Literature Review. *Mobile Media and Commun* (2022) 10(2):190–215. doi:10.1177/20501579211028647
 47. König D, Fellingner M, Pruckner N, Hinterbuchinger B, Dorffner G, Gleiss A, et al. Availability and Use of Mental Health Services in European Countries: Influence on National Suicide Rates. *J Affect Disord* (2018) 239:66–71. doi:10.1016/j.jad.2018.06.042

48. Michaud P-A, Visser A, Vervoort JPM, Kocken P, Reijneveld SA, Jansen DEMC. Availability and Accessibility of Primary Mental Health Services for Adolescents: An Overview of National Recommendations and Services in EU. *Eur J Public Health* (2020) 30(6):1127–33. doi:10.1093/eurpub/ckaa102
49. McGorry PD, Mei C, Chanen A, Hodges C, Alvarez-Jimenez M, Killackey E. Designing and Scaling up Integrated Youth Mental Health Care. *World Psychiatry* (2022) 21(1):61–76. doi:10.1002/wps.20938
50. Foulkes L, Andrews JL. Are Mental Health Awareness Efforts Contributing to the Rise in Reported Mental Health Problems? A Call to Test the Prevalence Inflation Hypothesis. *New Ideas Psychol* (2023) 69:101010. doi:10.1016/j.newideapsych.2023.101010
51. Kuosmanen T, Clarke AM, Barry MM. Promoting Adolescents' Mental Health and Wellbeing: Evidence Synthesis. *J Public Ment Health* (2019) 18(1):73–83. doi:10.1108/jpmh-07-2018-0036

Copyright © 2024 Schrijvers, Cosma, Potrebny, Thorsteinsson, Catunda, Reiss, Hulbert, Kostičová, Melkumova, Bersia, Klanšček, Gaspar and Dierckens. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.