



Kent Academic Repository

Maguire, K, Barnoux, Magali F.L., Collins, Josephine, Melvin, Clare L., Inkson, I, Alexander, Regi T., Devapriam, J, Duggan, C, Shepstone, Lee, Staufenberg, Ekkehart and others (2025) *The Factor Structure and Validity of the Psychopathy Checklist-Short Version When Used With Autistic Psychiatric Inpatients*. *Autism Research* . ISSN 1939-3792.

Downloaded from

<https://kar.kent.ac.uk/108721/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1002/aur.70004>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal** , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

RESEARCH ARTICLE OPEN ACCESS

The Factor Structure and Validity of the Psychopathy Checklist-Short Version When Used With Autistic Psychiatric Inpatients

Kate Maguire¹ | Magali Barnoux²  | Josie Collins³ | Clare L. Melvin⁴  | Ismay Inkson⁵ | Regi T. Alexander^{6,7}  | John Devapriam⁸ | Conor Duggan⁹ | Lee Shepstone¹⁰ | Ekkehart Staufenburg⁷ | Paul Thompson¹¹ | David Turner¹⁰ | Essi Viding¹² | Peter E. Langdon^{8,11,13} 

¹School of Health and Social Care, University of Essex, Colchester, UK | ²Tizard Centre, University of Kent, Kent, UK | ³Department of Law and Criminology, Royal Holloway, University of London, London, UK | ⁴Department of Psychology, University of East Anglia, Norwich, UK | ⁵Norfolk Community Health and Care NHS Trust, Norwich, UK | ⁶University of Hertfordshire, Hatfield, UK | ⁷Little Plumstead Hospital, Hertfordshire Partnership University NHS Foundation Trust, Norwich, UK | ⁸Herefordshire and Worcestershire Health and Care NHS Trust, Worcester, UK | ⁹Division of Psychiatry and Applied Psychology, School of Medicine, University of Nottingham, Nottingham, UK | ¹⁰Norwich Medical School, University of East Anglia, Norwich, UK | ¹¹Intellectual Disabilities Research Institute (IDRIS), University of Birmingham, Birmingham, UK | ¹²Division of Psychology and Language Sciences, University College London, London, UK | ¹³Birmingham Community Healthcare NHS Foundation Trust, Birmingham, UK

Correspondence: Peter E. Langdon (p.langdon@bham.ac.uk)

Received: 24 October 2024 | **Revised:** 28 January 2025 | **Accepted:** 9 February 2025

Funding: This study/project was funded by the NIHR Research for Patient Benefit Programme (NIHR: PB-PG-0214-33040). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Keywords: autism | crime | intellectual disabilities | psychiatric inpatient | psychopathy | risk

ABSTRACT

The Psychopathy Checklist Short Version (PCL:SV) is a brief measure of psychopathy. This study aimed to assess the reliability and validity of the PCL:SV with autistic adults detained in inpatient psychiatric care. Data were collected from 282 autistic adults at two time points separated by 12-months. Reliability and validity were investigated using omega, regression, receiver operating characteristic curves, and correlational analysis. PCL:SV Total, Factor 1, and Factor 2 had satisfactory to high reliability and construct validity. Higher PCL:SV scores were associated with poorer treatment progress, a longer length of stay, and previous criminal offending. Factor 1 was associated with a forensic history, detention under Part III of the Mental Health Act, and a personality disorder diagnosis, while Factor 2 was also associated with the absence of a forensic history, detention under Part II of the Mental Health Act, but not a personality disorder diagnosis. It was thought that Factor 2 most likely captured data associated with autism and/or intellectual disabilities (e.g., behaviors that challenge). Those with intellectual disabilities were less likely to have convictions, a history of violent offending, or a forensic history. They were also more likely to be detained under Part II of the Mental Health Act, and were more likely to have had a positive transfer 12-months later to a ward with lesser security. The PCL:SV correlated as expected with the HCR-20 and the START. This study provides preliminary evidence to support the use of the PCL:SV with autistic adults, including those with intellectual disabilities, within inpatient psychiatric hospitals.

Psychopathy is a construct characterized by shallow emotions, lack of empathy or remorse, callousness, and poor behavioral control (Cleckley 1941; Hare 1991). Prevalence in the general

population is estimated at 0.6% (Coid, Yang, Ullrich, Roberts, and Hare 2009), with a higher prevalence among offenders (Coid, Yang, Ullrich, Roberts, Moran, et al. 2009). It has long

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Autism Research* published by International Society for Autism Research and Wiley Periodicals LLC.

Summary

- Psychopathy is a construct that is characterized by shallow emotions, lack of empathy or remorse, callousness, and poor behavioral control.
- We recruited autistic psychiatric inpatients into our study, and their clinical teams completed the Psychopathy Checklist—Short Version (PCL:SV) to see how well this measure worked with autistic adults. Our results indicated that higher PCL:SV scores were associated with poorer treatment progress, a longer length of stay, and previous criminal offending and aggressive behavior 12-months later.

been associated with criminal and violent behavior and is a key predictor of recidivism (Azevedo et al. 2020; Hare 1999; Woodworth and Porter 2002).

Little is known about the co-existence of autism and psychopathy, and both conditions are characterized by difficulties with perspective-taking. Autistic individuals do come into contact with the criminal justice system (CJS), and prevalence estimates of criminal justice contact vary widely from 0.2% to 62.8% (Collins et al. 2022). These prevalence studies are fraught with methodological problems, and on the whole, autistic individuals present with a reduced risk of engaging in crime (Collins et al. 2022).

In an attempt to disentangle the relationship between autism and psychopathy, Rogers et al. (2006) proposed that while individuals can have both autism and psychopathy, the two are likely to be distinct and separate, which was referred to as the “double hit” hypothesis; that is, those with both autism and psychopathy have two separate conditions which represent a “double-hit.” Subsequent genetic and experimental data appear aligned with this hypothesis (e.g., Jones et al. 2010; O’Nions et al. 2014, 2015). In their systematic review, Maguire et al. (2024) investigated the relationship between autism and psychopathy. Across the included studies, a heightened prevalence of psychopathic traits among autistic individuals or those exhibiting high autistic traits was observed compared to the general population. But, there was evidence to support the view that cognitive empathy tends to be impaired among those with autism (e.g., Jones et al. 2010; Klapwijk et al. 2016), while affective empathy tends to be impaired among those scoring high on measures of psychopathy (e.g., Lockwood et al. 2013; Oliver et al. 2016), lending weight to the argument that autism and psychopathy are distinct. Those with both autism and psychopathy presented with difficulties with both cognitive and affective empathy, which has implications for risk assessment.

Maguire et al. (2024) also pointed out that there is a lack of validated measures for identifying psychopathic traits within autistic individuals. This is relevant, as both autism and psychopathy are associated with difficulties with empathy and measures of psychopathy should capture features of psychopathy, and not autism. Incorrectly labeling an autistic person as having psychopathy will cause substantial harm. Therefore, robust methods for measuring psychopathy with autistic individuals are needed to help researchers further develop an understanding of the relationship between autism and psychopathy, while

enabling clinicians to accurately assess clinical risk and develop risk mitigation strategies (Alexander et al. 2016; Barnoux et al. 2020; Melvin et al. 2017). Barnoux et al. (2020) argued that autistic individuals who have psychopathy are likely to present with an increased risk of committing crimes and require longer stays within forensic psychiatric and/or criminal justice settings due to the nature or degree of this risk. It is important that these risks and treatment needs are accurately identified among autistic people.

The Psychopathy Checklist—Revised (PCL-R) (Hare 1991) is one measure of psychopathy which is a widely used as a risk assessment tool within the CJS (Hare 2016) but can be lengthy to complete. Therefore, the shorter Psychopathy Checklist—Screening Version (PCL:SV) was derived (Hart et al. 1995). The PCL:SV is reported to be the strongest predictor of violence in psychiatric inpatient units when compared to other risk assessment tools (Doyle et al. 2002), and there is evidence that it predicts future violence amongst adults with intellectual disabilities (Gray et al. 2007), better than the PCL-R (Morrissey, Mooney, et al. 2007).

There is also evidence to indicate that psychopathy is associated with increased personality disorder symptomatology (Coid and Yang 2008; Coid and Ullrich 2010), including symptoms or a diagnosis of antisocial, histrionic, and borderline personality disorder, as well as paranoid personality disorder (Bergström et al. 2024). There is also evidence that autism and personality disorder may share overlapping features (e.g., Dudas et al. 2017) which may make accurate diagnosis challenging (Rinaldi et al. 2021). Considering antisocial personality disorder, Murphy (2006) reported that autistic inpatients and inpatients with schizophrenia performed more poorly on theory of mind tests than those with dissocial or borderline personality disorder within a high secure hospital, suggesting that those with autism or schizophrenia may have additional difficulties with perspective-taking. However, little is known about the relationship between personality disorder, autism, and psychopathy.

Within the current study, to investigate the reliability and validity of the PCL:SV when used with autistic inpatients detained within psychiatric hospitals, we captured data at two time points separated by 12-months. We initially examined whether the PCL:SV was associated with length of hospital stay, criminal history, violence offenses, forensic history, and diagnosis of personality disorder. Second, we investigated predictive validity by examining if the PCL:SV predicted: (a) moves across secure wards, and (b) aggressive or problematic behavior 12-months later. Finally, we investigated convergent validity by determining whether the PCL:SV was associated with other measures of clinical risk.

1 | Method

1.1 | Transparency and Openness

The study design and analysis were not preregistered. Data, analysis code, and research materials are not available. Data were analyzed using R (R Core Team 2023) and the package ggplot, version 3.2.1 (Wickham 2016).

1.2 | Design and Setting

This study utilized a prospective cohort design with two measurement points, separated by 12 months. Fifty-nine inpatient hospitals across 26 NHS Trusts and 7 hospitals from independent healthcare providers in England and Wales took part in this study. Sites were comprised of 22 low secure units, 13 medium secure units, two high secure units (low, medium and high secure units vary according to physical and procedural security and are predominantly for those with convictions, or behaviors likely to be criminal), 11 assessment and treatment units (for rehabilitation, acute mental health, behavioral problems), 12 locked hospital units (for rehabilitation, acute mental health, psychiatric intensive care or step-down services) and six open hospital units (for acute admissions, psychiatric services, and specialist residential services).

1.3 | Participants

Clinicians were asked to provide data about all their inpatients who met our eligibility criteria. Data were captured about 282 participants, who at the time of data collection were detained under the Mental Health Act, 1983, and/or subject to the Mental Capacity Act, 2005. All participants had a diagnosis of autism, including 251 males, 30 females, and one transgender person. Age ranged from 18 to 67 years, $M = 33.29$; $SD = 11.70$. The majority identified as Caucasian (88.6%), followed by mixed race (5%), Black African or Black Caribbean (4%), Asian (2%), and Chinese (0.4%). At baseline enrollment, most were single (98%), four were in relationships (1%), one participant was divorced (0.4%) and the majority did not have children (98%). Data about marital status was missing for one participant. Over half the sample had attended special educational needs schools (57%) and 43% were educated in mainstream schools. Data about educational status were missing for two participants. Forty-nine percent of the sample also had a diagnosis of intellectual disability. Regarding autism diagnosis, 47% had a diagnosis of childhood autism, 12% had a diagnosis of atypical autism, 39% had a diagnosis of Asperger syndrome, and 2% were diagnosed with pervasive developmental disorder-not otherwise specified. Fifty-one percent were detained under forensic sections, 44% were detained under civil sections as defined within the Mental Health Act, 1983, and a further 5% of participants were detained under the Mental Capacity Act, 2005. Data on section type was missing for one participant.

1.4 | Eligibility Criteria

Individuals were eligible for inclusion in the study if they were aged 18 years or older, had an ICD-10 diagnosis of an autism spectrum disorder made by a Clinical Psychologist, Psychiatrist, or other appropriately qualified professional, and were detained within a hospital using the Mental Health Act, 1983, or subject to the Mental Capacity Act, 2005. There were no specified exclusion criteria.

1.5 | Materials and Measures

A range of file-based information was collated from clinical records and staff working with the participants, including data on hospital admissions, an individual's detention and ward security level, behavioral factors, forensic history (a history of committing crime), and data on diagnoses of personality disorders as recorded by clinicians at the site. This information was anonymized at the site by clinicians.

1.6 | Aggressive/Problematic Behavior

Information regarding the frequency and type of aggressive or problematic behavior exhibited by participants was recorded by clinicians over the preceding 12-week period prior to 12-month census date. Clinicians were asked to document all instances of aggression during this period. Recorded data were sorted into eight frequency categories, as well as "overall presence/absence" of behaviors, and a category specifying if there was any evidence of clear violent intent exhibited by the participants across these behaviors, Table 1. Data were

TABLE 1 | Definitions of aggression or problematic behavior categories.

Physical aggression	Behaviors that lead to physical harm, such as hitting others.
Verbal aggression	Behaviors where individuals were verbally aggressive towards others, such as shouting or racial abuse.
Sexual behavior	Behaviors deemed inappropriately sexual in nature, such as masturbating in public.
Violence to self	Behaviors that led to self-injury, such as cutting or head banging.
Rule breaking	Behaviors that violated rules of the forensic mental health setting, such as absconding.
Threats of violence/aggression	Behaviors where individuals verbally threatened others, such as threatening to kill others.
Intimidating behavior	Behaviors where participants were physically threatening others through body language, such as raising fists.
Inappropriate behavior	Behaviors not considered socially acceptable behaviors, such as spitting/public defecation.
Overall presence	Overall presence of all recorded aggressive/problematic behaviors (Y/N).
Violent intent	Was there evidence of clear violent intent for behaviors? (Y/N)

categorized independently by two researchers and any discrepancies were discussed within the research team until consensus was reached, $k = 1$.

1.7 | The Psychopathy Checklist: Screening Version

The PCL:SV is a 12-item, two-factor tool designed for screening psychopathic traits and behaviors across forensic and non-forensic populations in individuals aged 16 years and older (Brazil and Forth 2016). Factor 1 assesses the interpersonal and affective features of psychopathy, such as deceitfulness, grandiosity, and lack of remorse and empathy, and Factor 2 assesses the socially deviant or antisocial behavior associated with psychopathy, such as impulsiveness and poor behavioral control. Items are scored on a three-point scale according to lifetime presence and severity of symptoms (0 = absent, 1 = possibly or partially present, and 2 = present). Individuals scoring 18 or over are considered as “psychopathic” and those scoring 13–17 are considered “maybe psychopathic” (Hart et al. 1995).

The PCL:SV was completed by senior clinicians (e.g., psychiatrist or psychologist) with responsibility for providing inpatient care. Each senior clinician had experience of interviewing the individual patient and was familiar with their history and records, including their diagnosis. While some clinicians had prior training in the PCL-R, all clinicians, regardless of experience, attended a 1-day group training programme for this study, which included training in the PCL-SV assessment procedure and practice scoring. This is not dissimilar from previous studies, including the MacArthur Violence Risk Assessment Study (Skeem and Mulvey 2001).

1.8 | Historical, Clinical, and Risk Management Tool

The HCR-20^{V3} (Douglas et al. 2013) is a 20-item tool to assess the risk of violence in 18–65-year-olds, containing three subscales: historical (10 items), clinical (5 items) and risk management (5 items), accounting for past, present, and future risk factors. Items are scored on a three-point scale (0 = absent, 1 = possibly or partially present, and 2 = definitely present) and a final summary rating of low, moderate, or high risk for violence is given. Although no cut-off points are provided, it is generally considered that the more risk factors present, the greater the risk. The HCR-20^{V2} has been subject to more than 200 empirical validations demonstrating its effectiveness as a risk assessment tool (Douglas et al. 2014) and versions 2 and 3 are strongly correlated (0.69–0.90) (Douglas and Belfrage 2014). The HCR-20^{V3} is routinely completed within secure services within England; the most recently completed version was taken from electronic patient records.

1.9 | Short-Term Assessment of Risk and Treatability (START)

The START (Webster et al. 2004) is a 20-item tool used to evaluate short-term risk of aggression, in individuals aged 16 and

above with psychiatric disorders. It assesses an individual's strengths and vulnerabilities, with items rated on a three-point scale (0 indicates no vulnerability/strength evident, 1 indicates moderate vulnerability/strength, and 2 indicates high vulnerability/strength). Raters then provide an overall risk rating (low, moderate, or high) about the likelihood of seven risk outcomes occurring: violence to others, self-harm, suicide, substance abuse, victimization, self-neglect, and unauthorized absence. Predictive validity of aggression has been demonstrated (Braithwaite et al. 2010; O'Shea et al. 2016). The START was also routinely completed by some included sites, and where this was the case, the most recent version was taken from electronic patient records. Where this was not the case, clinicians attended a 1-day group training programme for this study, which included training in the START assessment procedure and practice scoring.

1.10 | Analysis

All analyses were completed using R statistical software (R Core Team 2023). Descriptive statistics and the reliability coefficients, McDonald's Omega ($\omega =$) and Cronbach's alpha ($\alpha =$) were initially calculated. A confirmatory factor analysis was conducted on the PCL:SV items to determine if the two factor solution presented in the manual was appropriate for this sample, which was the case (Hart et al. 1995). All analyses were run using PCL:SV Total Scores, Factor 1, and Factor 2 separately due to collinearity. Data were not normally distributed, and alternative and appropriate tests were chosen.

For construct validity, regression was used to examine the relationship between the PCL:SV, length of time spent in hospital, and criminal offending. History of criminal offending was captured in two ways: (1) current convictions, cautions, and reprimands, which are a count of those associated with only the current hospital admission, and (2) total convictions, cautions, and reprimands, which are a count of all lifetime convictions, cautions, and reprimands. Negative binomial regressions were used for all count data as the assumptions governing linear and Poisson regression were violated. Age and/or diagnosis of intellectual disability (ID) were included in some models as appropriate covariates. Logistic regression models had appropriate fit and were run with the following binary outcome variables: (a) diagnosis of personality disorder (yes/no), (b) forensic background (contact with the CJS) (yes/no), and (c) Mental Health Act section (civil/forensic).

Multinomial logistic regression was used to examine the relationship between the PCL:SV and any changes in ward location (no change, transferred, discharge) at 12-month follow-up. Logistic regression was used to examine whether any ward transfers were positive or negative as an index of treatment progress at 12-months. Positive change was defined as moving down a security level (e.g., from high or medium to low security). Negative change was defined as moving up a security level (e.g., from low to medium or high security) or no change in security level due to too few cases in the negative change category. McFadden's Pseudo R^2 was reported for models as an approximation to variance explained (Harrell Jr. 2016).

Receiver operating characteristic curve (ROC) analysis was used to determine the predictive validity of the PCL-SV for aggressive or problematic behavior over the preceding 12-weeks at the 12-month follow-up. Values range from 0, a perfect negative classification, 0.5, a completely chance outcome to 1, a perfect prediction (Schmidt et al. 2006). Values below 0.70 are considered poor, 0.70–0.80 moderate, 0.80–0.90 good, and those higher than 0.90 are considered excellent (de Hond et al. 2022).

Correlations were used to measure the strength and direction of the relationship between the PCL:SV, HCR20^{V3}, and START. Finally, a sensitivity analysis for missing data using multiple imputation was conducted to ensure that results were robust to missing data assumptions (see [Supporting Information](#)). Additionally, Little's multivariate test (Little 1988) indicated that data were missing completely at random and unbiased ($p = 0.99$).

1.11 | Ethical Considerations

A favorable ethical opinion was granted by NHS Wales Research Ethics Committee 7 and associated Health Research Authority approval (REC Ref: 15-WA-0246; IRAS: 181659). Data were anonymized at the site prior to being shared with the research team, allowing for the continued confidentiality of participants.

2 | Results

2.1 | Descriptive Statistics and Reliability

Descriptive statistics for all our measures are found within Table 2. Within the sample, 13.48% ($n = 38$) met criteria for psychopathy using the cut-off score of ≥ 18 ; 18.79% ($n = 53$) met criteria for “maybe psychopathic” using the cut-off score of 13–17; the remaining 58.51% ($n = 165$) were categorized as “non

TABLE 2 | Descriptive statistics for included variables.

	<i>n</i>	<i>M</i>	<i>SD</i>	<i>Min–max</i>	<i>Frequency</i>
PCL:SV Total	257	10.70	5.83	0–24	—
Factor 1	257	4.81	3.23	0–12	—
Factor 2	256	5.89	3.30	0–12	—
CLoS	282	928.00	1534.23	1–17,934	—
Total days spent in SPC	271	2694.07	3332.57	7–20,805	—
Total previous CCR	271	4.46	16.96	0–235	—
Total current CCR	280	1.94	3.50	0–33	—
Total violent offenses	251	2.43	4.41	0–42	—
Forensic background	281	—	—	—	No—113 Yes—168
Diagnosis of PD	282	—	—	—	No—228 Yes—54
Diagnosis of LD	282	—	—	—	No—145 Yes—137
Mental health section	267	—	—	—	Forensic—144 Civil—123
Physical aggression	176	1.49	5.12	0–52	Present: $n = 60$ (34%)
Verbal aggression	176	1.41	2.66	0–17	Present: $n = 82$ (47%)
Sexual behavior	176	0.41	1.24	0–10	Present: $n = 32$ (18%)
Violence to self	176	0.59	3.23	0–38	Present: $n = 26$ (15%)
Rule breaking	176	0.70	2.04	0–15	Present: $n = 51$ (29%)
Threats of violence/aggression	176	0.75	2.44	0–20	Present: $n = 40$ (23%)
Intimidating behavior	176	0.95	2.86	0–22	Present: $n = 47$ (27%)
Inappropriate behavior	176	0.89	3.58	0–31	Present: $n = 31$ (18%)
Total frequency	176	7.19	12.93	0–84	Present: $n = 126$ (72%)
Violent intent	178	—	—	—	Present: $n = 100$ (56%)

Abbreviations: CCR = convictions, cautions, and reprimands, CLoS = current length of stay (days), PD = personality disorder, SPC = secure psychiatric care.

psychopathic.” Comparisons between these three groups across variables are found within Table 6 in [Supporting Information](#). PCL:SV data were missing for 26 participants. Structural reliability estimates indicated good to excellent reliability for the PCL:SV Total Score, $\omega=0.97$, 95% CI [0.93; 1.00], and $\alpha=0.87$, 95% CI [0.84; 0.89]. For Factor 1, the reliability estimates were also good to excellent, $\omega=0.93$, 95% CI [0.88; 0.98], $\alpha=0.83$, 95% CI [0.80; 0.86]. For Factor 2, the reliability estimates were satisfactory to good, $\omega=0.88$, 95% CI [0.82; 0.93], and $\alpha=0.77$, 95% CI [0.72; 0.81].

A confirmatory factor analysis was conducted on our data to confirm that a two-factor structure was appropriate for this sample. We found satisfactory model fit for our two-factor model after introducing residual covariances, $\chi^2(66)=2417.16$, $p<0.001$, CFI=0.98, TFI=0.96, RMSEA=0.071, and SMSR=0.067, Figure 1.

2.2 | Construct Validity

Controlling for age and diagnosis of intellectual disability, PCL:SV Total Score, Factor 1, and Factor 2 significantly and positively predicted current length of stay, total days spent in mental health hospitals, and total previous convictions, cautions, and reprimands, Table 3a. However, controlling for age and diagnosis of intellectual disability, PCL:SV Total Score, Factor 1, and Factor 2 did not significantly predict total *current* convictions, cautions, and reprimands; being older and having an intellectual disability was associated with fewer current convictions, cautions, and reprimands, Table 3a. PCL:SV Total Score, Factor 1, and Factor 2 significantly predicted the total number of violence offenses in the positive direction; having an intellectual disability was associated with fewer violent offenses, Table 3a. Results from missing data analysis were similar (see [Supporting Information](#)).

Increasing PCL:SV Total and Factor 1 was associated within significantly increased odds of having a forensic background, while Factor 2 was associated with significantly decreased odds of having a forensic background, Table 3b. A one-point increase in PCL:SV Total significantly increased the chances of having a forensic background by 1.06 times, or 6%, and a one-point increase in Factor 1 significantly increased the chances of

having a forensic background by 1.27 times, or 27%. However, the opposite was the case for Factor 2. With each of these models, having an intellectual disability was associated with a reduced odds of having a forensic background. Both PCL:SV Total and Factor 1 was significantly associated with increased odds of having a diagnosis of a personality disorder. A one-point increase in PCL:SV Total and Factor 1 significantly increased the chances of having a personality disorder diagnosis by 1.12 times, or 12%, and 1.30 times, or 30%, respectively. Factor 2 did not significantly increase the odds of having a personality disorder diagnosis. Turning to consider the type of Mental Health Act section, only Factor 1 and Factor 2 were associated with the odds of being detained under a forensic or civil section, in opposite directions. A one-point increase in Factor 1 was significantly associated with a 1.21 times (21%) greater chance of being detained under a forensic section, while Factor 2 was significantly associated with increased odds of being detained under a civil section. For both these models, having an intellectual disability was associated with increased odds of being detained under a civil section. All results remained consistent with the exception of the relationship between Factor 2 and the type of Mental Health Section, which was no longer significant, following our missing data sensitivity analysis (see [Supporting Information](#)).

2.3 | Predictive Validity: Transfer and Discharge From Secure Wards

PCL:SV Total score was a significant predictor of ward location 12-months later, Table 3c. These findings indicated that higher scores were associated with decreased odds of being discharged, relative to no change. Factor 2 was associated with a decreased odds of transfer to another ward, relative to no change, Table 3c.

There was a significant positive relationship between PCL:SV Total Score, Factor 1, and Factor 2 and the probability of experiencing no change or a negative change in ward security level at 12-months, Table 3d. A one-point increase in PCL:SV Total, Factor 1, and Factor 2 significantly increased the chances of experiencing a negative change or no change in ward security by 1.10 times (10%), 1.16, (16%), and 1.20 (20%) respectively. For the model including PCL:SV Total Score, and Factor 2, having an intellectual disability was associated with increased odds of experienced a positive change in ward security 12-months




FIGURE 1 | Path diagram for two factor confirmatory factor analysis.

TABLE 3 | (a) Negative binomial regression: The relationship between PCL:SV Total and Factor scores and length of stay and criminal offending history. (b) Logistic regression: PCL:SV and forensic background, personality disorder, and Mental Health Act section. (c) Multinomial logistic regression: PCL:SV and ward location at 12 months follow up. (d) Logistic regression: PCL:SV and changes in security ward at 12 months.

Panel (a)	β	SE	z	p	IRR (95% CI)	Pseudo R ² (McFaddon)
CLOS						
Model 1						
Age	0.02	0.01	2.806	<0.01***	1.02 [1.00–1.03]	0.0051
PCL:SV Total	0.04	0.01	3.6	<0.001**	1.04 [1.02–1.07]	
Intellectual disabilities	0.12	0.13	0.87	0.38	1.12 [0.86–1.46]	
Model 2						
Age	0.02	0.01	2.99	<0.01**	1.02 [1.01–1.03]	0.0041
Factor 1	0.06	0.02	2.95	<0.01**	1.06 [1.02–1.11]	
Intellectual disabilities	0.17	0.14	1.22	0.22	1.18 [0.9–1.54]	
Model 3						
Age	0.02	0.01	2.782	<0.01**	1.02 [1.00–1.03]	0.0053
Factor 2	0.07	0.02	3.519	<0.001***	1.07 [1.03–1.12]	
Intellectual disabilities	0.09	0.13	0.65	0.52	1.09 [0.84–1.42]	
Model 1						
Age	0.05	0.01	10.169	<0.001***	1.06 [1.04–1.07]	0.022
PCL:SV Total	0.04	0.01	3.51	<0.001***	1.04 [1.02–1.06]	
Intellectual disabilities	–0.09	0.13	–0.73	0.46	0.91 [0.71–1.17]	
Model 2						
Age	0.06	0.01	10.378	<0.001***	1.06 [1.05–1.07]	0.022
Factor 1	0.07	0.02	3.59	<0.001***	1.07 [1.03–1.12]	
Intellectual disabilities	–0.04	0.13	–0.3	0.77	0.96 [0.75–1.24]	
Model 3						
Age	0.06	0.01	10.248	<0.001***	1.06 [1.05–1.07]	0.022
Factor 2	0.06	0.02	3.07	<0.01**	1.06 [1.02–1.10]	
Intellectual disabilities	–0.1	0.13	–0.77	0.44	0.91 [0.71–1.16]	

(Continues)

TABLE 3 | (Continued)

Panel (a)	β	SE	z	p	IRR (95% CI)	Pseudo R ² (McFaddon)
Total previous CCR						
Model 1						0.036
Age	0.03	0.01	2.841	<0.01**	1.03 [1.01–1.06]	
PCL:SV Total	0.12	0.02	4.921	<0.001***	1.12 [1.07–1.18]	
Intellectual disabilities	–0.03	0.27	–0.12	0.91	1.03 [0.6–1.77]	
Model 2						0.035
Age	0.03	0.01	2.4235	0.02*	1.03 [1.01–1.05]	
Factor 1	0.21	0.04	4.8295	<0.001***	1.23 [1.13–1.34]	
Intellectual disabilities	0.08	0.28	0.29	0.77	1.09 [0.63–1.87]	
Model 3						0.029
Age	0.04	0.01	3.385	<0.001***	1.04 [1.02–1.07]	
Factor 2	0.17	0.04	4.048	<0.001***	1.19 [1.09–1.29]	
Intellectual disabilities	0.07	0.28	0.25	0.81	1.07 [0.62–1.85]	
Total current CCR						
Model 1						0.023
Age	–0.021	0.01	–2.01	0.04*	0.98 [0.97–1.00]	
PCL:SV Total	0.01	0.02	0.932	0.35	1.01 [0.98–1.04]	
Intellectual disabilities	–0.77	0.18	–4.36	<0.001***	0.46 [0.33–0.65]	
Model 2						0.025
Age	–0.021	0.01	–3.9	0.046*	0.98 [0.97–1.00]	
Factor 1	0.045	0.03	1.4678	0.1407	1.04 [0.99–1.1]	
Intellectual disabilities	–0.74	0.18	–4.15	<0.001***	0.48 [0.34–0.68]	
Model 3						0.023
Age	–0.021	0.01	–2.04	0.046*	0.98 [0.97–1.00]	
Factor 2	0.004	0.03	0.15	0.8863	1 [0.95–1.06]	
Intellectual disabilities	–0.78	0.18	–4.4	<0.001***	0.46 [0.33–0.65]	

(Continues)

TABLE 3 | (Continued)

Panel (a)	β	SE	z	p	IRR (95% CI)	Pseudo R ² (McFaddon)
Total violent offenses						
Model 1						
Age	0	0.01	0.134	0.69	1.00 [0.98–1.02]	0.012
PCL:SV Total	0.05	0.02	2.4557	0.01*	1.05 [1.01–1.09]	
Intellectual disabilities	–0.55	0.22	–2.49	0.01*	0.58 [0.37–0.89]	
Model 2						
Age	0	0.01	0.00422	0.9982	1.00 [0.98–1.02]	0.011
Factor 1	0.079	0.03	2.518	0.03*	1.08 [1.01–1.15]	
Intellectual disabilities	–0.52	0.23	–2.31	0.02*	0.59 [0.38–0.92]	
Model 3						
Age	0.001	0.01	0.2753	0.796	1.00 [0.98–1.02]	0.011
Factor 2	0.07	0.03	2.03	0.04*	1.07 [1.00–1.14]	
Intellectual disabilities	–0.6	0.22	–2.7	<0.01**	0.55 [0.36–0.85]	
Panel (b)						
Forensic background						
PCL:SV Total	0.06	0.02	2.658	0.01*	1.06 [1.02–1.11]	0.069
Intellectual disabilities	–1.07	0.27	–3.98	<0.001***	0.34 [0.20–0.58]	0.1
Factor 1	0.248	0.06	3.92	<0.001***	1.27 [1.13–1.44]	
Factor 2	–0.11	0.06	–1.98	0.047*	0.90 [0.80–1.00]	
Intellectual disabilities	–0.88	0.28	–3.14	<0.01*	0.42 [0.24–0.72]	
Diagnosis of PD						
PCL:SV Total	0.11	0.03	3.9285	<0.001***	1.12 [1.06–1.18]	0.069
Intellectual disabilities	0.43	0.33	1.33	0.18	1.54 [0.82–2.96]	
Factor 1	0.262	0.076	3.8348	<0.001***	1.30 [1.14–1.49]	0.093
Factor 2	–0.04	0.07	–0.62	0.53	0.96 [0.84–1.09]	
Intellectual disabilities	0.7	0.35	1.98	0.05	2.00 [1.02–4.07]	

(Continues)

TABLE 3 | (Continued)

Panel (b)	β	SE	z	p	OR (95% CI)	Pseudo R^2 (McFaddon)	
Mental Health Act section	0.04	0.02	1.6	0.109	0.96 [0.92–1.01]	0.045	
PCL:SV Total	-0.91	0.27	-3.43	<0.001***	0.40 [0.24–0.67]		
Intellectual disabilities	0.19	0.06	3.3	0.00**	1.21 [1.08–1.36]	0.074	
Factor 1	-0.11	0.05	-2.04	0.04*	0.89 [0.80–0.99]		
Factor 2	-0.74	0.28	-2.66	0.01*	0.48 [0.28–0.82]		
Intellectual disabilities							
Panel (c)	Location	β	SE	z	p	OR (95% CI)	Pseudo R^2 (McFaddon)
Model 1	PCL:SV Total	-0.01	0.03	-0.43	0.67	0.99 [0.93–1.04]	0.051
	Discharged	-0.13	0.03	-4.34	<0.001***	0.87 [0.82–0.93]	
	Transferred	-0.23	0.33	-0.68	0.5	0.80 [0.42–1.53]	
	Discharged	0.38	0.32	1.18	0.24	1.46 [0.78–2.73]	
Model 2	Transferred	-0.13	0.07	-1.85	0.06	0.98 [0.87–1.12]	0.052
	Discharged	-0.01	0.07	-0.27	0.79	0.88 [0.77–1.01]	
	Transferred	-0.14	0.07	-2.12	0.03*	0.99 [0.87–1.13]	
	Discharged	-0.008	0.07	-0.12	0.9	0.87 [0.77–0.99]	
	Transferred	0.36	0.33	1.06	0.29	0.78 [0.40–1.53]	
	Discharged	-0.25	0.35	-0.73	0.47	1.43 [0.74–2.74]	
Panel (d)	β	SE	z	p	OR (95% CI)	Pseudo R^2 (McFaddon)	
Changes in security ward	0.1	0.03	3.06	<0.01**	1.10 (1.04–1.18)	0.061	
PCL:SV Total	-0.061	0.35	-1.75	0.08	0.55 [0.27–1.07]		
Intellectual disabilities	0.145	0.06	2.465	0.01*	1.16 (1.03–1.30)	0.044	
Factor 1	-0.49	0.35	-1.42	0.16	0.61 [0.31–1.20]		
Intellectual disabilities	0.187	0.06	3.1904	<0.01**	1.20 (1.08–1.35)	0.065	
Factor 2	-0.71	0.35	-2.01	0.04*	0.49 [0.24–0.97]		
Intellectual disabilities							

Note: Reference categories logistic regressions (PCL:SV and characteristics associated with psychopathy) = No forensic background, no diagnosis of PD, forensic section. Location in secure psychiatric hospital based on changes from location at baseline (no change, transferred, discharged). Reference category for logistic regression (changes in security ward) = no change. Significance level, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Abbreviations: CCR = convictions, cautions, and reprimands, CLoS = current length of stay (days), IRR = incidence rate ratio, OR = odds ratio, PD = personality disorder, SE = standard error, SPC = secure psychiatric care.

later. All results remained consistent following missing data sensitivity analysis, with the exception that intellectual disability was not related to changes in ward security (see [Supporting Information](#)).

2.4 | Predictive Validity

In the 12-weeks prior to the 12-month follow-up, physical and verbal aggression were the most frequent types of aggressive/problematic behavior, while sexual behavior was the least frequent, Table 2. The PCL:SV (Total and Factor Scores) had an area under the curve (AUC) of less than 0.70 for all aggressive and problematic behaviors, indicating overall poor discriminatory ability (de Hond et al. 2022), Table 4. Despite this, the AUC estimates and logistic regression results were statistically significant for several specific behaviors. Factor 2 was the best predictor of aggressive or problematic behaviors, and AUC values showed that it was a significant predictor for all specified behaviors. The PCL:SV Total Score had a statistically significant AUC for all specified behaviors except physical aggression. Factor 1 significantly predicted all problematic behaviors with the exception of physical aggression and inappropriate behaviors. Some AUC results did not survive missing data analysis, with the AUC for Factor 2 and physical aggression changing from 0.59 ($p=0.03$) to 0.57 ($p=0.07$); however, changes were small and did not result in any other significant results becoming non-significant. The relationship between PCL:SV Total Score, Factor 1, and Factor 2 did not remain significant for Threats of Aggression and Intimidating Behavior following missing data analyses (see [Supporting Information](#)).

2.5 | Convergent Validity

There was a moderate degree of correlation between PCL:SV Total and HCR-20 Total Scores, with weaker correlations between Factor 1 and Factor 2, and HCR-20 Total Scores, Table 5. All correlations between PCL:SV Total and Factor Scores and Total and subscale scores on the HCR-20 were positive and significant, $p < 0.001$, except for the relationship between PCL:SV Total and the Imminent Violence subscale, $r = 0.19$, $p < 0.01$, as well as Factor 1 and the Imminent Violence subscale, $r = 0.14$, $p = 0.02$. Correlations between the PCL:SV and the START risk assessment revealed a moderate degree of correlation between the PCL:SV Total and Factor Scores and the Vulnerabilities subscale only, which were all significant, $p < 0.001$. No significant relationships between the PCL:SV Total or Factor Scores and the START Strengths subscale were observed. However, when performing missing data analysis, significant negative relationships between PCL:SV Total and Factor 2, and the START Strength subscale were observed, as would be expected.

3 | Discussion

Instruments like the PCL:SV are often used in populations that differ from those in which the measure was developed. This can be problematic, leading to invalid assessments and is particularly problematic within the CJS where labeling an

individual as a psychopath can influence decision-making and foster punitive actions (Blais and Forth 2014). Within this study, we aimed to explore the factor structure, reliability, and validity of the PCL:SV in autistic adults within psychiatric hospitals, a majority of whom were within forensic psychiatric hospitals. Their mean PCL:SV Total Score was similar to or slightly lower than that reported within studies that included forensic inpatients without autism (Bo et al. 2019; Ho et al. 2009; Pedersen et al. 2010), but higher than that reported within studies using non-forensic non-autistic inpatients (Doyle et al. 2002; Nicholls et al. 2004), including those within the MacArthur Violence Risk Assessment Study (Skeem and Mulvey 2001).

Turning to consider our factor analysis, we were able to fit a two-factor model that was similar to that originally reported for the PCL:SV (Hart et al. 1995). This model was chosen due to its similarity to the original two-factor PCL:SV structure and the associated scoring as found within the published manual. However, we included residual covariance between some variables within each factor to ensure a good fit. It is worth noting that there is some debate within the literature as to whether psychopathy is best represented by a two- or three-factor model. For example, Cooke and Michie (2001) proposed a three-factor model of psychopathy using PCL-R data incorporating: (a) arrogant and deceitful interpersonal style (glibness, superficial charm, grandiosity, pathologically lying, and conning/manipulative), (b) deficient affective experience (shallow affect and callousness, lack of empathy, lack of remorse, and failure to accept responsibility), and (c) impulsive and irresponsible behavior style (need for stimulation/boredom, impulsivity, irresponsibility, parasitic lifestyle, and lack of realistic goals). While our goal was not to test the validity of a two-factor versus a three-factor model of psychopathy using PCL:SV data, as there continues to be much debate, it is important to note that others have reported that a three-factor PCL:SV model is superior to a two-factor model using data captured from forensic psychiatric inpatients (Bo et al. 2019). While there is support for a three-factor PCL:SV relative to a unitary, two-factor, and four-factor model in some studies (Veal et al. 2021), this has not been consistent and a single, two-, three-, or four-factor model has been reported. Boduszek et al. (2016) reported a bifactor PCL:SV model with two-factors was superior to nine alternative PCL:SV models. It would be appropriate to undertake a future study investigating PCL:SV measurement invariance within autistic people and alternative and potentially better fitting factor structures.

Further, we found that the PCL:SV had good to excellent reliability and construct validity, particularly the PCL:SV Total and Factor 1 scores. Factor 2 showed a more intricate relationship with psychopathy-related traits, with some relationships reversed. Regarding predictive validity, higher PCL:SV scores were associated with factors indicating poorer treatment progress. Predictive validity for aggressive or problematic behaviors was limited. However, Factor 2 predicted aggressive and problematic behaviors more robustly, potentially influenced by observable behavior associated with autism and/or intellectual disabilities (e.g., behaviors that challenge). Generally, the results aligned with existing research and theoretical understanding of psychopathy and the PCL:SV.

TABLE 4 | Logistic regression and AUC analysis: PCL:SV scores as predictors of aggressive/problematic behaviors at 12 months.

		<i>B</i>	<i>SE</i>	<i>z</i>	<i>p</i>	OR (95% CI)	AUC (95% CI)	AUC <i>p</i>
Physical aggression	PCL:SV Total	0.03	0.03	0.94	0.35	1.03 [0.97–1.08]	0.57 [0.47–0.66]	0.08
	Intellectual disabilities	0.36	0.33	1.08	0.28	1.43 [0.75–2.76]		
	Factor 1	0	0.05	0.06	0.95	1.00 [0.91–1.11]	0.54 [0.45–0.63]	0.19
	Intellectual disabilities	0.33	0.33	0.97	0.33	1.39 [0.72–2.71]		
	Factor 2	0.1	0.05	1.93	0.05	1.10 [1.00–1.22]	0.60 [0.51–0.70]	0.01*
Verbal aggression	PCL:SV Total	0.12	0.03	3.97	<0.001***	1.12 [1.06–1.19]	0.67 [0.59–0.76]	<0.001***
	Intellectual disabilities	0.07	0.33	0.2	0.84	1.07 [0.56–2.04]		
	Factor 1	0.17	0.05	3.31	<0.001***	1.18 [1.07–1.31]	0.64 [0.56–0.73]	<0.01**
	Intellectual disabilities	0.2	0.33	0.6	0.55	1.22 [0.64–2.35]		
	Factor 2	0.22	0.05	4.13	<0.001***	1.25 [1.13–1.39]	0.69 [0.61–0.77]	<0.001***
Sexual behavior	PCL:SV Total	0.1	0.34	2.84	<0.01**	1.10 [1.03–1.18]	0.69 [0.61–0.77]	<0.01**
	Intellectual disabilities	−0.09	0.42	−0.216	0.83	0.91 [0.40–2.06]		
	Factor 1	0.15	0.06	2.4	0.01*	1.16 [1.03–1.30]	0.64 [0.54–0.75]	0.01*
	Intellectual disabilities	0.03	0.42	0.06	0.95	1.03 [0.44–2.36]		
	Factor 2	0.2	0.07	2.97	<0.01**	1.21 [1.07–1.39]	0.68 [0.58–0.77]	<0.01**
Violence towards self	PCL:SV Total	0.04	0.04	1.18	0.24	1.04 [0.97–1.12]	0.65 [0.55–0.75]	0.01*
	Intellectual disabilities	0.99	0.45	2.18	0.03*	2.69 [1.13–6.79]		
	Factor 1	0.03	0.07	0.5	0.62	1.03 [0.91–1.18]	0.63 [0.53–0.72]	0.02*
	Intellectual disabilities	0.98	0.46	2.12	0.03*	2.65 [1.10–6.81]		
	Factor 2	0.11	0.07	1.66	0.1	1.12 [0.98–1.27]	0.65 [0.55–0.76]	<0.01**
Rule breaking	PCL:SV Total	0.07	0.03	2.44	0.02*	1.08 [1.02–1.14]	0.65 [0.57–0.74]	<0.001*
	Intellectual disabilities	0.79	0.35	2.23	0.03*	2.2 [1.11–4.48]		
	Factor 1	0.13	0.05	2.35	0.02*	1.13 [1.02–1.26]	0.65 [0.56–0.73]	0.01*
	Intellectual disabilities	0.9	0.37	2.46	0.01*	2.46 [1.21–5.14]		
	Factor 2	0.13	0.05	2.38	0.02*	1.14 [1.03–1.27]	0.66 [0.57–0.74]	0.001*
Threats of aggression	PCL:SV Total	0.06	0.03	2.01	0.04*	1.06 [1.00–1.13]	0.61 [0.51–0.71]	0.02*
	Intellectual disabilities	0.144	0.37	0.39	0.7	1.16 [0.56–2.40]		
	Factor 1	0.1	0.06	1.79	0.07	1.10 [0.99–1.23]	0.61 [0.51–0.71]	0.02*
	Intellectual disabilities	0.22	0.38	0.59	0.56	1.25 [0.59–2.65]		
	Factor 2	0.12	0.06	2.18	0.03*	1.13 [1.01–1.27]	0.62 [0.52–0.72]	0.01*
	Intellectual disabilities	0.08	0.37	0.22	0.82	1.09 [0.52–2.26]		

(Continues)

TABLE 4 | (Continued)

		<i>B</i>	<i>SE</i>	<i>z</i>	<i>p</i>	OR (95% CI)	AUC (95% CI)	AUC <i>p</i>
Intimidating behavior	PCL:SV Total	0.09	0.03	3.02	<0.01**	1.09 [1.03–1.16]	0.65 [0.55–0.75]	<0.01**
	Intellectual disabilities	−0.38	0.37	−1.05	0.3	0.68 [0.32–1.39]		
	Factor 1	0.12	0.05	2.24	0.01*	1.13 [1.02–1.25]	0.62 [0.51–0.73]	<0.01**
	Intellectual disabilities	−0.3	0.37	−0.8	0.43	0.74 [0.35–1.53]		
	Factor 2	0.19	0.06	3.3	<0.001***	1.21 [1.08–1.36]	0.67 [0.57–0.77]	<0.001***
Inappropriate behavior	Intellectual disabilities	−0.52	0.37	−1.39	0.16	0.60 [0.28–1.22]		
	PCL:SV Total	0.06	0.04	1.6	0.11	1.06 [0.99–1.14]	0.62 [0.49–0.75]	0.02*
	Intellectual disabilities	0.61	0.43	1.42	0.16	1.83 [0.80–4.32]		
	Factor 1	0.06	0.06	0.89	0.37	1.06 [0.93–1.20]	0.58 [0.45–0.71]	0.09
	Intellectual disabilities	0.61	0.43	1.41	0.16	1.85 [0.79–4.42]		
Overall presence	Factor 2	0.16	0.07	2.4	0.02*	1.17 [1.03–1.35]	0.67 [0.54–0.79]	0.01*
	Intellectual disabilities	0.6	0.44	1.38	0.17	1.82 [0.78–4.37]		
	PCL:SV Total	0.09	0.03	2.76	<0.01*	1.09 [1.03–1.16]	0.65 [0.56–0.74]	<0.01**
	Intellectual disabilities	0.51	0.36	1.42	0.16	1.66 [0.83–3.40]		
	Factor 1	0.11	0.06	2.02	0.04*	1.12 [1.01–1.25]	0.62 [0.53–0.72]	<0.01*
Violent intent	Intellectual disabilities	0.57	0.36	1.58	0.12	1.77 [0.88–3.66]		
	Factor 2	0.19	0.06	3.19	<0.01**	1.20 [1.08–1.36]	0.67 [0.58–0.76]	<0.001***
	Intellectual disabilities	0.39	0.36	1.07	0.28	1.47 [0.73–3.01]		
	PCL:SV Total	0.09	0.03	3.02	<0.01**	1.09 [1.03–1.15]	0.66 [0.57–0.74]	<0.001***
	Intellectual disabilities	0.58	0.33	1.78	0.08	1.78 [0.95–3.42]		
Violent intent	Factor 1	0.09	0.05	1.89	0.06	1.10 [1.00–1.21]	0.61 [0.53–0.70]	<0.01*
	Intellectual disabilities	0.6	0.33	1.84	0.07	1.83 [0.97–3.52]		
	Factor 2	0.2	0.05	3.79	<0.001***	1.22 [1.11–1.36]	0.61 [0.61–0.77]	<0.001***
	Intellectual disabilities	0.47	0.33	1.43	0.15	1.60 [0.84–3.08]		

Note: Reference category = behavior not present. Significance level, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Abbreviations: CI = confidence intervals, SE = standard error.

3.1 | Reliability

Reliability analyses revealed robust estimates of internal consistency for the PCL:SV, with McDonald's Omega indicating excellent structural reliability. Reliability estimates were lowest for Factor 2, similar to prior research in different populations (Hart et al. 1995; Rogers et al. 2000; Žukauskienė et al. 2010). This may represent potential variability across populations, with different groups of people exhibiting a wider variation in the lifestyle/behavioral characteristics measured by Factor 2. Alternatively, this could be due to increased difficulty in measuring these behavioral characteristics.

3.2 | Construct Validity

Positive associations between the PCL:SV and variables that are thought to be associated with psychopathy were found, including length of stay, admissions, previous convictions, and

the number of violent offenses. Factor 1 was associated with an increased likelihood of having a forensic background, more violent offenses, and detention under a forensic section of the Mental Health Act, 1983. Factor 2 was also associated with having more violent offenses. This aligns with previous research on psychopathy, which has long been associated with criminal and violent behavior (Dhingra and Boduszek 2013; Hare 1999), longer offending trajectories, and an increased risk of re-institutionalization after release from prison (Porter et al. 2001).

PCL:SV Total and Factor 1 scores were significantly and positively related to having a personality disorder. Antisocial personality disorder (ASPD) is considered the closest diagnostic category to psychopathy (Ogloff 2006). Indeed, some research suggests that psychopathy is at the extreme end of ASPD (Coid and Ullrich 2010), with most psychopaths also meeting ASPD criteria (Hare 1996). No relationship was observed between Factor 2 and personality disorder. Factor 1 encompasses “core” personality features of psychopathy, capturing characteristics

TABLE 5 | Correlations between PCL:SV Total and Factor scores and HCR20 and START measures of clinical risk.

risk assessment tool	PCL:SV Total	PCL:SV Factor 1	PCL:SV Factor 2
HCR20			
Historical scale	$r=0.37, p<0.001^{***}$	$r=0.37, p<0.001^{***}$	$r=0.35, p<0.001^{***}$
Clinical scale	$r=0.41, p<0.001^{***}$	$r=0.35, p<0.001^{***}$	$r=0.41, p<0.001^{***}$
Risk management scale	$r=0.38, p<0.001^{***}$	$r=0.35, p<0.001^{***}$	$r=0.37, p<0.001^{***}$
Total score	$r=0.40, p<0.001^{***}$	$r=0.36, p<0.001^{***}$	$r=0.32, p<0.001^{***}$
Serious physical harm	$r=0.27, p<0.001^{***}$	$r=0.21, p<0.001^{***}$	$r=0.30, p<0.001^{***}$
Imminent violence	$r=0.19, p<0.01^{**}$	$r=0.14, p=0.02^*$	$r=0.22, p<0.001^{***}$
Future violence	$r=0.30, p<0.001^{***}$	$r=0.24, p<0.001^{***}$	$r=0.31, p<0.001^{***}$
START			
Strengths	$r=-0.01, p=0.83$	$r=0.06, p=0.35$	$r=-0.06, p=0.34$
Vulnerabilities	$r=0.45, p<0.001^{***}$	$r=0.32, p<0.001^{***}$	$r=0.50, p<0.001^{***}$

Note: Significance level, * $p<0.05$, ** $p<0.01$, *** $p<0.001$.

such as superficiality, deceitfulness, and lack of empathy, which may be more closely related to the central features of personality disorders than Factor 2 characteristics (Swales 2022).

In contrast to Factor 1, Factor 2 was associated with a greater likelihood of being detained under a civil section of the Mental Health Act, 1983, and a reduced likelihood of having a forensic background. In line with this, a systematic review by Collins et al. (2022) reported that autistic people who encounter the CJS are less likely to have a forensic history, with fewer previous convictions than non-autistic offenders. Consideration of the specific behavioral characteristics captured by Factor 2, such as impulsivity and poor behavioral control, this is unsurprising. Those presenting with behaviors that challenge, as seen more commonly amongst those with intellectual disabilities, may be more likely to score on Factor 2. It is worth noting that a substantial proportion (49%) of our participant population had an intellectual disability. Our findings across our models indicated that autistic inpatients with intellectual disabilities were less likely to have current convictions, cautions or reprimands, violent offenses, and a forensic background. They were more likely to be detained under a civil section of the Mental Health Act, and more likely to have had a positive transfer to a ward of lesser security 12-months later. Accurate levels of antisocial behavior may be hard to capture in adults with intellectual disability (Morrissey, Hogue, et al. 2007); carers often show a reluctance to report offending behaviors to the police (McBrien and Murphy 2006) and once reported, individuals are less likely to be charged for antisocial acts (Cockram 2005). It is also noteworthy that approximately 28% of autistic people have comorbid attention-deficit hyperactivity disorder (ADHD; Lai et al. 2019), which is characterized by inattention, hyperactivity, and impulsivity (World Health Organization 2021). Co-morbid ADHD could further contribute towards high scores on Factor 2. It has previously been reported that although the emotional features of psychopathy are not impaired in ADHD, the behavioral features of psychopathy are present (Eisenbarth et al. 2008), suggesting behavioral overlap between Factor 2 and ADHD.

No relationship was found between the PCL:SV and current convictions, cautions or reprimands, which may be due to two reasons, recognizing that those with intellectual disabilities were less likely to have current convictions, cautions or reprimands. First, most participants had current convictions, cautions or reprimands. However, we only had frequency data and not severity data. It may be expected that a proportion of those with higher PCL:SV scores would commit more severe, but less frequent offenses (e.g., murder) relative to those with lower PCL:SV scores. Second, detained individuals have restricted movement, increased supervision, and risk management plans; actions which collectively reduce opportunities to engage in offending behaviors while in hospital.

3.3 | Predictive Validity

PCL:SV Total score was significantly associated with decreased likelihood of being discharged, while Factor 2 was associated with decreased likelihood of transfer. PCL:SV Total and Factor Scores were also associated with increased likelihood of experiencing negative change/no change in security level at 12 months. This is unsurprising given that psychopathic traits are predictive of poor therapeutic progress and risk reduction (Olver et al. 2013), both of which are critical considerations when applying to transfer to a lower security ward (NHS England 2021). Research on the predictive validity of the PCL:SV and treatment progress is sparse. However, similar results have been reported with regards to the predictive validity of the PCL-R in offenders with intellectual disability (Morrissey, Mooney, et al. 2007) and in individuals with personality disorders detained in high-security settings (Tetley et al. 2010). In this study, Factor 1 and Factor 2 were both significant predictors of negative/no change in security level, indicating that both interpersonal and lifestyle factors can inhibit treatment progress. This contrasts to the existing PCL-R research where Factor 1 is reported as a stronger predictor of negative treatment progression (Morrissey, Mooney, et al. 2007; Olver et al. 2013; Tetley et al. 2010). Interpersonal factors such as superficiality and lack of remorse, likely

hinder therapeutic rapport building and insight development, resulting in reduced opportunities for treatment progress (Olver et al. 2013).

Differences in study design may contribute to varied findings regarding the influence of Factor 2. However, it is also necessary to consider the specific needs and behaviors of autistic adults within psychiatric hospitals. Not all patients in secure settings have a forensic background; some are admitted due to behaviors that challenge, which cannot be safely managed within general psychiatric services (Völlm et al. 2018), while not all our participants were detained with forensic psychiatric wards. Behaviors that challenge are common in autistic individuals and include behaviors such as aggression, destructive behavior, and self-injurious or stereotyped behavior (Matson and Rivet 2008). Managing these behaviors can be complex, posing challenges in maintaining a safe environment (Matson et al. 2011). Thus, behaviors that challenge may be driving Factor 2 as a stronger predictor of negative/no change in security level within the current sample.

The predictive validity of the PCL:SV for problematic or aggressive behaviors was explored but was limited. Nonetheless, Factor 2 was associated with significant AUC values for several behaviors and was the more robust predictor, relative to Factor 1, and PCL:SV Total Scores. Again, this may be influenced by the specific behavioral problems exhibited by autistic individuals. As the study looked at the presence versus absence of behaviors, rather than intensity or frequency, it is possible that individuals scoring highly on Factor 2 exhibited high frequency but low intensity behaviors (e.g., shouting and banging) rather than low frequency but higher intensity behaviors (e.g., sexual assault or violent attacks) which may be more associated with psychopathy than autism. Broadening the sample to encompass autistic individuals across different settings may enhance the predictive validity of the PCL:SV as the current sample is biased towards those prone to display a greater frequency of problematic or aggressive behaviors.

3.4 | Convergent Validity

The PCL:SV demonstrated convergent validity with other risk assessment tools. Significant and moderate correlations were observed between the PCL:SV, HCR-20, and START, except for the START strengths scale. Overall, the measures were expected to align but not be highly correlated as they focus on the measurement of different constructs or risk factors. However, it is important to note that there is a sparsity of research validating the HCR-20 and the START as risk assessment tools for use with autistic adults.

3.5 | Clinical Implications

Overall, these findings suggest that the PCL:SV has good reliability and validity when used with autistic adults within psychiatric hospitals. This allows for a more robust assessment of risk, as well as a more effective investigation of the “double hit” hypothesis (Rogers et al. 2006) in future studies. Individuals with both autism and psychopathy represent a small but clinically

significant subgroup of autistic offenders with unique treatment needs (Alexander et al. 2016). Barnoux et al. (2020) suggested that autistic individuals with a comorbid diagnosis of psychopathy present with increased forensic risk, requiring longer stays in secure psychiatric care, while those without a diagnosis of psychopathy may benefit from shorter stays and community-based placements. Validated psychometric tools such as the PCL:SV can aid in the accurate assessment and diagnosis of psychopathy in autistic individuals, ultimately enhancing treatment outcomes and risk management.

3.6 | Strengths and Limitations

The relatively large dataset from a range of inpatient services enhanced the generalizability of the study findings. The confirmatory factor analysis was consistent with the results derived from participants without autism, and missing data imputation showed that the data were largely unbiased, strengthening the integrity of the findings. While the intelligence quotient was unavailable for all participants, we were able to include whether participants had a diagnosis of an intellectual disability within our analysis. Considering limitations, data on aggressive/problematic behavior indexed the frequency of behaviors and not severity. Failure to account for this distinction complicates the differentiation between high-frequency, low-intensity behaviors potentially associated with autism rather than psychopathy (e.g., behaviors that challenge) and may contribute to the reduced predictive validity of the PCL:SV. Further, and arguably, it is possible that some autistic individuals may score on the PCL:SV due to the nature or degree of their autism. We were unable to investigate this possibility explicitly within our study because we did not capture data about the degree of autistic symptomatology. However, as PCL:SV Factor 1 captures difficulties with empathy, it could be argued that Factor 1 should overlap with autism and therefore predict violence poorly, relative to Factor 2. This was not the case as both Factors predicted history of violent offending and future aggressive and problematic behavior. However, Factor 1 was associated with having a forensic background, a diagnosis of personality disorder, and detention under a forensic section of the Mental Health Act, while the reverse was the case for Factor 2. Further, and while it is highly unlikely that those who completed the PCL:SV also diagnosed our participants with autism, we cannot guarantee this to be the case in all instances as we did not capture these data. Finally, the PCL:SV was used by clinicians within the intended setting, but we did not undertake reliability checks, which may have introduced some error; nevertheless, our findings indicated that the PCL:SV is an appropriate measure for use with autistic psychiatric inpatients.

3.7 | Constraints Upon Generality

There is evidence about the validity, reliability, and factor structure of the PCL:SV when used with community and psychiatric participants drawn from various countries (Veal et al. 2021), but no robust evidence supporting the use of the PCL:SV with autistic psychiatric inpatients. Within this study, we aimed to evaluate the performance of the PCL:SV

when used with autistic psychiatric inpatients in England and Wales. A majority were detained in secure psychiatric services, and some had intellectual disabilities, while the majority were men; we sampled from numerous hospitals helping to ensure that our sample is representative at two time points separated by 12-months. Researchers would need to sample in a similar way over a similar time period to generate comparable results. Our findings are limited to autistic psychiatric inpatients in England and Wales as we did not include autistic psychiatric inpatients from other countries, nor did we include autistic people from the community. Further, we did not include autistic prisoners. Our findings do not generalize to these additional autistic populations. We have no reason to believe that the results depend on other characteristics of the participants, materials, or context.

3.8 | Future Research

The findings from this study indicated that the PCL-SV can be used effectively with autistic psychiatric inpatients. Future studies should attempt to disentangle the relationship between autism and psychopathy further.

Ethics Statement

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

References

- Alexander, R., P. E. Langdon, V. Chester, M. Barnoux, I. Gunaratna, and S. Hoare. 2016. "Heterogeneity Within Autism Spectrum Disorder in Forensic Mental Health: The Introduction of Typologies." *Advances in Autism* 2, no. 4: 201–209. <https://doi.org/10.1108/AIA-08-2016-0021>.
- Azevedo, J., M. Vieira-Coelho, M. Castelo-Branco, R. Coelho, and M. Figueiredo-Bragaid. 2020. "Impulsive and Premeditated Aggression in Male Offenders With Antisocial Personality Disorder." *PLoS One* 15, no. 3: e0229876. <https://doi.org/10.1371/journal.pone.0229876>.
- Barnoux, M., R. Alexander, S. Bhaumik, et al. 2020. "The Face Validity of an Initial Sub-Typology of People With Autism Spectrum Disorders Detained in Psychiatric Hospitals." *Autism* 24, no. 7: 1885–1897. <https://doi.org/10.1177/1362361320929457>.
- Bergström, H., D. P. Farrington, and J. W. Coid. 2024. "Relationships Between Psychopathy and DSM Personality Disorders in Men and Women." *International Journal of Offender Therapy and Comparative Criminology* 68, no. 2–3: 278–295.
- Blais, J., and A. E. Forth. 2014. "Potential Labeling Effects: Influence of Psychopathy Diagnosis, Defendant Age, and Defendant Gender on

Mock Jurors' Decisions." *Psychology, Crime & Law* 20, no. 2: 116–134. <https://doi.org/10.1080/1068316X.2012.749473>.

Bo, S., L. Pedersen, K. B. Christensen, and K. Rasmussen. 2019. "Psychopathy in a Forensic Sample—The Factor Structure of the PCL:SV in a Danish Forensic Sample." *Scandinavian Journal of Psychology* 60: 447–455.

Boduszek, D., K. Dhingra, P. Hyland, and A. Debowska. 2016. "A Bifactorial Solution to the Psychopathy Checklist: Screening Version in a Sample of Civil Psychiatric Patients." *Criminal Behaviour and Mental Health* 26: 174–185. <https://doi.org/10.1002/cbm.1956>.

Braithwaite, E., Y. Charette, A. G. Crocker, and A. Reyes. 2010. "The Predictive Validity of Clinical Ratings of the Short-Term Assessment of Risk and Treatability (START)." *International Journal of Forensic Mental Health* 9, no. 4: 271–281. <https://doi.org/10.1080/14999013.2010.534378>.

Brazil, K. J., and A. E. Forth. 2016. "Psychopathy Checklist: Screening Version (PCL:SV)." In *Encyclopedia of Personality and Individual Differences*, 1–4. Springer International Publishing. https://doi.org/10.1007/978-3-319-28099-8_1101-1.

Cleckley, H. 1941. *The Mask of Sanity; an Attempt to Reinterpret the So-Called Psychopathic Personality*. Mosby.

Cockram, J. 2005. "Justice or Differential Treatment? Sentencing of Offenders With an Intellectual Disability." *Journal of Intellectual and Developmental Disability* 30, no. 1: 3–13. <https://doi.org/10.1080/13668250500033177>.

Coid, J., and S. Ullrich. 2010. "Antisocial Personality Disorder Is on a Continuum With Psychopathy." *Comprehensive Psychiatry* 51, no. 4: 426–433. <https://doi.org/10.1016/j.comppsy.2009.09.006>.

Coid, J., and M. Yang. 2008. "The Distribution of Psychopathy Among a Household Population: Categorical or Dimensional?" *Social Psychiatry and Psychiatric Epidemiology* 43: 773–781.

Coid, J., M. Yang, S. Ullrich, A. Roberts, and R. D. Hare. 2009. "Prevalence and Correlates of Psychopathic Traits in the Household Population of Great Britain." *International Journal of Law and Psychiatry* 32, no. 2: 65–73. <https://doi.org/10.1016/j.ijlp.2009.01.002>.

Coid, J., M. Yang, S. Ullrich, et al. 2009. "Psychopathy Among Prisoners in England and Wales." *International Journal of Law and Psychiatry* 32, no. 3: 134–141. <https://doi.org/10.1016/j.ijlp.2009.02.008>.

Collins, J., K. Horton, E. Gale-St Ives, G. Murphy, and M. Barnoux. 2022. "A Systematic Review of Autistic People and the Criminal Justice System: An Update of King and Murphy (2014)." *Journal of Autism and Developmental Disorders* 53, no. 8: 3151–3179. <https://doi.org/10.1007/s10803-022-05590-3>.

Cooke, D. J., and C. Michie. 2001. "Refining the Construct of Psychopathy: Towards a Hierarchical Model." *Psychological Assessment* 13, no. 2: 171.

de Hond, A. A. H., E. W. Steyerberg, and B. van Calster. 2022. "Interpreting Area Under the Receiver Operating Characteristic Curve." *Lancet Digital Health* 4, no. 12: e853–e855. [https://doi.org/10.1016/S2589-7500\(22\)00188-1](https://doi.org/10.1016/S2589-7500(22)00188-1).

Dhingra, K., and D. Boduszek. 2013. "Psychopathy and Criminal Behaviour: A Psychosocial Research Perspective." *Journal of Criminal Psychology* 3, no. 2: 83–107. <https://doi.org/10.1108/JCP-06-2013-0014>.

Douglas, K., and H. Belfrage. 2014. "Interrater Reliability and Concurrent Validity of the HCR-20 Version 3." *International Journal of Forensic Mental Health* 13, no. 2: 130–139. <https://doi.org/10.1080/14999013.2014.908429>.

Douglas, K., S. Hart, C. Webster, and H. Belfrage. 2013. *HCR-20V3: Assessing Risk for Violence—User Guide*. Mental Health, Law, and Policy Institute, Simon Fraser University.

Douglas, K. S., S. D. Hart, C. D. Webster, H. Belfrage, L. S. Guy, and C. M. Wilson. 2014. "Historical-Clinical-Risk Management-20, Version

- 3 (HCR-20 V3): Development and Overview.” *International Journal of Forensic Mental Health* 13, no. 2: 93–108. <https://doi.org/10.1080/14999013.2014.906519>.
- Doyle, M., M. Dolan, and J. McGovern. 2002. “The Validity of North American Risk Assessment Tools in Predicting In-Patient Violent Behaviour in England.” *Legal and Criminological Psychology* 7, no. 2: 141–154. <https://doi.org/10.1348/135532502760274756>.
- Dudas, R. B., C. Lovejoy, S. Cassidy, C. Allison, P. Smith, and S. Baron-Cohen. 2017. “The Overlap Between Autistic Spectrum Conditions and Borderline Personality Disorder.” *PLoS One* 12, no. 9: e0184447.
- Eisenbarth, H., G. W. Alpers, A. Conzelmann, C. P. Jacob, P. Weyers, and P. Pauli. 2008. “Psychopathic Traits in Adult ADHD Patients.” *Personality and Individual Differences* 45, no. 6: 468–472. <https://doi.org/10.1016/j.paid.2008.05.022>.
- Gray, N. S., S. Fitzgerald, J. Taylor, M. J. MacCulloch, and R. J. Snowden. 2007. “Predicting Future Reconviction in Offenders With Intellectual Disabilities: The Predictive Efficacy of VRAG, PCL-SV, and the HCR-20.” *Psychological Assessment* 19, no. 4: 474–479. <https://doi.org/10.1037/1040-3590.19.4.474>.
- Hare, R. 1991. *The Hare Psychopathy Checklist—Revised*. Multi-Health Systems.
- Hare, R. D. 1996. “Psychopathy and Antisocial Personality Disorder: A Case of Diagnostic Confusion.” *Psychiatric Times* 13, no. 2: 39–40.
- Hare, R. D. 1999. “Psychopathy as a Risk Factor for Violence.” *Psychiatric Quarterly* 70, no. 3: 181–197. <https://doi.org/10.1023/a:1022094925150>.
- Hare, R. D. 2016. “Psychopathy, the PCL-R, and Criminal Justice: Some New Findings and Current Issues.” *Canadian Psychology* 57, no. 1: 21–34. <https://doi.org/10.1037/cap0000041>.
- Harrell, F. E., Jr. 2016. *Regression Modeling Strategies*. Springer International Publishing.
- Hart, S., D. Cox, and R. Hare. 1995. *Manual for the Hare Psychopathy Checklist: Screening Version*. Multi-Health Systems.
- Ho, H., L. Thomson, and R. Darjee. 2009. “Violence Risk Assessment: The Use of the PCL-SV, HCR-20, and VRAG to Predict Violence in Mentally Disordered Offenders Discharged From a Medium Secure Unit in Scotland.” *Journal of Forensic Psychiatry & Psychology* 20, no. 4: 523–541.
- Jones, A. P., F. G. E. Happé, F. Gilbert, S. Burnett, and E. Viding. 2010. “Feeling, Caring, Knowing: Different Types of Empathy Deficit in Boys With Psychopathic Tendencies and Autism Spectrum Disorder.” *Journal of Child Psychology and Psychiatry* 51, no. 11: 1188–1197. <https://doi.org/10.1111/J.1469-7610.2010.02280.X>.
- Klapwijk, E. T., M. Aghajani, O. F. Colins, et al. 2016. “Different Brain Responses During Empathy in Autism Spectrum Disorders Versus Conduct Disorder and Callous-Unemotional Traits.” *Journal of Child Psychology and Psychiatry, and Allied Disciplines* 57, no. 6: 737–747. <https://doi.org/10.1111/jcpp.12498>.
- Lai, M., C. Kasse, R. Besney, et al. 2019. “Prevalence of Co-Occurring Mental Health Diagnoses in Autism Population: A Systematic Review and Meta-Analysis.” *Lancet Psychiatry* 6, no. 10: 819–829.
- Little, R. J. A. 1988. “A Test of Missing Completely at Random for Multivariate Data With Missing Values.” *Journal of the American Statistical Association* 83, no. 404: 1198–1202. <https://doi.org/10.1080/01621459.1988.10478722>.
- Lockwood, P. L., G. Bird, M. Bridge, E. Viding, J. Decety, and C. Frith. 2013. “Dissecting Empathy: High Levels of Psychopathic and Autistic Traits Are Characterized by Difficulties in Different Social Information Processing Domains.” *Frontiers in Human Neuroscience* 7: 760. <https://doi.org/10.3389/fnhum.2013.00760>.
- Maguire, K., H. Warman, F. Blumenfeld, and P. E. Langdon. 2024. “The Relationship Between Psychopathy and Autism: A Systematic Review and Narrative Synthesis.” *Frontiers in Psychiatry* 15: 1375170. <https://doi.org/10.3389/fpsy.2024.1375170>.
- Matson, J. L., and T. T. Rivet. 2008. “Characteristics of Challenging Behaviours in Adults With Autistic Disorder, PDD-NOS, and Intellectual Disability.” *Journal of Intellectual and Developmental Disability* 33, no. 4: 323–329. <https://doi.org/10.1080/13668250802492600>.
- Matson, J. L., M. Sipes, J. C. Fodstad, and M. E. Fitzgerald. 2011. “Issues in the Management of Challenging Behaviours of Adults With Autism Spectrum Disorder.” *CNS Drugs* 25, no. 7: 597–606. <https://doi.org/10.2165/11591700-000000000-00000>.
- McBrien, J., and G. Murphy. 2006. “Police and Carers’ Views on Reporting Alleged Offences by People With Intellectual Disabilities.” *Psychology, Crime & Law* 12, no. 2: 127–144. <https://doi.org/10.1080/10683160512331316262>.
- Melvin, C. L., P. E. Langdon, and G. H. Murphy. 2017. “Treatment Effectiveness for Offenders With Autism Spectrum Conditions: A Systematic Review.” *Psychology, Crime & Law* 23, no. 8: 748–776.
- Morrissey, C., T. Hogue, P. Mooney, et al. 2007. “Predictive Validity of the PCL-R in Offenders With Intellectual Disability in a High Secure Hospital Setting: Institutional Aggression.” *Journal of Forensic Psychiatry & Psychology* 18, no. 1: 1–15. <https://doi.org/10.1080/08990220601116345>.
- Morrissey, C., P. Mooney, T. E. Hogue, W. R. Lindsay, and J. L. Taylor. 2007. “Predictive Validity of the PCL-R for Offenders With Intellectual Disability in a High Security Hospital: Treatment Progress.” *Journal of Intellectual and Developmental Disability* 32, no. 2: 125–133. <https://doi.org/10.1080/13668250701383116>.
- Murphy, D. 2006. “Theory of Mind in Asperger’s Syndrome, Schizophrenia and Personality Disordered Forensic Patients.” *Cognitive Neuropsychiatry* 11, no. 2: 99–111.
- NHS England. 2021. “Service Specification: Medium Secure Mental Health Services (Adult).”
- Nicholls, T. L., J. R. P. Ogloff, and K. S. Douglas. 2004. “Assessing Risk for Violence Among Male and Female Civil Psychiatric Patients: The HCR-20, PCL:SV, and VSC.” *Behavioral Sciences & the Law* 22: 127–158.
- Ogloff, J. R. 2006. “Psychopathy/Antisocial Personality Disorder Conundrum.” *Australian & New Zealand Journal of Psychiatry* 40, no. 6–7: 519–528. <https://doi.org/10.1080/J.1440-1614.2006.01834.X>.
- Oliver, L. D., R. W. J. Neufeld, I. Dziobek, and D. G. V. Mitchell. 2016. “Distinguishing the Relationship Between Different Aspects of Empathic Responding as a Function of Psychopathic, Autistic, and Anxious Traits.” *Personality and Individual Differences* 99: 81–88. <https://doi.org/10.1016/j.paid.2016.04.058>.
- Olver, M. E., K. Lewis, and S. C. P. Wong. 2013. “Risk Reduction Treatment of High-Risk Psychopathic Offenders: The Relationship of Psychopathy and Treatment Change to Violent Recidivism.” *Personality Disorders, Theory, Research, and Treatment* 4, no. 2: 160–167. <https://doi.org/10.1037/a0029769>.
- O’Nions, E., C. L. Sebastian, E. McCrory, K. Chantiluke, F. Happé, and E. Viding. 2014. “Neural Bases of Theory of Mind in Children With Autism Spectrum Disorders and Children With Conduct Problems and Callous-Unemotional Traits.” *Developmental Science* 17, no. 5: 786–796. <https://doi.org/10.1111/DESC.12167>.
- O’Nions, E., B. Tick, F. Rijdsdijk, et al. 2015. “Examining the Genetic and Environmental Associations Between Autistic Social and Communication Deficits and Psychopathic Callous-Unemotional Traits.” *PLoS One* 10, no. 9: e0134331. <https://doi.org/10.1371/journal.pone.0134331>.
- O’Shea, L. E., M. M. Picchioni, and G. L. Dickens. 2016. “The Predictive Validity of the Short-Term Assessment of Risk and Treatability (START) for Multiple Adverse Outcomes in a Secure Psychiatric Inpatient

- Setting." *Assessment* 23, no. 2: 150–162. <https://doi.org/10.1177/1073191115573301>.
- Pedersen, L., C. Kunz, K. Rasmussen, and P. Elsass. 2010. "Psychopathy as a Risk Factor for Violent Recidivism: Investigating the Psychopathy Checklist Screening Version (PCL:SV) and the Comprehensive Assessment of Psychopathic Personality (CAPP) in a Forensic Psychiatric Setting." *International Journal of Forensic Mental Health* 9, no. 4: 308–315.
- Porter, S., A. R. Birt, and D. P. Boer. 2001. "Investigation of the Criminal and Conditional Release Profiles of Canadian Federal Offenders as a Function of Psychopathy and Age." *Law and Human Behavior* 25, no. 6: 647–661.
- R Core Team. 2023. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>. <https://www.R-project.org/>.
- Rinaldi, C., M. Attanasio, M. Valenti, M. Mazza, and R. Keller. 2021. "Autism Spectrum Disorder and Personality Disorders: Comorbidity and Differential Diagnosis." *World Journal of Psychiatry* 11, no. 12: 1366–1386.
- Rogers, J., E. Viding, R. J. Blair, U. Frith, and F. Happé. 2006. "Autism Spectrum Disorder and Psychopathy: Shared Cognitive Underpinnings or Double Hit?" *Psychological Medicine* 36, no. 12: 1789–1798. <https://doi.org/10.1017/S0033291706008853>.
- Rogers, R., R. Salekin, C. Hill, K. Sewell, M. Murdock, and C. Neumann. 2000. "The Psychopathy Checklist-Screening Version: An Examination of Criteria and Subcriteria in Three Forensic Samples." *Psychological Assessment* 7, no. 1: 1–15. <https://doi.org/10.1177/107319110000700101>.
- Schmidt, F., L. McKinnon, H. K. Chattha, and K. Brownlee. 2006. "Concurrent and Predictive Validity of the Psychopathy Checklist: Youth Version Across Gender and Ethnicity." *Psychological Assessment* 18, no. 4: 393–401. <https://doi.org/10.1037/1040-3590.18.4.393>.
- Skeem, J. L., and E. P. Mulvey. 2001. "Psychopathy and Community Violence Among Civil Psychiatric Patients: Results From the MacArthur Violence Risk Assessment Study." *Journal of Consulting and Clinical Psychology* 69, no. 3: 358–374. <https://doi.org/10.1037/0022-006X.69.3.358>.
- Swales, M. A. 2022. "Personality Disorder Diagnoses in ICD-11: Transforming Conceptualisations and Practice." *Clinical Psychology in Europe* 4: 1–18. <https://doi.org/10.32872/cpe.9635>.
- Tetley, A. C., T. E. Hogue, and S. Evershed. 2010. "Predictors of Progression From High to Medium Secure Services for Personality-Disordered Patients." *Personality and Mental Health* 4, no. 2: 108–120. <https://doi.org/10.1002/pmh>.
- Veal, R., C. Critchley, S. Luebbers, R. Cossar, and J. R. Ogloff. 2021. "Factor Structure of the Psychopathy Checklist: Screening Version (PCL: SV): A Systematic Review Using Narrative Synthesis." *Journal of Psychopathology and Behavioral Assessment* 43, no. 3: 565–582.
- Völm, B. A., R. Edworthy, N. Huband, et al. 2018. "Characteristics and Pathways of Long-Stay Patients in High and Medium Secure Settings in England; A Secondary Publication From a Large Mixed-Methods Study." *Frontiers in Psychiatry* 9: 140. <https://doi.org/10.3389/fpsy.2018.00140>.
- Webster, C., M. Martin, J. Brink, T. Nicholls, and C. Middleton. 2004. *Manual for the Short-Term Assessment of Risk and Treatability (START)*. St Joseph's Healthcare Hamilton, Ontario, Canada, and Forensic Psychiatric Services Commission.
- Wickham, H. 2016. "Getting Started With ggplot2." In *ggplot2: Elegant Graphics for Data Analysis*, 11–31. Springer International Publishing.
- Woodworth, M., and S. Porter. 2002. "In Cold Blood: Characteristics of Criminal Homicides as a Function of Psychopathy." *Journal of Abnormal Psychology* 111, no. 3: 436–445. <https://doi.org/10.1037/0021-843X.111.3.436>.
- World Health Organization. 2021. *International Classification of Diseases, Eleventh Revision (ICD-11)*. World Health Organization (WHO). <https://icd.who.int/en>.
- Žukauskienė, R., A. Laurinavičius, and I. Čėsniene. 2010. "Testing Factorial Structure and Validity of the PCL:SV in Lithuanian Prison Population." *Journal of Psychopathology and Behavioral Assessment* 32: 363–372. <https://doi.org/10.1007/s10862-009-9176-7>.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.