



Kent Academic Repository

Walker-Milne, Natasha, Elliott, Sophie Anne Marie, Wright, Peter J. and Bailey, David Mark (2025) *Seascape ecology of juvenile gadoid nursery areas*. Royal Society Open Science, 12 (9). ISSN 2054-5703.

Downloaded from

<https://kar.kent.ac.uk/112299/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/doi:10.1098/rsos.250705>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).



Research



Cite this article: Walker-Milne N, Elliott SAM, Wright PJ, Bailey DM. 2025 Seascape ecology of juvenile gadoid nursery areas. *R. Soc. Open Sci.* **12**: 250705.
<https://doi.org/10.1098/rsos.250705>

Received: 9 April 2025
Accepted: 4 August 2025

Subject Category:

Ecology, conservation, and global change biology

Subject Areas:

ecology, ecosystem

Keywords:

stereo baited remote underwater video, Atlantic cod (*Gadus morhua*), habitat modelling, species distribution modelling, haddock (*Melanogrammus aeglefinus*), whiting (*Merlangius merlangus*), heterogeneity, landscape

Author for correspondence:

Natasha Walker-Milne
e-mails: natasha.walker-milne@glasgow.ac.uk;
nwmilne@outlook.com

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.8014875>.

Seascape ecology of juvenile gadoid nursery areas

Natasha Walker-Milne¹, Sophie Anne Marie Elliott², Peter J. Wright³ and David Mark Bailey¹

¹School of Biodiversity, One Health & Veterinary Medicine, University of Glasgow, Glasgow, UK

²Game and Wildlife Conservation Trust, Fordingbridge, UK

³Marine Ecology and Conservation Consultancy, Ellon, UK

NW-M, 0000-0002-9575-0626; SAME, 0000-0001-6169-1560; PJW, 0000-0002-8402-5795; DMB, 0000-0002-0824-8823

Availability of juvenile fish habitat provision can impact recruitment. This study focused on identifying which environmental variables characterize the juvenile habitats of three commercially important gadoid species: Atlantic cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*) and whiting (*Merlangius merlangus*). Stereo baited remote underwater video surveys were conducted in the South Arran Marine Protected Area between 2013 and 2019 to collect presence/absence data on juvenile gadoids (>20 mm <120 mm) and demersal and epibenthic communities. Data were analysed using binomial generalized additive mixed models. The results revealed spatial segregation among species, each favouring distinct habitats. Predictive modelling suggests a substantial increase in presence probability from 0.25 to 0.75 as the inverse Simpson's diversity index increases, suggesting that biodiversity appears to be associated with species distribution. Boundary regions between seabed types were associated with variation in species distribution, underlining the importance of seascape heterogeneity. This study underscores the importance of conserving and restoring benthic and epibenthic biodiversity across spatially heterogeneous landscapes. Consequently, reducing benthic pressures could promote early survival for these species, thereby supporting broader ecosystem health and fisheries management goals.

1. Background

Understanding the habitat of juvenile fishes is essential for the management and conservation of populations, as juvenile life stages can have a large impact on recruitment success [1–4]. In species with a demersal or benthic settlement, protection of the early settlement stages could assist population recovery

[5,6]. With the vital role that suitable habitat plays, removal or degradation of these important post-settlement habitats may have a significant impact on populations [2,7,8]. Here, we define habitat as the combination of variables contributing to the characteristics of an area that influence where a species may occupy (e.g. depth, temperature, biodiversity, seabed type, etc.). In this study, seabed type refers to the composition of the substratum [9]. The definition of 'nursery' habitats has evolved in recent years, from being described as areas where higher densities of juveniles are found [10], to those that make a higher-than-average contribution to recruitment into the population [11]. Research has also been conducted to understand why some areas are more productive, considering aspects such as the complexities of patch connectivity, ontogenetic shifts and resource availability [12,13]. Habitats, which provide post-settlement refuge and foraging potential, can have a marked impact on the survivability of juvenile demersal and benthic fishes [14]. Understanding how these habitat attributes influence the early life stages of key demersal fishes, such as gadoids, is therefore important for effective management and conservation.

Juvenile fishes often do not rely on a single habitat [8,15], but rather a succession of differing ones, as their requirements change in response to prey availability and predation risk and through not only ontogenetic stages but also daily requirements [16]. Therefore, it is important to consider how a mosaic of habitat types and their arrangement influences post-settlement selection and survival. The 'seascape nursery' approach is valuable because it provides a framework for considering both spatial and temporal variation in habitat use, as well as the distinct requirements associated with different ontogenetic stages [13]. Given that survival through these early stages influences population growth [16,17], it is essential that the spatial and temporal scales reflect the ecological needs of the focal species, as requirements vary across species and their life-history phases.

For many species, nursery habitats are typically found in shallow coastal waters, where greater food availability, reduced physical stress and increased shelter enhance juvenile survival [18–21]. The quality of habitat, defined as the extent to which an area provides the species-specific physical, chemical and biological conditions necessary for optimal growth, survival and future reproductive success [9,22], is also important, and often indicated by higher biodiversity. Bottom-contacting mobile fishing gear can have long-term impacts on ecosystem health, including alteration of community structure [23], reductions in benthic biomass [24] and extended post-impact recovery times following removal [25]. This suggests that year-round protection of juvenile fish habitat rather than seasonal closures may be needed to promote recruitment while restoring and maintaining demersal and epibenthic biodiversity.

Epibenthos contributes to habitat complexity and provides food sources to lower trophic levels, subsequently supporting piscivorous predators. This interconnectivity across all trophic levels promotes ecosystem stability and faster recovery from disturbance [26–28]. Areas with high biodiversity are often areas of increased primary and secondary production and increased food sources, leading to higher growth, fecundity and resilience to invasive species [27,29,30]. A number of temperate coastal marine habitats have been associated with areas of increased biodiversity owing to their complex three-dimensional structures and increased foraging opportunities, including seagrass beds [31–33], maerl beds (*Phymatolithon calcareum*) [34–36] and kelp forests [37,38]. Predictor variables for biodiversity vary between biota; thus, each must be considered on a suitable landscape scale for each habitat [39]. Despite the many benefits of increased biodiversity, marine protected areas (MPAs) are often not designed with measurable improvements of metrics in biodiversity as a goal [40,41], and often focus on a specific species or habitat [42,43], despite biodiversity being a specific goal of the United Nations Convention on Biological Diversity [44]. To address this issue, some proposals have been put forward to pivot towards more 'whole-site' MPA management [41,45,46], with evidence that network-scale bioregional approaches are more effective for sustaining biodiversity than feature-specific designations [40]. If the focus remains on a selection of designated features such as those included in the Priority Marine Features approach used in Scotland (e.g. maerl beds, flame shell beds) [47], the types of seascapes protected may not be optimal for biodiversity protection or to support juvenile fishes.

Commercially important demersal species, such as Atlantic cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*) and whiting (*Merlangius merlangus*), have undergone population declines in several areas of the northeast Atlantic over recent decades, accompanied by changes in community structure [48,49]. Despite efforts to reduce fishing or enact spatial and temporal closures, long-term effects on population size and composition are evident [50–52]. These stocks are currently not achieving their productive potential, and rebuilding them requires a broad range of measures, including well-designed spatial measures [53]. Previous studies have demonstrated tendencies for each species

to favour certain juvenile gadoid conditions, such as differences in seabed composition [16,54,55] and depth [56]. All three species display some degree of site fidelity on settlement [14,57–59] and therefore, the availability of undisturbed and suitable nursery areas post-settlement may be a necessary component of successful recruitment and recovery. All three species display similar dietary niches when settlement first occurs and are in the transitional period from pelagic to demersal foraging [58]. As their size increases, there is divergence in dietary needs, although mid-sized *Mel. aeglefinus* and *Mer. merlangus* may still experience interspecific competition [58]. However, detailed dietary analyses indicate that resource partitioning among species and size classes reduces the likelihood of food limitation in nursery areas [60], although changing environmental conditions (e.g. climate-driven shifts in productivity) may increase the potential for competition in some systems. The use of shallower waters may not be a direct avoidance of predation, but rather a strategy to avoid dedicating as much time to vigilance, so more effort can be put into foraging [20].

The Firth of Clyde on the west coast of Scotland has a long history of commercial fishing; there is evidence that fisheries impact in the Clyde has been an ongoing concern for some time. The rise of industrial fishing in the nineteenth century was associated with a decline in most fish stocks [61]; the impacts of which were raised as concerns regarding bottom trawling on benthic spawners and the seabed as early as the 1880s. This heavy exploitation of the Firth of Clyde has led to changes in the community structure of populations, affecting taxonomic richness, evenness and maximum length [49]. Changes in Clyde fishery biomass showed that in the period 1927–1959, 13 taxa were responsible for 95% of biomass, which dropped to four taxa between 1995 and 2004, with *Mer. merlangus* alone accounting for greater than 80% of biomass [49].

In this study, we investigated how habitat variables (including proportional coverage of seabed types, depth, patch characteristics, distance to shore, current velocity) influence the presence of three commercially important juvenile gadoid species: *G. morhua*, *Mel. aeglefinus* and *Mer. merlangus*, at both patch and landscape scales, using stereo baited remote underwater video (SBRUV) surveys [62]. The aim of this study was to enhance the current understanding of juvenile habitat associations for all three species and to generate accurate predictions of species distribution, considering the effects of biodiversity and landscape composition. We focused on the Firth of Clyde in Scotland as a case study and hypothesized that juvenile individuals of each species would exhibit distinct habitat preferences influenced by factors such as foraging behaviour, dietary requirements and predator avoidance strategies.

2. Methods

Surveying was conducted in 2013, 2014, 2018 and 2019 around the Isle of Arran, the largest island located in the Firth of Clyde on the west coast of Scotland (55.5806° N, 5.2109° W). Surveying effort and spatial coverage varied between years owing to logistical constraints (equipment availability, vessel access and weather conditions). The dataset was designed to build cumulative spatial coverage rather than to support direct interannual comparison; therefore, year was not included as a variable in analyses. Stretching around Arran's southern coast is the South Arran Nature Conservation MPA. The South Arran MPA is a multi-use MPA with a series of fishing restriction measures (ranging from prohibition of dredging up to a no take zone) targeted at the protection of certain features, such as maerl beds and seagrass, while also allowing continued use of the area by both commercial and recreational human activities [62,63].

2.1. Survey methods and equipment

SBRUV surveys were carried out within the South Arran MPA using a 10.8 m research vessel (RV *Actinia*), 6.5 m rigid inflatable boat, and 10.0 m creel boat. Three SBRUV camera set-ups each consisted of a pair of Canon HF G25 video cameras encased in waterproof camera housings mounted on a steel frame 57 cm in height with an inward angle of approximately 8° and a basal separation of 58 cm [16,64], with two LED W38VR Archon lights providing illumination as per [16]. Frames were freshly baited with locally caught mackerel (*Scomber scombrus*) for each station. The camera systems remained on the seabed and recorded for 60 min per deployment with species recording commencing after a 3 min settlement time. Depths for SBRUV deployments ranged from 4.1 to 47.2 m and were conducted between 07.45 and 17.28 to avoid crepuscular behaviour. To maintain consistency in spatial coverage, SBRUV deployments deeper than 40 m were excluded from the analysis, as only a small proportion




Figure 1. Euclidean distance to the edge of seabed type patches as created from seabed patches produced in [62]. (a) Algae, (b) gravel, (c) pebble, (d) sand and (e) mud. Positive values denote within-habitat distances with increased values showing distance further into patches; mud is known to extend beyond the boundaries of the MPA [68]. Sampling locations are coloured by seabed type; red points indicate presence and black points indicate absence. Orange lines denote boundaries between MPA zones (numbered 1–5) used in generalized additive mixed models as a random effect.

of deployments exceeded this depth. The MPA was divided into five zones based on levels of fetch exposure, bathymetry and coastal characteristics [16,62]. Stations were selected within each zone using random points stratified by depth using a quantum geographic information system (QGIS) [65]. SBRUV camera equipment was used in conjunction with drop-down video carried out by NatureScot and the Scottish Environmental Protection Agency to examine and map seabed composition [62].

Videos were analysed using EVENTMEASURE™ software [64] with presence recorded for each fish species and a still image captured from each station for seabed analysis [62]. As SBRUV equipment was used in this survey, it was possible to take stereo measurements of fishes when visible in both cameras. Stereo video calibration and measurement protocols followed Elliott *et al.* [16]; length measurements with a root mean square error of greater than 2 cm or a precision greater than 0.5 cm were excluded. Owing to water clarity and identification requirements, most fishes measured were within 1–2 m of the camera system. This allowed for the confirmation of the 0-group gadoid status with sizes remaining

Table 1. Summary of environmental model variables, details of seabed classification [62].

variable	unit	description
mud	proportional coverage (0–1)	visual coverage of seabed type in each site from NatureScot and University of Glasgow surveys ^a
algae		
sand		
gravel		
pebble		
distance to shore	km	Euclidean distance to shore for each survey site
depth	m	bathymetry data 1 s arc
terrain ruggedness index (TRI)	in metres of elevation difference for grid points 1 arc-seconds apart	elevation differences between grid points calculated using the raster package in R
inverse Simpson's diversity index	$1/D = N(N-1)/\sum(n/N)$	biodiversity indices ^b
mean current velocity	ms^{-1}	mean current data [70]
distance to edge of seabed patch	m	Euclidean distance to patch edge for each survey site
patch perimeter	m	distance around patch area

^aGravel contains maerl, both dead and alive as live maerl was in too small quantities to be classified as separate seabed type.

^bPredicted demersal and epibenthic biodiversity calculated in R using rDiversity package [69].

less than 120 mm [35]; *G. morhua* larger than this were observed sporadically but were not recorded for this study as their size indicated 1-group (>200 mm) [66].

Environmental variables included proportional coverage of seabed type, classified as mud, sand, gravel (including both alive and dead maerl), pebble and algae (encompassing all types of macroalgae which were attached to the seabed), and seabed patch characteristics such as area (m^2), perimeter (m) and perimeter-to-area ratio. Euclidean distance to the edge of the patch (m) was also calculated using R statistical software (R v. 4.4.1) [62,67], and post-calculated analysis was conducted in QGIS. This resulted in a positive distance within the patch to the edge of the patch boundary and a negative value for distances outside the patches to allow for a metric of distance to the seabed type patch edges (figure 1). The other model variables were mean current velocity (ms^{-1}), distance to shore (km), depth (m) and terrain ruggedness index (TRI; table 1; electronic supplementary material, figure S2). Simpson's inverse diversity index was calculated for each site using species counts of epibenthic and demersal fishes, and macroinvertebrate species [62], hereafter referred to as biodiversity. Seagrass was excluded from the final models because of the small sample size. As noted previously, the year was not included as a variable in the analyses.

2.2. Statistical analysis

Collinearity checks were performed on all variables using multicollinearity diagnostic measures and Spearman's rank correlation coefficient [71]. Model selection was carried out using backward stepwise selection using Akaike information criterion model evaluation, and smoothing terms for the models were determined through model selection (electronic supplementary material, table S1). The model of best fit was tested against the null hypothesis using a log-likelihood ratio test. The months of sampling were combined into early (June/July, $n = 201$) and late (August/September, $n = 322$). This was included to investigate any interactions with depth and seabed type coverage to examine if there were significant ontogenetic shifts. The model results were checked for spatial autocorrelation using Moran's I statistic. There was initial evidence of spatial autocorrelation; therefore, the study area was partitioned into zones as described by Elliott *et al.* [16], which then accounted for spatial autocorrelation, and the MPA zone was included as a random effect.

The area under the receiver operating curve for fish species models was calculated for the model of best fit, as was the model's per cent correctly classified and a confusion matrix was calculated using the *PresenceAbsence* package [72]. Model accuracy testing was conducted on a 25% split of the data using the *caret* R package [73]. Predictions were mapped using a 100 m point grid of environmental variables

Table 2. Summary of model variables and outputs for 0-group gadoid presence including area under curve (AUC), per cent correctly classified (PCC). (Explanatory variables show proportional coverage of seabed type, depth (m), mean current velocity (ms^{-1}), inverse Simpson's diversity index, terrain ruggedness index (TRI) and distance from survey points to edge of seabed patch edge for sand, mud, algae and gravel (m), distance to shore (m), with MPA zone as a random effect. Coefficients and diagnostics (χ^2 and p -values) indicate the effect of each parameter, with variables of statistical significance in *italics*, * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.0001$.)

model: <i>G. morhua</i>	AUC: 0.85	PCC: 0.77	PCC s.d.: ± 0.02
	zero: 53%	deviance: 30.19	R^2: 0.34
variable	χ^2	P	
<i>algae</i>	63.49	<0.0001 ***	
<i>gravel</i>	21.06	<0.0001 ***	
<i>sand</i>	8.14	0.004 **	
<i>pebble</i>	2.86	0.091	
<i>TRI</i>	15.42	<0.0001 ***	
<i>Simpsons</i>	18.38	<0.0001 ***	
mean velocity	2.22	0.315	
<i>distance to sand edge</i>	11.35	0.024 *	
<i>distance to mud edge</i>	5.99	0.014 *	
model: <i>Mel. aeglefinus</i>	AUC: 0.89	PCC: 0.81	PCC s.d.: ± 0.02
	zero: 56%	deviance: 38.91	R^2: 0.43
variable	χ^2	P	
<i>algae</i>	39.07	<0.0001 ***	
<i>depth</i>	68.91	<0.0001 ***	
<i>mean velocity</i>	4.69	0.030 *	
<i>distance to shore</i>	6.25	0.012 *	
<i>Simpsons</i>	13.83	0.0002 ***	
model: <i>Mer. merlangus</i>	AUC: 0.89	PCC: 0.81	PCC s.d.: ± 0.02
	zero: 57%	deviance: 39.65	R^2: 0.44
variable	χ^2	P	
<i>depth</i>	44.25	<0.0001 ***	
<i>TRI</i>	8.68	0.003 **	
<i>mud</i>	3.37	0.066	
<i>mean velocity</i>	7.61	0.005 **	
<i>Simpsons</i>	24.87	<0.0001 ***	
<i>distance to algae edge</i>	21.48	0.0002 **	
<i>distance to gravel edge</i>	7.92	0.005 **	

(electronic supplementary material, figures S1 and S2) [62]. Error maps were produced by calculating the model fit for each grid point and subtracting the binomial presence/absence of the species. The predicted error of species presence/absence was mapped using ordinary kriging with *automap* [74], and the degree of extrapolation for model predictions was calculated using the multivariate environmental similarity surface index (electronic supplementary material, figure S3) using the R *dismo* package [75]. All figures were produced using the *ggplot2* package [76].




Figure 2. A combined generalized additive mixed model predicted the response of the influences of fixed biotic and abiotic variables on the probability of presence in 0-group fishes: (a) *G. morhua* – (i) proportional algae cover, (ii) proportional gravel cover, (iii) proportional sand cover, (iv) terrain ruggedness index (TRI), (v) Simpson's diversity index, (vi) distance to sand patch edge, (vii) distance to mud patch edge. (b) *Mel. aeglefinus* – (i) proportional algae cover, (ii) depth, (iii) mean current velocity, (iv) Simpson's diversity index, (v) distance to shore. (c) *Mer. merlangus* – (i) depth, (ii) distance to algae patch edge, (iii) distance to gravel patch edge, (iv) TRI, (v) mean current velocity, (vi) Simpson's diversity index. Distance values within and outside of patches, positive values are within patches, negative values are outside of patches. Grey shaded areas represent 95% confidence intervals, red dashed lines indicate the boundary of within and outside seabed patches.

3. Results

The surveys resulted in 544 SBRUV deployments across all years, of which 33 data points were excluded during initial review after deployment owing to technical issues (such as tipped frames or poor visibility owing to lighting failure), and a further 23 were removed during data cleaning (incomplete metadata, duplicate records), resulting in 488 data points. *Gadus morhua* was most likely to be present when the proportion of seabed covered by algae was approximately 0.5 ($\chi^2 = 6.49$, $p <$




Figure 3. Mapped presence predictions for 0-group gadoids: (a) *G. morhua*, (b) *Mel. aeglefinus* and (c) *Mer. merlangus* in the South Arran MPA. Species presence as recorded on stereo baited remote underwater video (SBRUV) surveys, black points absent and white points present.

0.0001; table 2). There was also a positive increase in *G. morhua* presence probability with increased gravel coverage, increased sand coverage, TRI and biodiversity.

Gadus morhua was also affected by the distance to the edge of the seabed type patch, with sand showing a positive relationship and mud showing a negative relationship (figure 2a(vi)(vii); table 2). Depth was not significant for *G. morhua*, unlike *Mel. aeglefinus* and *Mer. merlangus*, which were both affected by depth. *Melanogrammus aeglefinus* presence peaked between 20 and 30 m depth, whereas *Mer. merlangus* presence was influenced by depth, peaking at approximately 30 m (figure 2b(ii),c(i)). There was a negative relationship between *Mel. aeglefinus* presence and increases in algal coverage, mean current velocity and distance to shore (figure 2b(i)(iii)(iv)). *Merlangius merlangus* probability showed a negative relationship between the presence and increases in TRI and mean current velocity (figure 2c(iv)(v), respectively). The distance from algal patch edges influenced *Mer. merlangus*, resulting in decreases within algal patches and far outside of algal patch edges, but a peak in probability just outside algal patches (figure 2c(ii)). All three species showed a positive relationship with increased biodiversity, which was the only shared variable across the three species.

Gadus morhua were found in areas associated with high algae coverage to the southwest of the island and in areas of high sand coverage to the southeast (figure 3). *Melanogrammus aeglefinus* were found further away from the coast around the southwest of the island, avoiding areas of high algae coverage, but showed an increased presence in areas on the east coast of the island in deeper waters, both of which have a lower mean current velocity and are close to shore, providing increased shelter. *Merlangius merlangus* were predicted in a band around the island at increased depth away from areas of algal coverage but close to the borders of algal patches. This peaked in distance to the algae patch edge at approximately 500 m outside the patch areas (figures 2 and 3). Although *Mer. merlangus* showed a preference for increased depth and areas with less rugosity, the distance out into deeper water appeared to be limited by the distance from algal patches.

4. Discussion

This study demonstrated that the probability of the presence of all three gadoid species was positively associated with increased demersal and epibenthic biodiversity. Although each species responded differently to other environmental variables, biodiversity was the only variable consistently associated

with presence across all three species. These results for a temperate marine ecosystem build upon previous investigations into the impact of marine biodiversity on fish biomass and community regulation [77,78]. Identifying cross-species associations is important for informing multispecies management strategies, while recognizing that species-specific habitat requirements will vary between species. Areas with increased biodiversity have been demonstrated to have higher ecosystem services and greater community resilience to pressures and stochastic events [79,80]. This may be facilitated through factors including, but not limited to, genetic diversity and increased species interactions such as complementarity and facilitation [81,82].

The relationship between each species and seabed patch is interesting, as it is not only the degree of seabed coverage that plays a role but also the distance to or from patch edges, with *G. morhua* showing increased presence within sand patches and *Mer. merlangus* showing decreased presence inside gravel patches, as well as peaks in presence around the edges of algal patches (figure 2a(vi),c(ii)(iii)). Differences in the presence between species can be related to dietary requirements, predator avoidance strategies and size-specific segregation [83–86]. In the absence of predation pressure, juvenile *G. morhua* will select sand or gravel areas and will often forage more efficiently over sand, although this will leave them open to increased predation [55]. Despite this, juvenile *G. morhua* often settle in areas of higher complexity to increase survival [48,87,88]. As juvenile *G. morhua* rely on a predator avoidance strategy through camouflage rather than shoaling, it would be prudent to remain close or within areas of refuge [21]. The need to balance areas of concealment with areas of foraging may explain the preference for boundary areas of sand and areas where sporadic clumps of algae are present within sand patches. The presence of *G. morhua* in areas with a high degree of algal coverage far within a designated sand patch may also be owing to the usage of small clumps of unfastened algae that were not part of the main extent of an algae patch (figure 2a(i)(iii)). Sporadic algal patches have previously been observed to provide intermediary refuges for *G. morhua* transiting an area or indeed gaining access to foraging in open sand, but with a means of concealment, as such gap crossing is mediated by the provision of cover.

Predator avoidance is displayed in juvenile *G. morhua* activity compared with other more generalist species such as *Mer. merlangus*. This may be a result of boldness and more prominent scavenging behaviour exhibited by juvenile *Mer. merlangus*, resulting in a competitive advantage [86]. Such boldness may contribute to sampling bias associated with SBRUV equipment, as typically shy species will leave the field of view when larger or shoaling species arrive [86,89], and when encountering species with a high site fidelity, which may inflate perceived abundance in a small survey area [90]. Although all surveys were conducted within daylight hours owing to practical constraints (e.g. safety and tides), lights were included in the SBRUV set-up to add in species identification. Despite this limitation, SBRUVs have consistently demonstrated reliability in sampling carnivorous and scavenging species without negatively impacting the detection of herbivorous or omnivorous species [91,92]. However, considering the potential for diurnal bias, it is recommended that future studies also conduct nocturnal surveys for a more comprehensive understanding of species assemblages.

Merlangius merlangus were more likely to be present over areas with a lower TRI and were less likely to be observed closer to the edges of the gravel patch. The likelihood of the presence of *Mer. merlangus* further declined with increasing distance inside gravel patches. However, the peak in presence around algal patches (figure 2c(ii)(iii)) provides a noteworthy result. Although algal coverage itself does not provide the habitat necessary for juvenile *Mer. merlangus*, areas bordering it provide foraging opportunities that *Mer. merlangus* may find favourable [93]. Predators of juvenile *G. morhua*, such as older gadoids, including conspecifics, have been observed foraging on 0-group *G. morhua* along seagrass patch edges [94], and the same interaction may be present around areas of algae. Interestingly, given the results observed for juvenile *G. morhua*, if they are using algae for predator avoidance and foraging in areas surrounding algae, this could account for the presence of *Mer. merlangus* on this band bordering algal patches. *Merlangius merlangus* have been observed to predate on juvenile *G. morhua* [95]. The spike in presence outside of algal patches has been previously observed in other species, where patch edges attract predators [94]. *Merlangius merlangus* display numerous different habitat preferences [16,93,96]; as such, *Mer. merlangus* may display more plasticity in their preferred juvenile habitat depending on local conditions.

Because algae are only found in shallow water, this may limit the presence of juvenile *G. morhua* in some cases to shallower waters. While *Mer. merlangus* populations exhibit resilience and competitive advantages attributed to their extended settlement period and dominant behaviour, species such as *G. morhua* and *Mel. aeglefinus* display distinct depth preferences, as evidenced by presence patterns observed during this study. Specifically, *Mel. aeglefinus* presence was highly influenced by depth,

peaking at approximately 30 m. This is shallower than previous observations in the North Sea, where juvenile *Mel. aeglefinus* were found at much greater depths [56], and could be specific to some deeper inshore areas around Arran. *Melanogrammus aeglefinus* presence was also more likely in areas with low current velocity and close to shore. Juvenile *Mel. aeglefinus* in this study were predicted to be present mainly in areas with lower algal coverage, deeper than 20 m, with lower current velocity, higher biodiversity and close to shore. Although *Mel. aeglefinus* showed a relationship with the proportion of seabed type cover, characteristics such as distance to the seabed patch edge appeared to have no effect. Given the role that specific habitat characteristics, such as patch edges and areas of increased biodiversity, have on juvenile gadoids, ongoing anthropogenic impacts, such as those caused by bottom-contacting mobile fishing, pose a threat to these ecological components. Such activities are known to change community structure and damage the habitat complexity and biodiversity shown in this study to be important to juvenile fish presence.

While SBRUV deployments in this study were primarily limited to the nearshore and shallower areas (<40 m), this was based on both logistical constraints and the known ecology of juvenile gadoids, particularly *G. morhua* and *Mer. merlangus*, which predominantly occupy nearshore and shallow areas during their early demersal life stages [57]. While *Mel. aeglefinus* may show greater use of deeper water, SBRUV deployments in these areas were limited by logistical and safety constraints. The lack of offshore data may under-represent some aspects of spatial distribution, particularly for later stage *Mel. aeglefinus* and *Mer. merlangus*, and we recommend that future studies integrate additional sampling to address this. Nonetheless, our sampling strategy is consistent with both previous studies [16] and ecological understanding of juvenile habitat use [54,57,58] and is unlikely to have introduced substantial bias regarding the post-settlement habitats for the study species. Multivariate environmental similarity surface maps (electronic supplementary material, figure S3) indicate that statistical predictions are robust in the majority of nearshore and environmentally representative areas and decrease in predictive power in offshore regions.

5. Conclusion

It is the quality of habitat, through higher biodiversity, which is most strongly associated with the presence of all three species in conjunction with the benefits of boundary effects, which emphasize the importance of heterogeneous habitat mosaics. Our results emphasize the need for a seascape approach to understand habitat usage. It also highlights the importance of the conservation of habitats for commercial species, which would be best achieved through whole-site restrictions on harmful activities. Marine landscape composition and connectivity for mobile species, such as gadoids, would be better served by such whole-site management rather than protection of single features without consideration of their context [45,97–99].

Ethics. No aspect of this observational study required a UK Home Office license according to the University of Glasgow ethic committee, as all work undertaken comprised of in situ observations of species.

Data accessibility. The data and code that support the findings of this study are available from Dryad [100].

Supplementary material is available online [101].

Declaration of AI use. We have used AI-assisted technologies in creating this article.

Authors' contributions. N.W.-M.: conceptualization, data curation, formal analysis, investigation, methodology, visualization, writing—original draft; S.A.M.E.: formal analysis, investigation, writing—review and editing; P.J.W.: funding acquisition, project administration, supervision, writing—review and editing; D.B.: funding acquisition, methodology, project administration, supervision, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. This is contribution 2 from the importance of nursery habitat as a bottleneck to gadoid recruitment project. N.W.-M. was in receipt of a PhD studentship funded by the Ministry of Business, Innovation and Employment, in collaboration with the National Institute for Water and Atmospheric Research Ltd, New Zealand and Marine Scotland. S.A.M.E was in receipt of funding from the European Maritime and Fisheries Fund (EMFF), Blue Marine Foundation, Natural Environmental Research Council (NERC—grant NFSD/13/01), Sustainable Inshore Fisheries Trust (SIFT) and ClimateXChange.

Acknowledgements. The authors would like to extend their thanks and appreciation to the National Institute for Water and Atmospheric Research Ltd, New Zealand (NIWA) and Marine Scotland Science. Many thanks also to Dr Jo Clark, Dr Neil Burns and Dr Charlotte Hopkins for assistance with field work. We would also like to thank Marine Scotland Science, NatureScot, and all at the Community of Arran Seabed Trust (COAST) for their contribution and support.

References

1. Elliott SAM, Carpentier A, Feunteun E, Trancart T. 2020 Distribution and life history trait models indicate vulnerability of skates. *Prog. Oceanogr.* **181**, 102256. (doi:10.1016/j.pocean.2019.102256)
2. Caddy JF. 2008 The Importance of 'cover' in the life histories of demersal and benthic marine resources: a neglected issue in fisheries assessment and management. *Bull. Mar. Sci.* **83**, 7–52.
3. Wright P, Tobin D, Gibb F, Gibb I. 2010 Assessing nursery contribution to recruitment: relevance of closed areas to haddock *Melanogrammus aeglefinus*. *Mar. Ecol. Prog.* **400**, 221–232. (doi:10.3354/meps08384)
4. Laurel BJ, Knoth BA, Ryer CH. 2016 Growth, mortality, and recruitment signals in age-0 gadids settling in coastal Gulf of Alaska. *ICES J. Mar. Sci.* **73**, 2227–2237. (doi:10.1093/icesjms/fsw039)
5. Fodrie F, Levin L, Lucas A. 2009 Use of population fitness to evaluate the nursery function of juvenile habitats. *Mar. Ecol. Prog.* **385**, 39–49. (doi:10.3354/meps08069)
6. Lough RG. 2010 Juvenile cod (*Gadus morhua*) mortality and the importance of bottom sediment type to recruitment on Georges Bank. *Fish. Oceanogr.* **19**, 159–181. (doi:10.1111/j.1365-2419.2010.00535.x)
7. Lindholm J, Auster P, Kaufman L. 1999 Habitat-mediated survivorship of juvenile (0-year) Atlantic cod *Gadus morhua*. *Mar. Ecol. Prog. Ser.* **180**, 247–255. (doi:10.3354/meps180247)
8. Swadling DS, Knott NA, Taylor MD, Rees MJ, Cadiou G, Davis AR. 2024 Consequences of juvenile fish movement and seascape connectivity: does the concept of nursery habitat need a rethink? *Estuaries Coasts* **47**, 607–621. (doi:10.1007/s12237-023-01323-6)
9. Elliott S, Milligan R, Heath M, Turrell W, Bailey D. 2016 Disentangling habitat concepts for demersal marine fish management. *Oceanogr. Mar. Biol. Annu. Rev.* **54**, 173–192. (doi:10.1201/9781315368597-4)
10. Heck KL, Nadeau DA, Thomas R. 1997 The nursery role of seagrass beds. *Gulf Mex. Sci* **15**, 50–54. (doi:10.18785/goms.1501.08)
11. Beck MW *et al.* 2001 The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates. *Bioscience* **51**, 633–641. (doi:10.1641/0006-3568(2001)051[0633:TICAM0]2.0.CO;2)
12. Gillanders B, Able K, Brown J, Eggleston D, Sheridan P. 2003 Evidence of connectivity between juvenile and adult habitats for mobile marine fauna: an important component of nurseries. *Mar. Ecol. Prog.* **247**, 281–295. (doi:10.3354/meps247281)
13. Nagelkerken I, Sheaves M, Baker R, Connolly RM. 2015 The seascape nursery: a novel spatial approach to identify and manage nurseries for coastal marine fauna. *Fish Fish.* **16**, 362–371. (doi:10.1111/faf.12057)
14. Tupper M, Boutilier RG. 1995 Effects of habitat on settlement, growth, and postsettlement survival of Atlantic cod (*Gadus morhua*). *Can. J. Fish. Aquat. Sci.* **52**, 1834–1841. (doi:10.1139/f95-176)
15. Swadling DS, Knott NA, Taylor MD, Coleman MA, Davis AR, Rees MJ. 2022 Seascape connectivity of temperate fishes between estuarine nursery areas and open coastal reefs. *J. Appl. Ecol.* **59**, 1406–1416. (doi:10.1111/1365-2664.14157)
16. Elliott S, Turrell W, Heath M, Bailey D. 2017 Juvenile gadoid habitat and ontogenetic shift observations using stereo-video baited cameras. *Mar. Ecol. Prog.* **568**, 123–135. (doi:10.3354/meps12068)
17. Houde ED. 1987 Fish early life dynamics and recruitment variability. *Am. Fish. Soc. Symp.* **2**, 17–29.
18. Elliott SAM, Ahti PA, Heath MR, Turrell WR, Bailey DM. 2016 An assessment of juvenile Atlantic cod *Gadus morhua* distribution and growth using diver operated stereo-video surveys. *J. Fish Biol.* **89**, 1190–1207. (doi:10.1111/jfb.12998)
19. Gattuso JP, Gentili B, Duarte CM, Kleypas JA, Middelburg JJ, Antoine D. 2006 Light availability in the coastal ocean: impact on the distribution of benthic photosynthetic organisms and their contribution to primary production. *Biogeosciences* **3**, 489–513. (doi:10.5194/bg-3-489-2006)
20. Linehan JE, Gregory RS, Schneider DC. 2001 Predation risk of age-0 cod (*Gadus*) relative to depth and substrate in coastal waters. *J. Exp. Mar. Biol. Ecol.* **263**, 25–44. (doi:10.1016/S0022-0981(01)00287-8)
21. Lough R, Valentine P, Potter D, Auditory P, Bolz G, Neilson J, Perry R. 1989 Ecology and distribution of juvenile cod and haddock in relation to sediment type and bottom currents on eastern Georges Bank. *Mar. Ecol. Prog. Ser.* **56**, 1–12. (doi:10.3354/meps056001)
22. Gibson RN. 1994 Impact of habitat quality and quantity on the recruitment of juvenile flatfishes. *Neth. J. Sea Res.* **32**, 191–206. (doi:10.1016/0077-7579(94)90040-x)
23. Rijnsdorp AD *et al.* 2020 Mitigating seafloor disturbance of bottom trawl fisheries for North Sea sole *Solea solea* by replacing mechanical with electrical stimulation. *PLoS ONE* **15**, e0228528. (doi:10.1371/journal.pone.0228528)
24. Hiddink JG, Jennings S, Kaiser MJ. 2006 Indicators of the ecological impact of bottom-trawl disturbance on seabed communities. *Ecosystems* **9**, 1190–1199. (doi:10.1007/s10021-005-0164-9)
25. Rijnsdorp AD, Bolam SG, Garcia C, Hiddink JG, Hintzen NT, van Denderen PD, van Kooten T. 2018 Estimating sensitivity of seabed habitats to disturbance by bottom trawling based on the longevity of benthic fauna. *Ecol. Appl.* **28**, 1302–1312. (doi:10.1002/eap.1731)
26. Ziegler SL *et al.* 2023 Marine protected areas, marine heatwaves, and the resilience of nearshore fish communities. *Sci. Rep.* **13**, 1405. (doi:10.1038/s41598-023-28507-1)
27. Worm B *et al.* 2006 Impacts of biodiversity loss on ocean ecosystem services. *Science* **314**, 787–790. (doi:10.1126/science.1132294)
28. Ramírez-Ortiz G, Balart EF, Reyes-Bonilla H, Huato-Soberanis L, Cortés-Fuentes C, Micheli F. 2022 Greater resilience of reef fish assemblages in a no-take reserve compared to multi-use areas of the Gulf of California. *Prog. Oceanogr.* **204**, 102794. (doi:10.1016/j.pocean.2022.102794)
29. Palumbi SR, McLeod KL, Grünbaum D. 2008 Ecosystems in action: lessons from marine ecology about recovery, resistance, and reversibility. *BioScience* **58**, 33–42. (doi:10.1641/b580108)

30. Sala E, Knowlton N. 2006 Global marine biodiversity trends. *Annu. Rev. Environ. Resour.* **31**, 93–122. (doi:10.1146/annurev.energy.31.020105.100235))
31. Bertelli CM, Unsworth RKF. 2014 Protecting the hand that feeds us: seagrass (*Zostera marina*) serves as commercial juvenile fish habitat. *Mar. Pollut. Bull.* **83**, 425–429. (doi:10.1016/j.marpolbul.2013.08.011)
32. Hori M, Suzuki T, Monthum Y, Srisombat T, Tanaka Y, Nakaoka M, Mukai H. 2009 High seagrass diversity and canopy-height increase associated fish diversity and abundance. *Mar. Biol.* **156**, 1447–1458. (doi:10.1007/s00227-009-1184-3)
33. Jackson EL, Rowden AA, Attrill MJ, Bossey SJ, Jones MB. 2001 The importance of seagrass beds as a habitat for fishery species. In *Oceanography and marine biology* (eds RN Gibson, M Barnes, RJA Atkinson), pp. 269–304. London, UK: Taylor & Francis.
34. Abella E. 1998 Maerl grounds habitats of high biodiversity in European seas. In *Third European Marine Science and Technology Conf. (MAST Conf.)*, vol. 1, pp. 170–179, Lisbon, Portugal: European Commission.
35. Kamenos N, Moore P, Hall-Spencer J. 2004 Nursery-area function of maerl grounds for juvenile queen scallops *Aequipecten opercularis* and other invertebrates. *Mar. Ecol. Prog.* **274**, 183–189. (doi:10.3354/meps274183)
36. Riosmena-Rodríguez R. 2017 Natural history of rhodolith/maerl beds: their role in near-shore biodiversity and management. In *Rhodolith/maerl beds: a global perspective* (eds R Riosmena-Rodríguez, W Nelson, J Aguirre), pp. 3–26. Cham, Switzerland: Springer International Publishing. (doi:10.1007/978-3-319-29315-8_1)
37. Smale DA, Burrows MT, Moore P, O'Connor N, Hawkins SJ. 2013 Threats and knowledge gaps for ecosystem services provided by kelp forests: a northeast Atlantic perspective. *Ecol. Evol.* **3**, 4016–4038. (doi:10.1002/ece3.774)
38. Teagle H, Hawkins SJ, Moore PJ, Smale DA. 2017 The role of kelp species as biogenic habitat formers in coastal marine ecosystems. *J. Exp. Mar. Biol. Ecol.* **492**, 81–98. (doi:10.1016/j.jembe.2017.01.017)
39. Roland Pitcher C *et al.* 2012 Exploring the role of environmental variables in shaping patterns of seabed biodiversity composition in regional-scale ecosystems. *J. Appl. Ecol.* **49**, 670–679. (doi:10.1111/j.1365-2664.2012.02148.x)
40. Knott NA, Williams J, Harasti D, Malcolm HA, Coleman MA, Kelaher BP, Rees MJ, Schultz A, Jordan A. 2021 A coherent, representative, and bioregional marine reserve network shows consistent change in rocky reef fish assemblages. *Ecosphere* **12**, 1–22. (doi:10.1002/ecs2.3447)
41. Solandt JL, Mullier T, Elliott S, Sheehan E. 2020 Managing marine protected areas in Europe: moving from ‘feature-based’ to ‘whole-site’ management of sites. In *Marine protected areas* (eds J Humphreys, RWE Clark), pp. 157–181. Amsterdam, The Netherlands: Elsevier. (doi:10.1016/B978-0-08-102698-4.00009-5)
42. Hopkins CR, Burns NM, Brooker E, Dolman S, Devenport E, Duncan C, Bailey DM. 2020 Evaluating whether MPA management measures meet ecological principles for effective biodiversity protection. *Acta Oecologica* **108**, 103625. (doi:10.1016/j.actao.2020.103625)
43. Rice J *et al.* 2012 The role of MPAs in reconciling fisheries management with conservation of biological diversity. *Ocean Coast. Manag.* **69**, 217–230. (doi:10.1016/j.ocecoaman.2012.08.001)
44. United Nations. 1992 *Convention on biological diversity*. Rio de Janeiro, Brazil: United Nations.
45. Davies BFR, Holmes L, Attrill MJ, Sheehan EV. 2022 Ecosystem benefits of adopting a whole-site approach to MPA management. *Fish. Manag. Ecol.* **29**, 790–805. (doi:10.1111/fme.12581)
46. Davies BFR, Holmes L, Bicknell A, Attrill MJ, Sheehan EV. 2022 A decade implementing ecosystem approach to fisheries management improves diversity of taxa and traits within a marine protected area in the UK. *Divers. Distrib.* **28**, 173–188. (doi:10.1111/ddi.13451)
47. Scottish Natural Heritage and the Joint Nature Conservation Committee. 2012 Advice to the Scottish Government on the selection of Nature Conservation Marine Protected Areas (MPAs) for the development of the Scottish MPA network. Scottish Natural Heritage Commissioned Report No. 547
48. Bailey N *et al.* 2011 *The West of Scotland marine ecosystem: a review of scientific knowledge: Marine Scotland Science Report - Scottish Government*. Edinburgh, UK: The Scottish Government.
49. Heath MR, Speirs DC. 2012 Changes in species diversity and size composition in the Firth of Clyde demersal fish community (1927–2009). *Proc. R. Soc. B* **279**, 543–552. (doi:10.1098/rspb.2011.1015)
50. Hutchings JA. 2000 Collapse and recovery of marine fishes. *Nature* **406**, 882–885. (doi:10.1038/35022565)
51. Jackson JB *et al.* 2001 Historical overfishing and the recent collapse of coastal ecosystems. *Science* **293**, 629–637. (doi:10.1126/science.1059199)
52. Wright PJ, Trippel EA. 2009 Fishery-induced demographic changes in the timing of spawning: consequences for reproductive success. *Fish Fish.* **10**, 283–304. (doi:10.1111/j.1467-2979.2008.00322.x)
53. Worm B *et al.* 2009 Rebuilding global fisheries. *Science* **325**, 578–585. (doi:10.1126/science.1173146)
54. Kamenos NA, Moore PG, Hall-Spencer JM. 2004 Small-scale distribution of juvenile gadoids in shallow inshore waters; what role does maerl play? *ICES J. Mar. Sci.* **61**, 422–429. (doi:10.1016/j.icesjms.2004.02.004)
55. Gotceitas V, Brown JA. 1993 Substrate selection by juvenile Atlantic cod (*Gadus morhua*): effects of predation risk. *Oecologia* **93**, 31–37. (doi:10.1007/bf00321187)
56. Asjes A, González-Irusta J, Wright P. 2016 Age-related and seasonal changes in haddock *Melanogrammus aeglefinus* distribution: implications for spatial management. *Mar. Ecol. Prog. Ser.* **553**, 203–217. (doi:10.3354/meps11754)
57. Bastrikin DK, Gallego A, Millar CP, Priede IG, Jones EG. 2014 Settlement length and temporal settlement patterns of juvenile cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), and whiting (*Merlangius merlangus*) in a northern North Sea coastal nursery area. *ICES J. Mar. Sci.* **71**, 2101–2113. (doi:10.1093/icesjms/fsu029)

58. Demain DK, Gallego A, Jaworski A, Priede IG, Jones EG. 2011 Diet and feeding niches of juvenile *Gadus morhua*, *Melanogrammus aeglefinus* and *Merlangius merlangus* during the settlement transition in the northern North Sea. *J. Fish Biol.* **79**, 89–111. (doi:10.1111/j.1095-8649.2011.02997.x)
59. Rogers L, Olsen E, Knutsen H, Stenseth N. 2014 Habitat effects on population connectivity in a coastal seascape. *Mar. Ecol. Prog. Ser.* **511**, 153–163. (doi:10.3354/meps10944)
60. Rueda L, Massutí E, Alvarez-Berastegui D, Hidalgo M. 2015 Effect of intra-specific competition, surface chlorophyll and fishing on spatial variation of gadoid's body condition. *Ecosphere* **6**, 1–17. (doi:10.1890/es15-00087.1)
61. Thurstan RH, Brockington S, Roberts CM. 2010 The effects of 118 years of industrial fishing on UK bottom trawl fisheries. *Nat. Commun.* **1**, 15. (doi:10.1038/ncomms1013)
62. Walker-Milne NL, Elliott S, Wright PJ, Bailey DM. 2025 A novel use of stereo baited remote underwater video and drop-down video for biodiversity and marine landscape mapping and prediction. *PLoS ONE* **20**, e0319355. (doi:10.1371/journal.pone.0319355)
63. Scottish Government. 2015 *The South Arran Marine Conservation Order 2015*. SSA 2015/435. Edinburgh, UK.
64. SeaGIS Ltd. 2019 EventMeasure. Australia. See <https://www.seagis.com.au/>.
65. QGIS Development Team. 2009 *QGIS Geographic Information System*. Chicago, USA: Open Source Geospatial Foundation. See <https://qgis.org>.
66. Dunlop K, Staby A, van der Meeren T, Keeley N, Olsen EM, Bannister R, Skjærraasen JE. 2022 Habitat associations of juvenile Atlantic cod (*Gadus morhua* L.) and sympatric demersal fish communities within shallow inshore nursery grounds. *Estuar. Coast. Shelf Sci.* **279**, 108111. (doi:10.1016/j.ecss.2022.108111)
67. R Core Team. 2022 *R: a language and environment for statistical computing*, 4.2.2. Vienna, Austria: R Foundation for Statistical Computing.
68. Pace MC, Bailey DM, Donnan DW, Narayanaswamy BE, Smith HJ, Speirs DC, Turrell WR, Heath MR. 2021 Modelling seabed sediment physical properties and organic matter content in the Firth of Clyde. *Earth Syst. Sci. Data* **13**, 5847–5866. (doi:10.5194/essd-13-5847-2021)
69. Mitchell S, Reeve R, White T. 2020 *rdiversity*. R package. Glasgow. See <https://github.com/boydorr/rdiversity>.
70. Sabatino AD, O'Hara Murray RB, Hills A, Speirs DC, Heath MR. 2016 Modelling sea level surges in the Firth of Clyde, a fjordic embayment in south-west Scotland. *Nat. Hazards* **84**, 1601–1623. (doi:10.1007/s11069-016-2506-7)
71. Zuur AF, Ieno EN, Walker NJ, Savelieve AA, Smith GM. 2009 *Mixed effects models and extensions in ecology with R*. New York, NY: Springer.
72. Freeman EA, Moisen G. 2008 PresenceAbsence: an R package for presence-absence model analysis. *J. Stat. Softw.* **23**, 1–31. (doi:10.18637/jss.v023.i11)
73. Kuhn M. 2008 Building predictive models in R using the caret package. *J. Stat. Softw.* **28**, 1–26. (doi:10.18637/jss.v028.i05)
74. Hiemstra PH, Pebesma EJ, Twenhöfel CJW, Heuvelink GBM. 2009 Real-time automatic interpolation of ambient gamma dose rates from the Dutch radioactivity monitoring network. *Comput. Geosci.* **35**, 2008. (doi:10.1016/j.cageo.2008.10.011)
75. Hijmans RJ, Phillips S, Leathwick J, Elith J. 2024 *dismo*: Species Distribution Modeling. R package version 1.3-16. See <https://CRAN.R-project.org/package=dismo>.
76. Wickham H. 2016 *Ggplot2: elegant graphics for data analysis*. New York, NY: Springer.
77. Duffy JE, Lefcheck JS, Stuart-Smith RD, Navarrete SA, Edgar GJ. 2016 Biodiversity enhances reef fish biomass and resistance to climate change. *Proc. Natl Acad. Sci. USA* **113**, 6230–6235. (doi:10.1073/pnas.1524465113)
78. Stachowicz JJ, Bruno JF, Duffy JE. 2007 Understanding the effects of marine biodiversity on communities and ecosystems. *Annu. Rev. Ecol. Syst.* **38**, 739–766. (doi:10.1146/annurev.ecolsys.38.091206.095659)
79. Duarte CM. 2000 Marine biodiversity and ecosystem services: an elusive link. *J. Exp. Mar. Biol. Ecol.* **250**, 117–131. (doi:10.1016/S0022-0981(00)00194-5)
80. Allison G. 2004 The influence of species diversity and stress intensity on community resistance and resilience. *Ecol. Monogr.* **74**, 117–134. (doi:10.1890/02-0681)
81. Palumbi SR *et al.* 2009 Managing for ocean biodiversity to sustain marine ecosystem services. *Front. Ecol. Environ.* **7**, 204–211. (doi:10.1890/070135)
82. Hooper DU *et al.* 2005 Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. *Ecol. Monogr.* **75**, 3–35. (doi:10.1890/04-0922)
83. Skjærraasen JE, Fernö A, Folkvord A. 2003 Swimming, feeding and predator avoidance in cod larvae (*Gadus morhua* L.): trade-offs between hunger and predation risk. In *The big fish bang: Proc. of the 26th annual larval fish Conf.*, pp. 105–121. Bergen, Norway: Institute of Marine Research.
84. Gotcheitas V, Fraser S, Brown JA. 1995 Habitat use by juvenile Atlantic cod (*Gadus morhua*) in the presence of an actively foraging and non-foraging predator. *Mar. Biol.* **123**, 421–430. (doi:10.1007/BF00349220)
85. Shaw M, Diekmann R, Van DerKooij J, Milligan S, Bromley P, Righton D. 2008 Assessment of the diets of cod *Gadus morhua* and whiting *Merlangius merlangus* juveniles in a frontal region close to the Norwegian Trench: co-existence or competition? *J. Fish Biol.* **73**, 1612–1634. (doi:10.1111/j.1095-8649.2008.02035.x)
86. Elliott SAM, Allan BA, Turrell WR, Heath MR, Bailey DM. 2018 Survival of the fittest: explanations for gadoid imbalance in heavily fished seas. *Aquat. Conserv.* **28**, 1192–1199. (doi:10.1002/aqc.2926)
87. Juanes F. 2007 Role of habitat in mediating mortality during the post-settlement transition phase of temperate marine fishes. *J. Fish Biol.* **70**, 661–677. (doi:10.1111/j.1095-8649.2007.01394.x)
88. Laurel B, Gregory R, Brown J. 2003 Settlement and distribution of age-0 juvenile cod, *Gadus morhua* and *G. ogac*, following a large-scale habitat manipulation. *Mar. Ecol. Prog. Ser.* **262**, 241–252. (doi:10.3354/meps262241)

89. Willis T, Millar R, Babcock R. 2000 Detection of spatial variability in relative density of fishes: comparison of visual census, angling, and baited underwater video. *Mar. Ecol. Prog. Ser.* **198**, 249–260. (doi:[10.3354/meps198249](https://doi.org/10.3354/meps198249))
90. Stobart B, Diaz D, Álvarez F, Alonso C, Mallol S, Goñi R. 2015 Performance of baited underwater video: does it underestimate abundance at high population densities? *PLoS ONE* **10**, e0127559. (doi:[10.1371/journal.pone.0127559](https://doi.org/10.1371/journal.pone.0127559))
91. Harvey E, Cappo M, Butler J, Hall N, Kendrick G. 2007 Bait attraction affects the performance of remote underwater video stations in assessment of demersal fish community structure. *Mar. Ecol. Prog.* **350**, 92. (doi:[10.3354/meps07192](https://doi.org/10.3354/meps07192))
92. Watson DL, Harvey ES, Anderson MJ, Kendrick GA. 2005 A comparison of temperate reef fish assemblages recorded by three underwater stereo-video techniques. *Mar. Biol.* **148**, 415–425. (doi:[10.1007/s00227-005-0090-6](https://doi.org/10.1007/s00227-005-0090-6))
93. Atkinson CJL, Bergmann M, Kaiser MJ. 2004 Habitat selection in whiting. *Fish Biol.* **64**, 788–793. (doi:[10.1046/j.1095-8649.2003.00340.x](https://doi.org/10.1046/j.1095-8649.2003.00340.x))
94. Gorman AM, Gregory RS, Schneider DC. 2009 Eelgrass patch size and proximity to the patch edge affect predation risk of recently settled age 0 cod (*Gadus*). *J. Exp. Mar. Biol. Ecol.* **371**, 1–9. (doi:[10.1016/j.jembe.2008.12.008](https://doi.org/10.1016/j.jembe.2008.12.008))
95. Floeter J, Temming A. 2005 Analysis of prey size preference of North Sea whiting, saithe, and grey gurnard. *ICES J. Mar. Sci.* **62**, 897–907. (doi:[10.1016/j.icesjms.2005.03.004](https://doi.org/10.1016/j.icesjms.2005.03.004))
96. Pihl L, Baden S, Kautsky N, Rönnbäck P, Söderqvist T, Wennhage H. 2006 Shift in fish assemblage structure due to loss of seagrass *Zostera marina* habitats in Sweden. *Estuar. Coast. Shelf Sci.* **67**, 123–132. (doi:[10.1016/j.ecss.2005.10.016](https://doi.org/10.1016/j.ecss.2005.10.016))
97. Gallucci F, Christofoletti AR, Fonseca G, Dias MG. 2020 The effects of habitat heterogeneity at distinct spatial scales on hard-bottom-associated communities. *Diversity* **12**, 39. (doi:[10.3390/d12010039](https://doi.org/10.3390/d12010039))
98. Goetze JS *et al.* 2021 Increased connectivity and depth improve the effectiveness of marine reserves. *Glob. Chang. Biol.* **27**, 3432–3447. (doi:[10.1111/gcb.15635](https://doi.org/10.1111/gcb.15635))
99. Blampied SR, Rees SE, Attrill MJ, Binney FCT, Sheehan EV. 2022 Removal of bottom-towed fishing from whole-site marine protected areas promotes mobile species biodiversity. *Estuar. Coast. Shelf Sci.* **276**, 108033. (doi:[10.1016/j.ecss.2022.108033](https://doi.org/10.1016/j.ecss.2022.108033))
100. Walker-Milne N, Elliott S, Wright P, Bailey D. 2025 Seascape ecology of juvenile gadoid nursery areas [Dataset]. Dryad. (doi:[10.5061/dryad.fbg79cp7n](https://doi.org/10.5061/dryad.fbg79cp7n))
101. Walker-Milne N, Elliott SAM, Wright PJ, Bailey D. 2025 Supplementary material from: Seascape ecology of juvenile gadoid nursery areas. FigShare. (doi:[10.6084/m9.figshare.c.8014875](https://doi.org/10.6084/m9.figshare.c.8014875))