

Kent Academic Repository

Full text document (pdf)

Citation for published version

Paget, Rowena E. and Wildon, Mark (2011) Set families and Foulkes modules. *Journal of Algebraic Combinatorics*, 34 (3). pp. 525-544. ISSN 0925-9899.

DOI

<https://doi.org/10.1007/s10801-011-0282-8>

Link to record in KAR

<https://kar.kent.ac.uk/25196/>

Document Version

UNSPECIFIED

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

SET FAMILIES AND FOULKES MODULES

MARK WILDON AND ROWENA PAGET

ABSTRACT. We construct a new family of homomorphisms from Specht modules into Foulkes modules for the symmetric group. These homomorphisms are used to give a combinatorial description of the minimal partitions (in the dominance order) which label irreducible characters appearing as summands of the characters of Foulkes modules. The homomorphisms are defined using certain families of subsets of the natural numbers. These families are of independent interest; we prove a number of combinatorial results concerning them.

1. INTRODUCTION

The object of this paper is to construct a new family of homomorphisms from Specht modules into Foulkes modules, and to explore some of the consequences for the structure of Foulkes modules. Our main result gives a combinatorial description of the minimal partitions which label irreducible characters appearing as summands of the characters of Foulkes modules. (Here minimality is with respect to the dominance order on partitions, defined below.) The connection with Foulkes' long-standing conjecture [2, end of §1] on the symmetric group is explained later in the introduction.

Let S_r denote the symmetric group of degree r . For $m, n \in \mathbf{N}$, the *Foulkes module* $H^{(m^n)}$ is defined to be the permutation module for $\mathbf{Z}S_{mn}$ given by the action of the symmetric group S_{mn} on the collection of set partitions of a set of size mn into n sets each of size m . If λ is a partition of r , we denote by S^λ the associated Specht module for $\mathbf{Z}S_r$. The reader is referred to §2 of this paper for the definition of Specht modules and our notational conventions for Foulkes modules.

Our homomorphisms are defined using certain families of subsets of the natural numbers. We shall need the following combinatorial definitions.

Definition. Let $X = \{x_1, \dots, x_m\}$ and $Y = \{y_1, \dots, y_m\}$ be subsets of the natural numbers, written so that $x_1 < x_2 < \dots < x_m$ and $y_1 < y_2 < \dots < y_m$. We say that Y *majorizes* X , and write $X \preceq Y$, if $x_i \leq y_i$ for each i .

Date: June 2010.

The reader may find it helpful to refer to Figure 2 in §5.2 below, which shows part of the lattice of 4-subsets of \mathbf{N} under the majorization order.

Definition. A *set family of shape* (m^n) is a collection of n distinct m -subsets of the natural numbers. A set family \mathcal{P} is *closed* if whenever $Y \in \mathcal{P}$ and $X \prec Y$, then $X \in \mathcal{P}$.

In the following definition, λ' denotes the conjugate of the partition λ .

Definition. Let λ be a partition with largest part of size a . A set family has *type* λ if it has exactly λ'_i sets containing i for each $i \in \{1, \dots, a\}$.

It is easily seen that if \mathcal{P} is a closed set family then, for any $i \in \mathbf{N}$, at least as many sets in \mathcal{P} contain i as contain $i + 1$. Closed set families therefore have well-defined types. The reason for working with conjugate partitions will be seen in Theorem 1.1 below.

In the following definition we use the dominance order on partitions. Recall that if λ and μ are partitions of n then we say that λ *dominates* μ , and write $\lambda \succeq \mu$, if $\sum_{i=1}^j \lambda_i \geq \sum_{i=1}^j \mu_i$ for all $j \in \mathbf{N}$. (If i exceeds the number of parts of λ or μ , then the corresponding part size should be taken to be 0.)

Definition. Let \mathcal{P} be a set family of shape (m^n) and type λ . We say that \mathcal{P} is *minimal* if there is no set family \mathcal{Q} of shape (m^n) and type μ with $\mu \prec \lambda$.

It is an important fact that minimal set families are always closed; we shall prove this fact when we first need it, in Proposition 4.3 in §4 below.

We are now ready to state our main results. Let m be odd. In §2.2 we define for each closed set family \mathcal{P} of shape (m^n) and type λ , a homomorphism $f_{\mathcal{P}} : S^\lambda \rightarrow H^{(m^n)}$. One of our main results, which we prove in §3 below, is that these homomorphisms are well-defined.

Theorem 1.1. *Let m be odd and let $n \in \mathbf{N}$. Let \mathcal{P} be a closed set family of shape (m^n) and type λ . The map $f_{\mathcal{P}} : S^\lambda \rightarrow H^{(m^n)}$ is a well-defined injective homomorphism $S^\lambda \rightarrow H^{(m^n)}$.*

Let $\phi^{(m^n)}$ be the permutation character afforded by $H^{(m^n)}$. We shall refer to these characters as *Foulkes characters*. Let χ^λ be the irreducible character afforded by the Specht module S^λ . (More precisely, if λ is a partition of r , then χ^λ is the character of the $\mathbf{Q}S_r$ -module $S^\lambda \otimes_{\mathbf{Z}} \mathbf{Q}$.) It is well known that every irreducible character of a symmetric group is equal to a χ^λ : see, for instance, [4, Theorem 4.12].

If π is a character of S_r and λ is a partition of r , then we shall say that χ^λ is a *minimal constituent* of π if $\langle \pi, \chi^\lambda \rangle \geq 1$ and $\langle \pi, \chi^\mu \rangle = 0$ if $\mu \triangleleft \lambda$. In §4 we prove the following theorem which characterizes minimal constituents of Foulkes characters.

Theorem 1.2. *Let $m, n \in \mathbf{N}$.*

- (i) *If m is even then the unique minimal constituent of $\phi^{(m^n)}$ is $\chi^{(m^n)}$.*
- (ii) *If m is odd then χ^λ is a minimal constituent of $\phi^{(m^n)}$ if and only if there is a minimal set family of shape (m^n) and type λ .*

We also show that if m is even then $\langle \phi^{(m^n)}, \chi^{(m^n)} \rangle = 1$, and that part (ii) of the above theorem can be sharpened as follows.

Theorem 1.3. *Let m be odd and let $n \in \mathbf{N}$. Suppose that χ^λ is a minimal constituent of $\phi^{(m^n)}$. If $\mathcal{P}_1, \dots, \mathcal{P}_d$ are the set families of shape (m^n) and type λ , then $\mathcal{P}_1, \dots, \mathcal{P}_d$ are closed, and the homomorphisms $f_{\mathcal{P}_1}, \dots, f_{\mathcal{P}_d}$ are a \mathbf{Z} -basis for $\text{Hom}(S^\lambda, H^{(m^n)})$. In particular, $\langle \phi^{(m^n)}, \chi^\lambda \rangle = d$.*

Little is known about the constituents of Foulkes characters; indeed, it appears that the two preceding theorems give the strongest general results to date.

We pause to give a small example that will illustrate these theorems. We take $m = 3$ and $n = 4$. The three closed set families of shape (3^4) are

$$\begin{aligned} & \{\{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}, \{1, 2, 6\}\}, \\ & \{\{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}, \{1, 3, 4\}\}, \\ & \{\{1, 2, 3\}, \{1, 2, 4\}, \{1, 3, 4\}, \{2, 3, 4\}\}, \end{aligned}$$

of types $(6, 2, 2, 2)$, $(5, 4, 2, 1)$ and $(4, 4, 4)$ respectively. Since these partitions are incomparable in the dominance order, and (by Proposition 4.3) any minimal set family is closed, the set families above are minimal. It now follows from Theorems 1.2 and 1.3 that $\phi^{(3^4)}$ has $\chi^{(5,2,2,2)}$, $\chi^{(5,3,2,1)}$ and $\chi^{(4,4,4)}$ as summands, each with multiplicity 1. Moreover, if $\langle \phi^{(3^4)}, \chi^\mu \rangle \geq 1$ then μ dominates one of these partitions. The presence of such larger constituents cannot be detected by the homomorphisms $f_{\mathcal{P}}$.

Small examples of this kind are apt to create the false impression that every closed set family is minimal and is the unique set family of its type. In §5.1 we prove the following theorem which clarifies the relationship between these properties.

Theorem 1.4. *If \mathcal{P} is the unique set family of its type, then \mathcal{P} is minimal. If \mathcal{P} is a minimal set family then \mathcal{P} is closed. There exist closed set families that are not minimal, and minimal set families that are not unique for their type.*

The existence of minimal set families that are not unique for their type is of particular significance, since such families demonstrate that the multiplicity d in Theorem 1.3 can be > 1 .

Even with the help of Theorem 1.2, it appears to be a difficult matter to decide, when m is odd, whether a given partition of mn labels a minimal constituent of the Foulkes character $\phi^{(m^n)}$. In §5.2 we give a construction that gives some of these partitions. We prove that this construction gives every such partition if and only if $n \leq 5$.

We end in §6 by defining the generalized Foulkes characters ϕ^μ considered in [5] and showing how to obtain their minimal constituents from the minimal constituents of the $\phi^{(m^n)}$. The main part of this section may be read independently of the rest of the paper.

One motivation for Theorems 1.2 and 1.3 comes from Foulkes' Conjecture for the symmetric group. This conjecture, first made at end of §1 of [2], asserts that if m, n are natural numbers with $m < n$, then $\langle \phi^{(m^n)}, \chi^\lambda \rangle \geq \langle \phi^{(n^m)}, \chi^\lambda \rangle$ for every partition λ of mn .

Foulkes' Conjecture was already known to hold by the work of Thrall (see [8, Theorems III and IV]) in the case $m = 2$. It was proved when $m = 3$ by Dent and Siemons [1]. The most recent progress was made in 2008 by McKay [5], who proved it when $m = 3$ and $m = 4$. McKay's proof uses a family of maps $\psi_{(n^m)} : H^{(n^m)} \rightarrow H^{(m^n)}$ which were first defined by Wagner and Siemons¹ in 1986. McKay's main result is that if $\psi_{(m^m)}$ is invertible then $\psi_{(n^m)}$ is injective for any $n > m$. The maps $\psi_{(m^m)}$ for $m \leq 4$ were already known to be invertible by the work of Müller and Neunhöffer [6], so Foulkes' Conjecture is proved in these cases.

The main contribution of [6] was to show that $\psi_{(5^5)}$ has a non-zero kernel. It is however still possible that the maps $\psi_{(n^m)}$ will have a role in proving or disproving Foulkes' Conjecture. In particular, a conjecture of Johannes Siemons² implies that if the kernel of $\psi_{(n^m)}$ is non-zero, then there is a partition λ labelling a minimal constituent of $\phi^{(n^m)}$ and a homomorphism

¹See [5]. These maps were independently defined by Stanley: see the discussion following Problem 9 in [7].

²Seminar given at Castro Urdiales, October 2007.

$f : S^\lambda \rightarrow H^{(n^m)}$ such that $S^\lambda f \subseteq \ker \psi_{(n^m)}$. By our Theorem 1.3, any such homomorphism is an integral linear combination of the homomorphisms $f_{\mathcal{P}}$ for minimal set families \mathcal{P} of shape (n^m) and type λ .

Using Theorem 1.2, the authors have found all minimal constituents of the Foulkes characters $\phi^{(m^n)}$ for $m+n \leq 19$. The data and associated computer software are available from the second author's website: www.maths.bris.ac.uk/~mazzmjw.

Finally, we explain that we have chosen to work throughout with integral modules because this 'characteristic-free' setting is well-suited to our results and does not create any significant extra difficulties in their proofs. A subsequent paper by the authors will apply the results of this paper to study the behaviour of Foulkes modules over fields of prime characteristic.

2. SPECHT MODULES AND HOMOMORPHISMS

In this section we recall the definition of Specht modules as submodules of Young permutation modules and define the homomorphisms $f_{\mathcal{P}}$.

The following notation simplifies these definitions and will be found very useful in the proofs which follow. Given a partition λ of r with largest part of size a , let $A(\lambda)$ be the set consisting of the symbols i_j for $1 \leq i \leq a$ and $1 \leq j \leq \lambda'_i$. We say that i is the *number* and j is the *index* of the symbol i_j .

2.1. Specht modules. Let λ be a partition of r . A λ -*tableau* is an assignment of the elements of $A(\lambda)$ to the boxes of the Young diagram of λ . Given a λ -tableau t , we obtain the associated *tabloid* \mathbf{t} by disregarding the order of the elements within the rows of t . For example, if

$$t = \begin{array}{|c|c|c|c|} \hline 4_1 & 2_2 & 3_2 & 1_1 \\ \hline 2_1 & 1_2 & 3_1 & 4_2 \\ \hline 1_3 & & & \\ \hline \end{array} \quad \text{then} \quad \mathbf{t} = \frac{4_1 \ 2_2 \ 3_2 \ 1_1}{2_1 \ 1_2 \ 3_1 \ 4_2} = \frac{1_1 \ 2_2 \ 3_2 \ 4_1}{1_2 \ 2_1 \ 3_1 \ 4_2} = \dots \quad \text{etc.}$$

We may identify S_r with the symmetric group on $A(\lambda)$. After this identification is made, the natural permutation action of S_r on the set of λ -tableaux gives rise to a well-defined action of S_r on the set of λ -tabloids. We denote the associated permutation module for $\mathbf{Z}S_r$ by M^λ ; it is the *Young permutation module* corresponding to λ . For example, $M^{(r-1,1)}$ affords the natural integral representation of S_r as $r \times r$ permutation matrices.

Given a λ -tableau t , we let $C(t)$ be the subgroup of S_r consisting of those elements which fix setwise the columns of t . Define $b_t \in \mathbf{Z}S_r$ by

$$b_t = \sum_{\tau \in C(t)} \text{sgn}(\tau)\tau.$$

The *polytabloid* corresponding to t is the element $e_t \in M^\lambda$ defined by

$$e_t = \mathbf{t}b_t.$$

The *Specht module* S^λ is defined to be the submodule of M^λ spanned by the λ -polytabloids. An easy calculation shows that if $\sigma \in S_r$ then $(e_t)\sigma = e_{t\sigma}$, and so S^λ is cyclic, generated by any single polytabloid.

It follows from Theorem 4.12 of [4] that the rational $\mathbf{Q}S_n$ -modules $S^\lambda \otimes_{\mathbf{Z}} \mathbf{Q}$ for λ a partition of n are irreducible, and that they afford all the ordinary irreducible characters of S_n .

Let t_λ be the λ -tableau whose i -th column is $i_1, \dots, i_{\lambda'_i}$ when read from top to bottom. Note that the elements of $C(t_\lambda)$ permute the indices of symbols in $A(\lambda)$ while leaving the numbers unchanged.

2.2. Definition of the homomorphisms $f_{\mathcal{P}}$. Throughout this section, let $m, n \in \mathbf{N}$ and let λ be a partition of mn . After identifying S_{mn} with the symmetric group on the set $A(\lambda)$, the elements of the canonical permutation basis of $H^{(m^n)}$ are given by the following definition.

Definition. An *indexed set partition* of shape (m^n) and type λ is a set partition of $A(\lambda)$ into n sets each of size m .

Our notation allows us to pass easily from set families to indexed set partitions.

Definition. Let \mathcal{P} be a set family of shape (m^n) and type λ . Order the sets making up \mathcal{P} lexicographically, so that $\mathcal{P} = \{X_1, \dots, X_n\}$ where $X_1 < X_2 < \dots < X_n$. The *indexed set partition associated to \mathcal{P}* is the indexed set partition of type λ obtained by appending indices to the elements of the sets X_1, \dots, X_n so that the elements of X_1 all get the index 1, and an element $i \in X_r$ is given the smallest index not appended to any i appearing in X_1, \dots, X_{r-1} .

For example, the indexed set partition associated to the closed set family

$$\mathcal{Q} = \{\{1, 2, 3\}, \{1, 2, 4\}, \{1, 2, 5\}, \{1, 3, 4\}\}$$

of type $(5, 4, 2, 1)$ is

$$u = \{\{1_1, 2_1, 3_1\}, \{1_2, 2_2, 4_1\}, \{1_3, 2_3, 5_1\}, \{1_4, 3_2, 4_2\}\} \in H^{(3^4)}.$$

Since S^λ is generated by the polytabloid e_{t_λ} , any homomorphism from S^λ is determined by its effect on e_{t_λ} . Specifically, if $f : S^\lambda \rightarrow M$ is a homomorphism of S_{mn} -modules, t is a λ -tableau and $\sigma \in S_{mn}$ is such that $t_\lambda \sigma = t$, then $e_t f = (e_{t_\lambda} f)\sigma$.

Definition. Let \mathcal{P} be a set family of shape (m^n) and type λ . We define $f_{\mathcal{P}} : S^\lambda \rightarrow H^{(m^n)}$ by $e_{t_\lambda} f_{\mathcal{P}} = ub_{t_\lambda}$ where u is the indexed set partition associated to \mathcal{P} .

If \mathcal{Q} is as above then the homomorphism $f_{\mathcal{Q}} : S^{(5,4,2,1)} \rightarrow H^{(3^4)}$ is defined on the generator $e_{t_{(5,4,2,1)}}$ of $S^{(5,4,2,1)}$ by

$$e_{t_{(5,4,2,1)}} \mapsto \{\{1_1, 2_1, 3_1\}, \{1_2, 2_2, 4_1\}, \{1_3, 2_3, 5_1\}, \{1_4, 3_2, 4_2\}\} b_{t_{(5,4,2,1)}}.$$

We remark that while we have, for definiteness, given an explicit scheme for passing from set families to indexed set partitions, a different choice will at most lead to changes of sign in the maps $f_{\mathcal{P}}$. For example, if in our index appending scheme, the lexicographic order on sets is replaced with the colexicographic order, then the homomorphism above would instead be defined by

$$\begin{aligned} e_{t_{(5,4,2,1)}} &\mapsto \{\{1_1, 2_1, 3_1\}, \{1_2, 2_2, 4_1\}, \{1_3, 3_2, 4_2\}, \{1_4, 2_3, 5_1\}\} b_{t_{(5,4,2,1)}} \\ &= -\{\{1_1, 2_1, 3_1\}, \{1_2, 2_2, 4_1\}, \{1_3, 2_3, 5_1\}, \{1_4, 3_2, 4_2\}\} b_{t_{(5,4,2,1)}}. \end{aligned}$$

3. PROOF OF THEOREM 1.1

For technical reasons it will be useful to deal with the case $m = 1$ separately. The only closed set family of shape (1^n) is $\mathcal{P} = \{\{1\}, \{2\}, \dots, \{n\}\}$, which has type (n) . The homomorphism $f_{\mathcal{P}} : S^{(n)} \rightarrow H^{(1^n)}$ is defined by

$$e_{t_{(n)}} \mapsto \{\{1_1\}, \{2_1\}, \dots, \{n_1\}\}.$$

Since $S^{(n)}$ is the trivial FS_n -module, this map is clearly well-defined and injective.

To show that the homomorphisms $f_{\mathcal{P}}$ are well-defined when $m \geq 3$, we shall use the description of the Specht module S^λ given by Garnir relations. The following lemma states a suitable form of these relations in our numbers-and-indices notation.

Lemma 3.1. *Let U be a \mathbf{Z} -free $\mathbf{Z}S_r$ -module, let λ be a partition of r and let $t = t_\lambda$. If $u \in U$ is such that*

$$ub_t \sum_{\sigma \in S_{X \cup Y}} \sigma \operatorname{sgn}(\sigma) = 0$$

for every pair of subsets

$$\begin{aligned} X &\subseteq \{i_j : 1 \leq j \leq \lambda'_i\}, \\ Y &\subseteq \{(i+1)_j : 1 \leq j \leq \lambda'_{i+1}\} \end{aligned}$$

such that $|X| + |Y| > \lambda'_i$, then there is a homomorphism of $\mathbf{Z}S_r$ -modules $f : S^\lambda \rightarrow U$ such that $e_t f = ub_t$.

Proof. It follows from the remark at the top of page 102 of [3] that the kernel of the surjective map $\mathbf{Z}S_r \rightarrow S^\lambda$ defined by $x \mapsto e_t x$ is generated, as a right $\mathbf{Z}S_r$ -ideal, by elements of the following two types:

- $1 - (i_j, i_k)$ for $i_j, i_k \in A(\lambda)$;
- $G_{X,Y} = \sum \sigma \operatorname{sgn}(\sigma)$, where X and Y are as in the statement of the lemma and the sum is over a set of right-coset representatives for the cosets of $S_X \times S_Y$ in $S_{X \cup Y}$.

Clearly ub_t is killed by elements of the first type, so to prove the lemma, it will suffice to show that $ub_t G_{X,Y} = 0$ for each $G_{X,Y}$. As in the proof of Theorem 7.2 in [4], we set $\bar{S}_Z = \sum_{\sigma \in Z} \sigma \operatorname{sgn}(\sigma)$ for a subset Z of S_r . Note that $\bar{S}_X \bar{S}_Y G_{X,Y} = \bar{S}_{X \cup Y}$. By hypothesis $ub_t \bar{S}_{X \cup Y} = 0$, so we have

$$|X|! |Y|! ub_t G_{X,Y} = 0.$$

Since U is assumed to be free as a \mathbf{Z} -module, it follows that $ub_t G_{X,Y} = 0$, as required. \square

To show that the homomorphisms $f_{\mathcal{P}}$ are well-defined it suffices to check that $e_{t_\lambda} f_{\mathcal{P}}$ satisfies the relations in the previous lemma.

Proposition 3.2. *Let $m \geq 3$ be odd and let $n \in \mathbf{N}$. Suppose that \mathcal{P} is a closed set family of shape (m^n) and type λ and that u is the indexed set partition associated to \mathcal{P} . Let $t = t_\lambda$. If X and Y are as in the statement of Lemma 3.1 then*

$$ub_t \sum_{\sigma \in S_{X \cup Y}} \sigma \operatorname{sgn}(\sigma) = 0.$$

Proof. Let $\tau \in C(t)$. Suppose that there exist $i_x \in X$ and $(i+1)_y \in Y$ which appear in the same set in $u\tau$. Then $u\tau(1 - (i_x, (i+1)_y)) = 0$, and, taking coset representatives for $\langle (i_x, (i+1)_y) \rangle$ in $S_{X \cup Y}$, we see that $u\tau \sum_{\sigma \in S_{X \cup Y}} \sigma \operatorname{sgn}(\sigma) = 0$.

It therefore suffices to show that

$$u \sum_{\tau \in C'} \tau \operatorname{sgn}(\tau) \sum_{\sigma \in S_{X \cup Y}} \sigma \operatorname{sgn}(\sigma) = 0 \quad (1)$$

where C' is the set of $\tau \in C(t)$ such that no set in $u\tau$ meets both X and Y . We may assume that C' is non-empty.

Let $\vartheta \in C'$ and let $v = u\vartheta$. None of the $|Y|$ sets in v meeting Y can contain an element of X . At most $\lambda'_i - |X|$ of them can contain an element

of the complementary set $X' = \{i_{x'} : 1 \leq x' \leq \lambda'_i, i_{x'} \notin X\}$. By hypothesis $\lambda'_i - |X| < |Y|$. Hence the number of sets in v which meet Y is strictly greater than the number of sets in v which meet X' . It follows, on casting out the sets meeting both X' and Y , that there is a set

$$B = \{c(1)_{b(1)}, c(2)_{b(2)}, \dots, c(m-1)_{b(m-1)}, (i+1)_y\}$$

in v which meets Y but not X' . Moreover, since \mathcal{P} is closed, we may choose B so that, for some indices $a(1), \dots, a(m-1), x$, the set

$$A = \{c(1)_{a(1)}, c(2)_{a(2)}, \dots, c(m-1)_{a(m-1)}, i_x\}$$

also appears in v . Note that the indices $a(1), \dots, a(m-1), x$ are determined by the numbers $c(1), \dots, c(m-1), i$.

Let

$$\pi = (c(1)_{a(1)}, c(1)_{b(1)}) \cdots (c(m-1)_{a(m-1)}, c(m-1)_{b(m-1)}).$$

Since $B(i_x, (i+1)_y) = A\pi$ we have

$$v(i_x, (i+1)_y) = v\pi. \tag{2}$$

No set in $v\pi$ meets both X and Y , so since $v\pi = u\vartheta\pi$, we have $\vartheta\pi \in C'$. Thus $u\vartheta$ and $u\vartheta\pi$ are distinct summands of $u \sum_{\tau \in C'} \tau \operatorname{sgn}(\tau)$, appearing with the same sign. (This is the only point where we use our hypotheses on m .) If $\sigma_1, \dots, \sigma_s$ is a set of right coset representatives for the cosets of $\langle\langle i_x, (i+1)_y \rangle\rangle$ in $S_{X \cup Y}$ then, by (2),

$$(u\vartheta + u\vartheta\pi) \sum_{\sigma \in S_{X \cup Y}} \sigma \operatorname{sgn}(\sigma) = v(1 + \pi)(1 - (i_x, (i+1)_y)) \sum_{r=1}^s \sigma_r \operatorname{sgn}(\sigma_r) = 0.$$

Let H be the subgroup of $C(t)$ of elements that fix all the entries in columns i and $i+1$ of t . We have shown that given any $\vartheta \in C'$, there exists a non-identity even permutation $\pi_\vartheta \in H$ such that $\vartheta\pi_\vartheta \in C'$ and the contributions to (1) from $u\vartheta$ and $u\vartheta\pi_\vartheta$ cancel.

Let $C'_{i,i+1}$ be the subset of C' of elements which only move entries in columns i and $i+1$ of t . Let $\vartheta, \vartheta' \in C'_{i,i+1}$. If $\vartheta\pi_\vartheta = \vartheta'$ then $\vartheta'\vartheta^{-1} = \pi_\vartheta$; since π_ϑ fixes the entries in columns i and $i+1$ of ϑ , this implies that $\vartheta' = \vartheta$. Hence if the sets $\{\vartheta, \vartheta\pi_\vartheta\}, \{\vartheta', \vartheta'\pi_{\vartheta'}\}$ meet then $\vartheta = \vartheta'$. We may therefore pair up the elements of $C'_{i,i+1}$ to show that

$$u \sum_{\vartheta \in C'_{i,i+1}} \vartheta \operatorname{sgn}(\vartheta) \sum_{\sigma \in S_{X \cup Y}} \sigma \operatorname{sgn}(\sigma) = 0.$$

There exist $\tau_1, \dots, \tau_k \in H$ such that

$$C' = \tau_1 C'_{i,i+1} \cup \dots \cup \tau_k C'_{i,i+1}$$

where the union is disjoint. Hence the left-hand-side of (1) is

$$\begin{aligned} u \sum_{r=1}^k \tau_r \operatorname{sgn}(\tau_r) \sum_{\vartheta \in C'_{i,i+1}} \vartheta \operatorname{sgn}(\vartheta) \sum_{\sigma \in S_{X \cup Y}} \sigma \operatorname{sgn}(\sigma) = \\ \left(u \sum_{\vartheta \in C'_{i,i+1}} \vartheta \operatorname{sgn}(\vartheta) \sum_{\sigma \in S_{X \cup Y}} \sigma \operatorname{sgn}(\sigma) \right) \sum_{r=1}^k \tau_r \operatorname{sgn}(\tau_r) = 0, \end{aligned}$$

as we required. \square

To complete the proof of Theorem 1.1, we must show that the homomorphisms $f_{\mathcal{P}}$ are injective. This follows from the following general result.

Lemma 3.3. *Let λ be a partition of r and let M be a \mathbf{Z} -free $\mathbf{Z}S_r$ -module with character π . If $f : S^\lambda \rightarrow M$ is a non-zero homomorphism of $\mathbf{Z}S_r$ -modules then f is injective.*

Proof. The homomorphism f induces a non-zero homomorphism

$$f' : S^\lambda \otimes_{\mathbf{Z}} \mathbf{Q} \rightarrow M \otimes_{\mathbf{Z}} \mathbf{Q}.$$

Since $S^\lambda \otimes_{\mathbf{Z}} \mathbf{Q}$ is irreducible (see [4, Theorem 4.12]), f' is injective. Hence the original map f is also injective. \square

4. MINIMAL CONSTITUENTS OF FOULKES CHARACTERS

In this section we prove Theorems 1.2 and 1.3 on the minimal constituents of the Foulkes characters $\phi^{(m^n)}$.

4.1. Even case. Let m be even. To prove part (i) of Theorem 1.2, we must show that the unique minimal constituent of the Foulkes character $\phi^{(m^n)}$ is $\chi^{(m^n)}$. We do this using properties of the Young permutation modules defined in §2.1.

Let $q : M^{(m^n)} \rightarrow H^{(m^n)}$ be the map which sends an (m^n) -tabloid to the indexed set partition whose member sets are the rows of the tabloid. It is clear that q is a surjective homomorphism of $\mathbf{Z}S_{mn}$ -modules. It easily follows that if $\pi^{(m^n)}$ is the character of $M^{(m^n)}$ then

$$\langle \pi^{(m^n)}, \chi^\lambda \rangle \geq \langle \phi^{(m^n)}, \chi^\lambda \rangle$$

for every partition λ of mn .

The constituents of the character $\pi^{(m^n)}$ are given by Young's rule: see [4, Chapter 14]. Young's rule implies that $\chi^{(m^n)}$ is the unique minimal constituent of $\pi^{(m^n)}$, and that $\langle \pi^{(m^n)}, \chi^{(m^n)} \rangle = 1$. The following lemma implies that there is a non-zero homomorphism $S^\lambda \rightarrow H^{(m^n)}$, and hence $\langle \phi^{(m^n)}, \chi^{(m^n)} \rangle \geq 1$. Therefore $\chi^{(m^n)}$ is the unique minimal constituent of $\phi^{(m^n)}$ and $\langle \phi^{(m^n)}, \chi^{(m^n)} \rangle = 1$, as required.

Lemma 4.1. *The Specht module $S^{(m^n)}$ is not contained in the kernel of q .*

Proof. Let $t = t_{(m^n)}$ and let $\mathbf{t} \in M^{(m^n)}$ be the corresponding tabloid. By definition of q ,

$$\mathbf{t}q = \{ \{1_1, 2_1, \dots, m_1\}, \dots, \{1_n, 2_n, \dots, m_n\} \}.$$

Let $H \leq C(t)$ be the subgroup of the column permuting group of t that permutes as blocks for its action the rows of t . (As an abstract group, $H \cong S_n$.) For example, if $m = 4$ and $n = 3$, then H is generated by $(1_1, 1_2)(2_1, 2_2)(3_1, 3_2)(4_1, 4_2)$ and $(1_1, 1_2, 1_3)(2_1, 2_2, 2_3)(3_1, 3_2, 3_3)(4_1, 4_2, 4_3)$.

Note that since m is even, every element of H is an even permutation. Let $K \leq C(t)$ be the subgroup of permutations which fix the elements $1_1, \dots, 1_n$ in the first column of t . It is easy to see that $C(t) = HK$, and so

$$e_t q = (\mathbf{t}q) \sum_{\tau \in C(t)} \tau \operatorname{sgn}(\tau) = (\mathbf{t}q) \sum_{\pi \in H} \pi \sum_{\sigma \in K} \sigma \operatorname{sgn}(\sigma) = n! (\mathbf{t}q) \sum_{\sigma \in K} \sigma \operatorname{sgn}(\sigma).$$

The summands on the right-hand-side are distinct basis elements of $H^{(m^n)}$, hence $e_t q \neq 0$. \square

4.2. Odd case. We start with the following general form for a homomorphism from a Specht module into a Foulkes module $H^{(m^n)}$ when m is odd.

Proposition 4.2. *Let m be odd and let $n \in \mathbf{N}$. Let λ be a partition of mn and let $f : S^\lambda \rightarrow H^{(m^n)}$ be a homomorphism of $\mathbf{Z}S_{mn}$ -modules. Let $t = t_\lambda$. There exist set families $\mathcal{P}_1, \dots, \mathcal{P}_k$ of shape (m^n) and type λ and integers a_1, \dots, a_k such that*

$$e_t f = a_1 u_1 b_t + \dots + a_k u_k b_t$$

where u_i is the indexed set partition associated to \mathcal{P}_i .

Proof. Let

$$e_t f = \sum_u c_u u$$

where the sum is over all indexed set partitions u of shape (m^n) and type λ and $c_u \in \mathbf{Z}$.

Let u be such an indexed set partition. If the symbols i_j and i_k appear in the same set in u then, since $e_t(i_j, i_k) = -e_t$ and $u(i_j, i_k) = u$, we must have $c_u = 0$. Now suppose that u contains two sets

$$\begin{aligned} &\{c(1)_{a(1)}, c(2)_{a(2)}, \dots, c(m)_{a(m)}\}, \\ &\{c(1)_{b(1)}, c(2)_{b(2)}, \dots, c(m)_{b(m)}\} \end{aligned}$$

which become equal if the indices are removed. Let

$$\tau = (c(1)_{a(1)}, c(1)_{b(1)}) \dots (c(m)_{a(m)}, c(m)_{b(m)}).$$

Since m is odd, τ is an odd permutation in $C(t)$. Hence $e_t\tau = -e_t$ and $u\tau = u$, and again we have $c_u = 0$.

These remarks show that if $c_u \neq 0$ then removing indices from the symbols in the sets making up u leaves a set family of shape (m^n) and type λ . If removing indices from u and v gives the same set family then $v = u\tau$ for some $\tau \in C(t)$. Since $e_t\tau = \text{sgn}(\tau)e_t$ we have $c_v = \text{sgn}(\tau)c_u$. The proposition follows. \square

We also need a corollary of the following combinatorial proposition, which will be used again in §5 below.

Proposition 4.3. *If \mathcal{P} is a minimal set family then \mathcal{P} is closed.*

Proof. It will be necessary in this proof to extend the definition of type so that it applies to all set families. We define the *conjugate type* of a set family \mathcal{P} of shape (m^n) to be the composition α such that α_i is the number of sets in \mathcal{P} containing i . Note that if α is a partition, then α' is the type (in the usual sense) of \mathcal{P} .

Let \mathcal{P} be a set family of shape (m^n) and type λ . Suppose that \mathcal{P} is not closed. We may find $A \in \mathcal{P}$ and $i+1 \in \mathbf{N}$ such that $i+1 \in A$ and the set

$$B = A \setminus \{i+1\} \cup \{i\}$$

is not in \mathcal{P} . Let \mathcal{Q} be the set family obtained from \mathcal{P} by removing A and adding B . If $\alpha = \lambda'$, then the conjugate type of \mathcal{Q} is β where $\beta_i = \alpha_i + 1$, $\beta_{i+1} = \alpha_{i+1} - 1$ and $\beta_j = \alpha_j$ if $j \neq i, i+1$. Hence $\beta \triangleright \alpha$ (where \triangleright now refers to the dominance order on compositions). Iterating this construction, we will reach a closed set family \mathcal{R} of conjugate type γ where $\gamma \triangleright \alpha$. Since \mathcal{R} is closed, γ is a partition. If $\nu = \gamma'$ then, $\nu' \triangleright \lambda'$, and so $\nu \triangleleft \lambda$. Thus \mathcal{R} has smaller type than \mathcal{P} and so \mathcal{P} is not minimal. \square

Corollary 4.4. *If $\langle \phi^{(m^n)}, \chi^\lambda \rangle \geq 1$ then there is a minimal set family of shape (m^n) and type μ where $\mu \trianglelefteq \lambda$, and this set family is closed.*

Proof. The hypothesis implies that there is an injective homomorphism of $\mathbf{Q}S_{mn}$ -modules

$$f : S^\lambda \otimes_{\mathbf{Z}} \mathbf{Q} \rightarrow H^{(m^n)} \otimes_{\mathbf{Z}} \mathbf{Q}.$$

Let

$$e_{t_\lambda} f = \sum_u c_u u$$

where $c_u \in \mathbf{Q}$ and the sum is over all indexed set partitions u of shape (m^n) and type λ . For each such u , let $c_u = a_u/b_u$ where $a_u, b_u \in \mathbf{Z}$. Let m be the product of all the b_u . It is easy to see that the map $g : S^\lambda \rightarrow H^{(m^n)}$ defined by

$$e_{t_\lambda} g = \sum_u m c_u u$$

is a well-defined injective homomorphism of $\mathbf{Z}S_{mn}$ -modules.

Applying Proposition 4.2 to g , we see that there is a set family of shape (m^n) and type λ . Proposition 4.3 implies that a set family of minimal type $\triangleleft \lambda$ is closed. \square

We are now ready to prove part (ii) of Theorem 1.2. Suppose that χ^λ is a minimal constituent of $\phi^{(m^n)}$. By Corollary 4.4 there is a minimal set family \mathcal{Q} of type μ where $\mu \trianglelefteq \lambda$. Since \mathcal{Q} is closed, it follows from Theorem 1.1 that there is a non-zero homomorphism $S^\mu \rightarrow H^{(m^n)}$, and so $\langle \phi^{(m^n)}, \chi^\mu \rangle \geq 1$. Therefore $\mu = \lambda$ and \mathcal{Q} is a minimal set family of type λ .

Conversely, suppose that there is a minimal set family \mathcal{P} of type λ . By Proposition 4.3, \mathcal{P} is closed, and so it follows from Theorem 1.1 that χ^λ is a summand of $\phi^{(m^n)}$. If there is a summand χ^μ of $\phi^{(m^n)}$ with $\mu \triangleleft \lambda$, then by Proposition 4.2, there is a set family of type μ ; this contradicts the minimality of \mathcal{P} . Therefore χ^λ is a minimal constituent of $\phi^{(m^n)}$.

4.3. Proof of Theorem 1.3. Suppose that χ^λ is a minimal constituent of $\phi^{(m^n)}$. By Theorem 1.2, λ is the type of a minimal set family of shape (m^n) . Let u_1, \dots, u_d be the indexed set partitions associated to the set families $\mathcal{P}_1, \dots, \mathcal{P}_d$ of shape (m^n) and type λ . Proposition 4.3 implies that the \mathcal{P}_r are closed. By Proposition 4.2 we know that if $f : S^\lambda \rightarrow H^{(m^n)}$ is a homomorphism of $\mathbf{Z}S_{mn}$ -modules, then there exist $a_1, \dots, a_d \in \mathbf{Z}$ such that

$$e_t f = a_1 u_1 b_t + \dots + a_d u_d b_t$$

where $t = t_\lambda$. Hence

$$f = a_1 f_{\mathcal{P}_1} + \dots + a_d f_{\mathcal{P}_d}.$$

To show that homomorphisms $f_{\mathcal{P}_1}, \dots, f_{\mathcal{P}_d}$ are linearly independent it suffices to show that the images of e_t ,

$$e_t f_{\mathcal{P}_r} = \sum_{\tau \in C(t)} u_r \tau \operatorname{sgn}(\tau)$$

are linearly independent. Given $\tau \in C(t)$ and $r \in \{1, \dots, d\}$, we can easily recover \mathcal{P}_r from $u_r \tau$ by removing the indices from the symbols in the sets making up $u_r \tau$. Therefore each $e_t f_{\mathcal{P}_r}$ is a sum of distinct basis elements of $H^{(m^n)}$; as such, they are linearly independent.

5. SET FAMILIES AND PARTITIONS

5.1. Minimal and unique set families. We proved in Proposition 4.3 that minimal set families are closed. The following proposition implies that if \mathcal{P} is the unique set family of its shape and type, then \mathcal{P} is minimal. This establishes the following chain of implications on set families of a given shape:

$$\text{unique of its type} \implies \text{minimal} \implies \text{closed}. \quad (3)$$

Proposition 5.1. *If \mathcal{P} is a set family of shape (m^n) and type λ and $\mu \triangleright \lambda$ then there are two distinct set families of shape (m^n) and type μ .*

Proof. We may assume that λ and μ are neighbours in the dominance order, so μ is obtained from λ by moving a box upwards in its Young diagram. Suppose the box is moved from column i to column $j > i$. We have $\mu'_i = \lambda'_i - 1$, $\mu'_j = \lambda'_j + 1$ and $\mu'_k = \lambda'_k$ if $k \neq i, j$.

The sets in \mathcal{P} either contain both i and j , or i alone, or j alone, or neither. Since $\lambda'_i - \lambda'_j = \mu'_i - \mu'_j + 2$, at least two more sets contain i alone than contain j alone. Hence there are two sets

$$\begin{aligned} A &= \{x(1), x(2), \dots, x(m-1), i\} \in \mathcal{P}, \\ A' &= \{x'(1), x'(2), \dots, x'(m-1), i\} \in \mathcal{P} \end{aligned}$$

such that

$$\begin{aligned} B &= \{x(1), x(2), \dots, x(m-1), j\} \notin \mathcal{P}, \\ B' &= \{x'(1), x'(2), \dots, x'(m-1), j\} \notin \mathcal{P}. \end{aligned}$$

Let \mathcal{Q} be the set family obtained from \mathcal{P} by removing A and adding B , and let \mathcal{Q}' be the set family obtained from \mathcal{P} by removing A' and adding B' . Then \mathcal{Q} and \mathcal{Q}' are two different set families of type μ . \square

To complete the proof of Theorem 1.4, we must show that neither of the implications in the chain (3) is reversible. We shall use the following definition throughout the remainder of this section.

Definition. Let A be a subset of the natural numbers. The *downset* of A , denoted A^{\preceq} , is the set family consisting of all subsets X such that $X \preceq A$.

It is obvious that a set family is closed if and only if it is a union of downsets.

Our first example concerns the downset $\mathcal{P} = \{2, 4, 6, 8\}^{\preceq}$. This is a closed set family of shape (4^{42}) , but it is not unique for its type, since the set family

$$\mathcal{Q} = \mathcal{P} \setminus \{\{2, 4, 6, 8\}, \{1, 3, 5, 7\}\} \cup \{\{1, 2, 7, 8\}, \{3, 4, 5, 6\}\}$$

has the same shape and type as \mathcal{P} . Neither is \mathcal{P} minimal. This is most easily seen by noting that since \mathcal{Q} contains $\{1, 3, 5, 8\}$ but not $\{1, 3, 5, 7\}$, \mathcal{Q} is not closed. Following the proof of Proposition 4.3 leads one to (amongst others) the closed set family

$$\begin{aligned} \mathcal{R} &= \mathcal{Q} \setminus \{\{2, 4, 5, 8\}\} \cup \{\{1, 3, 5, 7\}\} \\ &= \mathcal{P} \setminus \{\{2, 4, 6, 8\}, \{2, 4, 5, 8\}\} \cup \{\{1, 2, 7, 8\}, \{3, 4, 5, 6\}\} \\ &= \{1, 2, 7, 8\}^{\preceq} \cup \{1, 4, 6, 8\}^{\preceq} \cup \{2, 3, 6, 8\}^{\preceq} \cup \{2, 4, 6, 7\}^{\preceq} \cup \{3, 4, 5, 6\}^{\preceq} \end{aligned}$$

which has smaller type than \mathcal{P} and \mathcal{Q} . (In fact, \mathcal{R} is minimal.)

The following example of a minimal set family that is not unique for its type was found by a computer search for minimal set families with prescribed shape and maximum entry. (A description of the algorithm used and accompanying source code is available from the second author's website, www.maths.bris.ac.uk/~mazmjw.) Let

$$\begin{aligned} \mathcal{S} &= \{1, 5, 9\}^{\preceq} \cup \{1, 6, 8\}^{\preceq} \cup \{2, 6, 7\}^{\preceq} \cup \{3, 4, 8\}^{\preceq} \cup \{3, 5, 6\}^{\preceq}, \\ \mathcal{S}' &= \{1, 4, 9\}^{\preceq} \cup \{1, 7, 8\}^{\preceq} \cup \{2, 3, 9\}^{\preceq} \cup \{2, 4, 8\}^{\preceq} \\ &\quad \cup \{2, 5, 7\}^{\preceq} \cup \{3, 4, 7\}^{\preceq} \cup \{4, 5, 6\}^{\preceq}. \end{aligned}$$

The set families \mathcal{S} and \mathcal{S}' both have type $\lambda = (24, 19, 17, 16, 13, 12, 10, 8, 4)'$ and shape (3^{41}) . Using the computer to enumerate all closed set families of shape (3^{41}) with maximum entry ≤ 9 confirms that there are no set families of shape (3^{41}) with type $\triangleleft \lambda$.

5.2. Constructing types of minimal set families. Let $m, n \in \mathbf{N}$. Given a partition ν of $n - 1$ with $k \leq m$ parts, we let $(m^n) \star \nu$ denote the partition obtained from (m^n) by deleting ν_i boxes from column $m + 1 - i$ and then

adding ν_i boxes to row i , for each i such that $1 \leq i \leq k$. This construction is illustrated in Figure 1 below. (If $\nu_1 = a$ then we add boxes to rows $1, 2, \dots, k$, and remove boxes from rows $n - a + 1, \dots, n$; since $k + a \leq n$, the partition $(m^n) \star \nu$ is well-defined.)




FIGURE 1. Construction of the partition $(5^8) \star (4, 2, 1)$.

Our first object in this section is to prove the following proposition.

Proposition 5.2. *Let ν be a partition of $n - 1$ with at most m parts. There is a unique set family of shape (m^n) and type $(m^n) \star \nu$.*

Proof. Suppose that ν has exactly k parts. Let $A = \{1, 2, \dots, m\}$. It is easily checked that

$$\mathcal{P} = \{A\} \cup \{A \setminus \{m - i + 1\} \cup \{m + j\} : 1 \leq i \leq k, 1 \leq j \leq \nu_i\}$$

is a set family of shape (m^n) and type $(m^n) \star \nu$.

Now suppose that \mathcal{Q} is a set family of this shape and type. We may write

$$\mathcal{Q} = \{A \setminus B_r \cup C_r : 1 \leq r \leq n\}$$

for some subsets $B_r, C_r \subseteq \mathbf{N}$ such that $B_r \cap C_r = \emptyset$ for $1 \leq r \leq n$. For each i such that $1 \leq i \leq k$, exactly ν_i of the sets B_1, \dots, B_n contain $m - i + 1$. It follows that $|B_1| + \dots + |B_n| = n - 1$, and so one of the sets B_r is empty, and the remaining $n - 1$ are singletons. Hence $A \in \mathcal{Q}$, and for each i such that $1 \leq i \leq k$, there are exactly ν_i sets in \mathcal{Q} of the form $A \setminus \{m - i + 1\} \cup \{c\}$ where $c \in \{m + 1, \dots, m + \nu_i\}$. Looking first at the case $i = 1$, we see that $A \setminus \{m\} \cup \{m + j\} \in \mathcal{Q}$ for each j such that $1 \leq j \leq \nu_1$. Iterating this argument for $i = 2, \dots, k$ shows that $\mathcal{Q} = \mathcal{P}$. \square

We isolate the following corollary of Proposition 5.2

Corollary 5.3. *Let m be odd and let $n \in \mathbf{N}$. If ν is a partition of $n - 1$ with at most m parts then*

$$\left\langle \phi^{(m^n)}, \chi^{(m^n) \star \nu} \right\rangle = 1$$

Moreover $\langle \phi^{(m^n)}, \chi^\mu \rangle = 0$ if $\mu \triangleleft \chi^{(m^n) \star \nu}$.

Proof. By Theorem 1.4, $(m^n) \star \nu$ is the type of a minimal set family of shape (m^n) . The result now follows from Theorem 1.3. \square

It is natural to ask when every minimal constituent $\phi^{(m^n)}$ arises from this construction. We shall show in Proposition 5.7 below that this is the case if and only if $n \leq 5$ or $m = 1$. We begin with the following straightforward lemma.

Lemma 5.4. *Let \mathcal{P} be a closed set family of shape (m^n) where $m \geq n$. If $X \in \mathcal{P}$ then $X \supseteq \{1, 2, \dots, m - n + 1\}$.*

Proof. Suppose that the smallest number not present in X is $m - t + 1$, so $X = \{1, 2, \dots, m - t, x(1), \dots, x(t)\}$ for some $x(r) > m - t + 1$. For each r such that $1 \leq r \leq t$, the set $X \setminus \{x(r)\} \cup \{m - t + 1\}$ is majorized by X , so must lie in \mathcal{P} . Hence $|\mathcal{P}| \geq t + 1$, and so $t \leq n - 1$. It follows that $m - t \geq m - n + 1$, as required. \square

In the proof of the following lemma, a further construction on partitions will be found useful: given a partition λ with exactly k parts, each of size $\geq c$, let $\lambda - (c^k)$ denote the partition obtained from $(\lambda_1 - c, \dots, \lambda_k - c)$ by removing any final parts of size zero.

Lemma 5.5. *Let $m \in \mathbf{N}$ and let $n \leq 5$. If \mathcal{P} is a closed set family of shape (m^n) then \mathcal{P} has type $(m^n) \star \nu$ for some partition ν of $n - 1$ with at most m parts.*

Proof. If $m < n$ then the result can be checked directly. For example, when $m = 4$ and $n = 5$, the set family \mathcal{P} must consist of 5 sets taken from the first 5 levels of the lattice of 4-subsets of \mathbf{N} under the majorization order. It is easily seen from Figure 2 below that there are 5 possibilities for \mathcal{P} , namely $\{1, 2, 3, 8\}^{\preceq}$, $\{1, 2, 3, 7\}^{\preceq} \cup \{\{1, 2, 4, 5\}\}$, $\{1, 2, 4, 6\}^{\preceq}$, $\{1, 3, 4, 5\}^{\preceq} \cup \{\{1, 2, 3, 6\}\}$ and $\{2, 3, 4, 5\}^{\preceq}$. The types of these set families are $(4^5) \star \nu$ where $\nu = (4), (3, 1), (2, 2), (2, 1, 1)$ and (1^4) respectively.

Now suppose that $m \geq n$. By Lemma 5.4, we know that every set in \mathcal{P} contains $\{1, 2, \dots, m - n + 1\}$. Let \mathcal{Q} be the set family obtained by removing the elements $1, 2, \dots, m - n + 1$ from every set in \mathcal{P} , and then subtracting

$m - n + 1$ from each remaining element. The shape of \mathcal{Q} is $((n - 1)^n)$. If \mathcal{P} has type λ , where λ has exactly k parts, then \mathcal{Q} has type $\lambda - ((m - n + 1)^k)$. By the result already proved, we know that

$$\lambda - ((m - n + 1)^k) = ((n - 1)^n) \star \nu$$

for some partition ν of $n - 1$. Since $\lambda - ((m - n + 1)^k)$ is a partition of $(n - 1)k$, we must have $k = n$. It is now easy to see that $\lambda = (m^n) \star \nu$, as required. \square




FIGURE 2. The first five levels of the lattice of 4-subsets of \mathbf{N} under the majorization order.

We now show that Lemma 5.5 is false when $n \geq 6$ and $m \geq 2$.

Lemma 5.6. *Let $n \geq 6$ and let $m \geq 2$. The unique set family of shape (m^n) and type $(m + n - 4, m + 2, m + 2, (m - 1)^{n-6}, (m - 2)^3)$ is*

$$\{1, 2, \dots, m - 2, m + 1, m + 2\}^{\preceq} \cup \{1, 2, \dots, m - 2, m - 1, m + n - 4\}^{\preceq}.$$

This type is not of the form $(m^n) \star \nu$ for any partition ν of $n - 1$ with at most m parts.

Proof. It is routine to check that this set family does have the claimed type. Suppose that \mathcal{P} is a set family of this type. Every set in \mathcal{P} contains $\{1, 2, \dots, m - 2\}$, so, as in the proof of Lemma 5.5, we may pass to a set family \mathcal{Q} of shape (2^n) and type $(n - 2, 4, 4, 1^{n-6})$ by removing the elements $1, 2, \dots, m - 2$ from every set, and then subtracting $m - 2$ from every remaining element. It suffices to show that $\mathcal{Q} = \{3, 4\}^{\preceq} \cup \{1, n - 2\}^{\preceq}$. Of the $n - 3$ sets in \mathcal{Q} which contain 1, at most one can contain 2, so 2 must appear

in two of the three remaining sets in \mathcal{Q} . A similar argument with 3 and 4 shows that $\{2, 3\}, \{2, 4\}, \{3, 4\} \in \mathcal{Q}$. It is now clear that \mathcal{Q} is as claimed.

If the final assertion in the lemma is false, then the type of Q is $(2^n) \star \nu$ where ν is a partition of $n-1$ with at most 2 parts. However, $(n-2, 4, 4, 1^{n-6})$ has three parts of size > 2 , so this is impossible. \square

We now use Lemmas 5.5 and 5.6 to prove the following proposition.

Proposition 5.7. *Let m be odd and let $n \in \mathbf{N}$. Every minimal constituent of $\phi^{(m^n)}$ is of the form $\chi^{(m^n) \star \nu}$ for some partition ν of $n-1$ with at most m parts if and only if $m = 1$ or $n \leq 5$. \square*

Proof. Since $\phi^{(1^n)} = \chi^{(n)}$ and $(n) = (1^n) \star (n-1)$, the proposition holds when $m = 1$. When $n \leq 5$ it follows from Lemma 5.5, Theorem 1.2 and Theorem 1.4. If $n \geq 6$ then, since the unique set family constructed in Lemma 5.6 is minimal by Theorem 1.4, there is a minimal constituent of $\phi^{(m^n)}$ not of the form $\chi^{(m^n) \star \nu}$ for any partition ν of $n-1$. \square

6. MINIMAL CONSTITUENTS OF GENERALIZED FOULKES CHARACTERS

We end by showing how to construct the minimal constituents of a wider class of permutation characters. Let μ be a partition of N with largest part of size a . If μ has exactly $n(i)$ parts of length i for each i such that $1 \leq i \leq a$, we define the *generalized Foulkes character* ϕ^μ to be the induced character

$$\phi^\mu = \left(\phi^{(1^{n(1)})} \times \phi^{(2^{n(2)})} \times \dots \times \phi^{(a^{n(a)})} \right) \uparrow_{S_{n(1)} \times S_{2n(2)} \times \dots \times S_{an(a)}}^{S_N}.$$

If $\mu = (m^n)$ for some $m, n \in \mathbf{N}$ then this definition agrees with the one given earlier.

Our main aim in this section is to prove Proposition 6.1 below describing the minimal constituents of generalized Foulkes characters. To state this result we need one final construction on partitions.

Definition. Given λ a partition of r and μ a partition of s , we denote by $\lambda \cup \mu$ the partition of $r+s$ whose multiset of parts is the union of the multisets of parts of λ and μ .

For example, $(5, 2) \cup (3, 2, 2, 1, 1) = (5, 3, 2, 2, 2, 1, 1)$.

Proposition 6.1. *Let $\mu = (m(1)^{n(1)}) \cup \dots \cup (m(t)^{n(t)})$ where the $m(k)$ are distinct. If χ^λ is a minimal constituent of ϕ^μ then*

$$\lambda = \nu(1) \cup \nu(2) \cup \dots \cup \nu(t)$$




FIGURE 3. Use of the Littlewood–Richardson rule to show that $(\chi^{(6,3,2)} \times \chi^{(4,3,1)}) \uparrow_{S_{10} \times S_8}^{S_{18}}$ has $\chi^{(6,3,2) \cup (4,3,1)} = \chi^{(6,4,3,3,2,1)}$ as a summand.

where $\nu(k)$ is a partition of $m(k)n(k)$ and $\chi^{\nu(k)}$ is a minimal constituent of $\phi^{(m(k)n(k))}$ for each k such that $1 \leq k \leq t$.

Our proof of Proposition 6.1 uses the following two general lemmas.

Lemma 6.2. *Let λ be a partition of r and let μ be a partition of s . The unique minimal constituent of $(\chi^\lambda \times \chi^\mu) \uparrow_{S_r \times S_s}^{S_{r+s}}$ is $\chi^{\lambda \cup \mu}$.*

Proof. Let $\vartheta = (\chi^\lambda \times \chi^\mu) \uparrow_{S_r \times S_s}^{S_{r+s}}$. It follows from the description of the Littlewood–Richardson rule given in [4, Chapter 16] that $\chi^{\lambda \cup \mu}$ is a constituent of the character ϑ . A typical example, which shows how the parts of $\lambda \cup \mu$ may be obtained by adding numbers to λ , is given in Figure 3. Note that at step j , the lowest μ_j positions that are eligible to be filled receive a j . For an explanation of the notation and method used, the reader is referred to [4, Chapter 16].

The remainder of the proof can be completed using the easier Young’s rule. The character

$$\psi = (\chi^\lambda \times (1_{S_\mu} \uparrow^{S_s})) \uparrow_{S_r \times S_s}^{S_{r+s}}$$

certainly contains all the constituents of ϑ , so to prove the lemma, it suffices to show that ψ has $\chi^{\lambda \cup \mu}$ as its least constituent. This follows by induction on the number of parts of μ if we rewrite ψ as

$$(\chi^\lambda \times 1_{\mu_1} \times \cdots \times 1_{\mu_k}) \uparrow_{S_r \times S_{\mu_1} \times \cdots \times S_{\mu_k}}^{S_{r+s}}$$

and then repeatedly apply Young’s rule (see [4, Chapter 14]). \square

Lemma 6.3. *Let π be a character of S_r and let ϑ be a character of S_s . If χ^ν is a minimal constituent of $(\pi \times \vartheta) \uparrow_{S_r \times S_s}^{S_{r+s}}$ then $\nu = \lambda \cup \mu$ where χ^λ is a minimal constituent of π and χ^μ is a minimal constituent of ϑ .*

Proof. Let $\psi = (\pi \times \vartheta) \uparrow_{S_r \times S_s}^{S_{r+s}}$. It follows from Lemma 6.2 that if χ^ν is a minimal constituent of ψ then there are partitions λ and μ such that $\langle \pi, \chi^\lambda \rangle \geq 1$, $\langle \vartheta, \chi^\mu \rangle \geq 1$ and $\nu = \lambda \cup \mu$.

Suppose that χ^λ is not a minimal constituent of π . Then there exists a partition λ^* such that $\lambda^* \triangleleft \lambda$ and $\langle \pi, \chi^{\lambda^*} \rangle \geq 1$. By Lemma 6.2 we have

$$\langle \psi, \chi^{\lambda^* \cup \mu} \rangle \geq 1.$$

It is easily seen that $\lambda^* \cup \mu \triangleleft \lambda \cup \mu$; this contradicts the minimality of χ^ν . Therefore χ^λ is a minimal constituent of π and similarly, χ^μ is a minimal constituent of ϑ . \square

We are now ready to prove Proposition 6.1. Let $N = m(1)n(1) + \dots + m(t)n(t)$. Since the $m(r)$ are distinct,

$$\phi^\mu = \left(\phi^{(m(1)n(1))} \times \dots \times \phi^{(m(t)n(t))} \right) \uparrow_{S_{m(1)n(1)} \times \dots \times S_{m(t)n(t)}}^{S_N}.$$

The proposition now follows by repeated applications of Lemma 6.3.

We finish with the observation that the converse to Proposition 6.1 (and to Lemma 6.3) is false. This can be demonstrated using Corollary 5.3 in §5.2. It follows from this corollary that $\phi^{(5^5)}$ has

$$\chi^{(5^5) \star (2,1,1)} = \chi^{(7,6,6,4,2)} \quad \text{and} \quad \chi^{(5^5) \star (1^4)} = \chi^{(6,6,6,6,1)}$$

as minimal constituents. Similarly, $\phi^{(3^5)}$ has

$$\chi^{(3^5) \star (4)} = \chi^{(7,2,2,2,2)} \quad \text{and} \quad \chi^{(3^5) \star (3,1)} = \chi^{(6,4,2,2,1)}$$

as minimal constituents. It is clear that

$$(6, 6, 6, 6, 1) \cup (7, 2, 2, 2, 2) \triangleright (7, 6, 6, 4, 2) \cup (6, 4, 2, 2, 1).$$

Hence $\chi^{(6,6,6,6,1) \cup (7,2,2,2,2)}$ is not a minimal constituent of $\phi^{(5^5, 3^5)}$, even though it arises from the \cup -construction applied to a minimal constituent of $\phi^{(5^5)}$ and a minimal constituent of $\phi^{(3^5)}$.

REFERENCES

- [1] Suzie C. Dent and Johannes Siemons. On a conjecture of Foulkes. *J. Algebra*, 226(1):236–249, 2000.
- [2] H. O. Foulkes. Concomitants of the quintic and sextic up to degree four in the coefficients of the ground form. *J. London Math. Soc.*, 25:205–209, 1950.
- [3] William Fulton. *Young tableaux*, volume 35 of *London Mathematical Society student texts*. CUP, 1997.
- [4] G. D. James. *The representation theory of the symmetric groups*, volume 682 of *Lecture Notes in Mathematics*. Springer, Berlin, 1978.

- [5] Tom McKay. On plethysm conjectures of Stanley and Foulkes. *J. Algebra*, 319(5):2050–2071, 2008.
- [6] Jurgen Müller and Max Neunhöffer. Some computations regarding Foulkes’ conjecture. *Experiment. Math.*, 14(3):277–283, 2005.
- [7] R. P. Stanley. Positivity problems and conjectures in algebraic combinatorics, in: *Mathematics: Frontiers and Perspectives*, 295–319. Amer. Math. Soc., 2000.
- [8] R. M. Thrall. On symmetrized Kronecker powers and the structure of the free Lie ring. *American Journal of Mathematics*, 64(2):371–388, 1942.