

Kent Academic Repository

Full text document (pdf)

Citation for published version

Clarkson, Peter and Jordaan, Kerstin and Kelil, Abey (2015) A Generalized Freud Weight. *Studies in Applied Mathematics*, 136 (3). pp. 288-320. ISSN 0022-2526.

DOI

<https://doi.org/10.1111/sapm.12105>

Link to record in KAR

<https://kar.kent.ac.uk/51095/>

Document Version

Pre-print

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

A Generalized Freud Weight

Peter A. Clarkson

School of Mathematics, Statistics and Actuarial Science,
University of Kent, Canterbury, CT2 7NF, UK
P.A.Clarkson@kent.ac.uk

Kerstin Jordaan and Abey Kelil

Department of Mathematics and Applied Mathematics,
University of Pretoria, Pretoria, 0002, South Africa
kerstin.jordaan@up.ac.za, abeysh2001@gmail.com

Abstract

We discuss the relationship between the recurrence coefficients of orthogonal polynomials with respect to a generalized Freud weight

$$w(x; t) = |x|^{2\lambda+1} \exp(-x^4 + tx^2), \quad x \in \mathbb{R},$$

with parameters $\lambda > -1$ and $t \in \mathbb{R}$, and classical solutions of the fourth Painlevé equation. We show that the coefficients in these recurrence relations can be expressed in terms of Wronskians of parabolic cylinder functions that arise in the description of special function solutions of the fourth Painlevé equation. Further we derive a second-order linear ordinary differential equation and a differential-difference equation satisfied by the generalized Freud polynomials.

1 Introduction

Let μ be a positive Borel measure defined on the real line for which all the moments

$$\mu_n = \int_{\mathbb{R}} x^n d\mu(x), \quad n \in \mathbb{N},$$

exist. The Hilbert space $L^2(\mu)$ contains the set of polynomials and hence Gram-Schmidt orthogonalization applied to the set $\{1, x, x^2, \dots\}$ yields a set of monic orthogonal polynomials on the real line defined by

$$\int_{\mathbb{R}} P_n(x)P_m(x) d\mu(x) = h_n \delta_{mn}, \quad m, n \in \mathbb{N},$$

with $h_n > 0$, δ_{mn} the Kronecker delta and $P_n(x)$ is a polynomial of exact degree n .

A family $\{P_n(x)\}_{n=0}^{\infty}$ of monic orthogonal polynomials satisfies a three-term recurrence relation of the form

$$xP_n(x) = P_{n+1}(x) + \alpha_n P_n(x) + \beta_n P_{n-1}(x), \quad n \in \mathbb{N}, \quad (1.1)$$

with $P_{-1}(x) = 0$ and $P_0(x) = 1$. Respectively multiplying (1.1) by $P_n(x)$ and $P_{n-1}(x)$ and integrating gives

$$\alpha_n = \frac{1}{h_n} \int_{\mathbb{R}} x P_n^2(x) d\mu(x), \quad \beta_n = \frac{1}{h_{n-1}} \int_{\mathbb{R}} x P_n(x) P_{n-1}(x) d\mu(x).$$

The converse statement, known as the spectral theorem for orthogonal polynomials, is often attributed to Favard [17] but was seemingly discovered independently, around the same time, by both Shohat [38, 39] and Natanson [34]. It is interesting to note that the result can be traced back to earlier work on continued fractions with a rudimentary form given already in 1894 by Stieltjes [42], see [12], also [30, 44]. The result also appears in the books by Stone [41] and Wintner [45], see [26]. A modern proof of the result is given by Beardon [2].

Theorem 1.1. *If a family of polynomials satisfies a three-term recurrence relation of the form*

$$xP_n(x) = P_{n+1}(x) + \alpha_n P_n(x) + \beta_n P_{n-1}(x),$$

with initial conditions $P_{-1}(x) = 0$ and $P_0(x) = 1$, where $\alpha_n, \beta_n \in \mathbb{R}$, then there exists a measure μ on the real line such that these polynomials are monic orthogonal polynomials satisfying

$$\int_{\mathbb{R}} P_n(x)P_m(x) d\mu(x) = h_n \delta_{mn}, \quad m, n \in \mathbb{N},$$

with $h_n > 0$.

Proof. See, for example, [12, 26]. □

Two important problems arise in the study of orthogonal polynomials.

1. *Direct problem:* Given a measure $\mu(x)$, what can be deduced about the recurrence coefficients $\{\alpha_n, \beta_n\}$, $n \in \mathbb{N}$?
2. *Inverse problem:* Given recurrence coefficients $\{\alpha_n, \beta_n\}$, $n \in \mathbb{N}$, what can be deduced about the orthogonality measure $\mu(x)$?

Given an orthogonality measure $\mu(x)$, several characteristic properties of the sequence $\{P_n(x)\}_{n=0}^{\infty}$ are determined by the nature of the measure. Extracting this information from the measure is an extension of the direct problem and one of the interesting and challenging problems in the study of systems of orthogonal polynomials. Properties typically studied include the Hankel determinants, the coefficients of the three-term recurrence relation, the coefficients of the differential-difference equation and the differential equation satisfied by the polynomials. For instance, the recurrence coefficients can be expressed in terms of Hankel determinants comprising the moments of the orthogonality measure. We refer to [12, 26, 43] for further information about orthogonal polynomials.

For classical orthogonal polynomials, namely the Hermite, Laguerre and Jacobi polynomials, the measure $\mu(x)$ is absolutely continuous and can be expressed in terms of a weight function $w(x)$ which is non-negative, with support on some interval $[a, b] \in \mathbb{R}$ (where $a = -\infty$ and $b = \infty$ are allowed), i.e. $d\mu(x) = w(x) dx$. The properties that these orthogonal polynomials satisfy are well known and include the fact that (cf. [26]):

- (a) their derivatives also form orthogonal polynomial sets;
- (b) they satisfy a Rodrigues' type formula

$$P_n(x) = \frac{1}{\kappa_n w(x)} \frac{d^n}{dx^n} \{w(x)\sigma^n(x)\},$$

where $w(x)$ is the weight function, $\sigma(x)$ is a polynomial in x independent of n and κ_n a constant;

- (c) they satisfy a non-linear equation of the form

$$\sigma(x) \frac{d}{dx} [P_n(x)P_{n-1}(x)] = (a_n x + b_n)P_n(x)P_{n-1}(x) + c_n P_n^2(x) + d_n P_{n-1}^2(x),$$

where a_n, b_n, c_n and d_n are independent of x ;

- (d) they satisfy a second order linear differential equation of the Sturm-Liouville type

$$\sigma(x) \frac{d^2 P_n}{dx^2} + \tau(x) \frac{d P_n}{dx} = \lambda_n P_n(x),$$

where $\sigma(x)$ is a polynomial of degree ≤ 2 , $\tau(x)$ is a linear polynomial, both independent of n , and λ_n is independent of x ;

- (e) they satisfy the differential-difference equation

$$\pi(x) \frac{d P_n}{dx} = (a_n x + b_n)P_n(x) + c_n P_{n-1}(x). \tag{1.2}$$

The converse, that any polynomial set which satisfies any one of the above properties must necessarily be one of the classical orthogonal polynomial sets, also holds. In particular, Al-Salam and Chihara [1] showed that orthogonal polynomial sets satisfying (1.2) must be either Hermite, Laguerre or Jacobi polynomials.

Askey raised the more general question of what orthogonal polynomial sets have the property that their derivatives satisfy a relation of the form

$$\pi(x) \frac{dP_n}{dx} = \sum_{k=n-t}^{n+s} a_{n,k} P_k(x).$$

This problem was solved by Shohat [40] and later, independently, by Freud [23], as well as Bonan and Nevai [6]. Maroni [31, 32] stated the problem in a different way, trying to find all orthogonal polynomial sets whose derivatives are quasi-orthogonal, and called such orthogonal polynomial sets semi-classical.

A useful characterization of classical orthogonal polynomials is the Pearson equation

$$\frac{d}{dx}[\sigma(x)w(x)] = \tau(x)w(x), \quad (1.3)$$

where $\sigma(x)$ and $\tau(x)$ are polynomials with $\deg(\sigma) \leq 2$ and $\deg(\tau) = 1$.

Semi-classical orthogonal polynomials are defined as orthogonal polynomials for which the weight function satisfies a Pearson equation (1.3) with $\deg(\sigma) \geq 2$ or $\deg(\tau) \neq 1$; see Hendriksen and van Rossum [25].

The relationship between semi-classical orthogonal polynomials and integrable equations dates back to Shohat [40] and Freud [23], see also Bonan and Nevai [6]. However it was not until the work of Fokas, Its and Kitaev [19, 20] that these equations were identified as discrete Painlevé equations. The relationship between semi-classical orthogonal polynomials and the Painlevé equations was discussed by Magnus [29] who, for example, showed that the coefficients in the three-term recurrence relation for the Freud weight [23]

$$w(x; t) = \exp(-x^4 + tx^2), \quad x \in \mathbb{R}, \quad (1.4)$$

with $t \in \mathbb{R}$ a parameter, can be expressed in terms of simultaneous solutions of the discrete equation

$$q_n(q_{n-1} + q_n + q_{n+1}) + 2tq_n = n, \quad (1.5)$$

which is discrete P_I (dP_I), as shown by Bonan and Nevai [6], and the differential equation

$$\frac{d^2 q_n}{dz^2} = \frac{1}{2q_n} \left(\frac{dq_n}{dz} \right)^2 + \frac{3}{2}q_n^3 + 4zq_n^2 + 2(z^2 + \frac{1}{2}n)q_n - \frac{n^2}{2q_n}, \quad (1.6)$$

which is a special case of the fourth Painlevé equation – see equation (1.8) below – with $n \in \mathbb{N}$. This connection between the recurrence coefficients for the Freud weight (1.4) and simultaneous solutions of (1.5) and (1.6) is due to Kitaev, see [19, 21]. The discrete equation (1.5) is also known as the “Freud equation” or the “string equation”.

It is known (cf. [26]) that polynomials orthogonal with respect to exponential weights $w(x) = \exp\{-Q(x)\}$ on \mathbb{R} for polynomial $Q(x)$ satisfy structural relations of the form

$$\frac{dP_n}{dx}(x) = A_n(x)P_{n-1}(x) + B_n(x)P_n(x).$$

Such structural relations and the three-term recurrence relation reveal that the orthogonal polynomials satisfy a second order differential equation.

It had been generally accepted that explicit expressions for the associated coefficients in the three-term recurrence relation and orthogonal polynomials were nonexistent for weights such as the Freud weight (1.4). To quote from the *Digital Library of Mathematical Functions* [36, §18.32]:

“A *Freud weight* is a weight function of the form

$$w(x) = \exp\{-Q(x)\}, \quad -\infty < x < \infty,$$

where $Q(x)$ is real, even, nonnegative, and continuously differentiable. Of special interest are the cases $Q(x) = x^{2m}$, $m = 1, 2, \dots$. No explicit expressions for the corresponding OP’s are available. However, for asymptotic approximations in terms of elementary functions for the OP’s, and also for their largest zeros, see Levin and Lubinsky [28] and Nevai [35]. For a uniform asymptotic expansion in terms of Airy functions for the OP’s in the case $Q(x) = x^4$ see Bo and Wong [4]. For asymptotic approximations to OP’s that correspond to Freud weights with more general functions $Q(x)$ see Deift *et al.* [15, 16], Bleher and Its [3], and Kriecherbauer and McLaughlin [27].”

In [13], the direct problem was studied for semi-classical Laguerre polynomials orthogonal with respect to the semi-classical Laguerre weight

$$w(x; t) = x^\lambda \exp(-x^2 + tx), \quad x \in \mathbb{R}^+, \quad \lambda > -1, \quad (1.7)$$

with $t \in \mathbb{R}$ a parameter, and it was shown that the coefficients in the three-term recurrence relation of these polynomials can be explicitly expressed in terms of Wronskians of parabolic cylinder functions which also arise in the description of special function solutions of the fourth Painlevé equation (P_{IV})

$$\frac{d^2 q}{dz^2} = \frac{1}{2q} \left(\frac{dq}{dz} \right)^2 + \frac{3}{2} q^3 + 4zq^2 + 2(z^2 - A)q + \frac{B}{q}, \quad (1.8)$$

where A and B are constants, and the second degree, second order equation satisfied by the associated Hamiltonian function, see equation (4.16).

Polynomials orthogonal with respect to a symmetric measure can be generated via quadratic transformation from the classical orthogonal polynomials, cf. [12]. For example, Laguerre polynomials generate a class of generalized Hermite polynomials while Jacobi polynomials give rise to a class of generalized Ultraspherical polynomials.

In this paper we are concerned with the positive, even weight function on the real line arising from a symmetrization of the semi-classical Laguerre weight (1.7), namely the generalized Freud weight

$$w(x; t) = |x|^{2\lambda+1} \exp(-x^4 + tx^2), \quad x \in \mathbb{R}, \quad (1.9)$$

with parameters $\lambda > -1$ and $t \in \mathbb{R}$.

We use two different methods to derive the differential-difference equation satisfied by generalized Freud polynomials $\{S_n(x; t)\}_{n=0}^\infty$, orthogonal with respect to the generalized Freud weight (1.9). In §2 we modify the ladder operator method (cf. [8, 9, 26]) to derive a general formula for the coefficients of the differential-difference as well as the second order differential equation satisfied by polynomials orthogonal with respect to a generalized Freud type weight which vanishes at a point. It is important to note that the second order differential equation is linear with coefficients that are rational functions of x with parameters t and λ involving parabolic cylinder functions. In §3 we derive the generalized Freud weight (1.9) through a symmetrization of the semi-classical Laguerre weight (1.7). In §4 we are concerned with specific results for generalized Freud weight (1.9). We show that the coefficient $\beta_n(t; \lambda)$ in the three-term recurrence relation

$$xS_n(x; t) = S_{n+1}(x; t) + \beta_n(t; \lambda)S_{n-1}(x; t),$$

satisfied by the polynomials associated with the weight (1.9) can be expressed in terms of Wronskians that arise in the description of special function solutions of P_{IV} which are expressed in terms of parabolic cylinder functions. Further we apply the results of §2 to the weight (1.9) to derive the differential-difference equation and the linear, second order differential equations satisfied by generalized Freud polynomials $\{S_n(x; t)\}_{n=0}^\infty$. A method due to Shohat [40], based on the concept of quasi-orthogonality and applied to the weight (1.9) is discussed in §4.5.

2 Semi-classical orthogonal polynomials

The coefficients $A_n(x; t)$ and $B_n(x; t)$ in the relation

$$\frac{dP_n}{dx}(x; t) = -B_n(x; t)P_n(x; t) + A_n(x; t)P_{n-1}(x; t), \quad (2.1)$$

satisfied by semi-classical orthogonal polynomials are of interest since differentiation of this differential-difference equation yields the second order differential equation satisfied by the orthogonal polynomials. Shohat [40] gave a procedure using quasi-orthogonality to derive (2.1) for weights $w(x; t)$ such that $w'(x; t)/w(x; t)$ is a rational function, which we apply to (1.9) later. This technique was rediscovered by several authors including Bonan, Freud, Mhaskar and Nevai approximately 40 years later, see [35, p. 126–132] and the references therein for more detail. The method of ladder operators was introduced by Chen and Ismail in [9]. Related work by various authors can be found in, for example, [10, 11, 18, 33] and a good summary of the technique is provided in [26, Theorem 3.2.1].

In [8], Chen and Feigin adapt the method of ladder operators to the situation where the weight function vanishes at one point. Our next result generalizes the work in [8], giving a more explicit expression for the coefficients in (2.1) when the weight function is positive on the real line except for one point.

Theorem 2.1. *Let*

$$w(x; t) = |x - k|^\gamma \exp\{-v(x; t)\}, \quad x, t, k \in \mathbb{R}, \quad (2.2)$$

where $v(x; t)$ is a continuously differentiable function on \mathbb{R} . Assume that the polynomials $\{P_n(x; t)\}_{n=0}^\infty$ satisfy the orthogonality relation

$$\int_{-\infty}^{\infty} P_n(x; t) P_m(x; t) w(x; t) dx = h_n \delta_{mn}.$$

Then, for $\gamma \geq 1$, $P_n(x; t)$ satisfy the differential-difference equation

$$(x - k) \frac{dP_n}{dx}(x; t) = -B_n(x; t) P_n(x; t) + A_n(x; t) P_{n-1}(x; t), \quad (2.3)$$

where

$$A_n(x; t) = \frac{x - k}{h_{n-1}} \int_{-\infty}^{\infty} P_n^2(y; t) \mathcal{K}(x, y) w(y; t) dy + a_n(x; t), \quad (2.4a)$$

$$B_n(x; t) = \frac{x - k}{h_{n-1}} \int_{-\infty}^{\infty} P_n(y; t) P_{n-1}(y; t) \mathcal{K}(x, y) w(y; t) dy + b_n(x; t), \quad (2.4b)$$

where

$$\mathcal{K}(x, y) = \frac{v'(x; t) - v'(y; t)}{x - y}, \quad (2.5)$$

with

$$a_n(x; t) = \frac{\gamma}{h_{n-1}} \int_{-\infty}^{\infty} \frac{P_n^2(y; t)}{y - k} w(y; t) dy, \quad (2.6a)$$

$$b_n(x; t) = \frac{\gamma}{h_{n-1}} \int_{-\infty}^{\infty} \frac{P_n(y; t) P_{n-1}(y; t)}{y - k} w(y; t) dy. \quad (2.6b)$$

Proof. Consider the generalized Freud-type weight (2.2). Since $\frac{dP_n}{dx}(x; t)$ is a polynomial of degree $n - 1$ in x , then it can be expressed in terms of the orthogonal basis as

$$\frac{dP_n}{dx}(x; t) = \sum_{j=0}^{n-1} c_{n,j} P_j(x; t). \quad (2.7)$$

Applying the orthogonality relation and integrating by parts, we obtain

$$\begin{aligned} c_{n,j} h_j &= \int_{-\infty}^{\infty} \frac{dP_n}{dy}(y; t) P_j(y; t) w(y; t) dy \\ &= \left[P_n(x; t) P_j(x; t) w(x; t) \right]_{-\infty}^{\infty} - \int_{-\infty}^{\infty} P_n(y; t) \left\{ \frac{dP_j}{dy}(y; t) w(y; t) + P_j(y; t) \frac{dw}{dy}(y; t) \right\} dy \\ &= - \int_{-\infty}^{\infty} P_n(y; t) P_j(y; t) \frac{dw}{dy}(y; t) dy \\ &= \int_{-\infty}^{\infty} P_n(y; t) P_j(y; t) \left[\frac{dv}{dy}(y; t) - \frac{\gamma}{y - k} \right] w(y; t) dy, \end{aligned}$$

provided that $\gamma \geq 1$.

Now, from (2.7), we can write

$$\begin{aligned}
\frac{dP_n}{dx}(x; t) &= \sum_{j=0}^{n-1} \frac{1}{h_j} \left\{ \int_{-\infty}^{\infty} P_n(y; t) P_j(y; t) \left[\frac{dv}{dy}(y; t) - \frac{\gamma}{y-k} \right] w(y; t) dy \right\} P_j(x; t) \\
&= \int_{-\infty}^{\infty} P_n(y; t) \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} \left[\frac{dv}{dy}(y; t) - \frac{\gamma}{y-k} \right] w(y; t) dy \\
&= \int_{-\infty}^{\infty} P_n(y; t) \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} \left[\frac{dv}{dy}(y; t) - \frac{dv}{dx}(x; t) \right] w(y; t) dy \\
&\quad + \frac{dv}{dx}(x; t) \int_{-\infty}^{\infty} P_n(y; t) \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} w(y; t) dy \\
&\quad - \gamma \int_{-\infty}^{\infty} \frac{P_n(y; t)}{y-k} \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} w(y; t) dy \\
&= \int_{-\infty}^{\infty} P_n(y; t) \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} \left[\frac{dv}{dy}(y; t) - \frac{dv}{dx}(x; t) \right] w(y; t) dy \\
&\quad - \gamma \int_{-\infty}^{\infty} \frac{P_n(y; t)}{y-k} \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} w(y; t) dy.
\end{aligned}$$

Next, using the orthogonality relation again, we obtain

$$\begin{aligned}
(x-k) \frac{dP_n}{dx}(x; t) &= (x-k) \int_{-\infty}^{\infty} P_n(y; t) \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} \left[\frac{dv}{dy}(y; t) - \frac{dv}{dx}(x; t) \right] w(y; t) dy \\
&\quad - \gamma \int_{-\infty}^{\infty} P_n(y; t) \left(\frac{x-k}{y-k} \right) \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} w(y; t) dy. \\
&= (x-k) \int_{-\infty}^{\infty} P_n(y; t) \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} \left[\frac{dv}{dy}(y; t) - \frac{dv}{dx}(x; t) \right] w(y; t) dy \\
&\quad - \gamma \int_{-\infty}^{\infty} P_n(y; t) \frac{x-y}{y-k} \sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} w(y; t) dy.
\end{aligned}$$

It now follows from the Christoffel-Darboux formula (cf. [26])

$$\sum_{j=0}^{n-1} \frac{P_j(y; t) P_j(x; t)}{h_j} = \frac{P_n(x; t) P_{n-1}(y; t) - P_n(y; t) P_{n-1}(x; t)}{(x-y) h_{n-1}},$$

that

$$\begin{aligned}
A_n(x; t) &= \frac{x-k}{h_{n-1}} \int_{-\infty}^{\infty} P_n^2(y; t) \mathcal{K}(x, y) w(y; t) dy + a_n(x; t) \\
B_n(x; t) &= \frac{x-k}{h_{n-1}} \int_{-\infty}^{\infty} P_n(y; t) P_{n-1}(y; t) \mathcal{K}(x, y) w(y; t) dy + b_n(x; t),
\end{aligned}$$

with $a_n(x; t)$ and $b_n(x; t)$ given by (2.6). □

Lemma 2.2. Consider the weight defined by (2.2) and assume that $v(x; t)$ is an even, continuously differentiable function on \mathbb{R} . Assume that the polynomials $\{P_n(x; t)\}_{n=0}^{\infty}$ satisfy the orthogonality relation

$$\int_{-\infty}^{\infty} P_n(x; t) P_m(x; t) w(x; t) dx = h_n \delta_{mn}.$$

and the three-term recurrence relation

$$P_{n+1}(x; t) = x P_n(x; t) - \beta_n(t; \lambda) P_{n-1}(x; t), \tag{2.8}$$

with $P_0 = 1$ and $P_1 = x$. Then the polynomials $P_n(x; t)$ satisfy

$$\int_{-\infty}^{\infty} \frac{P_n^2(y; t)}{y - k} w(y; t) dy = 0, \quad (2.9)$$

$$\int_{-\infty}^{\infty} \frac{P_n(y; t)P_{n-1}(y; t)}{y - k} w(y; t) dy = \frac{1}{2}[1 - (-1)^n] h_{n-1}, \quad (2.10)$$

where $n \in \mathbb{N}$ and

$$h_n = \int_{-\infty}^{\infty} P_n^2(y; t) w(y; t) dy.$$

Proof. Since $w(x; t)$ is even when $v(x; t)$ is assumed to be even, the integrand in (2.6a) is odd and hence $a_n(x; t) = 0$.

Furthermore, the monic orthogonal polynomials $P_n(x; t)$ satisfy the three-term recurrence relation (2.8), with $P_0 = 1$ and $P_1 = x$, hence

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{P_n(y; t)P_{n-1}(y; t)}{y - k} w(y; t) dy &= \int_{-\infty}^{\infty} \frac{[yP_{n-1}(y; t) - \beta_{n-1}P_{n-2}(y; t)]P_{n-1}(y; t)}{y - k} w(y; t) dy \\ &= \int_{-\infty}^{\infty} P_{n-1}^2(y; t) w(y; t) dy + k \int_{-\infty}^{\infty} \frac{P_{n-1}^2(y; t)}{y - k} w(y; t) dy \\ &\quad - \beta_{n-1} \int_{-\infty}^{\infty} \frac{P_{n-1}(y; t)P_{n-2}(y; t)}{y - k} w(y; t) dy \\ &= h_{n-1} - \beta_{n-1} \int_{-\infty}^{\infty} \frac{P_{n-1}(y; t)P_{n-2}(y; t)}{y - k} w(y; t) dy, \end{aligned}$$

since the integrand in the second integral is odd. Hence, if we define

$$I_n = \int_{-\infty}^{\infty} \frac{P_n(y; t)P_{n-1}(y; t)}{y - k} w(y; t) dy,$$

then I_n satisfies the recurrence relation

$$I_n = h_{n-1} - \beta_{n-1}I_{n-1} = h_{n-1} - \frac{h_{n-1}}{h_{n-2}}I_{n-1},$$

since $\beta_n = h_n/h_{n-1}$. Iterating this gives

$$I_n = \frac{h_{n-1}}{h_{n-3}}I_{n-2} = h_{n-1} - \frac{h_{n-1}}{h_{n-4}}I_{n-3} = \frac{h_{n-1}}{h_{n-5}}I_{n-4} = h_{n-1} - \frac{h_{n-1}}{h_{n-6}}I_{n-5},$$

and so on. Hence, by induction,

$$I_{2N} = \frac{h_{2N-1}}{h_1}I_2, \quad I_{2N+1} = h_{2N} - \frac{h_{2N}}{h_1}I_2,$$

and so, since

$$\begin{aligned} I_2 &= \int_{-\infty}^{\infty} \frac{P_2(y; t)P_1(y; t)}{y - k} w(y; t) dy \\ &= \int_{-\infty}^{\infty} P_2(y; t)w(y; t) dy + k \int_{-\infty}^{\infty} \frac{P_2(y; t)}{y - k} w(y; t) dy = 0, \end{aligned}$$

we have that

$$I_{2N} = 0, \quad I_{2N+1} = h_{2N},$$

as required. \square

Corollary 2.3. *Let*

$$w(x; t) = |x - k|^\gamma \exp\{-v(x; t)\}, \quad x, t, k \in \mathbb{R},$$

where $v(x; t)$ is an even, continuously differentiable function on \mathbb{R} . Assume that the polynomials $\{P_n(x; t)\}_{n=0}^\infty$ satisfy the orthogonality relation

$$\int_{-\infty}^{\infty} P_n(x; t)P_m(x; t)w(x; t) dx = h_n\delta_{mn}.$$

Then, for $\gamma \geq 1$, $P_n(x; t)$ satisfy the differential-difference equation

$$(x - k) \frac{dP_n}{dx}(x; t) = -B_n(x; t)P_n(x; t) + A_n(x; t)P_{n-1}(x; t),$$

where

$$A_n(x; t) = \frac{x - k}{h_{n-1}} \int_{-\infty}^{\infty} P_n^2(y; t) \mathcal{K}(x, y) w(y; t) dy, \quad (2.11a)$$

$$B_n(x; t) = \frac{x - k}{h_{n-1}} \int_{-\infty}^{\infty} P_n(y; t) P_{n-1}(y; t) \mathcal{K}(x, y) w(y; t) dy + \frac{1}{2} \gamma [1 - (-1)^n]. \quad (2.11b)$$

Proof. The result is an immediate consequence of Theorem 2.1 and Lemma 2.2. \square

Lemma 2.4. Let $w(x; t)$ be the weight defined by (2.2) with $v(x; t)$ an even, continuously differentiable function on \mathbb{R} and $A_n(x; t)$ and $B_n(x; t)$ defined by (2.11). Then

$$B_n(x; t) + B_{n+1}(x; t) = \frac{x A_n(x; t)}{\beta_n} + \gamma - (x - k) \frac{dv}{dx}(x; t),$$

when $\gamma \geq 1$.

Proof. From (2.11), (2.8) and the fact that $h_n = h_{n-1} \beta_n$ we have

$$\begin{aligned} B_n(x; t) + B_{n+1}(x; t) &= \frac{x - k}{h_{n-1}} \int_{-\infty}^{\infty} P_n(y; t) P_{n-1}(y; t) \mathcal{K}(x, y) w(y; t) dy \\ &\quad + \frac{x - k}{h_n} \int_{-\infty}^{\infty} P_n(y; t) P_{n+1}(y; t) \mathcal{K}(x, y) w(y; t) dy \\ &\quad + \frac{1}{2} \gamma [1 - (-1)^n] + \frac{1}{2} \gamma [1 - (-1)^{n+1}] \\ &= \frac{x - k}{h_n} \left\{ \int_{-\infty}^{\infty} P_n(y; t) [\beta_n P_{n-1}(y; t) + P_{n+1}(y; t)] \mathcal{K}(x, y) w(y; t) dy \right\} + \gamma \\ &= \frac{x - k}{h_n} \int_{-\infty}^{\infty} y P_n^2(y; t) \mathcal{K}(x, y) w(y; t) dy + \gamma \\ &= \frac{x - k}{h_n} \int_{-\infty}^{\infty} P_n^2(y; t) \left[\frac{dv}{dy}(y; t) - \frac{dv}{dx}(x; t) \right] w(y; t) dy \\ &\quad + \frac{x(x - k)}{h_n} \int_{-\infty}^{\infty} P_n^2(y; t) \mathcal{K}(x, y) w(y; t) dy + \gamma \\ &= \frac{x - k}{h_n} \int_{-\infty}^{\infty} P_n^2(y; t) \frac{dv}{dy}(y; t) w(y; t) dy \\ &\quad - \frac{x - k}{h_n} \frac{dv}{dx}(x; t) \int_{-\infty}^{\infty} P_n^2(y; t) w(y; t) dy + \frac{x}{h_n} h_{n-1} A_n(x; t) + \gamma \\ &= \frac{\gamma(x - k)}{h_n} \int_{-\infty}^{\infty} P_n^2(y; t) \frac{w(y; t)}{y - k} dy - \frac{x - k}{h_n} \int_{-\infty}^{\infty} P_n^2(y; t) \frac{dw}{dy}(y; t) dy \\ &\quad - (x - k) \frac{dv}{dx}(x; t) + \frac{x A_n(x; t)}{\beta_n} + \gamma, \end{aligned} \quad (2.12)$$

since

$$\frac{dw}{dy}(y; t) = \left[-\frac{dv}{dy}(y; t) + \frac{\gamma}{y - k} \right] w(y; t).$$

The first integral in (2.12) vanishes since the integrand is odd, hence it follows, using integration by parts, that

$$\begin{aligned} B_n(x; t) + B_{n+1}(x; t) &= -\frac{x - k}{h_n} \left\{ \left[P_n^2(x; t) w(x; t) \right]_{-\infty}^{\infty} - \int_{-\infty}^{\infty} 2P_n(y; t) \frac{dP_n}{dy}(y; t) w(y; t) dy \right\} \\ &\quad - (x - k) \frac{dv}{dx}(x; t) + \frac{x A_n(x; t)}{\beta_n} + \gamma, \end{aligned}$$

and the result follows from the orthogonality of P_n . \square

Lemma 2.5. Let $w(x; t)$ be the weight defined by (2.2) with $v(x; t)$ an even, continuously differentiable function on \mathbb{R} and $A_n(x; t)$ and $B_n(x; t)$ defined by (2.11). Then

$$\left\{ -(x-k) \frac{d}{dx} + B_n(x; t) + (x-k) \frac{dv}{dx}(x; t) - \gamma \right\} P_{n-1}(x; t) = \frac{A_{n-1}(x; t)}{\beta_{n-1}} P_n(x; t). \quad (2.13)$$

Proof. From the differential-difference equation (2.3) we have

$$\begin{aligned} (x-k) \frac{dP_{n-1}}{dx}(x; t) &= -B_{n-1}(x; t)P_{n-1}(x; t) + A_{n-1}(x; t)P_{n-2}(x; t) \\ &= -B_{n-1}(x; t)P_{n-1}(x; t) + \frac{A_{n-1}(x; t)}{\beta_{n-1}} [xP_{n-1}(x; t) - P_n(x; t)], \end{aligned}$$

using the recurrence relation (2.8). Hence, using Lemma 2.4,

$$\begin{aligned} \frac{A_{n-1}(x; t)}{\beta_{n-1}} P_n(x; t) &= -(x-k) \frac{dP_{n-1}}{dx}(x; t) - B_{n-1}(x; t)P_{n-1}(x; t) + \frac{x A_{n-1}(x; t)}{\beta_{n-1}} P_{n-1}(x; t) \\ &= -(x-k) \frac{dP_{n-1}}{dx}(x; t) - B_{n-1}(x; t)P_{n-1}(x; t) \\ &\quad + \left\{ B_n(x; t) + B_{n-1}(x; t) - \gamma + (x-k) \frac{dv}{dx}(x; t) \right\} P_{n-1}(x; t) \\ &= -(x-k) \frac{dP_{n-1}}{dx}(x; t) + B_n(x; t)P_{n-1}(x; t) + \left[(x-k) \frac{dv}{dx}(x; t) - \gamma \right] P_{n-1}(x; t) \\ &= \left\{ -(x-k) \frac{d}{dx} + B_n(x; t) + (x-k) \frac{dv}{dx}(x; t) - \gamma \right\} P_{n-1}(x; t). \end{aligned}$$

□

Theorem 2.6. Let $w(x; t) = |x-k|^\gamma \exp\{-v(x; t)\}$, for $x, t, k \in \mathbb{R}$, with $v(x; t)$ an even, continuously differentiable function on \mathbb{R} . Then

$$(x-k) \frac{d^2 P_n}{dx^2}(x; t) + R_n(x; t) \frac{dP_n}{dx}(x; t) + T_n(x; t) P_n(x; t) = 0, \quad (2.14a)$$

where

$$R_n(x; t) = \gamma - (x-k) \frac{dv}{dx}(x; t) - \frac{x-k}{A_n(x; t)} \frac{dA_n}{dx}(x; t) + 1, \quad (2.14b)$$

$$\begin{aligned} T_n(x; t) &= \frac{A_n(x; t)A_{n-1}(x; t)}{(x-k)\beta_{n-1}} + \frac{dB_n}{dx}(x; t) + \frac{\gamma B_n(x; t)}{x-k} \\ &\quad - B_n(x; t) \left[\frac{dv}{dx}(x; t) + \frac{B_n(x; t)}{x-k} \right] - \frac{B_n(x; t)}{A_n(x)} \frac{dA_n}{dx}(x; t), \end{aligned} \quad (2.14c)$$

with

$$\begin{aligned} A_n(x; t) &= \frac{x-k}{h_{n-1}} \int_{-\infty}^{\infty} P_n^2(y; t) \mathcal{K}(x, y) w(y; t) dy \\ B_n(x; t) &= \frac{x-k}{h_{n-1}} \int_{-\infty}^{\infty} P_n(y; t) P_{n-1}(y; t) \mathcal{K}(x, y) w(y; t) dy + \frac{1}{2} \gamma [1 - (-1)^n]. \end{aligned}$$

Proof. Differentiating both sides of (2.3) with respect to x we obtain

$$\begin{aligned} (x-k) \frac{d^2 P_n}{dx^2}(x; t) &= -[B_n(x; t) + 1] \frac{dP_n}{dx}(x; t) + \frac{dA_n}{dx}(x; t) P_{n-1}(x; t) \\ &\quad - \frac{dB_n}{dx}(x; t) P_n(x; t) + A_n(x; t) P_{n-1}(x; t). \end{aligned} \quad (2.15)$$

Substituting (2.13) into (2.15) yields

$$\begin{aligned} (x-k) \frac{d^2 P_n}{dx^2}(x; t) &= -[B_n(x; t) + 1] \frac{dP_n}{dx}(x; t) - \left[\frac{dB_n}{dx}(x; t) + \frac{A_n(x; t)A_{n-1}(x; t)}{\beta_{n-1}(x-k)} \right] P_n(x; t) \\ &\quad + \left\{ \frac{dA_n}{dx}(x; t) + A_n(x; t) \left[\frac{B_n(x; t)}{x-k} + \frac{dv}{dx}(x; t) - \frac{\gamma}{x-k} \right] \right\} P_{n-1}(x; t), \end{aligned} \quad (2.16)$$

and the results follows by substituting $P_{n-1}(x; t)$ in (2.16) using (2.3). □

3 Semi-classical Laguerre weight

We recall that the semi-classical Laguerre weight [13, 25] is given by

$$w(x; t) = x^\lambda \exp(-x^2 + tx), \quad x \in \mathbb{R}^+, \quad (3.1)$$

where $\lambda > -1$. The weight function satisfies the differential equation

$$x \frac{dw}{dx}(x; t) + (2x^2 - tx - \lambda)w(x; t) = 0,$$

which is the Pearson equation (1.3) with $\sigma(x) = x$ and $\tau(x; t) = -2x^2 + tx + \lambda + 1$.

Explicit expressions for the recurrence coefficients $\alpha_n(t; \lambda)$ and $\beta_n(t; \lambda)$ in the three-term recurrence relation

$$xP_n(x; t) = P_{n+1}(x; t) + \alpha_n(t; \lambda)P_n(x; t) + \beta_n(t; \lambda)P_{n-1}(x; t), \quad (3.2)$$

associated with the semi-classical Laguerre weight (3.1) were obtained in [13], and are given in the following theorem.

Theorem 3.1. *Suppose $\Psi_{n,\lambda}(z)$ is given by*

$$\Psi_{n,\lambda}(z) = \mathcal{W} \left(\psi_\lambda, \frac{d\psi_\lambda}{dz}, \dots, \frac{d^{n-1}\psi_\lambda}{dz^{n-1}} \right), \quad \Psi_{0,\lambda}(z) = 1,$$

where

$$\psi_\lambda(z) = \begin{cases} D_{-\lambda-1}(-\sqrt{2}z) \exp\left(\frac{1}{2}z^2\right), & \text{if } \lambda \notin \mathbb{N}, \\ \frac{d^m}{dz^m} \{ [1 + \operatorname{erf}(z)] \exp(z^2) \}, & \text{if } \lambda = m \in \mathbb{N}, \end{cases}$$

with $D_\nu(\zeta)$ is the parabolic cylinder function and $\operatorname{erfc}(z)$ the complementary error function

$$\operatorname{erfc}(z) = \frac{2}{\sqrt{\pi}} \int_z^\infty \exp(-t^2) dt, \quad (3.3)$$

Then coefficients $\alpha_n(t; \lambda)$ and $\beta_n(t; \lambda)$ in the recurrence relation (3.2) associated with the semi-classical Laguerre weight (3.1) are given by

$$\alpha_n(t; \lambda) = \frac{1}{2}q_n(z) + \frac{1}{2}t, \quad (3.4a)$$

$$\beta_n(t; \lambda) = -\frac{1}{8} \frac{dq_n}{dz} - \frac{1}{8}q_n^2(z) - \frac{1}{4}zq_n(z) + \frac{1}{4}\lambda + \frac{1}{2}n, \quad (3.4b)$$

with $z = \frac{1}{2}t$, where

$$q_n(z) = -2z + \frac{d}{dz} \ln \frac{\Psi_{n+1,\lambda}(z)}{\Psi_{n,\lambda}(z)},$$

which satisfies P_{IV} (1.8) with parameters $(A, B) = (2n + \lambda + 1, -2\lambda^2)$.

Proof. See [13]. □

3.1 Symmetrization of semi-classical Laguerre weight

In this section we show that symmetrizing the semi-classical Laguerre weight (3.1) gives rise to the generalized Freud weight (1.9).

Let $\{L_n^{(\lambda)}(x; t)\}$ denote the monic semi-classical Laguerre polynomials orthogonal with respect to semi-classical Laguerre weight (3.1) with

$$\int_0^\infty L_m^{(\lambda)}(x; t)L_n^{(\lambda)}(x; t) x^\lambda \exp(-x^2 + tx) dx = K_n(t) \delta_{mn}.$$

Define

$$S_{2m}(x; t) = L_m^{(\lambda)}(x^2; t); \quad S_{2m+1}(x; t) = xQ_n^{(\lambda)}(x^2; t), \quad (3.5)$$

where

$$Q_n^{(\lambda)}(x; t) = \frac{1}{x} \left[L_{n+1}^{(\lambda)}(x; t) - \frac{L_{n+1}^{(\lambda)}(0; t)}{L_n^{(\lambda)}(0; t)} L_n^{(\lambda)}(x; t) \right], \quad (3.6)$$

are also monic and of degree n , for $x \neq 0$.

Then

$$\begin{aligned} \int_0^\infty L_m^{(\lambda)}(x; t) L_n^{(\lambda)}(x; t) x^\lambda \exp(-x^2 + tx) dx &= 2 \int_0^\infty L_m^{(\lambda)}(x^2; t) L_n^{(\lambda)}(x^2; t) x^{2\lambda+1} \exp(-x^4 + tx^2) dx \\ &= \int_{-\infty}^\infty L_m^{(\lambda)}(x^2; t) L_n^{(\lambda)}(x^2; t) |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx \\ &= \int_{-\infty}^\infty S_{2m}(x; t) S_{2n}(x; t) |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx \\ &= K_n(t) \delta_{mn}. \end{aligned}$$

which implies that $\{S_{2m}(x; t)\}_{m=0}^\infty$ is a symmetric orthogonal sequence with respect to the even weight

$$w(x; t) = |x|^{2\lambda+1} \exp(-x^4 + tx^2),$$

on \mathbb{R} , i.e. the generalized Freud weight (1.9).

It is proved in [12, Thm 7.1] that the kernel polynomials $Q_m^L(x; t)$ are orthogonal with respect to $xw(x; t) = x^{\lambda+1} \exp(-x^2 + tx)$. Hence

$$\begin{aligned} K_n(t) \delta_{mn} &= \int_0^\infty Q_m^{(\lambda)}(x; t) Q_n^{(\lambda)}(x; t) x^{\lambda+1} \exp(-x^2 + tx) dx \\ &= 2 \int_0^\infty Q_m^{(\lambda)}(x^2; t) Q_n^{(\lambda)}(x^2; t) x^{2\lambda+3} \exp(-x^4 + tx^2) dx \\ &= \int_{-\infty}^\infty [x Q_m^{(\lambda)}(x^2; t)] [x Q_n^{(\lambda)}(x^2; t)] |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx \\ &= \int_{-\infty}^\infty S_{2m+1}(x; t) S_{2n+1}(x; t) |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx. \end{aligned}$$

Lastly, since in each case the integrand is odd, we have that

$$\int_{-\infty}^\infty S_{2m+1}(x; t) S_{2n}(x; t) |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx = 0, \quad m, n \in \mathbb{N},$$

and so we conclude that $\{S_n(x; t)\}_{n=0}^\infty$ is a sequence of polynomials orthogonal with respect to the even weight (1.9) on \mathbb{R} .

4 The generalized Freud weight

4.1 The moments of the generalized Freud weight

It is shown in [13] that the moments of the semi-classical weight provide the link between the weight and the associated Painlevé equation. The recurrence coefficients in the three-term recurrence relations associated with semi-classical orthogonal polynomials can often be expressed in terms of solutions of the Painlevé equations and associated discrete Painlevé equations.

For the generalized Freud weight (1.9), the first moment, $\mu_0(t; \lambda)$, can be obtained using the integral representation of a parabolic cylinder (Hermite-Weber) function $D_\nu(\xi)$. By definition

$$\begin{aligned} \mu_0(t; \lambda) &= \int_{-\infty}^\infty |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx \\ &= 2 \int_0^\infty x^{2\lambda+1} \exp(-x^4 + tx^2) dx \\ &= \int_0^\infty y^\lambda \exp(-y^2 + ty) dy \\ &= \frac{\Gamma(\lambda + 1)}{2^{(\lambda+1)/2}} \exp\left(\frac{1}{8}t^2\right) D_{-\lambda-1}\left(-\frac{1}{2}\sqrt{2}t\right). \end{aligned}$$

since the parabolic cylinder function $D_\nu(\xi)$ has the integral representation [36, §12.5(i)]

$$D_\nu(\xi) = \frac{\exp(-\frac{1}{4}\xi^2)}{\Gamma(-\nu)} \int_0^\infty s^{-\nu-1} \exp(-\frac{1}{2}s^2 - \xi s) ds, \quad \Re(\nu) < 0.$$

The even moments are

$$\begin{aligned} \mu_{2n}(t; \lambda) &= \int_{-\infty}^\infty x^{2n} |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx \\ &= \frac{d^n}{dt^n} \left(\int_{-\infty}^\infty |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx \right), \\ &= \frac{d^n}{dt^n} \mu_0(t; \lambda), \quad n \geq 1, \end{aligned}$$

whilst the odd ones are

$$\mu_{2n+1}(t; \lambda) = \int_{-\infty}^\infty x^{2n+1} |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx = 0, \quad n \in \mathbb{N},$$

since the integrand is odd.

We note that

$$\begin{aligned} \mu_{2n}(t; \lambda) &= \int_{-\infty}^\infty x^{2n} |x|^{2\lambda+1} \exp(-x^4 + tx^2) dx \\ &= \int_{-\infty}^\infty |x|^{2\lambda+2n+1} \exp(-x^4 + tx^2) dx \\ &= \mu_0(t; \lambda + n). \end{aligned}$$

Also, when $\lambda = n \in \mathbb{N}$, then

$$D_{-n-1}\left(-\frac{1}{2}\sqrt{2}t\right) = \frac{1}{2}\sqrt{2\pi} \frac{d^n}{dt^n} \left\{ \left[1 + \operatorname{erf}\left(\frac{1}{2}t\right)\right] \exp\left(\frac{1}{8}t^2\right) \right\},$$

where $\operatorname{erf}(z)$ is the error function [36, §12.7(ii)].

4.2 Recurrence coefficients of the generalized Freud polynomials

Monic orthogonal polynomials with respect to the symmetric generalized Freud weight (1.9) satisfy the three-term recurrence relation

$$xS_n(x; t) = S_{n+1}(x; t) + \beta_n(t; \lambda)S_{n-1}(x; t), \quad (4.1)$$

where $S_{-1}(x; t) = 0$ and $S_0(x; t) = 1$.

Our interest is to determine explicit expressions for the recurrence coefficients $\beta_n(t; \lambda)$ in the three-term recurrence relation (4.1). First we relate them to solutions of P_{IV} (1.8).

Lemma 4.1. *The recurrence coefficients $\beta_n(t; \lambda)$ in (4.1) satisfy the equation*

$$\frac{d^2\beta_n}{dt^2} = \frac{1}{2\beta_n} \left(\frac{d\beta_n}{dt} \right)^2 + \frac{3}{2}\beta_n^3 - t\beta_n^2 + \left(\frac{1}{8}t^2 - \frac{1}{2}A_n\right)\beta_n + \frac{B_n}{16\beta_n}, \quad (4.2)$$

where the parameters A_n and B_n are given by

$$\begin{aligned} A_{2n} &= -2\lambda - n - 1, & A_{2n+1} &= \lambda - n, \\ B_{2n} &= -2n^2, & B_{2n+1} &= -2(\lambda + n + 1)^2. \end{aligned}$$

Further $\beta_n(t; \lambda)$ satisfies the nonlinear difference equation

$$\beta_{n+1} + \beta_n + \beta_{n-1} = \frac{1}{2}t + \frac{2n + (2\lambda + 1)[1 - (-1)^n]}{8\beta_n}, \quad (4.3)$$

which is known as discrete P_I (dP_I).

Proof. See, for example, [18, Theorem 6.1]. □

Remarks 4.2.

1. Equation (4.2) is equivalent to P_{IV} (1.8) through the transformation $\beta_n(t; \lambda) = \frac{1}{2}w(z)$, with $z = -\frac{1}{2}t$. Hence

$$\beta_{2n}(t; \lambda) = \frac{1}{2}w(z; -2\lambda - n - 1, -2n^2), \quad (4.4a)$$

$$\beta_{2n+1}(t; \lambda) = \frac{1}{2}w(z; \lambda - n, -2(\lambda + n + 1)^2), \quad (4.4b)$$

with $z = -\frac{1}{2}t$, where $w(z; A, B)$ satisfies P_{IV} (1.8). The conditions on the P_{IV} parameters in (4.4) are precisely those for which P_{IV} has (1.8) has solutions expressible in terms of the parabolic cylinder function

$$\psi(z) = \mu_0(-2z; \lambda) = \frac{\Gamma(\lambda + 1)}{2^{(\lambda+1)/2}} \exp\left(\frac{1}{2}z^2\right) D_{-\lambda-1}(\sqrt{2}z),$$

[22, 37]; see also [13, Theorem 3.5].

2. The link between the differential equation (4.2) and the difference equation (4.3) is given by the Bäcklund transformations

$$\beta_{n+1} = \frac{1}{2\beta_n} \frac{d\beta_n}{dt} - \frac{1}{2}\beta_n + \frac{1}{4}t + \frac{\gamma_n}{4\beta_n}, \quad (4.5a)$$

$$\beta_{n-1} = -\frac{1}{2\beta_n} \frac{d\beta_n}{dt} - \frac{1}{2}\beta_n + \frac{1}{4}t + \frac{\gamma_n}{4\beta_n}, \quad (4.5b)$$

with $\gamma_n = \frac{1}{2}n + \frac{1}{4}(2\lambda + 1)[1 - (-1)^n]$. Letting $n \rightarrow n + 1$ in (4.5b) and substituting (4.5a) gives the differential equation (4.2), whilst eliminating the derivative yields the difference equation (4.3).

Lemma 4.3. *The recurrence coefficients $\beta_n(t; \lambda)$ in (4.1) are given by*

$$\beta_{2n}(t; \lambda) = \frac{d}{dt} \ln \frac{\tau_n(t; \lambda + 1)}{\tau_n(t; \lambda)}, \quad (4.6a)$$

$$\beta_{2n+1}(t; \lambda) = \frac{d}{dt} \ln \frac{\tau_{n+1}(t; \lambda)}{\tau_n(t; \lambda + 1)}, \quad (4.6b)$$

where $\tau_n(t; \lambda)$ is the Wronskian given by

$$\tau_n(t; \lambda) = \mathcal{W} \left(\phi_\lambda, \frac{d\phi_\lambda}{dt}, \dots, \frac{d^{n-1}\phi_\lambda}{dt^{n-1}} \right), \quad (4.7)$$

with

$$\phi_\lambda(t) = \mu_0(t; \lambda) = \frac{\Gamma(\lambda + 1)}{2^{(\lambda+1)/2}} \exp\left(\frac{1}{8}t^2\right) D_{-\lambda-1}\left(-\frac{1}{2}\sqrt{2}t\right).$$

Proof. From the parabolic cylinder solutions of P_{IV} (1.8) given in [13, Theorem 3.5], it is easily shown that the equation

$$\frac{d^2y}{dt^2} = \frac{1}{2y} \left(\frac{dy}{dt} \right)^2 + \frac{3}{2}y^3 - ty^2 + \left(\frac{1}{8}t^2 - \frac{1}{2}A \right)y + \frac{B}{16y}, \quad (4.8)$$

has the solutions $\left\{ y_n^{[j]}(t; A_n^{[j]}, B_n^{[j]}) \right\}_{j=1,2,3}$,

$$y_n^{[1]}(t; \lambda + 2n - 1, -2\lambda^2) = \frac{1}{2}t + \frac{d}{dt} \ln \frac{\tau_{n-1}(t; \lambda)}{\tau_n(t; \lambda)}, \quad (4.9a)$$

$$y_n^{[2]}(t; -2\lambda - n - 1, -2n^2) = \frac{d}{dt} \ln \frac{\tau_n(t; \lambda + 1)}{\tau_n(t; \lambda)}, \quad (4.9b)$$

$$y_n^{[3]}(t; \lambda - n + 1, -2(\lambda + n)^2) = \frac{d}{dt} \ln \frac{\tau_n(t; \lambda)}{\tau_{n-1}(t; \lambda + 1)}, \quad (4.9c)$$

where $\tau_n(t; \lambda)$ is the Wronskian (4.7). Comparing (4.4) and (4.9) gives the desired result. □

The first few recurrence coefficients $\beta_n(t; \lambda)$ are given by

$$\begin{aligned}\beta_1(t; \lambda) &= \Phi_\lambda, \\ \beta_2(t; \lambda) &= -\frac{2\Phi_\lambda^2 - t\Phi_\lambda - \lambda - 1}{2\Phi_\lambda}, \\ \beta_3(t; \lambda) &= -\frac{\Phi_\lambda}{2\Phi_\lambda^2 - t\Phi_\lambda - \lambda - 1} - \frac{\lambda + 1}{2\Phi_\lambda}, \\ \beta_4(t; \lambda) &= \frac{t}{2(\lambda + 2)} + \frac{\Phi_\lambda}{2\Phi_\lambda^2 - t\Phi_\lambda - \lambda - 1} + \frac{(\lambda + 1)(t^2 + 2\lambda + 4)\Phi_\lambda + (\lambda + 1)^2 t}{2(\lambda + 2)[2(\lambda + 2)\Phi_\lambda^2 - (\lambda + 1)t\Phi_\lambda - (\lambda + 1)^2]},\end{aligned}$$

where

$$\Phi_\lambda(t) = \frac{d}{dt} \ln \left\{ D_{-\lambda-1} \left(-\frac{1}{2}\sqrt{2}t \right) \exp \left(\frac{1}{8}t^2 \right) \right\} = \frac{1}{2}t + \frac{1}{2}\sqrt{2} \frac{D_{-\lambda} \left(-\frac{1}{2}\sqrt{2}t \right)}{D_{-\lambda-1} \left(-\frac{1}{2}\sqrt{2}t \right)}. \quad (4.10)$$

We note that

$$\Phi_\lambda(t) = \frac{d}{dt} \ln \phi_\lambda(t),$$

where $\phi_\lambda(t) = \exp(\frac{1}{8}t^2)D_{-\lambda-1}(-\frac{1}{2}\sqrt{2}t)$ satisfies

$$\frac{d^2\phi_\lambda}{dt^2} - \frac{1}{2}t \frac{d\phi_\lambda}{dt} - \frac{1}{2}(\lambda + 1)\phi_\lambda = 0,$$

and $\Phi_\lambda(t)$ satisfies the Riccati equation

$$\frac{d\Phi_\lambda}{dt} = -\Phi_\lambda^2 + \frac{1}{2}t\Phi_\lambda + \frac{1}{2}(\lambda + 1) = 0,$$

see Lemma 4.4.

Using the recurrence relation (4.1), the first few polynomials are

$$\begin{aligned}S_1(x; t, \lambda) &= x, \\ S_2(x; t, \lambda) &= x^2 - \Phi_\lambda, \\ S_3(x; t, \lambda) &= x^3 + \frac{t\Phi_\lambda + \lambda + 1}{2\Phi_\lambda} x, \\ S_4(x; t, \lambda) &= x^4 + \frac{2t\Phi_\lambda^2 - (t^2 + 2)\Phi_\lambda - (\lambda + 1)t}{2(2\Phi_\lambda^2 - t\Phi_\lambda - \lambda - 1)} x^2 - \frac{2(\lambda + 2)\Phi_\lambda^2 - (\lambda + 1)t\Phi_\lambda - (\lambda + 1)^2}{2(2\Phi_\lambda^2 - t\Phi_\lambda - \lambda - 1)}, \\ S_5(x; t, \lambda) &= x^5 - \frac{2(\lambda + 3)t\Phi_\lambda^2 - (\lambda + 1)(t^2 - 2)\Phi_\lambda - (\lambda + 1)^2 t}{4(\lambda + 2)\Phi_\lambda^2 - 2(\lambda + 1)t\Phi_\lambda - 2(\lambda + 1)^2} x^3 \\ &\quad - \frac{[2(\lambda + 2)^2 - t^2]\Phi_\lambda^2 - (\lambda + 1)(\lambda + 4)t\Phi_\lambda - (\lambda + 1)^2(\lambda + 3)}{4(\lambda + 2)\Phi_\lambda^2 - 2(\lambda + 1)t\Phi_\lambda - 2(\lambda + 1)^2} x.\end{aligned}$$

Lemma 4.4. *The function $\Phi_\lambda(t)$ defined by (4.10) satisfies the Riccati equation*

$$\frac{d\Phi_\lambda}{dt} = -\Phi_\lambda^2 + \frac{1}{2}t\Phi_\lambda + \frac{1}{2}(\lambda + 1), \quad (4.11)$$

and has the asymptotic expansion as $t \rightarrow \infty$

$$\Phi_\lambda(t) = \frac{1}{2}t + \sum_{n=1}^{\infty} \frac{a_n}{t^{2n-1}}, \quad (4.12)$$

where the constants a_n are given by the nonlinear recurrence relation

$$a_{n+1} = 2(2n - 1)a_n - 2 \sum_{j=1}^n a_j a_{n+1-j},$$

with $a_1 = \lambda$. In particular, as $t \rightarrow \infty$

$$\Phi_\lambda(t) = \frac{1}{2}t + \frac{\lambda}{t} + \frac{2\lambda(1 - \lambda)}{t^3} + \frac{4\lambda(\lambda - 1)(2\lambda - 3)}{t^5} + \mathcal{O}(t^{-7}). \quad (4.13)$$

Proof. Letting $\Phi_\lambda(t) = \frac{d}{dt} \ln \phi_\lambda(t)$ in (4.11) yields

$$\frac{d^2 \phi_\lambda}{dt^2} - \frac{1}{2}t \frac{d\phi_\lambda}{dt} - \frac{1}{2}(\lambda + 1)\phi_\lambda = 0,$$

which has solution

$$\phi_\lambda(t) = \left\{ C_1 D_{-\lambda-1} \left(-\frac{1}{2}\sqrt{2}t \right) + C_2 D_{-\lambda-1} \left(\frac{1}{2}\sqrt{2}t \right) \right\} \exp\left(\frac{1}{8}t^2\right),$$

with C_1 and C_2 arbitrary constants. Hence setting $C_1 = 1$ and $C_2 = 0$ gives the solution (4.10) and shows that $\Phi_\lambda(t)$ satisfies (4.11).

Substituting (4.12) into (4.11) gives

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{(2n-1)a_n}{t^{2n}} &= \frac{1}{2}t \sum_{n=1}^{\infty} \frac{a_n}{t^{2n-1}} + \left(\sum_{n=1}^{\infty} \frac{a_n}{t^{2n-1}} \right)^2 - \frac{1}{2}\lambda \\ &= a_1 - \lambda + \frac{1}{2} \sum_{n=1}^{\infty} \frac{a_{n+1}}{t^{2n}} + \sum_{n=1}^{\infty} \frac{1}{t^{2n}} \sum_{j=1}^n a_j a_{n+1-j}, \end{aligned}$$

hence, comparing coefficients of powers of t gives $a_1 = \lambda$ and

$$a_{n+1} = 2(2n-1)a_n - 2 \sum_{j=1}^n a_j a_{n+1-j},$$

as required. Hence

$$a_1 = \lambda, \quad a_2 = -2\lambda(\lambda-1), \quad a_3 = 4\lambda(\lambda-1)(2\lambda-3),$$

which gives (4.13) as required. \square

Lemma 4.5. Let $h_n(t; \lambda)$ be defined by

$$H_n(t; \lambda) = \frac{d}{dt} \ln \tau_n(t; \lambda), \quad (4.14)$$

where $\tau_n(t; \lambda)$ is the Wronskian given by

$$\tau_n(t; \lambda) = \mathcal{W} \left(\phi_\lambda, \frac{d\phi_\lambda}{dt}, \dots, \frac{d^{n-1}\phi_\lambda}{dt^{n-1}} \right),$$

with

$$\phi_\lambda(t) = \frac{\Gamma(\lambda+1)}{2^{(\lambda+1)/2}} \exp\left(\frac{1}{8}t^2\right) D_{-\lambda-1} \left(-\frac{1}{2}\sqrt{2}t \right).$$

Then $H_n(t; \lambda)$ satisfies the second-order, second-degree equation

$$\left(\frac{d^2 H_n}{dt^2} \right)^2 - \frac{1}{4} \left(t \frac{dH_n}{dt} - H_n \right)^2 + \frac{dH_n}{dt} \left(2 \frac{dH_n}{dt} - n \right) \left(2 \frac{dH_n}{dt} - n - \lambda \right) = 0. \quad (4.15)$$

Proof. Equation (4.15) is equivalent to S_{IV} , the P_{IV} σ -equation

$$\left(\frac{d^2 \sigma}{dz^2} \right)^2 - 4 \left(z \frac{d\sigma}{dz} - \sigma \right)^2 + 4 \frac{d\sigma}{dz} \left(\frac{d\sigma}{dz} + 2\theta_0 \right) \left(\frac{d\sigma}{dz} + 2\theta_\infty \right) = 0, \quad (4.16)$$

as shown in [13, Theorem 4.11]. Equation (4.15) is the same as equation (4.15) in [13]. Special function solutions of S_{IV} (4.16) in terms of parabolic cylinder functions have been classified in [22, 37]; see also [13, Theorem 3.5]. \square

We remark that equation (4.16), and hence also equation (4.15), is equivalent to equation SD-I.c in the classification of second order, second-degree differential equations with the Painlevé property by Cosgrove and Scoufis [14], an equation first derived and solved by Chazy [7].

Lemma 4.6. As $t \rightarrow \infty$, the recurrence coefficient $\beta_n(t)$ has the asymptotic expansion

$$\beta_{2n}(t; \lambda) = \frac{n}{t} - \frac{2n(2\lambda - n + 1)}{t^3} + \mathcal{O}(t^{-5}), \quad (4.17a)$$

$$\beta_{2n+1}(t; \lambda) = \frac{t}{2} + \frac{\lambda - n}{t} - \frac{2(\lambda^2 - 4\lambda n + n^2 - \lambda - n)}{t^3} + \mathcal{O}(t^{-5}), \quad (4.17b)$$

for $n \in \mathbb{N}$.

Proof. In terms of the function $H_n(t; \lambda)$ defined by (4.14), the recurrence coefficients are given as

$$\beta_{2n}(t; \lambda) = H_n(t; \lambda + 1) - H_n(t; \lambda), \quad (4.18a)$$

$$\beta_{2n+1}(t; \lambda) = H_{n+1}(t; \lambda) - H_n(t; \lambda + 1). \quad (4.18b)$$

As $t \rightarrow \infty$, $H_n(t; \lambda)$ has the asymptotic expansion

$$H_n(t; \lambda) = \frac{nt}{2} + \frac{n\lambda}{t} + \frac{2n\lambda(n - \lambda)}{t^3} + \mathcal{O}(t^{-5}), \quad (4.19)$$

for $n \in \mathbb{N}$, see [13, Lemma 5.2]; note that the functions $\Delta_n(t)$ and $S_n(t)$ in [13] are the same as our functions $\tau_n(t; \lambda)$ and $H_n(t; \lambda)$, respectively. Substituting (4.19) in (4.18) immediately gives the result. \square

4.3 The differential-difference equation satisfied by generalized Freud polynomials

In this section we derive a differential-difference equation satisfied by generalized Freud polynomials using our results in §2.

Lemma 4.7. For the generalized Freud weight (1.9) the monic orthogonal polynomials $S_n(x; t)$ satisfy

$$\int_{-\infty}^{\infty} \mathcal{K}(x, y) S_n^2(y; t) w(y; t) dy = 4(x^2 - \frac{1}{2}t + \beta_n + \beta_{n+1})h_n, \quad (4.20)$$

$$\int_{-\infty}^{\infty} \mathcal{K}(x, y) S_n(y; t) S_{n-1}(y; t) w(y; t) dy = 4xh_n, \quad (4.21)$$

where

$$\mathcal{K}(x, y) = \frac{v'(x; t) - v'(y; t)}{x - y},$$

with $v(x; t) = x^4 - tx^2$ and

$$h_n = \int_{-\infty}^{\infty} S_n^2(y; t) w(y; t) dy. \quad (4.22)$$

Proof. For the generalized Freud weight (1.9) we have

$$w(x; t) = |x|^{2\lambda+1} \exp(-x^4 + tx^2),$$

i.e. $v(x; t) = x^4 - tx^2$, and so

$$\mathcal{K}(x, y) = 4x^2 + 4xy + 4y^2 - 2t.$$

Hence for (4.20)

$$\begin{aligned} \int_{-\infty}^{\infty} \mathcal{K}(x, y) S_n^2(y; t) w(y; t) dy &= (4x^2 - 2t) \int_{-\infty}^{\infty} S_n^2(y; t) w(y; t) dy \\ &\quad + 4x \int_{-\infty}^{\infty} y S_n^2(y; t) w(y; t) dy + 4 \int_{-\infty}^{\infty} y^2 S_n^2(y; t) w(y; t) dy \\ &= (4x^2 - 2t)h_n + 4x \int_{-\infty}^{\infty} S_n(y; t) [S_{n+1}(y; t) + \beta_n S_{n-1}(y; t)] w(y; t) dy \\ &\quad + 4 \int_{-\infty}^{\infty} [S_{n+1}(y; t) + \beta_n S_{n-1}(y; t)]^2 w(y; t) dy \\ &= (4x^2 - 2t)h_n + 4h_{n+1} + 4\beta_n^2 h_{n-1} \\ &= 4(x^2 - \frac{1}{2}t + \beta_n + \beta_{n+1})h_n, \end{aligned}$$

since $\beta_n = h_n/h_{n-1}$, the monic orthogonal polynomials $S_n(x; t)$ satisfy the three-term recurrence relation (4.1), and are orthogonal, i.e.

$$\int_{-\infty}^{\infty} S_m(y; t) S_n(y; t) w(y; t) dy = 0, \quad \text{if } m \neq n. \quad (4.23)$$

Also for (4.21)

$$\begin{aligned} & \int_{-\infty}^{\infty} \mathcal{K}(x, y) S_n(y; t) S_{n-1}(y; t) w(y; t) dy \\ &= (4x^2 - 2t) \int_{-\infty}^{\infty} S_n(y; t) S_{n-1}(y; t) w(y; t) dy + 4x \int_{-\infty}^{\infty} y S_n(y; t) S_{n-1}(y; t) w(y; t) dy \\ & \quad + 4 \int_{-\infty}^{\infty} y^2 S_n(y; t) S_{n-1}(y; t) w(y; t) dy \\ &= 4x \int_{-\infty}^{\infty} S_n(y; t) [S_n(y; t) + \beta_{n-1} S_{n-2}(y; t)] w(y; t) dy \\ & \quad + 4 \int_{-\infty}^{\infty} [S_{n+1}(y; t) + \beta_n S_{n-1}(y; t)] [S_n(y; t) + \beta_{n-2} S_{n-2}(y; t)] w(y; t) dy \\ &= 4x h_n, \end{aligned}$$

using the recurrence relation (4.1) and orthogonality (4.23). \square

Theorem 4.8. *For the generalized Freud weight (1.9) the monic orthogonal polynomials $S_n(x; t)$ satisfy the differential-difference equation*

$$x \frac{dS_n}{dx}(x; t) = -B_n(x; t) S_n(x; t) + A_n(x; t) S_{n-1}(x; t), \quad (4.24)$$

where

$$A_n(x; t) = 4x\beta_n(x^2 - \frac{1}{2}t + \beta_n + \beta_{n+1}), \quad (4.25a)$$

$$B_n(x; t) = 4x^2\beta_n + (\lambda + \frac{1}{2})[1 - (-1)^n], \quad (4.25b)$$

with β_n the recurrence coefficient in the three-term recurrence relation (4.1).

Proof. Corollary 2.3 shows that monic orthogonal polynomials $S_n(x; t)$ with respect to the weight

$$w(x; t) = |x|^{2\lambda+1} \exp\{-v(x; t)\},$$

satisfy the differential-difference equation (4.24), where

$$A_n(x; t) = \frac{x}{h_{n-1}} \int_{-\infty}^{\infty} \mathcal{K}(x, y) S_n^2(y; t) w(y; t) dy, \quad (4.26a)$$

$$B_n(x; t) = \frac{x}{h_{n-1}} \int_{-\infty}^{\infty} \mathcal{K}(x, y) S_n(y; t) S_{n-1}(y; t) w(y; t) dy + (\lambda + \frac{1}{2})[1 + (-1)^n]. \quad (4.26b)$$

For the generalized Freud weight (1.9), using Lemma 4.7 yields the result. \square

4.4 The differential equation satisfied by generalized Freud polynomials

Now we derive a differential equation satisfied by generalized Freud polynomials.

Theorem 4.9. *For the generalized Freud weight (1.9) the monic orthogonal polynomials $S_n(x; t)$ satisfy the differential equation*

$$x \frac{d^2 S_n}{dx^2}(x; t) + R_n(x; t) \frac{dS_n}{dx}(x; t) + T_n(x; t) S_n(x; t) = 0, \quad (4.27)$$

where

$$R_n(x; t) = -4x^4 + 2tx^2 + 2\lambda + 1 - \frac{2x^2}{x^2 - \frac{1}{2}t + \beta_n + \beta_{n+1}}, \quad (4.28a)$$

$$\begin{aligned} T_n(x; t) &= 4nx^3 + 16x\beta_n(\beta_n + \beta_{n+1} - \frac{1}{2}t)(\beta_n + \beta_{n-1} - \frac{1}{2}t) \\ &\quad + 4x[1 + (2\lambda + 1)(-1)^n]\beta_n - \frac{8\beta_n x^3 + (2\lambda + 1)[1 - (-1)^n]x}{x^2 - \frac{1}{2}t + \beta_n + \beta_{n+1}} \\ &\quad + (2\lambda + 1)[1 - (-1)^n]x \left(t - \frac{1}{2x^2} \right). \end{aligned} \quad (4.28b)$$

Proof. In Theorem 2.6 we proved that the coefficients in the differential equation (2.14a) satisfied by polynomials orthogonal with respect to the weight

$$w(x; t) = |x - k|^\gamma \exp\{-v(x; t)\},$$

are given by (2.14b) and (2.14c). For the generalized Freud weight (1.9) we use (2.14b) and (2.14c) with $k = 0$, $\gamma = 2\lambda + 1$, $v(x; t) = x^4 - tx^2$, and A_n and B_n given by (4.25) to obtain the stated result. \square

Remark 4.10. We note that if $\{P_n(x)\}_{n=0}^\infty$ is a sequence of *classical* orthogonal polynomials (such as Hermite, Laguerre and Jacobi polynomials), then $P_n(x)$ satisfies second-order ordinary differential equation

$$\sigma(x) \frac{d^2 P_n}{dx^2} + \tau(x) \frac{d P_n}{dx} = \lambda_n P_n, \quad (4.29)$$

where $\sigma(x)$ is a monic polynomial with $\deg(\sigma) \leq 2$, $\tau(x)$ is a polynomial with $\deg(\tau) = 1$, and λ_n is a real number which depends on the degree of the polynomial solution, see Bochner [5]. For classical orthogonal polynomials, the polynomials $\sigma(x)$ and $\tau(x)$ are the same as in the associated Pearson equation (1.3). In contrast the coefficients in second-order ordinary differential equation satisfied by the polynomials for the generalized Freud weight given in Theorem 4.9 are not the same as the polynomials $\sigma(x) = x$ and $\tau(x) = -4x^4 + 2tx^2 + 2\lambda + 1$ in the Pearson equation (1.3) satisfied by the generalized Freud polynomials since

$$\frac{w'(x; t)}{w(x; t)} = \frac{\tau(x) - \sigma'(x)}{\sigma(x)} = -4x^3 + 2tx + \frac{2\lambda + 1}{x}, \quad x \in \mathbb{R} \setminus \{0\}.$$

4.5 An alternative method due to Shohat

It is shown in [35] and [40] that the monic orthogonal polynomials $S_n(x; t)$ orthogonal with respect to the generalized Freud weight (1.9) are quasi-orthogonal of order $m = 5$ and hence we can write

$$x \frac{dS_n}{dx}(x; t) = \sum_{k=n-4}^n c_{n,k} S_k(x; t), \quad (4.30)$$

where the coefficient $c_{n,k}$ is given by

$$c_{n,k} = \frac{1}{h_k} \int_{-\infty}^{\infty} x \frac{dS_n}{dx}(x; t) S_k(x; t) w(x; t) dx, \quad (4.31)$$

for $n - 4 \leq k \leq n$ and $h_k \neq 0$.

Integrating by parts, we obtain for $n - 4 \leq k \leq n - 1$,

$$\begin{aligned} h_k c_{n,k} &= \left[x S_k(x; t) S_n(x; t) w(x; t) \right]_{-\infty}^{\infty} - \int_{-\infty}^{\infty} \frac{d}{dx} [x S_k(x; t) w(x; t)] S_n(x; t) dx \\ &= - \int_{-\infty}^{\infty} \left[S_n(x; t) S_k(x; t) + x S_n(x; t) \frac{dS_k}{dx}(x; t) \right] w(x; t) dx - \int_{-\infty}^{\infty} x S_n(x; t) S_k(x; t) \frac{dw}{dx}(x; t) dx, \\ &= - \int_{-\infty}^{\infty} x S_n(x; t) S_k(x; t) \frac{dw}{dx}(x; t) dx \\ &= - \int_{-\infty}^{\infty} S_n(x; t) S_k(x; t) (-4x^4 + 2tx^2 + 2\lambda + 1) w(x; t) dx \\ &= \int_{-\infty}^{\infty} (4x^4 - 2tx^2) S_n(x; t) S_k(x; t) w(x; t) dx, \end{aligned} \quad (4.32)$$

since

$$x \frac{dw}{dx}(x; t) = (-4x^4 + 2x^2 + 2\lambda + 1)w(x; t).$$

Iterating the three-term recurrence relation (4.1), the following relations are obtained

$$x^2 S_n = S_{n+2} + (\beta_n + \beta_{n+1})S_n + \beta_n \beta_{n-1} S_{n-2}, \quad (4.33a)$$

$$\begin{aligned} x^4 S_n &= S_{n+4} + (\beta_n + \beta_{n+1} + \beta_{n+2} + \beta_{n+3})S_{n+2} \\ &\quad + [\beta_n(\beta_{n-1} + \beta_n + \beta_{n+1}) + \beta_{n+1}(\beta_n + \beta_{n+1} + \beta_{n+2})]S_n \\ &\quad + \beta_n \beta_{n-1}(\beta_{n-2} + \beta_{n-1} + \beta_n + \beta_{n+1})S_{n-2} + (\beta_n \beta_{n-1} \beta_{n-2} \beta_{n-3})S_{n-4}. \end{aligned} \quad (4.33b)$$

Substituting (4.33) into (4.32) yields the coefficients $\{c_{n,k}\}_{k=n-4}^{n-1}$ in (4.30).

$$c_{n,n-4} = 4\beta_n \beta_{n-1} \beta_{n-2} \beta_{n-3}, \quad (4.34a)$$

$$c_{n,n-3} = 0, \quad (4.34b)$$

$$c_{n,n-2} = 4\beta_n \beta_{n-1}(\beta_{n-2} + \beta_{n-1} + \beta_n + \beta_{n+1} - \frac{1}{2}t), \quad (4.34c)$$

$$c_{n,n-1} = 0. \quad (4.34d)$$

Lastly we consider the case when $k = n$. Integration by parts in (4.31) yields

$$\begin{aligned} h_n c_{n,n} &= \int_{-\infty}^{\infty} x \frac{dS_n}{dx}(x; t) S_n(x; t) w(x; t) dx \\ &= -\frac{1}{2} \int_{-\infty}^{\infty} S_n^2(x; t) \left[w(x; t) + x \frac{dw}{dx}(x; t) \right] dx \\ &= -\frac{1}{2} h_n + \int_{-\infty}^{\infty} S_n^2(x; t) (2x^4 - tx^2 - \lambda - \frac{1}{2}) w(x; t) dx \\ &= \int_{-\infty}^{\infty} (2x^4 - tx^2) S_n^2(x; t) w(x; t) dx - (\lambda + 1) h_n. \end{aligned} \quad (4.35)$$

From the three-term recurrence relation (4.1), we have

$$\begin{aligned} x^2 S_n^2 &= (S_{n+1} + \beta_n S_{n-1})^2 \\ &= S_{n+1}^2 + 2\beta_n S_{n+1} S_{n-1} + \beta_n^2 S_{n-1}^2 \\ x^4 S_n^2 &= x^2 (S_{n+1}^2 + 2\beta_n S_{n+1} S_{n-1} + \beta_n^2 S_{n-1}^2) \\ &= x^2 S_{n+1}^2 + 2\beta_n (x S_{n+1})(x S_{n-1}) + \beta_n^2 x^2 S_{n-1}^2 \\ &= (S_{n+2} + \beta_{n+1} S_n)^2 + 2\beta_n (S_{n+2} + \beta_{n+1} S_n)(S_n + \beta_{n-1} S_{n-2}) + \beta_n^2 (S_n + \beta_{n-1} S_{n-2})^2 \\ &= S_{n+2}^2 + 2(\beta_{n+1} + \beta_n) S_{n+2} S_n + (\beta_{n+1} + \beta_n)^2 S_n^2 + 2\beta_n \beta_{n-1} S_{n+2} S_{n-2} \\ &\quad + 2\beta_n \beta_{n-1} (\beta_n + \beta_{n+1}) S_n S_{n-2} + \beta_n^2 \beta_{n-1}^2 S_{n-2}^2 \end{aligned}$$

and so by orthogonality

$$\int_{-\infty}^{\infty} x^2 S_n^2(x; t) w(x; t) dx = h_{n+1} + \beta_n^2 h_{n-1} = (\beta_{n+1} + \beta_n) h_n, \quad (4.36)$$

$$\begin{aligned} \int_{-\infty}^{\infty} x^4 S_n^2(x; t) w(x; t) dx &= h_{n+2} + (\beta_{n+1} + \beta_n)^2 h_n + \beta_n^2 \beta_{n-1}^2 h_{n-2} \\ &= \beta_{n+2} \beta_{n+1} h_n + (\beta_{n+1} + \beta_n)^2 h_n + \beta_n \beta_{n-1} h_n \\ &= [(\beta_{n+1} + \beta_n + \beta_{n-1}) \beta_n + (\beta_{n+2} + \beta_{n+1} + \beta_n) \beta_{n+1}] h_n \\ &= \frac{1}{2} [t(\beta_{n+1} + \beta_n) + n + \lambda + 1] h_n, \end{aligned} \quad (4.37)$$

using $h_{n+1} = \beta_{n+1} h_n$ and dP_I (4.3). Hence from (4.35), (4.36) and (4.37) we have

$$\begin{aligned} c_{n,n} &= t(\beta_{n+1} + \beta_n) + n + \lambda + 1 - t(\beta_{n+1} + \beta_n) - (\lambda + 1) \\ &= n. \end{aligned} \quad (4.38)$$

Combining (4.34) with (4.30), we write

$$x \frac{dS_n}{dx}(x; t) = c_{n,n-4} S_{n-4}(x; t) + c_{n,n-2} S_{n-2}(x; t) + c_{n,n} S_n(x; t). \quad (4.39)$$

In order to express S_{n-4} and S_{n-2} in (4.39) in terms of S_n and S_{n-1} , we iterate (4.1) to obtain

$$S_{n-2} = \frac{xS_{n-1} - S_n}{\beta_{n-1}}, \quad (4.40)$$

$$S_{n-3} = \frac{xS_{n-2} - S_{n-1}}{\beta_{n-2}} = \frac{x^2 - \beta_{n-1}}{\beta_{n-1}\beta_{n-2}} S_{n-1} - \frac{x}{\beta_{n-1}\beta_{n-2}} S_n, \quad (4.41)$$

$$S_{n-4} = \frac{xS_{n-3} - S_{n-2}}{\beta_{n-3}} = \frac{x^3 - (\beta_{n-1} + \beta_{n-2})x}{\beta_{n-1}\beta_{n-2}\beta_{n-3}} S_{n-1} - \frac{x^2 - \beta_{n-2}}{\beta_{n-1}\beta_{n-2}\beta_{n-3}} S_n. \quad (4.42)$$

Substituting (4.34), (4.38), (4.40) and (4.42) into (4.39) yields

$$x \frac{dS_n}{dx}(x; t) = -B_n(x) S_n(x; t) + A_n(x; t) S_{n-1}(x; t), \quad (4.43)$$

where $A_n(x)$ and $B_n(x)$ are given by (4.25).

5 Conclusion

In this paper, for the generalized Freud weight (1.9) we have obtained explicit expressions for the coefficients of the three-term recurrence relation and differential-difference equation satisfied by generalized Freud polynomials. We also proved that the generalized Freud polynomials satisfy a linear ordinary differential equation. We note that the closed form expressions for the coefficients provided allow investigation of other properties, including properties of the zeros such as monotonicity, convexity and inequalities satisfied by the zeros. However, although the expressions for the coefficients given in this paper are explicit, they are rather complicated and given in terms of special function solutions of the fourth Painlevé equation which does not necessarily lead to elegant results in applications. For this reason, a natural extension of this work would be an investigation of asymptotic properties using limiting relations satisfied by the polynomials as the parameters t and/or λ tend to ∞ .

Acknowledgments

We thank the London Mathematical Society for support through a ‘‘Research in Pairs’’ grant. PAC thanks Alexander Its, Ana Loureiro, and Walter van Assche for their helpful comments and illuminating discussions. KJ thanks the School of Mathematics, Statistics & Actuarial Science at the University of Kent for their hospitality during her visit when some of this research was done. The research by KJ was partially supported by the National Research Foundation of South Africa.

References

- [1] W.A. Al Salam and T.S. Chihara, Another characterization of the classical orthogonal polynomials, *SIAM J. Math. Anal.*, **18** (1987) 228–242.
- [2] A.F. Beardon, The theorems of Stieltjes and Favard, *Comput. Methods Funct. Theory*, **11** (2011) 247–262.
- [3] P. Bleher and A. Its, Semiclassical asymptotics of orthogonal polynomials, Riemann-Hilbert problem, and universality in the matrix model, *Ann. Math.*, **150** (1999) 185–266.
- [4] R. Bo and R. Wong, A uniform asymptotic formula for orthogonal polynomials associated with $\exp(-x^4)$, *J. Approx. Theory*, **98** (1999) 146–166.
- [5] S. Bochner, Über Sturm-Liouvillesche Polynomsysteme, *Math. Z.*, **29** (1929) 730–736.
- [6] S. Bonan and P. Nevai, Orthogonal polynomials and their derivatives. I, *J. Approx. Theory*, **40** (1984) 134–147.
- [7] J. Chazy, Sur les équations différentielles du troisième ordre et d’ordre supérieur dont l’intégrale générale a ses points critiques fixes, *Acta Math.*, **34** (1911) 317–385.
- [8] Y. Chen and M.V. Feigin, Painlevé IV and degenerate Gaussian Unitary Ensembles, *J. Phys. A: Math. Gen.*, **39** (2006) 12381–12393.

- [9] Y. Chen and M.E.H. Ismail, Ladder operators and differential equations for orthogonal polynomials, *J. Phys. A*, **30** (1997) 7817–7829.
- [10] Y. Chen and A. Its, Painlevé III and a singular statistics in Hermitian random matrix ensembles, I, *J. Approx. Theory*, **162** (2010) 270–297.
- [11] Y. Chen and L. Zhang, Painlevé VI and the Unitary Jacobi Ensembles, *Stud. Appl. Math.*, **125** (2010) 91–112.
- [12] T.S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, 1978. [Reprinted by Dover Publications, 2011.]
- [13] P.A. Clarkson and K. Jordaan, The relationship between semi-classical Laguerre polynomials and the fourth Painlevé equation, *Constr. Approx.*, **39** (2014) 223–254.
- [14] C.M. Cosgrove and G. Scoufis, Painlevé classification of a class of differential equations of the second order and second-degree, *Stud. Appl. Math.*, **88** (1993) 25–87.
- [15] P. Deift, T. Kriecherbauer, K.T.-R. McLaughlin, S. Venakides, and X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, *Comm. Pure Appl. Math.*, **52** (1999) 1491–1552.
- [16] P. Deift, T. Kriecherbauer, K.T.-R. McLaughlin, S. Venakides, and X. Zhou, Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory, *Comm. Pure Appl. Math.*, **52** (1999) 1335–1425.
- [17] J. Favard, Sur les polynômes de Tchebicheff., *C. R. Acad. Sci., Paris*, **200** (1935) 2052–2053.
- [18] G. Filipuk, W. van Assche, and L. Zhang, The recurrence coefficients of semi-classical Laguerre polynomials and the fourth Painlevé equation, *J. Phys. A*, **45** (2012) 205201.
- [19] A.S. Fokas, A.R. Its, and A.V. Kitaev, Discrete Painlevé equations and their appearance in quantum-gravity, *Commun. Math. Phys.*, **142** (1991) 313–344.
- [20] A.S. Fokas, A.R. Its, and A.V. Kitaev, The isomonodromy approach to matrix models in 2D quantum-gravity, *Commun. Math. Phys.*, **147** (1992) 395–430.
- [21] A.S. Fokas, A.R. Its, and X. Zhou, Continuous and discrete Painlevé equations, in *Painlevé Transcendents. Their Asymptotics and Physical Applications*, D. Levi and P. Winternitz (Editors), NATO Adv. Sci. Inst. Ser. B Phys., vol. 278, pp. 33–47 (Plenum, New York, 1992).
- [22] P.J. Forrester and N.S. Witte, Application of the τ -function theory of Painlevé equations to random matrices: PIV, PII and the GUE, *Commun. Math. Phys.*, **219** (2001) 357–398.
- [23] G. Freud, On the coefficients in the recursion formulae of orthogonal polynomials, *Proc. R. Irish Acad., Sect. A*, **76** (1976) 1–6.
- [24] P. Gochhayat, K. Jordaan, K. Raghavendar, and A. Swaminathan, Interlacing properties and bounds for zeros of ${}_2\phi_1$ hypergeometric and little q -Jacobi polynomials, *Ramanujan Journal*, doi.org/10.1007/s11139-015-9709-5 (2015).
- [25] E. Hendriksen and H. van Rossum, Semi-classical orthogonal polynomials, in *Polynômes Orthogonaux et Applications*, C. Brezinski, A. Draux, A.P. Magnus, P. Maroni, and A. Ronveaux (Editors), Lect. Notes Math., Vol. 1171 pp. 354–361, Springer-Verlag, Berlin, 1985.
- [26] M.E.H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, Encyclopedia of Mathematics and its Applications, Vol. 98, Cambridge University Press, Cambridge, 2005.
- [27] T. Kriecherbauer and K.T.-R. McLaughlin, Strong asymptotics of polynomials orthogonal with respect to Freud weights, *Internat. Math. Res. Notices*, **6** (1999) 299–333.
- [28] E. Levin and D. S. Lubinsky, *Orthogonal Polynomials for Exponential Weights*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, Vol. 4, Springer-Verlag, New York, 2001.
- [29] A.P. Magnus, Painlevé-type differential equations for the recurrence coefficients of semi-classical orthogonal polynomials, *J. Comput. Appl. Math.*, **57** (1995) 215–237.
- [30] F. Marcellán and R. Álvarez-Nodarse, On the “Favard theorem” and its extensions, *J. Comput. Appl. Math.*, **127** (2001) 231–254.
- [31] P. Maroni, Une caractérisation des polynômes orthogonaux semi-classique, *Ann. Mat. pura et Appl.*, **149(4)** (1987) 165–184.
- [32] P. Maroni, Le calcul des formes lineaires et les polynômes orthogonaux semiclassiques, in *Orthogonal Polynomials and their Applications*, M. Alfaro, J.S. Dehesa, F.J. Marcellán, J.L. Rubio de Francia, and J. Vinesa (Editors), Lecture Notes Math., Vol. 1329 pp. 279–290, Springer, Berlin, 1988.
- [33] H.N. Mhaskar, Bounds for certain Freud-type orthogonal polynomials, *J. Approx. Theory*, **63** (1990) 238–254.
- [34] I.P. Natanson, *Konstruktive Funktionentheorie*, Akademie-Verlag, Berlin, 1955.
- [35] P. Nevai, Géza Freud, Orthogonal Polynomials and Christoffel Functions. A case study, *J. Approx. Th.*, **48** (1986) 3–167.

- [36] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, and C.W. Clark (Editors), *DLMF Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, 2010.
- [37] K. Okamoto, Studies on the Painlevé equations III. Second and fourth Painlevé equations, P_{II} and P_{IV} , *Math. Ann.*, **275** (1986) 221–255.
- [38] J. Shohat, The Relation of the Classical Orthogonal Polynomials to the Polynomials of Appell, *Amer. J. Math.*, **58** (1936) 453–464.
- [39] J. Shohat, Sur les polynômes orthogonaux généralisés, *C. R. Acad. Sci., Paris*, **207** (1938) 556–558.
- [40] J. Shohat, A differential equation for orthogonal polynomials, *Duke Math. J.*, **5** (1939) 401–417.
- [41] M.H. Stone, *Linear transformations in Hilbert space and their applications to analysis*, AMS Colloquium Publications, Vol. 15, Amer. Math. Soc., Providence RI, 1932.
- [42] T.J. Stieltjes, Recherches sur les fractions continues, *Ann. Fac. Sci. Toulouse Sci. Math. Sci. Phys.*, **8:J1–J122 (1894); 9** (1895) A5–A47.
- [43] G. Szegő, *Orthogonal Polynomials*, AMS Colloquium Publications, Vol. 23, Amer. Math. Soc., Providence RI, 1975.
- [44] W. Van Assche, The impact of Stieltjes work on continued fractions and orthogonal polynomials, in *Collected Papers*, G. van Dijk (Editor), pp. 5–37, Springer-Verlag, Berlin, 1993.
- [45] A. Wintner, *Spektraltheorie der unendlichen Matrizen*, S. Hirzel, Leipzig, 1929.