



Kent Academic Repository

Pech, Clelia and Rietsch, Konstanze (2018) *A comparison of Landau-Ginzburg models for odd-dimensional Quadrics*. Bulletin of the Institute of Mathematics Academia Sinica, 13 (3). pp. 249-291. ISSN 2304-7909.

Downloaded from

<https://kar.kent.ac.uk/51113/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.21915/BIMAS.2018301>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

A COMPARISON OF LANDAU-GINZBURG MODELS FOR ODD DIMENSIONAL QUADRICS

CLELIA PECH AND KONSTANZE RIETSCH

This paper is dedicated to George Lusztig on his 70th birthday.

ABSTRACT. In [Rie08], the second author defined a Landau-Ginzburg model for homogeneous spaces G/P . In this paper, we reformulate this LG model in the case of the odd-dimensional quadric $X = Q_{2m-1}$. Namely we introduce a regular function \mathcal{W}_{can} on a variety $\check{X}_{\text{can}} \times \mathbb{C}^*$, where \check{X}_{can} is the complement of a particular anticanonical divisor in the projective space $\mathbb{C}\mathbb{P}^{2m-1} = \mathbb{P}(H^*(X, \mathbb{C})^*)$. Firstly we prove that the Jacobi ring associated to \mathcal{W}_{can} is isomorphic to the quantum cohomology ring of the quadric, and that this isomorphism is compatible with the identification of homogeneous coordinates on $\check{X}_{\text{can}} \subset \mathbb{C}\mathbb{P}^{2m-1}$ with elements of $H^*(X, \mathbb{C})$. Secondly we find a very natural Laurent polynomial formula for \mathcal{W}_{can} by restricting it to a ‘Lusztig torus’ in \check{X}_{can} . Thirdly we show that the Dubrovin connection on $H^*(X, \mathbb{C}[q])$ embeds into the Gauss-Manin system associated to \mathcal{W}_{can} and deduce a flat section formula in terms of oscillating integrals. Finally, we compare $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ with previous Landau-Ginzburg models defined for odd quadrics. Namely, we prove that it is a partial compactification of Givental’s original LG model [Giv96]. We show that our LG model is isomorphic to the Lie-theoretic LG model from [Rie08]. Moreover it is birationally equivalent to an LG model introduced by Gorbounov and Smirnov [GS13], and it is algebraically isomorphic to Gorbounov and Smirnov’s mirror for Q_3 , implying a tameness property in that case.

1. INTRODUCTION

The geometric Satake correspondence [Lus83, Gin95, MV07] constructs representations of a reductive algebraic group G in terms of geometry of the affine Grassmannian of the Langlands dual group G^\vee . It has its origins in the seminal paper of Lusztig [Lus83]. In this paper we describe the mirror symmetry partner of a smooth, odd-dimensional complex quadric $X = Q_{2m-1}$ from the point of view of its automorphism group $G^\vee = \text{SO}_{2m+1}(\mathbb{C})$, Langlands duality and the geometric Satake correspondence.

Recall that the Langlands dual group of $\text{SO}_{2m+1}(\mathbb{C})$ is the symplectic group $\text{Sp}_{2m}(\mathbb{C})$. The geometric Satake correspondence provides us with a ‘Langlands dual’ interpretation of the cohomology $H^*(X, \mathbb{C})$ of the smooth quadric $X = Q_{2m-1}$ as follows. The quadric X appears as one of the simplest Schubert varieties inside the affine Grassmannian of $\text{SO}_{2m+1}(\mathbb{C})$,

$$X \hookrightarrow \text{Gr}_{G^\vee} = \text{SO}_{2m+1}(\mathbb{C}((t)))/\text{SO}_{2m+1}(\mathbb{C}[[t]]).$$

Namely this Schubert variety is associated to the first fundamental coweight of $\text{SO}_{2m+1}(\mathbb{C})$. The geometric Satake correspondence reinterprets this coweight as a dominant *weight* for the Langlands dual group, $\text{Sp}_{2m}(\mathbb{C})$. Moreover the intersection cohomology of the associated Schubert variety X is then understood to be the representation of $\text{Sp}_{2m}(\mathbb{C})$ with that highest weight. In our setting, since the quadric is smooth, the intersection cohomology coincides

2000 *Mathematics Subject Classification.* 14N35, 14M17, 14J33, 57T15.

Key words and phrases. Mirror Symmetry, quadrics, Lie theory, Gromov-Witten theory, quantum cohomology, Landau-Ginzburg model, Gauss-Manin system.

This work was supported by the Leverhulme Trust grant n° F07040AW - A Lie theoretic approach to derived categories of flag varieties G/P .

with the usual cohomology of X and we obtain the interpretation,

$$H^*(X, \mathbb{C}) = \mathbb{C}^{2m} = V_{\omega_1},$$

of the cohomology of X , where V_{ω_1} is the defining representation of $G = \mathrm{Sp}_{2m}(\mathbb{C})$.

In mirror symmetry a ‘mirror dual’ construction of the *quantum* cohomology ring of X is sought, along with other structures involving Gromov-Witten invariants of X , see [Bea95, CK99]. In the setting of the quadric X , the (small) quantum cohomology ring is a 1-parameter deformation of $H^*(X, \mathbb{C})$, whose structure constants are Gromov-Witten invariants that count 3-pointed genus 0 holomorphic curves in X subject to certain constraints. The result is a commutative algebra structure on $H^*(X, \mathbb{C}) \otimes \mathbb{C}[q]$ which recovers the usual cohomology ring when $q \rightarrow 0$. For explicit formulas in the case of quadrics we refer to [CMP07].

There are various previous mirror constructions that apply to odd quadrics, which we recall in Section 2. Some of these already recover the quantum cohomology ring, and one construction is already in terms of the Langlands dual group $\mathrm{Sp}_{2m}(\mathbb{C})$. The main new construction we introduce in this paper combines the geometric Satake correspondence with the ‘Langlands dual group’ construction of the mirror. As a result we construct a mirror for the quadric X that is expressed in terms of coordinates which are naturally identified with cohomology classes of the quadric. An analogous construction was carried out for Grassmannians in [MR13], and then for Lagrangian Grassmannians in [PR13].

The mirror of the quadric X takes the form of a *Landau-Ginzburg model* or *LG model*, that is, of a pair $(\check{X}_{\mathrm{can}}, \mathcal{W}_{\mathrm{can}})$, where \check{X}_{can} is an affine Calabi-Yau variety and $\mathcal{W}_{\mathrm{can}}$ is a regular function $\check{X}_{\mathrm{can}} \rightarrow \mathbb{C}$. In our construction \check{X}_{can} is the complement of a particular anticanonical divisor in projective space,

$$\check{X}_{\mathrm{can}} := \mathbb{C}\mathbb{P}^{2m-1} = \mathbb{P}(H^*(X, \mathbb{C})^*) = \mathrm{Proj}(\mathbb{C}[p_0, \dots, p_{2m-1}]).$$

Here $\mathbb{C}\mathbb{P}^{2m-1}$ is viewed as the homogeneous space $\mathbb{P}(V_{\omega_1}^*)$ for the symplectic group $G = \mathrm{Sp}_{2m}(\mathbb{C})$. Thus the first equality is by the geometric Satake correspondence. In the second equality, the variables or homogeneous coordinates, $p_0, p_1, \dots, p_{2m-1}$ are identified with the Schubert basis, $\sigma_0, \sigma_1, \dots, \sigma_{2m-1} \in H^*(X, \mathbb{C})$, which has one element in each even degree.

To give a concrete example, in the case of $X = Q_3$ our formula reads

$$(1) \quad \mathcal{W}_{\mathrm{can}, q} = \frac{p_1}{p_0} + \frac{p_2^2}{p_1 p_2 - p_0 p_3} + q \frac{p_1}{p_3}$$

in terms of the homogeneous coordinates $(p_0 : p_1 : p_2 : p_3)$ on $\check{X}_{\mathrm{can}} = \mathbb{C}\mathbb{P}^3$, which are identified with the Schubert classes of Q_3 .

In the case of $X = Q_5$ our formula reads

$$(2) \quad \mathcal{W}_{\mathrm{can}, q} = \frac{p_1}{p_0} + \frac{p_2 p_4}{p_1 p_4 - p_0 p_5} + \frac{p_3^2}{p_2 p_3 - p_1 p_4 + p_0 p_5} + q \frac{p_1}{p_5}$$

in terms of the homogeneous coordinates $(p_0 : p_1 : p_2 : p_3 : p_4 : p_5)$ on $\check{X}_{\mathrm{can}} = \mathbb{C}\mathbb{P}^5$, which are identified with the Schubert classes of Q_5 .

We compare our formula with previous constructions of Landau-Ginzburg models [Giv96, Rie08, GS13], and obtain various mirror theorems for our LG model.

1.1. Quantum Cohomology. The LG model $(\check{X}_{\mathrm{can}}, \mathcal{W}_{\mathrm{can}})$ provides the following Jacobi ring description of the small quantum cohomology ring of the quadric X . Let δ_i denote the i^{th} quadratic denominator of the superpotential $\mathcal{W}_{\mathrm{can}}$ when $i = 1 \dots m-1$, and $\delta_m = p_{2m-1}$.

Then

$$(3) \quad \mathrm{qH}^*(X, \mathbb{C})[q^{-1}] = \mathbb{C}[\check{X}_{\mathrm{can}} \times \mathbb{C}_q^*] / \left(\frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_1}, \frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_2}, \dots, \frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_{2m-1}} \right) \\ = \mathbb{C}[p_1, \dots, p_{2m-1}, \delta_1^{-1}, \dots, \delta_m^{-1}, q^{\pm 1}] / \left(\frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_1}, \frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_2}, \dots, \frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_{2m-1}} \right),$$

where p_i is identified with the (unique) Schubert class generator σ_i in $\mathrm{qH}^{2i}(X, \mathbb{C})$ and we have set $p_0 = 1$. In particular, the equality (3) says that the p_i span the right hand side as a free $\mathbb{C}[q, q^{-1}]$ -module, and the multiplicative structure constants compute 3-point genus 0 Gromov-Witten invariants of the quadric X .

1.2. The Dubrovin connection and flat sections. The next more sophisticated mirror theorem says that the Gauss-Manin connection defined using $\mathcal{W}_{\mathrm{can},q}$ recovers Dubrovin's connection on the free $\mathbb{C}[q^{\pm 1}]$ -module, $H^*(X, \mathbb{C}[q^{\pm 1}])$, which is defined using the small quantum cup product \star_q , see [Dub96, CK99]. The precise statement is formulated in Section 11. Namely we have a natural embedding of the Dubrovin connection into the Gauss-Manin system, Theorem 11.1, in which the coordinates p_i on the Gauss-Manin side match up with the Schubert classes σ_i on the Dubrovin connection side. The theorem in particular implies an integral formula for a global flat section of the connection, which is stated in Corollary 11.2. Much earlier Givental [Giv96] constructed a flat section for this connection without using mirror symmetry, as a power series with coefficients given by descendent 2-point Gromov-Witten invariants of X . Our Corollary 11.2 implies integral formulas for these invariants, via a comparison with Givental's formula. See the sequel paper [PRW16].

To illustrate the flat section formulas, let $X = Q_3$ and ω be a meromorphic 3-form on $\mathbb{C}\mathbb{P}^3$ with simple poles along the divisor

$$D = \{p_0 = 0\} \cup \{p_3 = 0\} \cup \{p_1 p_2 - p_0 p_3 = 0\}.$$

Suppose Γ is a real 3-dimensional cycle in $\mathbb{C}\mathbb{P}^3 \setminus D$ for integrating over; e.g. the compact 3-torus used in Section 1.3 below. Then our result implies that the $H^*(X, \mathbb{C})$ -valued function in q ,

$$S(q) = \left(\int_{\Gamma} e^{\mathcal{W}_{\mathrm{can},q} p_3} \omega \right) \sigma_0 + \left(\int_{\Gamma} e^{\mathcal{W}_{\mathrm{can},q} p_2} \omega \right) \sigma_1 + \left(\int_{\Gamma} e^{\mathcal{W}_{\mathrm{can},q} p_1} \omega \right) \sigma_2 + \left(\int_{\Gamma} e^{\mathcal{W}_{\mathrm{can},q}} \omega \right) \sigma_3,$$

satisfies the 'flat section' differential equation

$$q \frac{d}{dq} S = \sigma_1 \star_q S.$$

Here we have set $p_0 = 1$ and used p_1, p_2, p_3 as coordinates on $\mathbb{C}\mathbb{P}^3 \setminus D$, as in Section 1.1.

We can improve on Theorem 11.1 in the special case of $X = Q_3$ using a paper of Gorbounov and Smirnov. Namely in [GS13] Gorbounov and Smirnov construct their own ad hoc partial compactification of the original Givental-mirror [Giv96] for Q_{2m-1} . They prove with Sabbah and Nemethi that their superpotential $\mathcal{W}_{\mathrm{GS}}$ is cohomologically tame, which implies that the associated Gauss-Manin system reconstructs the Dubrovin connection without needing to pass to a submodule, see [Sab99]. We compare the Gorbounov-Smirnov partial compactification of Givental's mirror to the canonical LG model and show that in the case of Q_3 they are isomorphic. Together with the result of [GS13], we obtain in this case an isomorphism of the Gauss-Manin system of $(\check{X}_{\mathrm{can}}, \mathcal{W}_{\mathrm{can}})$ with the Dubrovin connection, see Theorem 11.1.

For dimension greater than three the canonical LG model is only birationally isomorphic to the Gorbounov-Smirnov mirror. However it still has the expected number of critical points. We therefore conjecture that the canonical and the Lie-theoretic superpotentials

are also cohomologically tame, and the isomorphism statement of Theorem 11.1 extends to all quadrics.

1.3. The Lie-theoretic superpotential and Lusztig coordinates. The flat sections $S(q)$ discussed above can also be written in terms of $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$, the Lie-theoretic LG model which was defined in [Rie08]. In this case the top degree coefficient of $S(q)$ (with respect to the grading on cohomology) takes on the form

$$(4) \quad \langle S(q), \sigma_0 \rangle = \int_{\Gamma_{\text{Lie}} \subset \check{X}_{\text{Lie}}} e^{\mathcal{W}_{\text{Lie}, q} \omega}.$$

Here \check{X}_{Lie} is a $(2m - 1)$ -dimensional affine subvariety of the full flag variety Sp_{2m}/B (it is a Schubert variety intersected with an opposite big cell), and ω is a particular holomorphic volume form on \check{X}_{Lie} . This formula (4) which we prove here was conjectured in [Rie08, Conjecture 8.1].

Crucial to our proofs is a new Laurent-polynomial LG-model. From the point of view of Lie theory, \check{X}_{can} can be described as the image inside $\check{X}_{\text{can}} = \mathbb{C}\mathbb{P}^{2m-1}$ of the intersection of two opposite Bruhat cells. It is therefore natural to restrict $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ to a ‘Lusztig torus’ \check{X}_{Lus} inside \check{X}_{can} . That is we consider the same torus which would be used by Lusztig to parametrise the totally positive part of \check{X}_{can} viewed as an Sp_{2m} -homogeneous space, in the theory of total positivity [Lus94]. After restriction of \mathcal{W}_{can} to this torus we obtain a very nice Laurent polynomial, which is reminiscent of the standard superpotential for projective space $\mathbb{C}\mathbb{P}^n$,

$$z_1 + \dots + z_n + q \frac{1}{z_1 \dots z_n},$$

and which can be used to compute the integral (4).

For example in the case of $X = Q_3$ we have a 3-dimensional torus with coordinates a, b, c , and

$$(5) \quad \mathcal{W}_{\text{Lus}, q} = a + b + c + q \frac{a + b}{abc}.$$

In the case of $X = Q_5$ we have a 5-dimensional torus with coordinates a_1, a_2, b_1, b_2, c , and

$$(6) \quad \mathcal{W}_{\text{Lus}, q} = a_1 + a_2 + b_1 + b_2 + c + q \frac{a_1 + b_1}{a_1 a_2 b_1 b_2 c}.$$

We may rewrite our integral formulas in terms of these coordinates. For example if $X = Q_3$ we obtain the easily computable integral

$$(7) \quad \langle S(q), \sigma_0 \rangle = \oint \oint \oint e^{(a+b+c+q \frac{a+b}{abc})} \frac{da}{a} \wedge \frac{db}{b} \wedge \frac{dc}{c}.$$

Finally, we note that analogous results in the parallel case of even quadrics are worked out in [PRW16]. For even-dimensional quadrics, the Langlands dual homogeneous space is another even-dimensional quadric, thus the canonical mirror $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ looks quite different from the one in the odd quadrics case. However when we restrict to the Lusztig torus in that setting, the formula is a straightforward generalisation of (5), (6), etc., to an even number of coordinates.

Acknowledgements : The second author thanks George Lusztig for his great PhD supervision and for introducing her to the theory of total positivity, which turns out to have so many beautiful connections. The second author also thanks Dale Peterson for his inspiring lectures on quantum cohomology.

2. OVERVIEW OF EARLIER LG MODELS

We begin by recalling various earlier constructions of mirror Landau-Ginzburg models which are relevant in our setting.

The Givental mirror. The earliest Landau-Ginzburg model construction which applies to odd quadrics is due to Givental [Giv96], who wrote down an LG model for any complex projective hypersurface $Y \hookrightarrow \mathbb{C}\mathbb{P}^N$. Givental's LG model is a regular function \mathcal{W}_{Giv} on a hypersurface inside an N -dimensional torus.

The odd quadric Q_{2m-1} is a hypersurface inside $\mathbb{C}\mathbb{P}^{2m}$, and Givental's mirror is a regular function on a particular hypersurface in a $2m$ -dimensional torus, namely

$$\left\{ (x_1, \dots, x_{2m}) \mid x_{2m} + \frac{q}{\prod_{i=1}^{2m} x_i} = 1 \right\}.$$

In a more symmetric formulation the Givental mirror of Q_{2m-1} is

$$\check{X}_{\text{Giv}} = \left\{ (\nu_1, \dots, \nu_{2m+1}) \in (\mathbb{C}^*)^{2m+1} \mid \prod_{i=1}^{2m+1} \nu_i = q, \nu_{2m} + \nu_{2m+1} = 1 \right\},$$

$$\mathcal{W}_{\text{Giv}} = \nu_1 + \dots + \nu_{2m-1}.$$

Additionally, \check{X}_{Giv} comes with a holomorphic volume form. But we do not include it here as it will not be used later.

The Przyjalkowski mirror. We use the notation $(\check{X}_{\text{Prz}}, \mathcal{W}_{\text{Prz}})$ for a Laurent polynomial mirror written down in [Prz07] which extends Givental's mirror from \check{X}_{Giv} to a $(2m-1)$ -dimensional torus containing it. In the case where Y is the smooth quadric Q_3 in \mathbb{P}^4 the Przyjalkowski mirror is given by

$$\check{X}_{\text{Prz}} := (\mathbb{C}^*)^3, \quad \mathcal{W}_{\text{Prz},q} := Y_1 + Y_2 + \frac{(Y_3 + q)^2}{Y_1 Y_2 Y_3}.$$

More generally for a quadric Q_{2m-1} the formula reads

$$\check{X}_{\text{Prz}} = (\mathbb{C}^*)^{2m-1}, \quad \mathcal{W}_{\text{Prz},q} = Y_1 + \dots + Y_{2m-2} + \frac{(Y_{2m-1} + q)^2}{Y_1 \dots Y_{2m-1}}.$$

\mathcal{W}_{Prz} is obtained from \mathcal{W}_{Giv} via the change of coordinates described in Section 10.

One issue with both $(\check{X}_{\text{Giv}}, \mathcal{W}_{\text{Giv}})$ and $(\check{X}_{\text{Prz}}, \mathcal{W}_{\text{Prz}})$ is that the superpotential does not in general have the expected number of critical points (at fixed generic value of q). Namely the expected number of critical points should be equal to $\dim H^*(Q_{2m-1}, \mathbb{C}) = 2m$. The analogous problem in the case of the even quadric Q_4 was already observed in [EHX97]. In this example [EHX97] constructed a partial compactification to solve this problem, albeit in an ad hoc fashion.

The following LG models, described in detail later in the paper, are partial compactifications of Givental's mirror $(\check{X}_{\text{Giv}}, \mathcal{W}_{\text{Giv}})$ which are known to have the correct number of critical points.

The Lie-theoretic mirror. The quadratic hypersurfaces Q_{2m-1} have large symmetry groups. Indeed Q_{2m-1} can be viewed as the Grassmannian of isotropic lines in \mathbb{C}^{2m} for a fixed non-degenerate quadratic form. In this way the quadric Q_{2m-1} is identified as a cominuscule homogeneous space for the group $G^\vee = \text{SO}_{2m+1}$ or its universal cover $\text{Spin}_{2m+1}(\mathbb{C})$.

For any projective homogeneous space of a complex algebraic group there is a Landau-Ginzburg model which was defined by the second author using a Lie-theoretic construction. Namely, in [Rie08] a conjectural LG model $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$ is constructed for any projective homogeneous space $X = G^\vee/P^\vee$ of a simple complex algebraic group G^\vee , as a regular function on an affine subvariety of the Langlands dual group G . This affine variety is generally larger than a torus. It is shown in [Rie08] that this Lie-theoretic LG model has the correct number of critical points. Namely its Jacobi ring is shown to recover the Peterson variety presentation [Pet97] of the quantum cohomology of X .

The Gorbounov-Smirnov mirror. For odd-dimensional quadrics Q_{2m-1} a recent paper [GS13] of Gorbounov and Smirnov directly constructs a partial compactification $(\check{X}_{\text{GS}}, \mathcal{W}_{\text{GS}})$ of the Givental mirror, without making use of [Rie08]. Moreover a version of mirror symmetry is proved, which identifies the initial data of the Frobenius manifold associated to their LG model with that constructed out of the quantum cohomology. In particular the Gauss-Manin connection associated to $(\check{X}_{\text{GS}}, \mathcal{W}_{\text{GS}})$ is shown to be isomorphic to the small Dubrovin connection.

3. PLAN OF THE PAPER

We begin in Section 4 by setting up notation and giving a careful definition of the Lie-theoretic LG model $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$. The domain is an open subvariety of a $(2m - 1)$ -dimensional Schubert variety \check{X}_{Lie} of the full flag variety of $\text{Sp}_{2m}(\mathbb{C})$.

Our first result is that \mathcal{W}_{Lie} restricted to a certain torus recovers the Laurent polynomial superpotential \mathcal{W}_{Lus} . This is proved in Section 5.

We introduce the canonical LG model in Section 6. It has domain \check{X}_{can} equal to the complement of an anti-canonical divisor in $\check{X}_{\text{can}} = \mathbb{C}\mathbb{P}^{2m}$, where $\mathbb{C}\mathbb{P}^{2m}$ is viewed as right homogeneous space for $\text{Sp}_{2m}(\mathbb{C})$. We then describe a birational map (depending on q)

$$(8) \quad \check{X}_{\text{Lie}} \dashrightarrow \check{X}_{\text{can}}.$$

This birational map sends the torus used above isomorphically to the Lusztig torus \check{X}_{Lus} in \check{X}_{can} . We express the Lusztig coordinates in terms of the homogeneous coordinates of \check{X}_{can} and show that the formula for \mathcal{W}_{Lus} transforms to the formula for the canonical superpotential \mathcal{W}_{can} . Therefore we see that $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$ and $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ are birationally isomorphic (to each other as well as to $(\check{X}_{\text{Lus}}, \mathcal{W}_{\text{Lus}})$).

Next, in Section 7 we show that the birational map (8) restricts to an isomorphism

$$\check{X}_{\text{Lie}} \rightarrow \check{X}_{\text{can}}.$$

It follows that the canonical and the Lie-theoretic LG models are isomorphic.

Then in Section 9 we deduce an isomorphism which identifies the Jacobi ring associated to $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ with the quantum cohomology ring of X . We show that under this isomorphism the homogeneous coordinates $\{p_i\}$ map to the Schubert classes $\{\sigma_i\}$, and the quadratic denominators of \mathcal{W}_{can} each map to either q or σ_{2m-1} .

In Section 10 we show that the following LG models are all birationally equivalent:

- the Givental mirror $(\check{X}_{\text{Giv}}, \mathcal{W}_{\text{Giv}})$ from [Giv96],
- the Przyjalkowski mirror $(\check{X}_{\text{Prz}}, \mathcal{W}_{\text{Prz}})$ from [Prz07],
- the Gorbounov-Smirnov mirror $(\check{X}_{\text{GS}}, \mathcal{W}_{\text{GS}})$ from [GS13],
- and the canonical mirror $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$, or equivalently $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$ from [Rie08].

In the case of $X = Q_3$ we also show that the Gorbounov-Smirnov mirror and the canonical mirror are isomorphic.

Section 11 is devoted to the Gauss-Manin system of $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$. Inside this Gauss-Manin system we identify a free $\mathbb{C}[q^{\pm 1}]$ -submodule with connection, which is then shown to be isomorphic to the Dubrovin connection on $H^*(X, \mathbb{C}[q^{\pm 1}])$. From this result we deduce integral formulas for flat sections of the Dubrovin connection.

Using results from [GS13] we deduce that the Dubrovin connection on $H^*(X, \mathbb{C}[q^{\pm 1}])$ is isomorphic to the Gauss-Manin system of $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ (and not just to a submodule) in the special case of the quadric Q_3 . In the final section we collect together and write out explicitly the formulas in the example of Q_3 .

The 1-parameter subgroups given by the x_i generate U_+ , and those given by the y_i generate U_- . We define the following additive characters on U_+ and U_- , respectively,

$$(9) \quad \begin{aligned} e_i^* : U_+ &\rightarrow \mathbb{C}, & e_i^*(x_j(m)) &= m\delta_{i,j}, \\ f_i^* : U_+ &\rightarrow \mathbb{C}, & f_i^*(y_j(m)) &= m\delta_{i,j}. \end{aligned}$$

Recall that we realised our quadric Q_{2m-1} as the homogeneous space $G^\vee/P_{\omega_1}^\vee$ for $SO_{2m+1}(\mathbb{C})$. We now consider the *dual parabolic subgroup* $P = P_{\omega_1}$ of $G = \mathrm{Sp}_{2m}(\mathbb{C})$ associated with the first fundamental weight. Explicitly it is the subgroup whose Lie algebra is generated by all of the Chevalley generators e_i together with f_2, \dots, f_m , leaving out f_1 . We let W_P denote the subgroup of the Weyl group W associated with P_{ω_1} , namely $W_P = \langle s_2, \dots, s_m \rangle$. We write w_P for the longest element in W_P , W^P for the set of minimal length coset representatives for W/W_P , and $w^P \in W^P$ for the minimal length coset representative of w_0 .

4.2. Definition of the Lie-theoretic LG model. In this section we follow [Rie08], adapting the results there to our special case. We first introduce the domain, $\check{X}_{\mathrm{Lie}} \subset G/B_-$, of the Lie-theoretic mirror, namely

$$\check{X}_{\mathrm{Lie}} := (B_+ \dot{w}_P B_- \cap B_- \dot{w}_0 B_-) / B_-.$$

Its closure in G/B_- is the Schubert variety

$$\overline{\check{X}}_{\mathrm{Lie}} := \overline{B_+ \dot{w}_P B_-} / B_-.$$

To write down the superpotential $\mathcal{W}_{\mathrm{Lie}} : \check{X}_{\mathrm{Lie}} \times \mathbb{C}_q^* \rightarrow \mathbb{C}$, we introduce a variety $Z \subset G$ which is a covering of $\check{X}_{\mathrm{Lie}} \times \mathbb{C}_q^*$. Let T^{W_P} be the W_P -fixed part of the maximal torus T . Since P is a maximal parabolic, this is a one-dimensional torus, and we have that

$$\begin{aligned} \alpha_1 : T^{W_P} &\rightarrow \mathbb{C}_q^*, \\ t &\mapsto \alpha_1(t) \end{aligned}$$

is a double cover. We set

$$(10) \quad Z := B_- \dot{w}_0 \cap U_+ T^{W_P} \dot{w}_P U_-,$$

and define a map

$$\begin{aligned} \pi_{\mathrm{Lie}} : \quad Z &\rightarrow \check{X}_{\mathrm{Lie}} \times \mathbb{C}_q^* \\ z = u_1 t \dot{w}_P \bar{u}_2 &\mapsto (z B_-, \alpha_1(t)). \end{aligned}$$

which is again a double cover. Note that if we were to quotient out Z by the action of the centre, $\{\pm 1\}$, of $\mathrm{Sp}_{2m}(\mathbb{C})$, then the map would be an isomorphism. This would be the convention taken in [Rie08].

We define a regular function on Z by

$$(11) \quad \begin{aligned} \mathcal{F} : \quad Z &\rightarrow \mathbb{C} \\ z = u_1 t \dot{w}_P \bar{u}_2 &\mapsto \sum e_i^*(u_1) + \sum f_i^*(\bar{u}_2). \end{aligned}$$

From [Rie08] it follows that \mathcal{F} is well-defined and descends to a regular function $\mathcal{W}_{\mathrm{Lie}} : \check{X}_{\mathrm{Lie}} \times \mathbb{C}_q^* \rightarrow \mathbb{C}$ such that the diagram

$$\begin{array}{ccc} Z & & \\ \pi_{\mathrm{Lie}} \downarrow & \searrow \mathcal{F} & \\ \check{X}_{\mathrm{Lie}} \times \mathbb{C}_q^* & \xrightarrow{\mathcal{W}_{\mathrm{Lie}}} & \mathbb{C} \end{array}$$

commutes. The corresponding map for fixed q , is denoted

$$\mathcal{W}_{\text{Lie},q} : \check{X}_{\text{Lie}} \rightarrow \mathbb{C}, \quad u_1 \dot{w}_P B_- \mapsto \mathcal{W}_{\text{Lie}}(u_1 \dot{w}_P B_-, q).$$

5. THE LAURENT POLYNOMIAL SUPERPOTENTIAL \mathcal{W}_{Lus} .

We continue using all of the notations from the previous section. We want to restrict the Lie-theoretic superpotential \mathcal{W}_{Lie} to a well-chosen torus to obtain a particularly nice Laurent polynomial. Instead of constructing the torus inside \check{X}_{Lie} , we will use Z , the double cover of $\check{X}_{\text{Lie}} \times \mathbb{C}^*$. Recall that $Z \subset B_- \dot{w}_0$ consists of those elements z which can be written in the form

$$z = u_1 t \dot{w}_P \bar{u}_2$$

for $u_1 \in U_+, t \in T^{W_P}$ and $\bar{u}_2 \in U_-$. However, the factors u_1 and \bar{u}_2 in this factorisation are not uniquely determined. We can make them uniquely determined for example by restricting the domain of \bar{u}_2 , which is what we will do now. Let

$$(12) \quad U_-^P := U_- \cap B_+ (\dot{w}^P)^{-1} B_+.$$

We have the following proposition.

Proposition 5.1. *For any $z \in Z$ there exists a unique $\bar{u}_2 = \bar{u}_2(z) \in U_-^P$ such that z has a factorisation of the form*

$$z = u_1 t \dot{w}_P \bar{u}_2.$$

We also write $t(z) = t$ if z is factored as above. The map $\theta : Z \rightarrow U_-^P \times T^{W_P}$ defined by

$$\theta : z \mapsto (\bar{u}_2(z), t(z))$$

is an isomorphism of affine varieties.

Remark. Note that for any $z \in Z$ with factorisation $z = u_1 t \dot{w}_P \bar{u}_2$, the factor $t \in T^{W_P}$ is uniquely determined and we may write $t = t(z)$. Under the conditions of the proposition, $\bar{u}_2 = \bar{u}_2(z) \in U_-^P$ is uniquely determined by z . This implies that the first factor, $u_1 \in U_+$, is also uniquely determined (if we keep the condition on \bar{u}_2). Namely $u_1(z) = z \bar{u}_2(z)^{-1} \dot{w}_P^{-1} t(z)^{-1}$.

The above proposition is proved using the twist map of Berenstein and Zelevinsky.

Theorem 5.2 ([BZ97, Theorem 1.2]). *Let G be a semisimple algebraic group. Let B_+, B_- be opposite Borel subgroups in G and U_+, U_- their unipotent radicals. Denote by $\dot{w} \in G$ a choice of representative for an element w of the Weyl group W as in Section 4.1. Consider $y \in U_- \cap B_+ \dot{w}^{-1} B_+$. There exists a unique $x \in U_+ \cap B_- \dot{w} B_-$ such that $U_+ \cap B_- \dot{w} y = \{x\}$, and the resulting map*

$$\tilde{\eta}_w : U_- \cap B_+ \dot{w}^{-1} B_+ \rightarrow U_+ \cap B_- \dot{w} B_-, y \mapsto x$$

is an isomorphism. In particular there exists an inverse isomorphism

$$\epsilon_w : U_+ \cap B_- \dot{w} B_- \rightarrow U_- \cap B_+ \dot{w}^{-1} B_+.$$

Proof of Proposition 5.1. The map $z \mapsto (t(z), \bar{u}_2(z))$ is constructed as the composition

$$Z \rightarrow (U_- \dot{w}_0 \cap B_+ \dot{w}_P U_-) \times T^{W_P} \rightarrow (U_+ \cap B_- \dot{w}^P B_-) \times T^{W_P} \rightarrow U_-^P \times T^{W_P},$$

where the rightmost map is defined using the isomorphism ϵ_{w^P} from Theorem 5.2, the middle map is defined using left multiplication by \dot{w}_0^{-1} , which is also an isomorphism, and the leftmost map is defined to be

$$z = b_- \dot{w}_0 \mapsto ([b_-]_0^{-1} b_- \dot{w}_0, t(z)).$$

Here $[b_-]_0$ denotes the torus part of the Borel group element $b_- \in TU_-$. This latter map is also an isomorphism with inverse $(b_+ \dot{w}_P u_-, t) \rightarrow t [b_+]_0^{-1} b_+ \dot{w}_P u_-$. \square

5.1. **The intermediate LG model.** We now have the commutative diagram

$$\begin{array}{ccc}
 \check{X}_{\text{Lie}} \times \mathbb{C}_q^* & \xleftarrow{\pi_{\text{Lie}}} & Z \\
 \searrow \mathcal{W}_{\text{Lie}} & & \downarrow \mathcal{F}_\theta \\
 & \mathbb{C} & \\
 \swarrow \mathcal{W}_\theta & & \downarrow \mathcal{F}_\theta \\
 U_-^P \times \mathbb{C}_q^* & \xleftarrow{\text{id} \times \alpha_1} & U_-^P \times T^{W^P}
 \end{array}$$

defining the maps $\mathcal{F}_\theta : U_-^P \times T^{W^P} \rightarrow \mathbb{C}$, and $\mathcal{W}_\theta : U_-^P \times \mathbb{C}_q^* \rightarrow \mathbb{C}$. Note that we may invert π_{Lie} and $\text{id} \times \alpha_1$, if we quotient out by the action of $\{\pm 1\}$ on Z and T^{W^P} . Therefore we may think of $(U_-^P, \mathcal{W}_\theta)$ as being an isomorphic LG model to $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$; there is an isomorphism $U_-^P \times \mathbb{C}_q^* \rightarrow \check{X}_{\text{Lie}} \times \mathbb{C}_q^*$ which is the identity on the second factor, and under which \mathcal{W}_{Lie} pulls back to \mathcal{W}_θ .

5.2. **The Laurent polynomial LG model.** We define an open dense torus inside U_-^P as follows. The Weyl group element $w^P \in W^P$ has the reduced expression

$$w^P = s_1 s_2 \dots s_{m-1} s_m s_{m-1} \dots s_2 s_1.$$

As a consequence of this and the Bruhat lemma, a generic element \bar{u}_2 in U_-^P can be written as a product of elements of 1-parameter subgroups as follows,

$$(13) \quad \bar{u}_2 = y_1(a_1) \dots y_{m-1}(a_{m-1}) y_m(c) y_{m-1}(b_{m-1}) \dots y_1(b_1),$$

where $a_i, c, b_j \neq 0$. Thus we define the torus $\mathcal{T} \subset U_-^P$ to be

$$\mathcal{T} := \{y_1(a_1) \dots y_{m-1}(a_{m-1}) y_m(c) y_{m-1}(b_{m-1}) \dots y_1(b_1) \mid a_i, c, b_i \in \mathbb{C}^*\}.$$

Before working out the restriction of the superpotential to this torus, we note that it is natural to think of \mathcal{T} as embedded in the homogeneous space

$$(14) \quad \check{X}_{\text{can}} := P \backslash Sp_{2m}(\mathbb{C}) \cong \mathbb{C}\mathbb{P}^{2m},$$

via $\bar{u}_2 \mapsto P_{w_1} \bar{u}_2$, setting the stage for the *canonical superpotential* to be introduced in the next section.

Thus we make the following definition.

Definition 5.3. We denote the image of the torus \mathcal{T} in \check{X}_{can} by \check{X}_{Lus} , and denote the coordinates on \check{X}_{Lus} in the same way as those on \mathcal{T} , by $a_i, c, b_j \neq 0$. Explicitly,

$$\check{X}_{\text{Lus}} := \{P y_1(a_1) \dots y_{m-1}(a_{m-1}) y_m(c) y_{m-1}(b_{m-1}) \dots y_1(b_1) \mid a_i, c, b_i \in \mathbb{C}^*\}.$$

The restriction of \mathcal{W}_θ to \mathcal{T} defines a map

$$\mathcal{W}_{\text{Lus}} : \check{X}_{\text{Lus}} \times \mathbb{C}_q^* \rightarrow \mathbb{C}.$$

Theorem 5.4. *In terms of the coordinates a_i, b_i, c on \check{X}_{Lus} ,*

$$(15) \quad \mathcal{W}_{\text{Lus}} = a_1 + \dots + a_{m-1} + c + b_{m-1} + \dots + b_1 + q \frac{a_1 + b_1}{a_1 \dots a_{m-1} c b_{m-1} \dots b_1}.$$

Proof of Theorem 5.4. Consider an element $\bar{u}_2 \in \mathcal{T} \subset U_-^P$ and choose a $t \in T^{W^P}$ such that $\alpha_1(t) = q$. By definition, \bar{u}_2 admits a factorisation

$$\bar{u}_2 = y_1(a_1) \dots y_{m-1}(a_{m-1}) y_m(c) y_{m-1}(b_{m-1}) \dots y_1(b_1).$$

Let $z := \theta^{-1}(\bar{u}_2, t)$, where θ is the isomorphism from Proposition 5.1. Then z can be written as $z = u_1 t \dot{w}_P \bar{u}_2$ for some unique $u_1 \in U_+$, and

$$\mathcal{W}_\theta(\bar{u}_2, q) = \mathcal{F}(u_1 t \dot{w}_P \bar{u}_2) = \sum_{i=1}^m e_i^*(u_1) + \sum_{i=1}^m f_i^*(\bar{u}_2).$$

The theorem now follows from the lemma below. \square

Lemma 5.5. *If u_1 and \bar{u}_2 are as above then we have the following identities*

$$(16) \quad f_i^*(\bar{u}_2) = \begin{cases} a_i + b_i & \text{if } 1 \leq i \leq m, \\ c & \text{otherwise.} \end{cases}$$

$$(17) \quad e_i^*(u_1) = \begin{cases} 0 & \text{if } 2 \leq i \leq m, \\ q \frac{a_1 + b_1}{a_1 \dots a_{m-1} c b_{m-1} \dots b_1} & \text{if } i = 1. \end{cases}$$

Proof. Equation (16) is obtained immediately from the definition of \bar{u}_2 . For Equation (17), let $v_{\omega_i}^-$ and $v_{\omega_i}^+$ denote a lowest, respectively highest, weight vector in V_{ω_i} and notice that

$$\begin{aligned} e_i^*(u_1) &= \frac{\langle u_1^{-1} \cdot v_{\omega_i}^-, e_i \cdot v_{\omega_i}^- \rangle}{\langle u_1^{-1} \cdot v_{\omega_i}^-, v_{\omega_i}^- \rangle} \\ &= \frac{\langle t \dot{w}_P \bar{u}_2 \cdot v_{\omega_i}^+, e_i \cdot v_{\omega_i}^- \rangle}{\langle t \dot{w}_P \bar{u}_2 \cdot v_{\omega_i}^+, v_{\omega_i}^- \rangle}. \end{aligned}$$

Assume $2 \leq i \leq m$. Then $e_i^*(u_1) = 0$ if and only if $\langle \bar{u}_2 \cdot v_{\omega_i}^+, \dot{w}_P^{-1} e_i \cdot v_{\omega_i}^- \rangle = 0$. Now the vector $w_P^{-1} e_i \cdot v_{\omega_i}^-$ is in the μ -weight space of the i -th fundamental representation, where $\mu = w_P^{-1} s_i(-\omega_i)$. Moreover, $\bar{u}_2 \in B_+(\dot{w}^P)^{-1} B_+$, hence $\bar{u}_2 \cdot v_{\omega_i}^+$ can have non-zero components only down to the weight space of weight $(w^P)^{-1}(\omega_i) = w_P^{-1}(-\omega_i)$. Since $\ell(w_P^{-1} s_i) > \ell(w_P^{-1})$ for $2 \leq i \leq m$, this is higher than μ , which proves that $e_i^*(u_1) = 0$.

Now assume $i = 1$. We have

$$\begin{aligned} e_1^*(u_1) &= \frac{\langle t \dot{w}_P \bar{u}_2 \cdot v_{\omega_1}^+, e_1 \cdot v_{\omega_1}^- \rangle}{\langle t \dot{w}_P \bar{u}_2 \cdot v_{\omega_1}^+, v_{\omega_1}^- \rangle} \\ &= (\omega_1 + \alpha_1 - \omega_1)(t) \frac{\langle \bar{u}_2 \cdot v_{\omega_1}^+, \dot{w}_P^{-1} e_1 \cdot v_{\omega_1}^- \rangle}{\langle \bar{u}_2 \cdot v_{\omega_1}^+, \dot{w}_P v_{\omega_1}^- \rangle} \\ &= q \frac{\langle \bar{u}_2 \cdot v_{\omega_1}^+, \dot{w}_P^{-1} e_1 \cdot v_{\omega_1}^- \rangle}{\langle \bar{u}_2 \cdot v_{\omega_1}^+, v_{\omega_1}^- \rangle}. \end{aligned}$$

First look at the denominator. The only way to go from the highest weight vector $v_{\omega_1}^+$ of the first fundamental representation to the lowest $v_{\omega_1}^-$ is to apply $g \in B_+ w B_+$ for $w \geq (w^P)^{-1}$. Since $\bar{u}_2 \in B_+(\dot{w}^P)^{-1} B_+$, it follows that we need to use all factors of \bar{u}_2 , and normalising $v_{\omega_1}^-$ appropriately, we get

$$\langle \bar{u}_2 \cdot v_{\omega_1}^+, v_{\omega_1}^- \rangle = a_1 \dots a_{m-1} c b_{m-1} \dots b_1.$$

Finally, we look at the numerator $\langle \bar{u}_2 \cdot v_{\omega_1}^+, \dot{w}_P^{-1} e_1 \cdot v_{\omega_1}^- \rangle$. Let ϵ_i denote the weight of the basis vector $v_i \in V_{\omega_1}$ when $1 \leq i \leq m$. The vector $\dot{w}_P^{-1} e_1 \cdot v_{\omega_1}^-$ has weight

$$\mu' = \dot{w}_P^{-1} s_1(-\omega_1) = \dot{w}_P^{-1}(-\epsilon_2) = \epsilon_2.$$

Indeed, $\dot{w}_P^{-1} e_1 \cdot v_{\omega_1}^- = v_2$. From the definition of \bar{u}_2 , it follows that $\langle \bar{u}_2 \cdot v_{\omega_1}^+, v_2 \rangle = a_1 + b_1$, which concludes the proof of the lemma and of the theorem. \square

In this section we have re-expressed the LG model $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$ in terms of a regular function on a subvariety of U_- , namely we introduced the intermediate LG model $(U_-^P, \mathcal{W}_\theta)$. Then we restricted to a natural choice of torus inside U_-^P to find a simple Laurent polynomial expression, leading us to $(\check{X}_{\text{Lus}}, \mathcal{W}_{\text{Lus}})$.

We are now ready to introduce the canonical mirror.

6. CONSTRUCTION OF THE CANONICAL LG MODEL $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$.

We now construct the canonical LG model $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ and state our main comparison theorem.

6.1. \check{X}_{can} and its affine subvariety \check{X}_{can} . Recall the definition of \check{X}_{can} from (14), as the right homogeneous space for $\text{Sp}_{2m}(\mathbb{C})$,

$$\check{X}_{\text{can}} = P_{\omega_1} \backslash \text{Sp}_{2m}(\mathbb{C}).$$

If $V = \mathbb{C}^{2m}$ is the defining representation of $\text{Sp}_{2m}(\mathbb{C})$ as in Section 4.1, then \check{X}_{can} may equivalently be described as $\mathbb{P}(V^*)$, viewed as an orbit of $\text{Sp}_{2m}(\mathbb{C})$ acting from the right.

Remark. We note that on V^* we have both the action from the right (matrix multiplication from the right on the vector space of row vectors), and the action from the left (dual representation of V). Namely these are related by $g \cdot v^* = v^* \cdot g^{-1}$ for $v^* \in V^*$.

We want to choose fixed coordinates on \check{X}_{can} . Let $v^0 = (0, \dots, 0, 1)$ in V^* , so that the line $\langle v^0 \rangle_{\mathbb{C}} \in \mathbb{P}(V^*)$ has stabiliser $P = P_{\omega_1}$. We let $w_{(k)} \in W$ be defined by

$$w_{(k)} = \begin{cases} s_1 s_2 \dots s_k & \text{if } k \leq m, \\ s_1 s_2 \dots s_{m-1} s_m s_{m-1} \dots s_{2m-k} & \text{if } m+1 \leq k \leq 2m-1. \end{cases}$$

This defines a total ordering, $w_{(0)} = e < w_{(1)} < \dots < w_{(2m-1)}$, on the minimal length coset representatives for $W_P \backslash W$. It gives rise to a basis $\{v^0, \dots, v^{2m-1}\}$ of V^* where $v^k := v^0 \cdot w_{(k)}$. Explicitly,

$$(18) \quad v^0 = (0, \dots, 0, 1), v^1 = (0, \dots, 1, 0), \dots, v^{2m-1} = (1, 0, \dots, 0).$$

We can now introduce notation for the homogeneous coordinates of an element $x \in \check{X}_{\text{can}}$, described as a coset $x = Pg$, by using the identification $\check{X}_{\text{can}} = \mathbb{P}(V^*)$ and the basis (18).

Definition 6.1. For $0 \leq k \leq 2m-1$ and $g \in \text{Sp}_{2m}(\mathbb{C})$ define the homogeneous coordinates $p_k(g)$ for the coset $Pg \in \check{X}_{\text{can}}$ by

$$p_k(g) = \langle v^0 \cdot g, v^k \rangle.$$

Here the angle brackets refer to the coefficient with respect to the basis $\{v^0, \dots, v^{2m-1}\}$.

Applying this definition, the homogeneous coordinates for a coset $Pg \in \check{X}_{\text{can}}$ are just given by the bottom row entries of the matrix g read from right to left. We note that, if as before we write g as $g = u_1 t \dot{w}_P \bar{u}_2$, then

$$(p_0(g) : \dots : p_{2m-1}(g)) = (p_0(\bar{u}_2) : \dots : p_{2m-1}(\bar{u}_2)),$$

since $Pg = P\bar{u}_2$. Changing the coset representative only rescales all of the homogeneous coordinates by a common factor.

Finally, note that the basis elements are of the form $v^k = v^0 \cdot w_{(k)}$ and these homogeneous coordinates can also be interpreted as generalised minors.

Definition 6.2. We define an affine subvariety of \check{X}_{can} by

$$(19) \quad \check{X}_{\text{can}} := \check{X}_{\text{can}} \setminus D,$$

where $D := D_0 \cup D_1 \cup \dots \cup D_{m-1} \cup D_m$, the divisors D_i being given by

$$\begin{aligned} D_0 &:= \{p_0 = 0\}, \\ D_\ell &:= \left\{ p_\ell p_{2m-1-\ell} - p_{\ell-1} p_{2m-\ell} + \dots + (-1)^\ell p_0 p_{2m-1} = 0 \right\} \text{ for } 1 \leq \ell \leq m-1, \\ D_m &:= \{p_{2m-1} = 0\}. \end{aligned}$$

Since the index of projective space $\check{X}_{\text{can}} = \mathbb{C}\mathbb{P}^{2m-1}$ is $2m$ which is also the degree of the divisor D , it follows that D is an anticanonical divisor. We may also use the notation δ_ℓ for the quadratic expression

$$\delta_\ell := p_\ell p_{2m-1-\ell} - p_{\ell-1} p_{2m-\ell} + \dots + (-1)^\ell p_0 p_{2m-1},$$

where $0 \leq \ell \leq m-1$.

6.2. The superpotential \mathcal{W}_{can} and the isomorphism theorem. In the previous section we defined \check{X}_{can} and \check{X}_{can} and the homogeneous coordinates p_i . To define the ‘canonical’ LG model $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ it remains to define the superpotential \mathcal{W}_{can} .

Definition 6.3. \mathcal{W}_{can} is defined to be the regular map $\mathcal{W}_{\text{can}} : \check{X}_{\text{can}} \times \mathbb{C}_q^* \rightarrow \mathbb{C}$ expressed in terms of the homogeneous coordinates of \check{X}_{can} by

$$(20) \quad \mathcal{W}_{\text{can}} = \frac{p_1}{p_0} + \sum_{\ell=1}^{m-1} \frac{p_{\ell+1} p_{2m-1-\ell}}{p_\ell p_{2m-1-\ell} - p_{\ell-1} p_{2m-\ell} + \dots + (-1)^\ell p_0 p_{2m-1}} + q \frac{p_1}{p_{2m-1}}.$$

If q is fixed we use the notation $\mathcal{W}_{\text{can},q}$. We refer to the pair $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ as the ‘canonical’ LG model.

The next two sections will be devoted to proving the following comparison theorem between $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$ and $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$. Recall the definition of the subvarieties $Z \subset B_- \dot{u}_0$ and $U_P^- \subset U_-$ from (10) and (12), respectively.

By Proposition 5.1 and Section 5.1 we have an isomorphism $\check{X}_{\text{Lie}} \times \mathbb{C}_q^* \xrightarrow{\sim} U_P^- \times \mathbb{C}_q^*$. We use it to define a ‘comparison’ map

$$(21) \quad \Psi : \check{X}_{\text{Lie}} \times \mathbb{C}_q^* \xrightarrow{\sim} U_P^- \times \mathbb{C}_q^* \xrightarrow{\pi_{\text{can}}} \check{X}_{\text{can}} \times \mathbb{C}_q^*$$

where the right hand side map is simply defined by $(\bar{u}_2, q) \mapsto (P\bar{u}_2, q)$. All in all the map Ψ is given by

$$(22) \quad (gB_-, q) = (u_1 t \dot{w}_P \bar{u}_2 B_-, q) \mapsto (\bar{u}_2, q) \mapsto (P\bar{u}_2, q),$$

where $z = u_1 t \dot{w}_P \bar{u}_2$ is the uniquely (up to ± 1) determined element of Z for which $\alpha_1(t) = q$ and $zB_- = gB_-$.

Theorem 6.4. *The map Ψ from (21) has image $\check{X}_{\text{can}} \times \mathbb{C}_q^*$ and defines an isomorphism between $\check{X}_{\text{Lie}} \times \mathbb{C}_q^*$ and $\check{X}_{\text{can}} \times \mathbb{C}_q^*$ such that the following diagram commutes*

$$\begin{array}{ccc} \check{X}_{\text{Lie}} \times \mathbb{C}_q^* & \xrightarrow{\Psi} & \check{X}_{\text{can}} \times \mathbb{C}_q^* \\ & \searrow \mathcal{W}_{\text{Lie}} & \swarrow \mathcal{W}_{\text{can}} \\ & & \mathbb{C} \end{array}$$

In other words, we have an isomorphism $\check{X}_{\text{Lie}} \times \mathbb{C}_q^ \rightarrow \check{X}_{\text{can}} \times \mathbb{C}_q^*$ which is the identity on the second factor, and under which \mathcal{W}_{can} pulls back to \mathcal{W}_{Lie} .*

7. THE SUPERPOTENTIAL \mathcal{W}_{can} AND THE LAURENT POLYNOMIAL \mathcal{W}_{Lus}

In this section we prove a birational version of Theorem 6.4. To do this we make use of the Laurent polynomial LG model $(\check{X}_{\text{Lus}}, \mathcal{W}_{\text{Lus}})$ which is birational to $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$ by construction.

Proposition 7.1. *Consider the canonical superpotential \mathcal{W}_{can} as the rational function on $\check{X}_{\text{can}} \times \mathbb{C}_q^*$ given by,*

$$\mathcal{W}_{\text{can},q} = \frac{p_1}{p_0} + \sum_{\ell=1}^{m-1} \frac{p_{\ell+1}p_{2m-1-\ell}}{p_{\ell}p_{2m-1-\ell} - p_{\ell-1}p_{2m-\ell} + \cdots + (-1)^{\ell}p_0p_{2m-1}} + q \frac{p_1}{p_{2m-1}}.$$

The restriction of \mathcal{W}_{can} to the Lusztig torus $\check{X}_{\text{Lus}} \subset \check{X}_{\text{can}}$ is regular and agrees with \mathcal{W}_{Lus} , see Definition 5.3 and Theorem 5.4.

Note that the Laurent polynomial superpotential \mathcal{W}_{Lus} was in fact obtained from \mathcal{W}_{Lie} by restriction of \mathcal{W}_{Lie} to the torus $\Psi^{-1}(\check{X}_{\text{Lus}} \times \mathbb{C}_q^*)$, compare Section 5. Thus Proposition 7.1 has the following Corollary.

Corollary 7.2. *We have a commutative diagram of (rational) maps*

$$\begin{array}{ccccc} \check{X}_{\text{Lie}} \times \mathbb{C}_q^* & \longrightarrow & \check{X}_{\text{Lus}} \times \mathbb{C}_q^* & \hookrightarrow & \check{X}_{\text{can}} \times \mathbb{C}_q^* \\ & \searrow \mathcal{W}_{\text{Lie}} & \downarrow \mathcal{W}_{\text{Lus}} & \swarrow \mathcal{W}_{\text{can}} & \\ & & \mathbb{C} & & \end{array}$$

□

Proof of Proposition 7.1. Consider an element $x = P\bar{u}_2 \in \check{X}_{\text{Lus}}$. By definition of \check{X}_{Lus} , \bar{u}_2 admits a factorisation

$$\bar{u}_2 = y_1(a_1) \cdots y_{m-1}(a_{m-1})y_m(c)y_{m-1}(b_{m-1}) \cdots y_1(b_1).$$

Recall from Definition 6.1 that the homogeneous coordinates are given by

$$p_k(x) = \langle v^0 \cdot \bar{u}_2, v^k \rangle = \langle v^0 \cdot \bar{u}_2, v^0 \cdot \dot{w}_{(k)} \rangle.$$

Using the factorisation of \bar{u}_2 the following result is immediate.

Lemma 7.3. *Let $0 \leq k \leq 2m-1$ be an integer. Then if \bar{u}_2 possesses a factorisation of the form (13) we have*

$$p_k(\bar{u}_2) = \begin{cases} 1 & \text{if } k = 0, \\ a_1 \cdots a_{k-1}(a_k + b_k) & \text{if } 1 \leq k \leq m-1, \\ a_1 \cdots a_{m-1}cb_{m-1} \cdots b_{2m-k} & \text{otherwise.} \end{cases} \quad \square$$

By an easy computation it follows that

$$\mathcal{W}_{\text{can},q}(x) = \frac{p_1}{p_0} + \sum_{\ell=1}^{m-1} \frac{p_{\ell+1}p_{2m-1-\ell}}{p_{\ell}p_{2m-1-\ell} - p_{\ell-1}p_{2m-\ell} + \cdots + (-1)^{\ell}p_0p_{2m-1}} + q \frac{p_1}{p_{2m-1}} = \mathcal{W}_{\text{Lus},q}(x).$$

Since \check{X}_{Lus} is open dense in \check{X}_{can} this completes the proof of the proposition. □

In the next section we will study the locus \check{X}_{can} where $\mathcal{W}_{\text{can},q}$ is regular and prove that Ψ is an isomorphism $\check{X}_{\text{Lie}} \times \mathbb{C}_q^* \rightarrow \check{X}_{\text{can}} \times \mathbb{C}_q^*$. This will complete the proof of Theorem 6.4.

8. THE CANONICAL MIRROR VARIETY

Recall that an element in $\check{X}_{\text{can}} = P \backslash SL_{2m}(\mathbb{C})$ has projective coordinates $(p_0 : p_1 : \dots : p_{2m-1})$ which were introduced in Definition 6.1. Then the affine subvariety \check{X}_{can} was defined as the complement of a particular anticanonical divisor $D \subset \check{X}_{\text{can}}$ expressed in these coordinates. Namely

$$D := D_0 + D_1 + \dots + D_{m-1} + D_m,$$

with $D_0 = \{p_0 = 0\}$, $D_m = \{p_{2m-1} = 0\}$ and each remaining D_ℓ of the form $D_\ell = \{\delta_\ell = 0\}$ for a particular quadratic polynomial δ_ℓ , see Definition 6.2. We let $\delta_0 = p_0 p_{2m-1}$.

The goal of this section is to prove the following proposition.

Proposition 8.1. *The map from (21) defines an isomorphism $\Psi : \check{X}_{\text{Lie}} \times \mathbb{C}_q^* \rightarrow \check{X}_{\text{can}} \times \mathbb{C}_q^*$.*

Proof. By construction, the canonical superpotential $\mathcal{W}_{\text{can},q}$ is regular on \check{X}_{can} and this is the whole regular locus. Now from Corollary 7.2 it follows that Ψ maps $\check{X}_{\text{Lie}} \times \mathbb{C}_q^*$ into the regular locus of \mathcal{W}_{can} . Therefore Ψ must have its image in $\check{X}_{\text{can}} \times \mathbb{C}_q^*$. We will now prove the result by constructing an inverse map $\Psi^{-1} : \check{X}_{\text{can}} \times \mathbb{C}_q^* \rightarrow \check{X}_{\text{Lie}} \times \mathbb{C}_q^*$.

It suffices to set $q = 1$ and construct a map $\Psi_{q=1}^{-1} : \check{X}_{\text{can}} \rightarrow \check{X}_{\text{Lie}}$ for which

$$\Psi_{q=1}^{-1}(P\bar{u}_2) = zB_-$$

where $z = u_1 \dot{w}_P \bar{u}_2 \in Z$. Then Ψ^{-1} is constructed out of $\Psi_{q=1}^{-1}$ by setting

$$\Psi^{-1} : (P\bar{u}_2, q) \mapsto t(q)\Psi_{q=1}^{-1}(P\bar{u}_2) = t(q)zB_-,$$

where for any q we let $t(q) \in T^{WP}$ be a torus element for which $\alpha_1(t(q)) = q$.

To construct $\Psi_{q=1}^{-1}$ we consider the morphism

$$\Phi : \check{X}_{\text{can}} \rightarrow B_- \dot{w}_0, \quad (p_0 : p_1 : \dots : p_{2m-1}) \rightarrow \Phi(p_0 : p_1 : \dots : p_{2m-1}),$$

where $\Phi(p_0 : p_1 : \dots : p_{2m-1})$ is the matrix of the linear map which to the basis element v_j of $V \cong \mathbb{C}^{2m}$ associates

$$\begin{cases} p_{2m-1}v_{2m} & \text{if } j = 1, \\ (-1)^{j-1} \frac{\delta_{j-1}}{\delta_{j-2}} v_{2m+1-j} + p_{2m-j} \left(\sum_{\ell=1}^{j-2} (-1)^\ell \frac{p_\ell}{\delta_{\ell-1}} v_{2m-\ell} + v_{2m} \right) & \text{if } 2 \leq j \leq m, \\ (-1)^{j-1} \frac{\delta_{2m-1-j}}{\delta_{2m-j}} v_{2m+1-j} \\ + p_{2m-j} \left(\sum_{\ell=m+1}^{j-1} (-1)^{\ell-1} \frac{p_{\ell-1}}{\delta_{2m-\ell}} v_{2m+1-\ell} + \sum_{\ell=1}^{m-1} (-1)^\ell \frac{p_\ell}{\delta_{\ell-1}} v_{2m-\ell} + v_{2m} \right) & \text{if } m+1 \leq j \leq 2m-1, \\ \frac{-1}{\delta_0} v_1 + \sum_{\ell=1}^{m-1} (-1)^{\ell+1} \frac{p_{2m-1-\ell}}{\delta_\ell} v_{\ell+1} + \sum_{\ell=1}^{m-1} (-1)^\ell \frac{p_\ell}{\delta_{\ell-1}} v_{2m-\ell} + v_{2m} & \text{if } j = 2m. \end{cases}$$

Here $p_0 = 1$. Let $\Omega \subset \check{X}_{\text{can}}$ be the open dense subset where the coordinates $p_m, p_{m+1}, \dots, p_{2m-2}$ do not vanish. This subset is isomorphic to the Lusztig torus \check{X}_{Lus} via the change of coordinates

$$a_i = \frac{p_{2m-1}\delta_i}{p_{2m-1-i}\delta_{i-1}}, \quad b_i = \frac{p_{2m-1}}{p_{2m-1-i}} \text{ for all } 1 \leq i \leq m-1, \quad c = \frac{p_m^2}{\delta_{m-1}}.$$

Lemma 8.2. *For any element $(p_0 : p_1 : \dots : p_{2m-1}) \in \Omega$, $\Phi(p_0 : p_1 : \dots : p_{2m-1})$ factorizes as $u_1 \dot{w}_P \bar{u}_2$, where*

$$\bar{u}_2 = y_1(a_1) \dots y_{m-1}(a_{m-1}) y_m(c) y_{m-1}(b_{m-1}) \dots y_1(b_1)$$

and u_1 is given by the matrix

$$\begin{pmatrix} 1 & \frac{-(a_1+b_1)}{a_1 \dots a_{m-1} c b_{m-1} \dots b_1} & \dots & \frac{-(a_{m-1}+b_{m-1})}{a_1 \dots a_{m-1} c b_{m-1}} & \frac{-1}{a_1 \dots a_{m-1}} & \dots & \frac{-1}{a_1} & \frac{-1}{a_1 \dots a_{m-1} c b_{m-1} \dots b_1} \\ & 1 & & & & & \frac{1}{a_1} & \\ & & \ddots & & & & \vdots & \\ & & & 1 & & & \frac{(-1)^m}{a_1 \dots a_{m-1}} & \\ & & & & 1 & & (-1)^{m-1} \frac{a_{m-1}+b_{m-1}}{a_1 \dots a_{m-1} c b_{m-1}} & \\ & & & & & \ddots & \vdots & \\ & & & & & & 1 & \frac{-(a_1+b_1)}{a_1 \dots a_{m-1} c b_{m-1} \dots b_1} \\ & & & & & & & 1 \end{pmatrix}$$

Proof of Lemma 8.2. Using the definition of the y_i , it is easy to check that $\bar{u}_2 \cdot v_j$ is equal to

$$\begin{cases} v_j + \sum_{\ell=0}^{m-1-j} (a_{j+\ell} + b_{j+\ell}) \left(\prod_{r=0}^{\ell-1} b_r \right) v_{j+\ell+1} + \sum_{\ell=0}^{m-1} \left(\prod_{r=1}^{\ell} a_{m-r} \right) c b_{m-1} \dots b_j v_{m+1+\ell} & \text{if } 1 \leq j \leq m-1, \\ v_m + \sum_{k=0}^{m-1} a_{m-k} \dots a_{m-1} c v_{m+1+k} & \text{if } j = m, \\ v_j + (a_{2m-j} + b_{2m-j}) \sum_{\ell=0}^{2m-1-j} \left(\prod_{r=1}^{\ell-1} a_{2m-1-j-r} \right) v_{j+1+\ell} & \text{if } m+1 \leq j \leq 2m. \end{cases}$$

Now a straightforward computation shows that $\Phi(p_0 : p_1 : \dots : p_{2m-1}) = u_1 \dot{w}_P \bar{u}_2$. \square

Lemma 8.2 shows that $\Phi(\Omega)$ is contained in $Z_{q=1} := B_- \dot{w}_0 \cap U_+ \dot{w}_P U_-$. In the next lemma we prove that the entire image of Φ is contained in $Z_{q=1}$.

Lemma 8.3. *The image $\Phi(p_0 : p_1 : \dots : p_{2m-1})$ of any element $(p_0 : p_1 : \dots : p_{2m-1}) \in \check{X}_{\text{can}}$ lies in $Z_{q=1}$.*

Proof of Lemma 8.3. Since Ω is open dense in \check{X}_{can} we have that $\Phi(\check{X}_{\text{can}}) \subset B_- \dot{w}_0 \cap \overline{U_+ \dot{w}_P U_-}$. Suppose indirectly there exists $x = (p_0 : p_1 : \dots : p_{2m-1}) \in \check{X}_{\text{can}}$ such that $\Phi(x) \notin U_+ \dot{w}_P U_-$. Then from the Bruhat decomposition, we get $\Phi(x) \dot{w}_0^{-1} \in U_+ \dot{w} U_+$ with $w < w_P w_0$. It follows that we must have

$$(23) \quad 0 = \langle \Phi(x) \dot{w}_0^{-1} \cdot v_{\omega_1}^+, v_{\omega_1}^- \rangle = \langle \Phi(x) \cdot v_{\omega_1}^-, v_{\omega_1}^- \rangle$$

in the representation V_{ω_1} of G , where $v_{\omega_1}^+ = v_1$ and $v_{\omega_1}^- = v_{2m}$, compare Section 4.1. However we have seen that the lower right hand corner of the matrix representing $\Phi(x)$ is equal to 1, contradicting (23). Thus we must have $\Phi(x) \in Z_{q=1}$. \square

We have thus shown that the map Φ is a regular morphism $\check{X}_{\text{can}} \rightarrow Z_{q=1}$. Moreover by construction the composition with the coset map, $\pi : Z_{q=1} \rightarrow \check{X}_{\text{Lie}}, z \mapsto z B_-$, gives the map

$$\Psi_{q=1}^{-1} : \check{X}_{\text{can}} \rightarrow \check{X}_{\text{Lie}}$$

we were looking for. This concludes the proof of Proposition 8.1. \square

We have now proved our main comparison result, Theorem 6.4.

9. THE JACOBI RING PRESENTATION OF $\mathrm{qH}^*(X, \mathbb{C})[q^{-1}]$

The main result of [Rie08] was to show that there is an isomorphism between the quantum cohomology of X and the Jacobi ring of $(\check{X}_{\mathrm{Lie}}, \mathcal{W}_{\mathrm{Lie}})$ (for the case of the quadric X as well as for a general homogeneous space $X = G^\vee/P^\vee$). This result made use of the remarkable Peterson presentation [Pet97] of $\mathrm{qH}^*(G^\vee/P^\vee)$, which identifies the quantum cohomology ring with the coordinate ring of an associated affine variety $\mathcal{Y}_P \subset G/B$. The variety \mathcal{Y}_P is called the ‘Peterson variety’ associated to the parabolic subgroup P .

Now that we have proved that the canonical LG model $(\check{X}_{\mathrm{can}}, \mathcal{W}_{\mathrm{can}})$ is isomorphic to the Lie-theoretic LG model $(\check{X}_{\mathrm{Lie}}, \mathcal{W}_{\mathrm{Lie}})$, Theorem 6.4, we may apply the result from [Rie08] to deduce that there is an isomorphism between the quantum cohomology of $X = Q_{2m-1}$ and the Jacobi ring of $(\check{X}_{\mathrm{can}}, \mathcal{W}_{\mathrm{can}})$,

$$(24) \quad \mathbb{C}[\check{X}_{\mathrm{can}} \times \mathbb{C}_q^*] / \left(\frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_1}, \dots, \frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_{2m-1}} \right) \cong \mathrm{qH}^*(X, \mathbb{C})[q^{-1}].$$

In this section we prove that the isomorphism sends the homogeneous coordinate p_i to the Schubert class σ_i .

Theorem 9.1. *The isomorphism*

$$(25) \quad \mathbb{C}[\check{X}_{\mathrm{can}} \times \mathbb{C}_q^*] / (\partial \mathcal{W}_{\mathrm{can},q}) \rightarrow \mathrm{qH}^*(X)[q^{-1}]$$

defined using [Rie08], identifies the coordinate q with the quantum parameter q and sends p_i to the Schubert class $\sigma_i \in H^{2i}(X, \mathbb{C})$.

We first prove the following lemma. Note that when we write p_0 we mean 1, since this homogeneous coordinate has been fixed as $p_0 = 1$.

Lemma 9.2. *In the Jacobi ring $\mathbb{C}[\check{X}_{\mathrm{can}} \times \mathbb{C}_q^*] / (\partial \mathcal{W}_{\mathrm{can},q})$ of $\mathcal{W}_{\mathrm{can}}$, the element $\delta_0 = p_0 p_{2m-1}$ has the property*

$$(26) \quad \delta_0^2 = p_{2m-1}^2 = q^2, \quad \text{and} \quad \delta_0 p_i = q p_i, \quad \text{for } i = 1, \dots, 2m-1.$$

For the elements δ_ℓ with $1 \leq \ell \leq m-1$ we have

$$(27) \quad \delta_\ell = p_\ell p_{2m-1-\ell} - p_{\ell-1} p_{2m-\ell} + \dots + (-1)^\ell p_0 p_{2m-1} = \begin{cases} q & \text{if } \ell \text{ is odd} \\ p_{2m-1} & \text{if } \ell \text{ is even,} \end{cases}$$

in the Jacobi ring.

Proof. The equations (26) and (27) for p_i replaced by σ_i are a straightforward consequence of quantum Schubert calculus on the quadric (which can be deduced from the quantum Chevalley formula [FW04] in this case, see also (33)). It is not hard to check by a direct calculation that the relations

$$\frac{\partial \mathcal{W}_{\mathrm{can},q}}{\partial p_i} = 0$$

imply that the p_i in the Jacobi ring satisfy all the same relations as the σ_i do in the quantum cohomology ring.

Alternatively, particularly to prove (26), one can check that the coordinates of the critical points of $\mathcal{W}_{\mathrm{can},q}$ satisfy the equations. They are all non-degenerate; the quantum cohomology ring is semisimple. These critical points are worked out explicitly in [PRW16, Proposition 2.3]. \square

Proof of Theorem 9.1. While we already noted in the preceding proof that the p_i satisfy the relations of quantum Schubert calculus in the Jacobi ring of $\mathcal{W}_{\mathrm{can}}$, the statement of the theorem is about a specific isomorphism. It remains to check that this isomorphism does indeed send p_i to σ_i .

The isomorphism,

$$(28) \quad \mathrm{qH}^*(X, \mathbb{C})[q^{-1}] \longrightarrow \mathbb{C}[\check{X}_{\mathrm{Lie}} \times \mathbb{C}_q^*] / (\partial \mathcal{W}_{\mathrm{Lie},q}),$$

coming from [Rie08] and involving the Peterson presentation [Pet97] takes the following form. The Schubert class σ_k is associated to the element $w^{(k)} \in W^P$ defined by

$$w^{(k)} = \begin{cases} s_k s_{k-1} \cdots s_1 & \text{if } k \leq m, \\ s_{2m-k} \cdots s_{m-1} s_m s_{m-1} \cdots s_1 & \text{if } m+1 \leq k \leq 2m-1. \end{cases}$$

Note that $w^{(k)} = w_{(k)}^{-1}$, compare Section 6.1. The isomorphism (28) is explicitly given by $\sigma_k \mapsto f_k$ where f_k is the regular function

$$f_k(zB_-) := \frac{\langle z \cdot v_{\omega_1}^-, \dot{w}^{(k)} \cdot v_{\omega_1}^- \rangle}{\langle z \cdot v_{\omega_1}^-, v_{\omega_1}^- \rangle}.$$

Here $v_{\omega_1}^- = v_{2m}$ is the lowest weight vector of the representation $V = V_{\omega_1}$. We may assume that $z = u_1 t(q) \dot{w}_P \bar{u}_2$, as in Section 4.2. We need to show that

$$f_k(zB_-) = p_k(\bar{u}_2),$$

whenever $P\bar{u}_2$ is a critical point of $\mathcal{W}_{\text{can},q}$ with $q = \alpha_1(t)$. We use the notations from the proof of Proposition 8.1. Recall that there we have a map $\Phi : \check{X}_{\text{can}} \rightarrow B_- \dot{w}_0$ given explicitly in terms of the coordinates p_i , for which

$$t(q) \Phi(p_1(\bar{u}_2), \dots, p_{2m-1}(\bar{u}_2)) = z = u_1 t(q) \dot{w}_P \bar{u}_2 \in Z.$$

We can now work out $f_k(zB_-)$ by looking at the entries of the last column of the matrix $z = t(q) \Phi(p_1(\bar{u}_2), \dots, p_{2m-1}(\bar{u}_2))$. Namely we get

$$f_k(zB_-) = \begin{cases} 1, & \text{if } k = 0, \\ \frac{qp_k}{\delta_{k-1}}, & \text{if } 1 \leq k \leq m, \\ \frac{qp_k}{\delta_{2m-k-1}}, & \text{if } m+1 \leq k \leq 2m-1, \\ \frac{q^2}{\delta_0}, & \text{if } k = 2m-1. \end{cases}$$

Applying the relations (26) and (27) we get the identity $f_k(zB_-) = p_k(\bar{u}_2)$ as required. \square

It is interesting to note that by the isomorphism from Theorem 9.1 and the quantum Pieri rule, the summands of \mathcal{W}_{can} map to σ_1 or $2\sigma_1$, and \mathcal{W}_{can} maps to the anticanonical class of X in the quantum cohomology ring. Indeed for $k = 2, \dots, m-2$ it follows from the relations that the k -th summand of $\mathcal{W}_{\text{can}} = W_1 + W_2 + \dots + W_{m-1} + qW_m$ in the Jacobi ring simplifies to

$$W_k = \frac{p_{k+1} p_{2m-1-k}}{\delta_k} = 2p_1,$$

while $k = 1, m-1, m$ gives

$$W_1 = p_1, \quad W_{m-1} = \frac{p_m^2}{\delta_m} = p_1, \quad W_m = q \frac{p_1}{p_{2m-1}} = p_1.$$

In total we have $\mathcal{W}_{\text{can}} = (2m-1)p_1$ in $\mathbb{C}[\check{X}_{\text{can}} \times \mathbb{C}_q^*]/(\partial \mathcal{W}_{\text{can},q})$ and \mathcal{W}_{can} represents the anticanonical class $(2m-1)\sigma_1$ of $X = Q_{2m-1}$ via the isomorphism (25).

10. COMPARISON WITH OTHER LG MODELS

Let us now see how our canonical mirror $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ and the corresponding Laurent polynomial mirror $(\check{X}_{\text{Lus}}, \mathcal{W}_{\text{Lus}})$ compare with previous Landau-Ginzburg models for odd quadrics presented in Section 2. From Theorem 6.4 we already know that $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ is isomorphic to the Lie-theoretic mirror $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$, so it only remains to consider the Givental, Przyjalkowski mirrors and the partial compactification by Gorbounov and Smirnov.

Comparison with the Givental mirror and Przyjalkowski Laurent polynomial.

First recall the definition of the Givental mirror of Q_{2m-1} ,

$$\check{X}_{\text{Giv}} = \left\{ (\nu_1, \dots, \nu_{2m+1}) \in (\mathbb{C}^*)^{2m+1} \mid \prod_{i=1}^{2m+1} \nu_i = q, \nu_{2m} + \nu_{2m+1} = 1 \right\},$$

$$\mathcal{W}_{\text{Giv}} = \nu_1 + \dots + \nu_{2m-1}.$$

The Laurent polynomial ‘extension’ of this mirror, written down in [Prz07], is as follows

$$\check{X}_{\text{Prz}} = (\mathbb{C}^*)^{2m-1}, \quad \mathcal{W}_{\text{Prz}} = Y_1 + \dots + Y_{2m-2} + \frac{(Y_{2m-1} + q)^2}{Y_1 \dots Y_{2m-1}}.$$

It is obtained via the change of coordinates

$$Y_i = \begin{cases} \nu_{i+1} & \text{if } 1 \leq i \leq 2m-2, \\ q \frac{\nu_{2m}}{\nu_{2m+1}} & \text{if } i = 2m-1. \end{cases}$$

The torus \check{X}_{Prz} is slightly larger than \check{X}_{Giv} , as

$$\check{X}_{\text{Giv}} \cong \check{X}_{\text{Prz}} \setminus \{Y_{2m-1} + q = 0\}.$$

Our canonical mirror is a partial compactification of the Givental mirror, as shown by the following result, whose proof is immediate.

Proposition 10.1. *The change of coordinates (dependant on q)*

$$Y_i = \begin{cases} \frac{p_i}{p_{i-1}} & \text{if } 1 \leq i \leq m-1, \\ \frac{p_{2m-1-i} \delta_{2m-3-i}}{p_{2m-2-i} \delta_{2m-2-i}} & \text{if } m \leq i \leq 2m-3, \\ q \frac{p_1}{p_{2m-1}} & \text{if } i = 2m-2, \\ q \frac{\delta_{2m-2}}{\delta_{m-1}} & \text{if } i = 2m-1, \end{cases}$$

induces an isomorphism between \check{X}_{Prz} and the torus inside \check{X}_{can} where the Plücker coordinates p_i for $0 \leq i \leq m-1$ are all non-zero.

Thus we have two distinguished tori inside the canonical mirror variety \check{X}_{can} , namely the Lusztig torus, \check{X}_{Lus} , and the image of \check{X}_{Prz} under the map from Proposition 10.1. They are however distinct, since \check{X}_{Lus} is the torus inside \check{X}_{can} where the Plücker coordinates p_i for $m \leq i \leq 2m-1$ are all non-zero. In fact, \check{X}_{can} is an example of a cluster variety, see [GLS08a, Section 12], so it contains multiple tori. We will describe the cluster structure in detail in Section 11 where we will also see that the two distinguished tori are cluster tori.

Comparison with the Gorbounov-Smirnov mirror. The Landau-Ginzburg model $(\check{X}_{\text{GS}}, \mathcal{W}_{\text{GS}})$ from [GS13] goes as follows

$$\check{X}_{\text{GS}} = \{(x; y_1, \dots, y_{m-1}; z_1, \dots, z_{m-1}) \in \mathbb{C}^m \times (\mathbb{C}^*)^{m-1} \mid xy_1 \dots y_{m-1} - 1 \neq 0\},$$

$$(29) \quad \mathcal{W}_{\text{GS}} = \sum_{i=1}^{m-1} y_i(1 + z_i) + q \frac{x^2}{(xy_1 y_2 \dots y_{m-1} - 1) z_1 z_2 \dots z_{m-1}}.$$

Consider the change of coordinates:

$$y_i = \frac{p_i}{p_{i-1}} \quad \text{for } 1 \leq i \leq m-1;$$

$$z_i = \frac{\delta_{i-2}}{\delta_{i-1}} \quad \text{for } 2 \leq i \leq m-1;$$

$$z_1 = q \frac{p_0}{p_{2m-1}}, \quad x = \frac{p_0 p_m}{\delta_{m-1}}.$$

Proposition 10.2. *The change of coordinates $\{p_i\} \mapsto \{x, y_i, z_i\}$ above defines an isomorphism between the torus $\{p_1 \dots p_{m-1} \neq 0\}$ and $\{y_1 \dots y_{m-1} \neq 0\}$ inside \check{X}_{GS} and the torus inside \check{X}_{can} , which pulls back the Gorbounov-Smirnov superpotential \mathcal{W}_{GS} to \mathcal{W}_{can} . Moreover when $m = 2$ the change of coordinates is a well-defined isomorphism between \check{X}_{can} and \check{X}_{GS} .*

Proof. We have $y_1(1 + z_1) = p_1 + q \frac{p_1}{p_{2m-1}}$, and $y_i(1 + z_i) = \frac{p_i p_{2m-i}}{\delta_{i-1}}$ for $2 \leq i \leq m-1$. Moreover

$$xy_1 \dots y_{m-1} - 1 = \frac{\delta_{m-2}}{\delta_{m-1}}, \quad z_1 \dots z_{m-1} = \frac{q}{\delta_{m-2}}, \quad x^2 = \frac{p_m^2}{(\delta_{m-1})^2},$$

which gives

$$q \frac{x^2}{(xy_1 y_2 \dots y_{m-1} - 1) z_1 z_2 \dots z_{m-1}} = \frac{p_m^2}{\delta_{m-1}},$$

hence the change of coordinates maps \mathcal{W}_{GS} to \mathcal{W}_{can} . Finally, the change of coordinates is well-defined on the tori, and so is its inverse,

$$p_0 = 1, \quad p_i = \prod_{j=1}^{m-1} y_j \quad \text{for } 1 \leq i \leq m-1, \quad p_m = \frac{q}{z_1 \dots z_{m-1}} \frac{x}{xy_1 \dots y_{m-1} - 1}$$

$$p_{2m-1-i} = \frac{q(1 + z_{i+1})}{y_1 z_1 \dots y_i z_i} \quad \text{for } 1 \leq i \leq m-2, \quad p_{2m-1} = \frac{q}{z_1},$$

which concludes the proof. \square

It follows from Proposition 10.2 that the LG models $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ and $(\check{X}_{\text{GS}}, \mathcal{W}_{\text{GS}})$ are isomorphic in the case of the three-dimensional quadric Q_3 , but in the general case we only get a birational equivalence.

11. THE A-MODEL AND B-MODEL CONNECTIONS

Recall that we have proved that the canonical LG model $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ is isomorphic to the Lie-theoretic LG model $(\check{X}_{\text{Lie}}, \mathcal{W}_{\text{Lie}})$; hence using [Rie08] we deduced that there is an isomorphism between the quantum cohomology of $X = Q_{2m-1}$ and the Jacobi ring of $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$,

$$\text{qH}^*(X, \mathbb{C}) [q^{-1}] \cong \mathbb{C} [\check{X}_{\text{can}} \times \mathbb{C}_q^*] / \left(\frac{\partial \mathcal{W}_{\text{can}, q}}{\partial p_1}, \dots, \frac{\partial \mathcal{W}_{\text{can}, q}}{\partial p_{2m-1}} \right).$$

Furthermore we have proved that the isomorphism is given by mapping the Schubert class $\sigma_i \in H^{2i}(X, \mathbb{Z})$ to the Plücker coordinate p_i .

We may now prove a more detailed mirror theorem by comparing two flat connections, one related to $X = Q_{2m-1}$ and one constructed from $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$. Let \mathcal{H}_A be the sheaf of regular functions of the trivial vector bundle with fibre $H^*(X, \mathbb{C})$ over $\mathbb{C}_\hbar^* \times \mathbb{C}_q^*$, the two-dimensional complex torus with coordinates \hbar and q . The *A-model connection* ${}^A\nabla$, also known as the Dubrovin connection, is defined on \mathcal{H}_A by

$${}^A\nabla_{q\partial_q} = q \frac{\partial}{\partial q} + \frac{1}{\hbar} p_1 \star_q \bullet,$$

$${}^A\nabla_{\hbar\partial_\hbar} = \hbar \frac{\partial}{\partial \hbar} + \text{gr} - \frac{1}{\hbar} c_1(TX) \star_q \bullet,$$

where gr is a diagonal operator on $H^*(X, \mathbb{C})$ given by $\text{gr}(\alpha) = k\alpha$ for $\alpha \in H^{2k}(X, \mathbb{C})$. Here we are using the conventions of [Iri09]. Let \mathcal{H}_A^\vee be the vector bundle on $\mathbb{C}_\hbar^* \times \mathbb{C}_q^*$ defined by

$\mathcal{H}_A^\vee = j^* \mathcal{H}_A$ for $j : (\hbar, q) \mapsto (-\hbar, q)$. This vector bundle with the pulled back connection ${}^A \nabla^\vee = j^* ({}^A \nabla)$ is dual to $(\mathcal{H}_A, {}^A \nabla)$ via the flat non-degenerate pairing,

$$\langle \sigma_j, \sigma_k \rangle = (2\pi i \hbar)^{2m-1} \int_{[X]} \sigma_j \cup \sigma_k = (2\pi i \hbar)^{2m-1} \delta_{j+k, 2m-1}.$$

The dual A-model connection ${}^A \nabla^\vee$ defines a system of differential equations which we call the (small) quantum differential equations

$$(30) \quad {}^A \nabla_{q \partial_q}^\vee S = 0.$$

Let us now define a $\mathbb{C}[\hbar^{\pm 1}, q^{\pm 1}]$ -module

$$\mathcal{G} := \Omega^N(\check{X}_{\text{can}})[\hbar^{\pm 1}, q^{\pm 1}] / \left(d - \frac{1}{\hbar} d\mathcal{W}_{\text{can}, q} \wedge \bullet \right) \Omega^{N-1}(\check{X}_{\text{can}})[\hbar^{\pm 1}, q^{\pm 1}],$$

where $\Omega^k(\check{X}_{\text{can}})$ is the space of holomorphic k -forms on \check{X}_{can} . We denote by \mathcal{H}_B the sheaf with global sections \mathcal{G} . The B -model connection, or Gauss-Manin connection, on \mathcal{H}_B is given by

$$\begin{aligned} {}^B \nabla_{q \partial_q}[\eta] &= q \frac{\partial}{\partial q}[\eta] + \frac{1}{\hbar} \left[q \frac{\partial \mathcal{W}_{\text{can}, q}}{\partial q} \eta \right], \\ {}^B \nabla_{\hbar \partial_{\hbar}}[\eta] &= \hbar \frac{\partial}{\partial \hbar}[\eta] - \frac{1}{\hbar} [\mathcal{W}_{\text{can}, q} \eta]. \end{aligned}$$

Since \check{X}_{can} is a cluster variety there exists a unique up to a scalar non-vanishing N -form on \check{X}_{can} with simple poles along the boundary, which we denote by ω_{can} , compare [LS16]. Explicitly, in terms of homogeneous coordinates we write

$$\omega_{\text{can}} := \bigwedge_{i=1}^{m-1} \frac{dp_i}{p_i} \wedge \bigwedge_{i=1}^{m-1} \frac{d\delta_i}{\delta_i} \wedge \frac{dp_{2m-1}}{p_{2m-1}}.$$


In the isomorphic case of \check{X}_{Lie} this is the form introduced in [Rie08, Section 7].

Theorem 11.1. *The map*

$$(31) \quad \begin{aligned} \Psi : (\mathcal{H}_A, {}^A \nabla) &\rightarrow (\mathcal{H}_B, {}^B \nabla) \\ \sigma_j &\mapsto [p_j \omega_{\text{can}}] \end{aligned}$$

is an injective homomorphism of bundles with connection. Moreover, in the case of the three-dimensional quadric Q_3 it is an isomorphism.

Proof. We use the cluster variety structure of the mirror of Q_{2m-1} . Namely, the coordinate ring $\mathbb{C}[\check{X}_{\text{can}}]$ has a cluster algebra structure of type A_1^{m-1} , which is described in detail in [GLS08b, Section 2] and [GLS08a, Section 12]. Consider the following initial quiver:



Here the initial cluster variables correspond to the vertices in the top row of the quiver, while the frozen variables (or coefficients) correspond to the vertices in the bottom row. In particular, it is of finite type A_1^{m-1} , and there are 2^{m-1} different clusters, consisting of

- the cluster variables r_1, \dots, r_{m-1} , where $r_i \in \{p_i, p_{2m-1-i}\}$;
- the frozen variables (or coefficients) $\delta_1, \dots, \delta_{m-1}$, and p_{2m-1} .

Moreover we have set $p_0 = 1$. The exchange relations are

$$(32) \quad p_i p_{2m-1-i} = \begin{cases} p_{2m-1} + \delta_1 & \text{for } i = 1; \\ \delta_{i-1} + \delta_i & \text{for } 2 \leq i \leq m-1. \end{cases}$$

For the injectivity of Ψ the proof agrees with the proof of the analogous lemma [MR13, Lemma 9.3]. We note that the only ingredients used in [MR13, Lemma 9.3] are the isomorphism of the Jacobi ring with the quantum cohomology ring, see Theorem 9.1, and the fact that \mathcal{W}_{can} has isolated critical points, see [PRW16, Proposition 2.3]. It remains to prove that Ψ maps the A -model connection to the B -model connection. We use a change of coordinates to reduce the problem to checking only the action of $q\partial_q$. Namely, this follows by replacing (p_i, q, \hbar) with $(\mathbf{p}_i, \mathbf{q}, \hbar)$, where

$$\mathbf{p}_i = \hbar^{-i} p_i, \quad \mathbf{q} = \hbar^{1-2m} q, \quad \hbar = \hbar,$$

and observing that written in these coordinates the Gauss-Manin system for $\frac{1}{\hbar}\mathcal{W}_{\text{can},q}$ no longer involves the \hbar .

Now we check that the map Ψ preserves the action of $q\partial_q$. We consider the following identities in $q\mathbb{H}^*(Q_{2m-1}, \mathbb{C})$, which are a special case of results in [FW04]:

$$(33) \quad \sigma_1 \star_q \sigma_i = \begin{cases} \sigma_{i+1} & \text{for } 0 \leq i \leq m-2 \text{ or } m \leq i \leq 2m-3; \\ 2\sigma_m & \text{for } i = m-1; \\ \sigma_{2m-1} + q & \text{for } i = 2m-2; \\ q\sigma_1 & \text{for } i = 2m-1, \end{cases}$$

We need to prove that there are similar identities on the B side:

$$(34) \quad \left[q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} p_i \omega_{\text{can}} \right] = \begin{cases} [p_{i+1} \omega_{\text{can}}] & \text{for } 0 \leq i \leq m-2 \text{ or } m \leq i \leq 2m-3; \\ [2p_{m-1} \omega_{\text{can}}] & \text{for } i = m-1; \\ [(p_{2m-1} + q) \omega_{\text{can}}] & \text{for } i = 2m-2; \\ [qp_1 \omega_{\text{can}}] & \text{for } i = 2m-1, \end{cases}$$

where ω_{can} is the canonical $(2m-1)$ -form on \check{X}_{can} . The proof of these identities on the B side proceeds by constructing closed $(2m-2)$ -forms ν_i such that the relation corresponding to p_i will follow from the fact that

$$[d\mathcal{W}_{\text{can},q} \wedge \nu_i] = [(d + d\mathcal{W}_{\text{can},q} \wedge -)\nu_i] = 0.$$

(The first equality above comes from the fact that ν_i is closed, and the second comes from the definition of the B -model.)

Concretely, we will pick a cluster \mathcal{C} containing a particular Plücker coordinate, say p_i , and use the following Ansatz for constructing ν_i . We define a vector field

$$\xi_i = p_i \left(\sum_{c \in \mathcal{C} \setminus \{p_i\}} m_c c \partial_c \right)$$

and define an associated $(2m-2)$ -form by insertion $\nu_i = \iota_{\xi_i} \omega_{\text{can}}$. Here the m_c 's are constants and ι is the interior product.

To see that these $(2m-2)$ -forms are closed, write $\omega_{\text{can}} = \bigwedge_{p \in \mathcal{C}} \frac{dp}{p}$. For $c \in \mathcal{C}$, we have $\iota_{c\partial_c} \omega_{\text{can}} = \bigwedge_{p \in \mathcal{C} \setminus \{c\}} \frac{dp}{p}$, and so ν_i is a \mathbb{C} -linear combination of terms of the form $p_i \bigwedge_{p \in \mathcal{C} \setminus \{c\}} \frac{dp}{p}$ for $c \neq p_i$. Such a term is closed, because p_i lies in $\mathcal{C} \setminus \{c\}$. Using the fact that $d\mathcal{W}_{\text{can},q} \wedge \omega_{\text{can}} = 0$, we get $d\mathcal{W}_{\text{can},q} \wedge \nu_i = \pm d\mathcal{W}_{\text{can},q}(\xi_i) \omega_{\text{can}}$. It follows that

$$d\mathcal{W}_{\text{can},q} \wedge \nu_i = p_i \left(\sum_{c \in \mathcal{C} \setminus \{p_i\}} m_c c \frac{\partial \mathcal{W}_{\text{can},q}}{\partial c} \right) \omega_{\text{can}}.$$

Therefore e.g. in order to prove that $\left[q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} p_i \omega_{\text{can}} \right] - [p_{i+1} \omega_{\text{can}}] = 0$, we will show that $q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} p_i - p_{i+1}$ has the form $p_i \left(\sum_{c \in \mathcal{C} \setminus \{p_i\}} m_c c \frac{\partial \mathcal{W}_{\text{can},q}}{\partial c} \right)$, for some choice of coefficients m_c .

To prove these identities, we will work with two clusters:

- the initial cluster $\mathcal{C}_1 = \{p_1, \dots, p_{m-1}, \delta_1, \dots, \delta_{m-1}, p_{2m-1}\}$;
- the cluster $\mathcal{C}_2 = \{p_{2m-2}, \dots, p_m, \delta_1, \dots, \delta_{m-1}, p_{2m-1}\}$.

Let us first start with \mathcal{C}_1 and express $\mathcal{W}_{\text{can},q}$ in terms of it using the exchange relations (32). To simplify our notation we let δ_0 denote p_{2m-1} .

$$\mathcal{W}_{\text{can},q} = p_1 + \sum_{\ell=1}^{m-1} \left(\frac{p_{\ell+1} \delta_{\ell-1}}{p_\ell \delta_\ell} + \frac{p_{\ell+1}}{p_\ell} \right) + q \frac{p_1}{\delta_0}.$$

The partial derivatives of $\mathcal{W}_{\text{can},q}$ are:

$$\begin{aligned} q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} &= q \frac{p_1}{\delta_0}, \\ p_1 \frac{\partial \mathcal{W}_{\text{can},q}}{\partial p_1} &= p_1 - \frac{p_2 \delta_0}{p_1 \delta_1} - \frac{p_2}{p_1} + q \frac{p_1}{\delta_0}, \\ p_i \frac{\partial \mathcal{W}_{\text{can},q}}{\partial p_i} &= \frac{p_i \delta_{i-2}}{p_{i-1} \delta_{i-1}} + \frac{p_i}{p_{i-1}} - \frac{p_{i+1} \delta_{i-1}}{p_i \delta_i} - \frac{p_{i+1}}{p_i} \text{ for } 2 \leq i \leq m-1, \\ \delta_0 \frac{\partial \mathcal{W}_{\text{can},q}}{\partial \delta_0} &= \frac{p_2 \delta_0}{p_1 \delta_1} - q \frac{p_1}{\delta_0}, \\ \delta_i \frac{\partial \mathcal{W}_{\text{can},q}}{\partial \delta_i} &= -\frac{p_{i+1} \delta_{i-1}}{p_i \delta_i} + \frac{p_{i+2} \delta_i}{p_{i+1} \delta_{i+1}} \text{ for } 1 \leq i \leq m-1. \end{aligned}$$

Hence

$$q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} p_i - p_{i+1} = -p_i \left(\sum_{j=i+1}^{m-1} p_j \frac{\partial \mathcal{W}_{\text{can},q}}{\partial p_j} + \sum_{j=0}^{m-1} \delta_j \frac{\partial \mathcal{W}_{\text{can},q}}{\partial \delta_j} + \sum_{j=i}^{m-1} \delta_j \frac{\partial \mathcal{W}_{\text{can},q}}{\partial \delta_j} \right)$$

for $0 \leq i \leq m-2$, and

$$q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} p_{m-1} - 2p_m = -p_{m-1} \left(\sum_{j=0}^{m-1} \delta_j \frac{\partial \mathcal{W}_{\text{can},q}}{\partial \delta_j} \right).$$

Since the right-hand sides of the above equations have the form $p_i \left(\sum_{c \in \mathcal{C} \setminus \{p_i\}} m_c c \frac{\partial \mathcal{W}_{\text{can},q}}{\partial c} \right)$, this proves identity (34) for $0 \leq i \leq m-2$.

To prove the remaining identities, we use the cluster \mathcal{C}_2 . In this cluster chart, $\mathcal{W}_{\text{can},q}$ takes the following form:

$$\mathcal{W}_{\text{can},q} = \frac{\delta_0}{p_{2m-2}} + \frac{\delta_1}{p_{2m-2}} + \sum_{\ell=1}^{m-1} \left(\frac{p_{2m-1-\ell}}{p_{2m-2-\ell}} + \frac{p_{2m-1-\ell} \delta_{\ell+1}}{p_{2m-2-\ell} \delta_\ell} \right) + \frac{q}{p_{2m-2}} + \frac{q \delta_1}{p_{2m-2} \delta_0}.$$

Working out the partial derivatives of $\mathcal{W}_{\text{can},q}$ as before, we get

$$q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} p_i - p_{i+1} = p_i \left(- \sum_{j=i+1}^{2m-2} p_j \frac{\partial \mathcal{W}_{\text{can},q}}{\partial p_j} - \sum_{j=0}^{2m-2-i} \delta_j \frac{\partial \mathcal{W}_{\text{can},q}}{\partial \delta_j} \right) \text{ for } m \leq i \leq 2m-3.$$

Recall that δ_0 is p_{2m-1} . The final two relations are

$$\begin{aligned} q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} p_{2m-2} - (p_{2m-1} + q) &= -p_{2m-2} \delta_0 \frac{\partial \mathcal{W}_{\text{can},q}}{\partial \delta_0} \quad \text{and} \\ q \frac{\partial \mathcal{W}_{\text{can},q}}{\partial q} p_{2m-1} - q p_1 &= 0. \end{aligned}$$

This gives us the identities (34) for $m - 1 \leq i \leq 2m - 1$, which concludes the proof of the homomorphism. To deduce that the map is injective as claimed we use the same strategy as in [MR13, Lemma 9.3]. Namely, observe that the relations in the Gauss-Manin system recover the relations of the Jacobi ring as \hbar tends to zero. On the other hand as we already proved, the Jacobi ring is isomorphic to quantum cohomology with the homogeneous coordinates p_i playing the role of the Schubert basis. Therefore we see that the $[p_i \omega_{can}] \in \mathcal{G}$ are linearly independent in the $\hbar \rightarrow 0$ limit. Hence they must be linearly independent already in \mathcal{G} .

In the case of Q_3 the added surjectivity result is a consequence of the fact that $(\check{X}_{can}, \mathcal{W}_{can})$ is isomorphic to the Gorbounov-Smirnov mirror in that case, see Proposition 10.2. Indeed \mathcal{W}_{GS} is cohomologically tame [GS13], hence so is \mathcal{W}_{can} . Therefore \mathcal{G} is a free $\mathbb{C}[\hbar^{\pm 1}, q^{\pm 1}]$ -module of rank $2m$ (cf. [Sab99]), and \mathcal{H}_B a trivial vector bundle of that dimension. \square

Let Γ_0 be a compact oriented real $(2m - 1)$ -dimensional submanifold of \check{X}_{can} representing a cycle in $H^{2m-1}(\check{X}_{can}, \mathbb{Z})$ dual to ω_{can} , in the sense that $\frac{1}{(2i\pi)^{2m-1}} \int_{\Gamma_0} \omega_{can} = 1$. From Theorem 11.1 we deduce the following formula.

Corollary 11.2. *The integral formula*

$$S_0(\hbar, q) = \frac{1}{(2i\pi\hbar)^{2m-1}} \sum_{j=0}^{2m-1} \left(\int_{\Gamma_0} e^{\frac{\mathcal{W}_{can,q}}{\hbar}} p_j \omega_{can} \right) \sigma_{2m-1-j}$$

describes a solution to the quantum differential equation (30). \square

The corollary follows as in [MR13, Theorem 4.2]. If we replace $(\check{X}_{can}, \mathcal{W}_{can})$ by the isomorphic LG model $(\check{X}_{Lie}, \mathcal{W}_{Lie})$ the above corollary implies a special case of [Rie08, Conj. 8.1] for odd-dimensional quadrics.

12. THE MIRROR TO Q_3

In this section we work out in detail the example of the three-dimensional quadric, Q_3 , to illustrate our main results.

The Laurent polynomial mirror $(\check{X}_{Lus}, \mathcal{W}_{Lus})$. Recall from Section 4 the definition of the variety $Z \subset G$,

$$Z := B_- \dot{w}_0 \cap U_+ T^{W_P} \dot{w}_P U_-.$$

A generic element $g \in Z$ can be written as $g = u_1 t \dot{w}_P \bar{u}_2$, where

$$\bar{u}_2 = y_1(a) y_2(c) y_1(b),$$

and a, b, c are non-zero, i.e.

$$\bar{u}_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ a+b & 1 & 0 & 0 \\ cb & c & 1 & 0 \\ acb & ac & a+b & 1 \end{pmatrix}.$$

From this the expression of the Laurent polynomial mirror follows, namely,

$$\check{X}_{Lus} = (\mathbb{C}^*)_{a,b,c}^3, \quad \mathcal{W}_{Lus} = a + b + c + \frac{a+b}{abc}.$$

This illustrates Theorem 5.4 in the case of Q_3 .

The canonical mirror. The map $Z_t \rightarrow \check{X}_{\text{can}} \cong \mathbb{CP}^3$ takes $z = u_1 t \dot{w}_P \bar{u}_2$ to $Pz = P\bar{u}_2$. This may be interpreted as taking z to the span of the reverse row vector corresponding to the last row of \bar{u}_2 after the identification $\check{X}_{\text{can}} \cong \mathbb{CP}^3$. The homogeneous coordinates of \bar{u}_2 are given by $p_0 = 1, p_1 = a + b, p_2 = ac, p_3 = acb$.

The image of $Z_{q=q_0}$ in \mathbb{CP}^3 is independent of q_0 , so we may choose q_0 to be 1, and restrict our attention to $Z_{q=1} := B_- \dot{w}_0 \cap U_+ \dot{w}_P U_-$. The image of $Z_{q=1}$ is obtained in coordinates $(p_0 : p_1 : p_2 : p_3)$ by removing the anticanonical divisor

$$D := \{p_0 = 0\} \cup \{p_3 p_0 - p_2 p_1 = 0\} \cup \{p_3 = 0\}.$$

Thus

$$\check{X}_{\text{can}} = \{(p_0 : p_1 : p_2 : p_3) \in \mathbb{CP}^3 \mid p_0(p_1 p_2 - p_0 p_3) p_3 \neq 0\}$$

In terms of the homogeneous coordinates, the canonical superpotential from Equation (20) is given by

$$\mathcal{W}_{\text{can},q} = \frac{p_1}{p_0} + \frac{p_2^2}{p_1 p_2 - p_0 p_3} + q \frac{p_1}{p_3}.$$

Comparison with the Gorbounov-Smirnov-mirror. The mirror $(\check{X}_{\text{GS}}, \mathcal{W}_{\text{GS}})$ for Q_3 is given by

$$\check{X}_{\text{GS}} = \{(x, y, z) \in \mathbb{C}^3 \mid (xy - 1)z \neq 0\}, \mathcal{W}_{\text{GS}} = y(1 + z) + q \frac{x^2}{(xy - 1)z}.$$

It corresponds to $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ via the change of coordinates (setting $p_0 = 1$):

$$x = \frac{p_2}{p_1 p_2 - p_3}; \quad y = p_1; \quad z = \frac{q}{p_3}.$$

This change of coordinates is well-defined on \check{X}_{can} , and its inverse,

$$p_1 = y; \quad p_2 = \frac{qx}{(xy - 1)z}; \quad p_3 = \frac{q}{z}$$

is well-defined on \check{X}_{GS} . This illustrates the result of Proposition 10.2, namely that for Q_3 the LG models $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ and $(\check{X}_{\text{GS}}, \mathcal{W}_{\text{GS}})$ are isomorphic, even though that is not the case in higher dimension.

Isomorphism between the quantum cohomology and the Jacobi ring. Recall that the cohomology of Q_3 is generated by the Schubert classes $\sigma_i \in H^{2i}(Q_3, \mathbb{C})$ for $i = 0, \dots, 3$. Moreover it has a presentation:

$$H^*(Q_3, \mathbb{C}) = \mathbb{C}[\sigma_1, \sigma_2] / (\sigma_1^4, \sigma_1^2 - 2\sigma_2),$$

and the quantum cohomology of Q_3 is presented as follows,

$$\text{qH}^*(Q_3, \mathbb{C}) = \mathbb{C}[\sigma_1, \sigma_2, q] / (\sigma_1^4 - q\sigma_1, \sigma_1^2 - 2\sigma_2).$$

Now the Jacobi ring of $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ is

$$\mathbb{C}[\check{X}_{\text{can}} \times \mathbb{C}_q^*] / (p_3^2 - qp_3, p_1 p_2 - 2p_3, p_2^2 - qp_1)$$

The map

$$p_1 \mapsto \sigma_1, \quad p_2 \mapsto \sigma_2, \quad p_3 \mapsto \frac{1}{2}\sigma_1^3$$

defines an isomorphism between the Jacobi ring of $(\check{X}_{\text{can}}, \mathcal{W}_{\text{can}})$ and $\text{qH}^*(Q_3, \mathbb{C})$ as in Theorem 9.1.

The quantum differential equations. Recall from Section 11 that the dual A -model connection ${}^A\nabla^\vee$ defines a system of differential equations called the quantum differential equations,

$${}^A\nabla_{q\partial q}^\vee S = 0.$$

In the case of Q_3 our mirror result, Theorem 11.1, tells us that the map $(\mathcal{H}_A, {}^A\nabla) \rightarrow (\mathcal{H}_B, {}^B\nabla)$ given by

$$\sigma_i \mapsto [p_i \omega_{can}],$$

where $\omega_{can} = \frac{dp_1}{p_1} \wedge \frac{d\delta_1}{\delta_1} \wedge \frac{dp_3}{p_3}$, is an isomorphism. If Γ is a real 3-dimensional cycle in $\mathbb{C}\mathbb{P}^3 \setminus D$ this implies in particular that the function

$$S_\Gamma(q) := \left(\int_\Gamma e^{\mathcal{W}_{can,q} p_3} \omega_{can} \right) \sigma_0 + \left(\int_\Gamma e^{\mathcal{W}_{can,q} p_2} \omega_{can} \right) \sigma_1 + \left(\int_\Gamma e^{\mathcal{W}_{can,q} p_1} \omega_{can} \right) \sigma_2 + \left(\int_\Gamma e^{\mathcal{W}_{can,q}} \omega_{can} \right) \sigma_3$$

satisfies ${}^A\nabla_{q\partial q}^\vee S_\Gamma(q) = 0$.

REFERENCES

- [Bea95] Arnaud Beauville. Quantum cohomology of complete intersections. *Mat. Fiz. Anal. Geom.*, 3–4(2):384–398, 1995.
- [BZ97] Arkady Berenstein and Andrei Zelevinsky. Total positivity in Schubert varieties. *Comment. Math. Helv.*, 72(1):128–166, 1997.
- [CMP07] Chaput, P.-E., L. Manivel, and N. Perrin. Quantum Cohomology of Minuscule Homogeneous Spaces II Hidden Symmetries, International Mathematics Research Notices, Vol. 2007, Article ID rnm107, 29 pages. doi:10.1093/imrn/rnm107.
- [CK99] David A. Cox and Sheldon Katz. Mirror symmetry and algebraic geometry. *Mathematical Surveys and Monographs*, 68:xxii+469, American Mathematical Society, Providence, RI, 1999.
- [Dub96] Boris Dubrovin. Geometry of 2D topological field theories. In *Integrable systems and quantum groups (Montecatini Terme, 1993)*, volume 1620 of *Lecture Notes in Math.*, pages 120–348. Springer, Berlin, 1996.
- [EHX97] Tohru Eguchi, Kentaro Hori, and Chuan-Sheng Xiong. Gravitational quantum cohomology. *Internat. J. Modern Phys. A*, 12(9):1743–1782, 1997.
- [FW04] W. Fulton and C. Woodward. On the quantum product of Schubert classes. *J. Algebraic Geom.*, 13(4):641–661, 2004.
- [Gin95] V. Ginzburg. Perverse sheaves on a Loop group and Langlands’ duality. arXiv:9511007, 1995.
- [Giv95] Alexander Givental. Homological geometry and mirror symmetry. In *Proceedings of the International Congress of Mathematicians*, Vol. 1, 2 (Zürich, 1994), pages 472–480, Birkhäuser, Basel, 1995.
- [Giv96] Alexander Givental. Equivariant Gromov-Witten invariants. *Internat. Math. Res. Notices* 13:613–663, 1996.
- [GLS08a] Christof Geiß, Bernard Leclerc, and Jan Schröer. Partial flag varieties and preprojective algebras. *Ann. Inst. Fourier (Grenoble)* 58(3):825–876, 2008.
- [GLS08b] Christof Geiß, Bernard Leclerc, and Jan Schröer. Preprojective algebras and cluster algebras, Trends in representation theory of algebras and related topics. In *EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich* 253–283, 2008.
- [GS13] Vassily Gorbounov and Maxim Smirnov. Some remarks on Landau-Ginzburg potentials for odd-dimensional quadrics. *Glasgow Mathematical Journal*, Volume 57, Issue 3, DOI: <http://dx.doi.org/10.1017/S0017089514000433>, pp. 481–507, September 2015.
- [HV00] Kentaro Hori and Cumrun Vafa. Mirror symmetry. *arXiv preprint hep-th/0002222*, 2000.
- [Iri09] Hiroshi Iritani. An integral structure in quantum cohomology and mirror symmetry for toric orbifolds. *Adv. Math.*, 222(3):1016–1079, 2009.
- [LS16] Thomas Lam and David Speyer. Cohomology of cluster varieties. I. Locally acyclic case. *arXiv:1604.06843v1 [math.AG]*, 2016
- [Lus83] George Lusztig. Singularities, character formulas, and a q -analog of weight multiplicities. In *Analysis and topology on singular spaces, II, III (Luminy, 1981)*, volume 101 of *Astérisque*, pages 208–229. Soc. Math. France, Paris, 1983.

- [Lus94] George Lusztig. Total positivity in reductive groups. In *Lie theory and geometry*, volume 123, *Progr. Math.*, pages 531–568, Birkhäuser Boston, 1994.
- [MR13] R. Marsh and K. Rietsch. The B -model connection and T -equivariant mirror symmetry for Grassmannians. arXiv:1307.1085v2 [math.AG].
- [MV07] I. Mirković and K. Vilonen. Geometric Langlands duality and representations of algebraic groups over commutative rings. *Ann. of Math. (2)*, 166(1):95–143, 2007.
- [Pet97] D. Peterson. Quantum cohomology of G/P . Lecture Course, MIT, Spring Term, 1997.
- [PR13] C. Pech and K. Rietsch. A Landau–Ginzburg model for Lagrangian Grassmannians, Langlands duality and relations in quantum cohomology. arXiv:1304.4958, 2013.
- [PRW16] C. Pech, K. Rietsch, and L. Williams. On Landau–Ginzburg models for quadrics and flat sections of Dubrovin connections. *Adv. Math.*, 300:275–319, 2016.
- [Prz07] V. V. Przhlyakovskiĭ. Gromov-Witten invariants of Fano threefolds of genera 6 and 8. *Mat. Sb.*, 198(3):145–158, 2007.
- [Rie08] Konstanze Rietsch. A mirror symmetric construction of $qH_T^*(G/P)_{(q)}$. *Adv. Math.*, 217(6):2401–2442, 2008.
- [Sab99] Claude Sabbah. Hypergeometric period for a tame polynomial. *C. R. Acad. Sci. Paris Sér. I Math.*, 328(7):603–608, 1999.

UNIVERSITY OF KENT, CANTERBURY, CT2 7NZ,

KING’S COLLEGE LONDON, STRAND, LONDON, WC2R 2LS