

Kent Academic Repository

Full text document (pdf)

Citation for published version

Waterstraat, Nils (2013) A remark on the space of metrics having non-trivial harmonic spinors. *Journal of Fixed Point Theory and Applications*, 13 . pp. 143-149. ISSN 1661-7738.

DOI

<https://doi.org/10.1007/s11784-013-0096-5>

Link to record in KAR

<http://kar.kent.ac.uk/51390/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

A remark on the space of metrics having non-trivial harmonic spinors

Nils Waterstraat

Abstract

Let M be a closed spin manifold of dimension $n \equiv 3 \pmod{4}$. We give a simple proof of the fact that the space of metrics on M with invertible Dirac operator is either empty or it has infinitely many path components.

1 Introduction

Let M be an n -dimensional closed spin manifold and let $R(M)$ be the space of all Riemannian metrics on M . For any choice of a metric $g \in R(M)$, we can build the associated spinor bundle $\Sigma_g M$ and obtain a natural first order operator D_g acting on sections of $\Sigma_g M$ and which we call the *Dirac operator*. Elements of $\ker D_g$ are called *harmonic spinors* and their existence has been studied for a long time. While one can show that on S^2 no non-trivial harmonic spinors exist for any choice of g (cf. [Ba92]), it is conjectured that on every closed spin manifold of dimension $n \geq 3$ there exists a Riemannian metric g such that $\ker D_g \neq 0$. The conjecture has been proved by N. Hitchin in [Hi74] if $n \equiv 0, \pm 1 \pmod{8}$ and by C. Bär in [Ba96] if $n \equiv 3 \pmod{4}$.

As a more general question, one may ask how many metrics exist on M such that the corresponding Dirac operator has non-trivial kernel. A possible way to study this question is to consider the complementary set of metrics $R^{\text{inv}}(M)$ consisting of all metrics $g \in R(M)$ such that $\ker D_g = 0$. M. Dahl showed in [Da08] that elements of $R^{\text{inv}}(M)$ can be extended to $R^{\text{inv}}(W)$ if W is the trace of a surgery of codimension at least 3 on M . By using the Atiyah-Singer index theorem and special metrics on the spheres originating from the study of positive scalar curvature, he concluded from this result that $R^{\text{inv}}(M)$ is in all dimensions $n \geq 5$ which were considered by Hitchin and Bär either empty or disconnected. Moreover, in the case $n \equiv 3 \pmod{4}$, $n \geq 7$, he even obtained that, if non-empty, $R^{\text{inv}}(M)$ has infinitely many path components. Recently he improved this conclusion in collaboration with N. Grosse to dimension 3 by studying extensions of metrics to attached handles (cf. [DG12]).

The aim of this article is to show that the existence of infinitely many connected components of $R^{\text{inv}}(M)$ in all dimensions $n \equiv 3 \pmod{4}$ can be derived easily from Bär's results in [Ba96] by using spectral flow and rather elementary homotopy arguments.

Finally, we want to mention that Bär improved his theorem in [Ba97] to twisted Dirac operators. Note that for any fixed pair (F, ∇) of a bundle F over M and a connection ∇ on F , we obtain a family $D^{(F, \nabla)}$ of twisted Dirac operators which is again parametrised by the space of Riemannian metrics $R(M)$ on M . We believe that one can extend our argument here to this case by using the results from [Ba97] instead of [Ba96]. Accordingly, we conjecture that the corresponding space $R_{(F, \nabla)}^{\text{inv}}(M) = \{g \in R(M) : \ker D_g^{(F, \nabla)} = 0\}$ is either empty or it has infinitely many path components.

2010 Mathematics Subject Classification: Primary 58J30; Secondary 53C27

Acknowledgement: I want to thank Jacobo Pejsachowicz for encouraging me to write this paper and for several helpful suggestions improving its presentation.

2 Preliminaries: Dirac operators

In this section we recall briefly the definition of spinor bundles and their Dirac operators. Among the many references for these topics we want to mention [Hij01] and [Am01], on which we base our exposition. In order to simplify the presentation we assume throughout that M is an oriented closed manifold of odd dimension $n \geq 3$.

We denote by $\widetilde{\text{GL}}^+(M)$ the principal $\widetilde{\text{GL}}^+(n; \mathbb{R})$ -bundle of oriented bases over M and recall that $\widetilde{\text{GL}}^+(n; \mathbb{R})$ has a unique connected 2-fold covering $\Theta : \widetilde{\text{GL}}^+(n; \mathbb{R}) \rightarrow \text{GL}^+(n; \mathbb{R})$ since the fundamental group of $\text{GL}^+(n; \mathbb{R})$ is of order two. A *spin structure* on M is a pair $(\widetilde{\text{GL}}^+(M), \vartheta)$, where $\widetilde{\text{GL}}^+(M)$ is a principal $\widetilde{\text{GL}}^+(n; \mathbb{R})$ -bundle over M and $\vartheta : \widetilde{\text{GL}}^+(M) \rightarrow \text{GL}^+(M)$ is a 2-fold covering such that

$$\pi \circ \vartheta = \tilde{\pi} \quad \text{and} \quad \vartheta(u \cdot v) = \vartheta(u) \cdot \Theta(v), \quad \text{for all } v \in \widetilde{\text{GL}}^+(n; \mathbb{R}), u \in \widetilde{\text{GL}}^+(M),$$

where π and $\tilde{\pi}$ denote the corresponding projections of the bundles. Henceforth we assume that M is a *spin manifold*, that is, M is oriented and a spin structure on M is given. Note that so far we have not required that M is endowed with a Riemannian metric.

Let now g be a Riemannian metric on M and denote by $\text{SO}(M, g)$ the associated principal $\text{SO}(n)$ -bundle of positively oriented orthonormal bases. Then $\text{Spin}(M, g) := \vartheta^{-1}(\text{SO}(M, g))$ is a principal $\text{Spin}(n)$ -bundle over M , where $\text{Spin}(n) := \Theta^{-1}(\text{SO}(n))$ is the unique connected 2-fold covering of $\text{SO}(n)$. Let $\rho : \mathbb{C}l_n \rightarrow \text{End}(\Sigma_n)$ denote the usual irreducible representation of the complex Clifford algebra, where Σ_n is the space of complex spinors. We fix an inner product $\langle \cdot, \cdot \rangle$ on Σ_n such that $\langle \rho(x)\sigma_1, \rho(x)\sigma_2 \rangle = \langle \sigma_1, \sigma_2 \rangle$ for all $x \in \mathbb{R}^n$, $\|x\| = 1$, and $\sigma_1, \sigma_2 \in \Sigma_n$. If now $\rho' : \text{Spin}(n) \rightarrow \text{Aut}(\Sigma_n)$ denotes the complex spinor representation of $\text{Spin}(n)$, which is obtained by restricting ρ to $\text{Spin}(n) \subset \mathbb{C}l_n$, then the *spinor bundle* $\Sigma_g M$ of M with respect to g is defined as the associated vector bundle $\text{Spin}(M, g) \times_{\rho'} \Sigma_n$.

The representation ρ induces a *Clifford multiplication* on $\Sigma_g M$, that is, a complex linear vector bundle homomorphism

$$m : T^*M \otimes \Sigma_g M \rightarrow \Sigma_g M, \quad X^\flat \otimes \varphi \mapsto X \cdot \varphi$$

such that $X \cdot (Y \cdot \varphi) + Y \cdot (X \cdot \varphi) = -2g(X, Y)\varphi$ for all $X, Y \in TM$ and $\varphi \in \Sigma_g M$. Moreover, the inner product on Σ_n gives rise to an Hermitian structure on the bundle $\Sigma_g M$ such that $\langle X \cdot \varphi, \psi \rangle = -\langle \varphi, X \cdot \psi \rangle$ for all $X \in TM$ and $\varphi, \psi \in \Sigma_g M$. Finally, the Levi-Civita connection on TM induces a connection on $\text{SO}(M, g)$ and this connection lifts in a canonical way to a connection on $\text{Spin}(M, g)$. The associated covariant derivative $\nabla : C^\infty(M, \Sigma_g M) \rightarrow C^\infty(M, T^*M \otimes \Sigma_g M)$ on the spinor bundle has the properties

$$X \langle \varphi, \psi \rangle = \langle \nabla_X \varphi, \psi \rangle + \langle \varphi, \nabla_X \psi \rangle \quad \text{and} \quad \nabla_X (Y \cdot \varphi) = (\nabla_X^{TM} Y) \cdot \varphi + Y \cdot (\nabla_X \varphi)$$

for vector fields X, Y and a spinor field φ .

Now the Dirac operator with respect to the metric g is defined by

$$D_g = m \circ \nabla : C^\infty(M, \Sigma_g M) \rightarrow C^\infty(M, \Sigma_g M)$$

and is an elliptic, essentially selfadjoint differential operator of first order.

3 The Proof

We assume from now on that M is a closed spin manifold of dimension $n \equiv 3 \pmod{4}$. We denote by $R(M)$ the space of all Riemannian metrics on M with the C^1 -topology and note that it is obviously contractible. Moreover, we define

$$R^{\text{inv}}(M) = \{g \in R(M) : \ker D_g = 0\} \subset R(M)$$

and recall that our aim is to show that this set has infinitely many path components if it is not empty. Accordingly, we assume henceforth that $R^{\text{inv}}(M) \neq \emptyset$ and now we conclude in three steps the announced disconnectedness of this space.

Step 1: The spectral flow

Since our operators D_g , $g \in R(M)$, are essentially selfadjoint, they have real spectra. Moreover, by ellipticity their spectra are discrete and consist entirely of eigenvalues of finite multiplicity. We define for any compact interval $[a, b] \subset \mathbb{R}$ a non-negative integer by

$$m(g, [a, b]) = \sum_{\lambda \in [a, b]} \dim \ker(D_g - \lambda \cdot id).$$

Next we quote the following stability result for the spectra of the operators D_g that can be found in [Ba96, Prop. 7.1].

Theorem 3.1. *Let (M, g) be a closed Riemannian spin manifold with Dirac operator D_g . Let $\varepsilon > 0$ and let $\Lambda > 0$ such that $-\Lambda, \Lambda \notin \sigma(D_g)$. Write*

$$\sigma(D_g) \cap (-\Lambda, \Lambda) = \{\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k\}.$$

Then there exists a neighbourhood of g in the C^1 -topology such that for any metric \tilde{g} in this neighbourhood with Dirac operator $D_{\tilde{g}}$ the following holds:

- $\sigma(D_{\tilde{g}}) \cap (-\Lambda, \Lambda) = \{\mu_1 \leq \mu_2 \leq \dots \leq \mu_k\}$,
- $|\lambda_i - \mu_i| < \varepsilon$, $i = 1, \dots, k$.

The eigenvalues λ_i and μ_i are repeated according to their multiplicities.

We obtain immediately the following corollary.

Corollary 3.2. *For all $g_0 \in R(M)$ and $\Lambda > 0$ such that $\pm\Lambda \notin \sigma(D_{g_0})$ there exists an open neighbourhood $N(g_0, \Lambda) \subset R(M)$ such that $\pm\Lambda \notin \sigma(D_g)$ and $m(g, [-\Lambda, \Lambda]) = m(g_0, [-\Lambda, \Lambda])$ for all $g \in N(g_0, \Lambda)$.*

Let now $\gamma : I \rightarrow R(M)$ be a path of metrics. Because of corollary 3.2 we can find a decomposition $0 = t_0 < t_1 < \dots < t_N = 1$ and positive numbers a_1, \dots, a_N such that the functions $[t_{i-1}, t_i] \ni t \mapsto m(\gamma(t), [-a_i, a_i])$ are constant. We define

$$\Gamma(\gamma) = \sum_{i=1}^N m(\gamma(t_i), [0, a_i]) - m(\gamma(t_{i-1}), [0, a_i]) \in \mathbb{Z} \quad (1)$$

and note that, roughly speaking, $\Gamma(\gamma)$ counts the number of negative eigenvalues of $D_{\gamma(t)}$ that become positive as the parameter t travels from 0 to 1 minus the number of positive eigenvalues of $D_{\gamma(t)}$ that become negative; i.e., the net number of eigenvalues which cross zero. The formula (1) corresponds precisely to the definition of the spectral flow for paths of selfadjoint Fredholm operators acting on a fixed Hilbert space which can be found for example in [Phi96] and [BLP05]. Accordingly, one can show verbatim as in [Phi96] that $\Gamma(\gamma)$ indeed does only depend on the path γ and not on the choices of the $t_i, a_i, i = 1, \dots, N$. Moreover, if $\gamma, \tilde{\gamma} : I \rightarrow R(M)$ are two paths of metrics, then the following properties hold:

- i) $\Gamma(\gamma) = 0$ if $\gamma(t) \in R^{\text{inv}}(M)$ for all $t \in [0, 1]$,
- ii) $\Gamma(\gamma * \tilde{\gamma}) = \Gamma(\gamma) + \Gamma(\tilde{\gamma})$, whenever the concatenation $\gamma * \tilde{\gamma}$ exists,
- iii) $\Gamma(\gamma^{-1}) = -\Gamma(\gamma)$, where $\gamma^{-1}(t) = \gamma(1 - t), t \in I$,
- iv) $\Gamma(\gamma) = \Gamma(\tilde{\gamma})$ if $\gamma \simeq \tilde{\gamma}$ through a homotopy having ends in $R^{\text{inv}}(M)$.

Note that the first three properties are immediate consequences of the definition. The homotopy invariance can be obtained again verbatim as in [Phi96].

Step 2: The range of Γ

Our argument in this section is based on results from [Ba96] which we introduce before we proceed with the proof. At first, we need the existence of the following metrics on the sphere S^n , that were constructed in [Ba96, §3].

Proposition 3.3. *For $n \equiv 3 \pmod{4}$ and any integer $m > 0$, there exists a path of metrics $g_t^m, t \in [0, 1]$, on S^n such that the following holds for the associated Dirac operators \mathcal{D}_t^m :*

- there is $\lambda(t) \in \sigma(\mathcal{D}_t^m)$ such that $\lambda(0) = -1$ and $\lambda(1) = 1$,
- $\lambda(t)$ depends linearly on t ,
- the multiplicity of $\lambda(t)$ is constant in t and greater than m ,
- $\lambda(t)$ is the only eigenvalue of \mathcal{D}_t^m in the interval $[-2, 2]$.

Bär combined in [Ba96] proposition 3.3 and a general gluing theorem for Dirac operators [Ba96, theorem B] to conclude the existence of non-trivial harmonic spinors in dimensions $n \equiv 3 \pmod{4}$. Actually, in order to find the spinors he just needed a special case of his gluing theorem which reads as follows.

Theorem 3.4. *Let (M, g) be a closed Riemannian spin manifold of odd dimension $n \geq 3$. Let D_g be the corresponding Dirac operator and let \mathcal{D} denote the Dirac operator on S^n with respect to some Riemannian metric. Finally, let $\Lambda > 0$ be such that $\pm\Lambda \notin \sigma(D_g) \cup \sigma(\mathcal{D})$. Write*

$$(\sigma(D_g) \cup \sigma(\mathcal{D})) \cap (-\Lambda, \Lambda) = \{\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k\}.$$

Then for any $\varepsilon > 0$ there exists a Riemannian metric \tilde{g} on M such that the corresponding Dirac operator $D_{\tilde{g}}$ has the following properties:

- i) $\pm\Lambda \notin \sigma(D_{\tilde{g}})$,*
- ii) $\sigma(D_{\tilde{g}}) \cap (-\Lambda, \Lambda) = \{\mu_1 \leq \mu_2 \leq \dots \leq \mu_k\}$*
- iii) $|\lambda_j - \mu_j| < \varepsilon$ for $j = 1, \dots, k$.*

The eigenvalues λ_i and μ_i are repeated according to their multiplicities.

We now take some metric $g_0 \in R^{\text{inv}}(M)$. Because of the conformal covariance of the Dirac operator (cf. [Hij01, Prop. 5.13]), we can assume that $[-2, 2] \cap \sigma(D_{g_0}) = \emptyset$ simply by rescaling the metric if necessary.

Let $m > 0$ be an integer and consider the operators \mathcal{D}_t^m on S^n from proposition 3.3. Recall that we denote by $\lambda(t)$ the unique eigenvalue of \mathcal{D}_t^m in the interval $[-2, 2]$ and that $\lambda(t)$ depends linearly on t with $\lambda(0) = -1$, $\lambda(1) = 1$.

We now apply theorem 3.4 for $\Lambda = 2$ and $\varepsilon = \frac{1}{2}$ to D_{g_0} and the operators \mathcal{D}_t^m , $t \in [0, 1]$, on S^n . We obtain for any $t \in [0, 1]$ a metric \tilde{g}_t on M such that each eigenvalue of $D_{\tilde{g}_t}$ in the interval $[-2, 2]$ is of distance less than $\frac{1}{2}$ to $\lambda(t)$. In particular, $D_{\tilde{g}_0}$ and $D_{\tilde{g}_1}$ are invertible and hence $\{\tilde{g}_t\}_{t \in [0, 1]}$ defines a path $\gamma : (I, \partial I) \rightarrow (R(M), R^{\text{inv}}(M))$. Moreover, the function $t \mapsto m(\gamma(t), [-2, 2])$ is constant on the whole interval $[0, 1]$. Hence we finally obtain from the definition of Γ

$$\Gamma(\gamma) = m(\tilde{g}_1, [0, 2]) - m(\tilde{g}_0, [0, 2]) = m(\tilde{g}_1, [0, 2]) = \dim \ker(\mathcal{D}_1^m - id) > m.$$

To sum up, we have shown that the set

$$\{\Gamma(\gamma) : \gamma : (I, \partial I) \rightarrow (R(M), R^{\text{inv}}(M)) \text{ continuous}\} \subset \mathbb{Z}$$

is not bounded from above.

Step 3: The final argument

We fix some $g_0 \in R^{\text{inv}}(M)$. Our first aim of this final step is to construct inductively a sequence of paths $\gamma_k : (I, \partial I) \rightarrow (R(M), R^{\text{inv}}(M))$, $k \in \mathbb{N}$, such that $\gamma_k(0) = g_0$ for all $k \in \mathbb{N}$ and $\Gamma(\gamma_i) \neq \Gamma(\gamma_j)$ for all $i \neq j$.

Let γ_1 be the constant path $\gamma_1 \equiv g_0 \in R^{\text{inv}}(M)$. Assume that we have already constructed $\gamma_1, \dots, \gamma_k : (I, \partial I) \rightarrow (R(M), R^{\text{inv}}(M))$ such that $\gamma_i(0) = g_0$, $i = 1, \dots, k$, and $\Gamma(\gamma_i) \neq \Gamma(\gamma_j)$ for all $i \neq j$.

According to the second step of our proof we can find a path $\tilde{\gamma} : (I, \partial I) \rightarrow (R(M), R^{\text{inv}}(M))$ such that

$$\Gamma(\tilde{\gamma}) > \max_{1 \leq i, j \leq k} |\Gamma(\gamma_i) - \Gamma(\gamma_j)|. \quad (2)$$

Moreover, we choose a path $\hat{\gamma} : (I, \partial I) \rightarrow (R(M), R^{\text{inv}}(M))$ such that $\hat{\gamma}(0) = g_0$ and $\hat{\gamma}(1) = \tilde{\gamma}(0)$. Then $\hat{\gamma} * \tilde{\gamma} : (I, \partial I) \rightarrow (R(M), R^{\text{inv}}(M))$ and we set $\gamma_{k+1} = \hat{\gamma} * \tilde{\gamma}$ if $\Gamma(\hat{\gamma} * \tilde{\gamma}) \neq \Gamma(\gamma_j)$ for all $j = 1, \dots, k$.

If, on the other hand, $\Gamma(\hat{\gamma} * \tilde{\gamma}) = \Gamma(\gamma_j)$ for some $j = 1, \dots, k$, then we set $\gamma_{k+1} = \hat{\gamma}$. In order to justify this choice, assume that also $\Gamma(\hat{\gamma}) = \Gamma(\gamma_i)$ for some $1 \leq i \leq k$. Then we obtain

$$\Gamma(\gamma_j) = \Gamma(\hat{\gamma} * \tilde{\gamma}) = \Gamma(\hat{\gamma}) + \Gamma(\tilde{\gamma}) = \Gamma(\gamma_i) + \Gamma(\tilde{\gamma}),$$

which contradicts (2). Hence we indeed obtain a sequence $\{\gamma_k\}_{k \in \mathbb{N}}$ with the required properties. We now finish our proof by claiming that the metrics $\gamma_k(1)$, $k \in \mathbb{N}$, all lie in different path components of $R^{\text{inv}}(M)$. Assume on the contrary that we can find $i, j \in \mathbb{N}$, $i \neq j$, and a path $\tilde{\gamma} : I \rightarrow R^{\text{inv}}(M)$ such that $\tilde{\gamma}(0) = \gamma_i(1)$ and $\tilde{\gamma}(1) = \gamma_j(1)$. Then $\gamma_i * \tilde{\gamma} * \gamma_j^{-1}$ is a closed path with initial point $g_0 \in R^{\text{inv}}(M)$. Since $R(M)$ is contractible, $\gamma_i * \tilde{\gamma} * \gamma_j^{-1}$ is homotopic to the constant path $\gamma_1 \equiv g_0$ through a g_0 -preserving homotopy. We obtain from the properties of Γ

$$0 = \Gamma(\gamma_1) = \Gamma(\gamma_i * \tilde{\gamma} * \gamma_j^{-1}) = \Gamma(\gamma_i) + \Gamma(\tilde{\gamma}) + \Gamma(\gamma_j^{-1}) = \Gamma(\gamma_i) + \Gamma(\gamma_j^{-1})$$

and hence $\Gamma(\gamma_i) = \Gamma(\gamma_j)$ contradicting the construction of the sequence $\{\gamma_k\}_{k \in \mathbb{N}}$.

References

- [Am01] B. Ammann, **Spectral estimates on 2-tori**, arXiv:math/0101061v1, 2001
- [Ba92] C. Bär, **Lower eigenvalue estimates for Dirac operators**, Math. Ann. **293**, 39-46, 1992
- [Ba96] C. Bär, **Metrics with Harmonic Spinors**, Geom. Funct. Anal. **6**, 1996, 899-942
- [Ba97] C. Bär, **Harmonic Spinors for Twisted Dirac Operators**, Math. Ann. **309**, 1997, 225-246, arXiv:dg-ga/9706016v1
- [BLP05] B. Boock-Bavnbek, M. Lesch, J. Phillips, **Unbounded Fredholm Operators and Spectral Flow**, Canad. J. Math. **57**, 2005, 225-250, arXiv:math/0108014v3
- [Da08] M. Dahl, **On the Space of Metrics with Invertible Dirac Operator**, Comment. Math. Helv. **83**, 2008, 451-469, arXiv:math/0603018v1
- [DG12] M. Dahl, N. Grosse, **Invertible Dirac operators and handle attachments on manifolds with boundary**, arXiv:1203.3637v1
- [Hij01] O. Hijazi, **Spectral properties of the Dirac operator and geometrical structures**, Proceedings of the Summer School on Geometric Methods in Quantum Field Theory, Villa de Leyva, Colombia, July 12-30, (1999), 2001
- [Hi74] N. Hitchin, **Harmonic Spinors**, Adv. Math. **14**, 1-55, 1974

[Phi96] J. Phillips, **Self-Adjoint Fredholm Operators and Spectral Flow**, *Canad. Math. Bull.* **39**, 1996, 460-467

Dipartimento di Scienze Matematiche
Politecnico di Torino
Corso Duca degli Abruzzi, 24
10129 Torino, Italy
E-mail: waterstraat@daad-alumni.de