

Kent Academic Repository

Full text document (pdf)

Citation for published version

Wildman, Scott S.P. and King, Brian F. (2008) P2X receptors: epithelial ion channels and regulators of salt and water transport. *Nephron Physiology*, 108 (3). p60-67. ISSN 1660-8151.

DOI

<https://doi.org/10.1159/000122028>

Link to record in KAR

<http://kar.kent.ac.uk/53053/>

Document Version

Publisher pdf

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

P2X Receptors: Epithelial Ion Channels and Regulators of Salt and Water Transport

Scott S.P. Wildman^a Brian F. King^b

^aDepartment of Basic Veterinary Sciences, The Royal Veterinary College, and ^bDepartment of Physiology (Hampstead Campus), Centre for Nephrology, University College London, London, UK

Key Words

P2X receptor · Ligand-gated ion channel · Nucleotide · Adenosine 5'-triphosphate · Epithelial cells, renal · ATP · P2Y

Abstract

When the results from electrophysiological studies of renal epithelial cells are combined with data from *in vivo* tubule microperfusion experiments and immunohistochemical surveys of the nephron, the accumulated evidence suggests that ATP-gated ion channels, P2X receptors, play a specialized role in the regulation of ion and water movement across the renal tubule and are integral to electrolyte and fluid homeostasis. In this short review, we discuss the concept of P2X receptors as regulators of salt and water salvage pathways, as well as acknowledging their accepted role as ATP-gated ion channels.

Copyright © 2008 S. Karger AG, Basel

Ion Channels in Polarized Renal Epithelial Cells

Polarized renal epithelial cells serve to maintain electrolyte and fluid balance in the body by regulating the movement of ions and water across the surface of the kidney tubule. Epithelial cells are involved in either the net absorption or secretion of ions, or absorption of water – a role dependent on the concerted actions of channels,

co-transporters, exchangers and pumps that are selectively distributed to the apical and basolateral membranes. These primary structures are regulated, in turn, by a complex series of intracellular and extracellular signaling pathways which come under genomic and non-genomic control. Voltage-independent ion channels (e.g., ENaC, ROMK1 and CFTR), voltage-dependent ion channels (e.g., the CLC family) and cAMP-dependent water channels (i.e., the AQP protein family) exert a considerable influence on net ion and fluid transport. Their importance in solute and water salvage is apparent from a number of inheritable human diseases that disrupt the expression of normal channels in the kidney (e.g., Liddle's and Dent's diseases, cystic fibrosis, Bartter's syndrome types II and III, pseudohypoaldosteronism type I, autosomal dominant polycystic kidney disease type II, and autosomal nephrogenic diabetes insipidus).

Voltage-independent and voltage-dependent ion channels are not the only types of ion channels found in epithelial cells. Additionally, a number of ligand-gated ion channels (LGICs) have been identified and they are activated by substances more commonly associated with neurotransmission (e.g., NMDA, GABA_A and nACh receptors). This list of LGICs now includes the surface receptors known as P2X receptors, which are activated principally by extracellular adenosine 5'-triphosphate (ATP). Experimental observations have led us to propose that P2X receptors regulate the opening and trafficking

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2008 S. Karger AG, Basel
1660–2137/08/1083–0060\$24.50/0

Accessible online at:
www.karger.com/nep

Dr. Scott S.P. Wildman
Department of Veterinary Basic Sciences, Royal Veterinary College
Royal College Street
London NW1 0TU (UK)
Tel. +44 20 7121 1903, Fax +44 20 7468 5204, E-Mail swildman@rvc.ac.uk

of ion and water channels in the kidney tubule. Evidence for this advanced role is also supported by findings from *in vivo* microperfusion of the kidney tubule with ATP and its analogues, which have been shown to alter the degree of electrolyte and water salvage in salt-depleted animals. In this short review, we discuss a novel role of P2X receptors as regulators of epithelial cell transport, in addition to their widely accepted role as membrane pathways for Ca^{2+} and Na^{+} influx.

Introduction to P2X Receptors

P2X receptors represent the LGIC subdivision of the P2 receptor family and are activated by purine nucleotides such as ATP, GTP and, to a lesser extent, by related pyrimidine nucleotides such as CTP and UTP. Another subdivision of P2 receptors is represented by the G protein-coupled P2Y receptors, of which there are eight members – P2Y_{1,2,4,6,11,12,13,14} receptors – activated by purine or pyrimidine nucleotides (and sometimes both). The function of P2Y receptors in polarized epithelial cells has been discussed in an earlier review [1].

Seven genes encode the P2X receptor subunits in mammals; otherwise, simpler vertebrates (e.g., zebra fish) may possess up to nine genes. Examples of P2X receptor genes have also been found in invertebrates [2]. In heterologous expression systems, the seven mammalian subunits form functional homomeric assemblies (called P2X₁₋₇; fig. 1). Seven functional heteromeric assemblies formed by the union of two different P2X subunits (P2X_{1/2,1/4,1/5,2/3,2/6,4/6,4/7}) have also been established from a larger cohort of 13 heteromeric assemblies (that further includes P2X_{1/3,1/6,2/5,3/5,4/5,5/6}) first identified in immunoprecipitation studies of co-expressed subunits. Established P2X receptor subtypes have distinct pharmacological and operational profiles but, fundamentally, all are ATP receptors and all are potently activated by this energy molecule (fig. 1). There is a paucity of selective agonists and antagonists for each of the known P2X receptor subtypes and, accordingly, establishing a correspondence between native P2X receptors and known recombinant P2X receptor subtypes is, in equal measure, arduous and laborious. Such studies typically require the careful cross-correlation of the potency and efficacy of agonists at native and recombinant P2X receptors, their modulation by a range of substances and by the capacity of a limited number of antagonists to block P2X receptor subtypes (table 1).




Fig. 1. P2X receptors. The P2X receptors comprise a series of seven homomeric assemblies (P2X₁–P2X₇) and a second series of seven heteromeric assemblies (P2X_{1/2}, P2X_{1/4}, P2X_{1/5}, P2X_{2/3}, P2X_{2/6}, P2X_{4/6} and P2X_{4/7}; dark blue circles). A further series of heteromeric assemblies has been identified from immunoprecipitation studies, but none has yet been established as functional ion channels (pale blue circles). The series of P2X₁–P2X₇ receptors have different affinities for ATP which, as shown in the accompanying graph, is more potent at certain assemblies. The potency indices for ATP are given in table 1.

It is generally accepted that P2X receptor assemblies are comprised of 3 glycosylated subunits [2]. Accordingly, three ATP molecules are required for receptor activation. Each subunit protein has two transmembrane-spanning domains, with intracellular N and C termini and a highly folded extracellular loop held together by a number of disulfide bridges. The binding sites for ATP are thought

Table 1. Homomeric P2X receptor subtypes

	P2X ₁	P2X ₂	P2X ₃	P2X ₄	P2X ₅	P2X ₆	P2X ₇
Gene (human)	<i>17p13</i>	<i>12q24.33</i>	<i>11q12</i>	<i>12q24.32</i>	<i>17p13</i>	<i>22q11.21</i>	<i>12q24.31</i>
Protein (rat)	399 aa	472 aa	397 aa	388 aa	445 aa	379 aa	595 aa
Natural ligands (pEC ₅₀)	ATP (7.0) CTP (4.4) Ap ₆ A (6.0) Ap ₆ G (5.7)	ATP (5.3) Ap ₄ A (4.8)	ATP (5.9) UTP (4.0) Ap ₆ A (5.9) Ap ₃ A (6.0)	ATP (5.4) CTP (3.5) Ap ₄ A (5.5) ^a	ATP (6.4) GTP (4.6) Ap ₄ A (6.6)	ATP (6.3)	ATP (3.4) ADP (2.7) (m) ^b
Other agonists (pEC ₅₀)	2MeSATP (7.0) ATPγS (6.2) αβmeATP (5.5) BzATP (4.6)	2MeSATP (5.1) ATPγS (5.1) αβmeATP (3.0)	2MeSATP (6.7) ATPγS (5.9) αβmeATP (5.7) BzATP (7.1)	2MeSATP (3.6) ATPγS (~5.0) αβmeATP (4.2) ^a	2MeSATP (6.4) ATPγS (6.5) αβmeATP (6.0) BzATP (5.9)	αβmeATP (6.2)	2MeSATP (5.0) BzATP (5.2)
Antagonist – selective (pIC ₅₀)	NF449 (9.5) Ip ₅ I (8.5)		A317491 (7.6)				A740003 (7.75) BBG (8.0) KN-62 (7.4) (h) ^b
Antagonist – nonselective (pIC ₅₀)	TNP-ATP (9.0) PPADS (6.9) Suramin (5.7) RB-2 (5.7)	TNP-ATP (5.9) PPADS (5.8) Suramin (5.0) RB-2 (6.4)	TNP-ATP (9.5) PPADS (6.7) Suramin (5.4) RB-2 (4.3)	TNP-ATP (4.8) PPADS (<4.0) Suramin (<4.0) BBG (3.9)	TNP-ATP (6.3) PPADS (6.7) Suramin (5.8) RB-2 (4.7)	TNP-ATP (6.1) PPADS (4.7)	TNP-ATP (~4.3) PPADS (4.3) Suramin (<4.0)
Signaling	Cation channel P _{Ca} /P _{Na} ~4	Cation channel P _{Ca} /P _{Na} ~2	Cation channel P _{Ca} /P _{Na} ~4	Cation channel P _{Ca} /P _{Na} ~4	Cation channel P _{Ca} /P _{Na} ~1.5	Cation channel ND	Channel-to-pore Pass size <700 Da

The pharmacological and signaling properties of homomeric P2X receptors, listing the natural and synthetic agonists and selective and nonselective antagonists for each P2X receptor subtype. Data are given as $-\log_{10}$ EC₅₀ (pEC₅₀) and $-\log_{10}$ IC₅₀ (pIC₅₀). Data are given for rat isoforms. Adapted from Bailey et al. [35].

^a Data are given for partial agonists in some cases.

^b Data are also given for human (h) and mouse (m) isoforms in a few cases.

to lie between (rather than within) the extracellular loops. The integral ion channel formed by the close association of six transmembrane-spanning domains (i.e., two from each of three subunits) is permeable to Na⁺, K⁺ and Ca²⁺ ions and, in a few examples, also to Cl⁻ ions. Upon activation, P2X receptors open within milliseconds and cause cell depolarization by a predominant cationic influx. Via Ca²⁺ influx, P2X receptors also activate intracellular signaling cascades (including transcription pathways) and can regulate the activity of a broad series of intracellular kinases [3]. Ca²⁺ influx makes up some 3–15% of the total current carried by P2X receptor ion channels [4].

The current carrying capacity differs between the known P2X receptor subtypes. Some (including P2X₁, P2X_{1/2}, P2X_{1/5} and P2X₃) activate, then rapidly inactivate, in the continued presence of extracellular ATP. These P2X subtypes can produce membrane currents that are large in amplitude yet brief in duration, and they are referred to as ‘rapidly desensitizing’ P2X receptors. They may serve a role for intermittent, yet intense, cell signaling. Others (including P2X₂, P2X_{2/6}, P2X₄ and

P2X_{4/6}) activate, then slowly inactivate, over a longer period of time (10–100 s), and these are referred to as ‘slowly desensitizing’ P2X receptors. Such P2X subtypes may play a role in prolonged, and probably tonic, cell signaling. Lastly, the activated P2X₇ receptor, in sharp contrast to all other P2X subtypes, progresses to a non-inactivating conductance state that involves the opening of an ancillary membrane channel [5]. This may bring about profound changes to the cell and may limit its lifespan.

There are also striking differences in the amplitude of currents carried by P2X receptors, a phenomenon related both to receptor number and the efficiency of receptor insertion into the surface membrane. Homomeric P2X₅ and P2X₆ receptors carry only small amplitude membrane currents, because these assemblies traffic poorly to the cell surface and numbers there are low. On the other hand, P2X₁, P2X₂, P2X₃ and P2X₄ receptors yield large amplitude membrane currents because they are transported more efficiently to the cell surface. Of note, some P2X subtypes (P2X₂, P2X_{2/3}, P2X₄, P2X₅ and P2X₇) undergo a time-dependent change in ion channel con-

ductance that alters the pass size of ion channel permeants – a phenomenon called pore dilatation. P2X₇ receptors first undergo pore dilatation and, subsequently, activate a much larger ancillary pore [5].

P2X Receptor Distribution

Immunohistochemical, RT-PCR and pharmacological studies have demonstrated the presence of P2X receptors in all types of cells, including neurons, glia, muscle, bone and hemopoietic tissues, endothelial and epithelial cells [6]. In spite of the widespread distribution of P2X subunits in cells and tissues, much of the research on P2X receptors has focused on just a few key roles, such as smooth muscle contraction (involving P2X₁ and P2X₅ subunits), neuronal excitation (involving P2X₂, P2X₃, P2X₄ and P2X₆ subunits), inflammation and cell death (involving P2X₇ subunits). With gene targeting in mice, P2X receptor subtypes have been linked to less traditional roles such as autoregulation of renal blood flow (P2X₁^{-/-} data), to pain, visceral sensation, ventilatory responses during hypoxia (P2X₂^{-/-} and P2X₃^{-/-} data), bone growth and macrophage function (P2X₇^{-/-} data) [7].

Investigators have reported the surface expression of nearly all P2X (and P2Y) proteins in epithelial cells of the kidney, lung, gastrointestinal tract and accessory exocrine glands [8–11]. P2X (and P2Y) receptors occur on the apical and basolateral membranes of native epithelial cells and related cell lines, with some evidence for polarity in their distribution [12]. Investigators have also shown that ATP can be released from epithelial cells *in vitro* by a series of stimuli such as membrane distortion, osmotic swelling, hypoxia and acidosis [13]. ATP extrusion pathways have been linked to the CFTR protein, to P-glycoprotein-based organic anion transporters, stretch-activated cationic channels (and Ca²⁺-dependent exocytosis of ATP) and also to hemiconnexin channels [7, 14]. In turn, regulated ATP release has been linked to disparate cell functions, such as cell defense, hydration of the mucus barrier, enhanced ciliary beating, cell volume regulation, ischemic protection and cell division [10].

P2X Receptors in the Kidney

All seven P2X subunits have been localized by immunohistochemical techniques in kidney tissue, with identifiable patterns of distribution along the rat nephron [11, 15, 16]. P2X_{1–3} subunits have been found mainly in vas-

cular smooth muscle, while P2X₂ and P2X₇ subunits were found in endothelial and mesangial cells, as well as podocytes. High levels of P2X₄ and P2X₆ proteins occur in renal tubule cells, sometimes accompanied by relatively low levels of P2X₁, P2X₂, P2X₅ and P2X₇ protein. The variety and abundance of P2X subunits is greatest in the distal segments of the renal tubule, with P2X_{1,2,4,5,6,7} subunits present in the apical membrane of collecting duct (CD) principal cells in the rat, and P2X₄, P2X₆ and P2X₇ subunits in the basolateral membrane. Using real-time PCR, mRNA levels in the CD are highest for P2X₄, with P2X₆ some 10-fold lower [16]. Transcripts for P2X₁, P2X₂ and P2X₄ are found in similar quantities as P2X₆ in the distal tubule. In salt-depleted rats (maintained on 0.01% NaCl for 8–10 days), there is a significant upregulation of P2X₁, P2X₄ and P2X₆ transcripts in CD cells [16].

With respect to the vasculature, activation of P2X receptors typically causes vasoconstriction. This applies especially to the afferent arteriole, where ATP in the interstitium and probably originating from the macula densa can profoundly regulate glomerular perfusion and intraglomerular pressure. With gene deletion in P2X₁^{-/-} mice, this effect is largely ablated [17]. Perfusion of the renal vasculature with nucleotides results in both short- and long-lasting increases in the muscle tone of blood vessels. This dual action points to heterogeneity amongst the endogenous P2X receptors. By itself, the rapidly desensitizing P2X₁ cannot comfortably explain the long-term actions of perfused nucleotides and, more likely, a mixture of homomeric and heteromeric P2X receptors are involved. The molecular identity of P2X receptors responsible for tonic vasoconstriction is a key area for future research.

In the nephron, nucleotides in the luminal fluid may reach sufficiently high concentrations to activate P2X receptors on the apical membrane [18]. At one time, it was suggested that luminal nucleotides took their origin from the blood and were filtered freely by the glomerulus, thereafter to be concentrated in the tubular fluid. However, kidney micropuncture studies support an alternative hypothesis that principally involves ATP secretion in the proximal convoluted tubule with a minimal contribution towards the total ATP concentration by glomerular filtration [19]. The molecular basis of tubular ATP secretion is presently unknown, but mechanoreception via the central cilium may play a key role here [20].

For a number of reasons, most reports on the regulation of ion transport by extracellular ATP have focused on the involvement of P2Y receptors, to the exclusion P2X receptors. Traditionally, signaling by P2X receptors has




Fig. 2. Extracellular ATP actions in renal epithelium. A summary of effects of extracellular ATP in the nephron and epithelial cell lines derived from specific nephron segments [15, 16, 27, 29, 32]. In most cases, the activated P2R subtypes have not been identified. Many of the effects evoked by extracellular ATP in distal nephron epithelia are attributed to the activation of P2X receptors.

been associated with short-term phenomena and, accordingly, transient effects have been considered to have no practicality in the renal tubule. P2Y receptors, on the other hand, appear to be abundant in the apical membrane of the renal tubule, especially P2Y₂-like receptors in mouse tissues and mouse-derived cell lines which are used commonly in renal research. Consistently, signaling by P2Y₂ receptors has been associated with profound and long-term actions wherever they are found. For example, P2Y₂ receptors have been associated with cell growth in

epithelial and non-epithelial cells and, in part, found to exert their effects genomically. However, our immunohistochemical data (unpublished) from rat and human renal tubules place P2Y₂ receptors primarily on intercalated cells of the CD [11, 16] or the basolateral membrane of tubules (unpublished). We can say with some authority that, instead, P2Y₄ receptors appear in the apical membrane of principal cells of rat CDs [16]. However, P2Y₂ and P2Y₄ receptors are almost identical in their pharmacological profiles as far as rodent isoforms are concerned

[21, 22] and it is easy to see why P2Y₂ receptors were implicated in past studies. The involvement of apical P2Y₂ receptors in regulating kidney function probably has been overrepresented through misguided pharmacology. The actions of extracellular ATP in the renal epithelium are listed in figure 2.

Many of the effects evoked by extracellular ATP in the distal nephron (and derived cell lines) can be attributed to the activation of P2X receptors, particularly where transport of Mg²⁺, Na⁺ and Cl⁻ ions is concerned [23–27] and movement of water is considered [28, 29]. For Mg²⁺ transport, P2X receptor activation in a mouse distal convoluted tubule cell line resulted in the inhibition of magnesium absorption by a presumed action on TRPM6 channels [24]. This inhibition occurred via an increase in intracellular Ca²⁺ ions which may enter through as yet unidentified P2X receptors involving P2X₁, P2X₂, P2X₃, P2X₄ and P2X₅ subunit proteins which were found in this cell line [24]. Similarly, P2X-mediated calcium influx involving P2X₁ and/or P2X₄ subunits has been reported in isolated perfused mouse medullary CD preparation [27]. Further to these observations, it should be noted that P2X₁ subunits are capable of polymerization with P2X₂, P2X₃, P2X₄ and P2X₅ subunits, to form a number of subtypes of P2X receptors.

For Na⁺ transport, tubular perfusion of ATP inhibits Na⁺ uptake in the isolated mouse CCD even though ENaC channels are insensitive to nucleotides [30]. Several independent studies link this phenomenon in part to P2X receptor activation. First, Na⁺ absorption and Cl⁻ secretion appears to involve P2X₃ and P2X₄ receptors in mIMCD-K2 cells, an immortalized mouse CD cell line [23]. Second, the recovery of ²²Na⁺ ions in the urine of Na⁺-restricted living rats is increased in distal tubules microperfused with the nonhydrolysable nucleotide, ATPγS [25]. This effect was linked to the activation of P2X_{4/6} receptors [31]. Third, the sequential co-expression of rat ENaC channels with rat P2X receptor subtypes in *Xenopus* oocytes showed that a single ATP challenge irreversibly inhibited ENaC channels when either P2X₂, P2X₄, P2X_{2/6} or P2X_{4/6} receptors were activated [28]. Inhibition occurred by trafficking ENaC subunits out of the cell membrane and, mechanistically, was associated with Na influx through P2X₂ and P2X_{2/6} receptors and Na⁺ and Ca²⁺ influx through P2X₄ and P2X_{4/6} receptors. Recently, patch clamp experiments in the microdissected CD of the rat kidney showed that nucleotide activation of P2X₄ and possibly P2X_{4/6} receptors on the apical surface of principal cells in situ can either inhibit or potentiate an amiloride-sensitive inward current, presumably involv-

ing ENaC channels, depending on the concentration of extracellular sodium [16]. These extended observations have led to the proposal that P2X₄ and P2X_{4/6} receptors may function as sodium sensors responsible for local regulation of ENaC activity in the rat CD in response to changes in luminal sodium concentration. Potentiation of ENaC-mediated currents by P2X₄ receptors, albeit those expressed on the basolateral membrane, has also been reported in A6 cells [32].

For the movement of water, sequential co-expression of rat AQP2 channels with known rat P2X receptor subtypes in *Xenopus* oocytes showed a diminished responsiveness in a swelling assay, but only when P2X₂ receptors were present, and they were activated by ATP [28]. Furthermore, dDAVP treatment of cultured mpkCCD(c14) cells not only increased AQP2-positive immunostaining, but also P2X₁ and P2X₂ subunit immunostaining in the apical membrane; secondly, activation of these apical P2X receptors with ATPγS removed immunopositive AQP2 protein from the apical membrane [29]. This apart, P2Y₂ receptors on the basolateral membrane of principal cells are also known to regulate water movement in the CD.

Are P2X Receptors Also Directly Involved in the Net Transport of Solutes?

It is not unreasonable to suppose that, as apical membrane channels, P2X receptors can assist in transporting sodium and calcium across the kidney tubule. We have calculated the charge transfer during the maximal activation of P2X₂ receptors and estimated the net increase in intracellular sodium to be 0.5 μmol/min for a model cell (*Xenopus* oocyte, 1 μl cell volume) [26]. We have calculated the charge transfer for activated ENaC channels, which yielded tonic inward currents of comparable amplitude to that evoked by P2X₂ receptors, and estimated the net increase in intracellular sodium to be 5 μmol/min. Even if channel numbers were matched in vivo, the lower amount of sodium carried by P2X₂ receptors compared to ENaC channels suggests a minor role, at best, for the former in sodium transport.

P2X Receptors as Regulators of Other LGICs

Functional interactions between P2X receptors and other LGICs have been described in excitable tissues and in expression systems [2]. Thus far, cross-inhibition ap-

appears to be the norm when P2X receptors are co-expressed, or co-localized, with members of the Cys-loop superfamily of LGICs including GABA_A, GABA_C, nACh and 5-HT₃ receptors. Use of fluorescence resonance energy transfer and total internal reflection fluorescence suggests that P2X₂ receptors and nACh (α4β2) receptors physically interact and the activation of one can lead to the internalization of both. As indicated above, it has also been postulated that TRPM6 channels are also regulated by P2X receptors, but the mechanism behind this regulation is unknown [24]. Otherwise, TRPV1 was shown to exert a one-way inhibition on P2X_{2/3} and P2X₃ receptors again by an unknown mechanism [33, 34]. As yet, it remains to be shown how and where channel–channel interactions occur naturally in the kidney tubule.

Conclusions

For the kidney, the role of P2X receptors in polarized epithelial cells has been largely neglected. However, our research on kidney cell function over the past 5 years, including our supporting work of modeling in expression

systems, now points to a regulatory role of P2X receptors in the movement of salt and water across the wall of the distal tubule and CDs. We find no credible evidence for a role of P2X receptors in punctate cell signaling, which is the conventional role for P2X receptors outside the kidney and in excitable tissues. Instead, P2X receptor activation and resultant cationic currents appear to change the rate of internalization and reclamation of salt and water channels. Uppermost in our minds is the question of the physiological relevance of these phenomena. Nature, typically, is economical; complex mechanisms tend to evolve into efficient regulatory processes rather than linger as redundant systems. Thus, we are working towards a better understanding of this regulatory mechanism and, in writing this review, we hope that our opinions will stimulate a debate and lead to new research directions.

Acknowledgments

We gratefully acknowledge the financial support of the British Heart Foundation and the St Peter's Trust for Kidney, Bladder and Prostate Research.

References

- Dubyak GR: Knock-out mice reveal tissue-specific roles of P2Y receptor subtypes in different epithelia. *Mol Pharmacol* 2003;63:773–776.
- Nicke A, King BF: Heteromerization of P2X receptors; in Arias HR (ed): *Biological Aspects of Ligand-Gated Ion Channel Receptor Superfamilies*. Kerala, Research SignPost, 2006, pp 383–417.
- North RA: Molecular physiology of P2X receptors. *Physiol Rev* 2002;82:1013–1067.
- Egan TM, Khakh BS: Contribution of calcium ions to P2X channel responses. *J Neurosci* 2004;24:3413–3420.
- Pelegrin P, Surprenant A: Pannexin-1 couples to maitotoxin- and nigericin-induced interleukin-1β release through a dye uptake-independent pathway. *J Biol Chem* 2007;282:2386–2394.
- Ralevic V, Burnstock G: Receptors for purines and pyrimidines. *Pharmacol Rev* 1998;50:413–492.
- Boeynaems JM, Communi D, Gonzales NS, Robaye B: Overview of the P2 receptors. *Semin Thromb Hemost* 2005;31:139–149.
- Taylor AL, Schwiebert LM, Smith JL, King C, Jones JR, Sorscher EJ, Schwiebert EM: Epithelial P2X purinergic receptor channel expression and function. *J Clin Invest* 1999;104:875–884.
- Bucheimer RE, Linden J: Purinergic regulation of epithelial transport. *J Physiol* 2003;555:311–321.
- Leipziger J: Control of epithelial transport via luminal P2 receptors. *Am J Physiol Renal Physiol* 2003;284:F419–F432.
- Turner CM, Vonend O, Chan C, Burnstock G, Unwin RJ: The pattern of distribution of selected ATP-sensitive P2 receptor subtypes in normal rat kidney: an immunohistological study. *Cells Tissues Organs* 2003;175:105–117.
- Wolff SC, Qi AD, Harden TK, Nicholas RA: Polarized expression of human P2Y receptors in epithelial cells from kidney, lung, and colon. *Am J Physiol Cell Physiol* 2005;288:C624–C632.
- Schwiebert EM, Zsembery A: Extracellular ATP as a signaling molecule for epithelial cells. *Biochim Biophys Acta* 2003;1615:7–32.
- Lazarowski ER, Boucher RC, Harden TK: Mechanisms of release of nucleotides and integration of their action as P2X- and P2Y-receptor activating molecules. *Mol Pharmacol* 2003;64:785–795.
- Unwin RJ, Bailey MA, Burnstock G: Purinergic signaling along the renal tubule: current state of play. *News Physiol Sci* 2003;18:237–241.
- Wildman SS, Turner CM, Marks J, Shirley DG, King BF, Wang W, Unwin RJ: Sodium-dependent regulation of ENaC by apical P2 receptors in rat renal collecting duct. *J Am Soc Nephrol* 2008, Jan. 30, E-pub.
- Inscho EW, Cook AK, Imig JD, Vial C, Evans RJ: Renal autoregulation in P2X₁ knockout mice. *Acta Physiol Scand* 2004;181:445–453.
- Schwiebert EM: ATP release mechanisms, ATP receptors and purinergic signalling along the nephron. *Clin Exp Pharmacol Physiol* 2001;28:340–350.
- Vekaria R, Unwin RJ, Shirley DJ: Intraluminal ATP concentrations in rat renal tubules. *J Am Soc Nephrol* 2006;17:1841–1847.
- Hovater MB, Olteanu D, Hanson EL, Cheng NL, Siroky B, Fintha A, Komiosi P, Liu W, Satlin LM, Bell PD, Yoder BK, Schwiebert EM: Loss of apical monocilia on collecting duct principal cells impairs ATP secretion across the apical cell surface and ATP-dependent and flow-induced calcium signals. *Purinergic Signal* 2008, in press.

- 21 Lazarowski ER, Rochelle LG, O'Neal WK, Ribeiro CM, Grubb BR, Zhang V, Harden TK, Boucher RC: Cloning and functional characterization of two murine uridine nucleotide receptors reveal a potential target for correcting ion transport deficiency in cystic fibrosis gallbladder. *J Pharmacol Exp Ther* 2001;297:43–49.
- 22 Wildman SS, Unwin RJ, King BF: Extended pharmacological profiles of rat P2Y₂ and rat P2Y₄ receptors and their sensitivity to extracellular H⁺ and Zn²⁺ ions. *Br J Pharmacol* 2003;140:1177–1186.
- 23 McCoy DE, Taylor AL, Kudlow BA, Karlson K, Slattery MJ, Schwiebert LM, Schwiebert EM, Stanton BA: Nucleotides regulate NaCl transport in mIMCD-K2 cells via P2X and P2Y purinergic receptors. *Am J Physiol* 1999; 277:F552–F559.
- 24 Dai LJ, Kang HS, Kerstan D, Ritchiw G, Quamme GA: ATP inhibits Mg²⁺ uptake in MDCT cells via P2X purinoceptors. *Am J Physiol Renal Physiol* 2001;281:F833–F840.
- 25 Shirley DG, Bailey MA, Unwin RJ: In vivo stimulation of apical P2 receptors in collecting ducts: evidence for inhibition of sodium reabsorption. *Am J Physiol Renal Physiol* 2005;288:F1243–F1248.
- 26 Wildman SS, Marks J, Churchill LJ, Peppiatt CM, Chraibi A, Shirley DG, Horisberger JD, King BF, Unwin RJ: Regulatory interdependence of cloned epithelial Na⁺ channels and P2X receptors. *J Am Soc Nephrol* 2005;16: 2686–2597.
- 27 Li L, Lynch IJ, Zheng W, Cash MN, Teng X, Wingo CS, Verlander JW, Xia S-L: Apical P2XR contribute to [Ca²⁺]_i signaling and I_{sc} in mouse renal MCD. *Biochem Biophys Res Commun* 2007;359:438–444.
- 28 Wildman SS, Peppiatt CM, Boone M, Konings I, Marks J, Churchill LJ, Turner CM, Shirley DG, King BF, Deen PMT, Unwin RJ: Potential role of apical P2 receptors in modulating aquaporin-2-mediated water reabsorption in the collecting duct. *Proc Physiol Soc* 2006;2:PC4.
- 29 Wildman SS, Boone M, Peppiatt CM, Churchill LJ, Shirley DG, King BF, Deen PMT, Unwin RJ: Immunofluorescent labeling reveals dDAVP-dependent P2 receptor expression and apical P2 receptor-mediated inhibition of AQP2 expression in mpkCCD(cl4) cultures. *Proc Physiol Soc* 2006;3:PC3.
- 30 Lehrmann H, Thomas J, Kim SJ, Jacobi C, Leipziger J: Luminal P2Y₂ receptor-mediated inhibition of Na⁺ absorption in isolated perfused mouse CCD. *J Am Soc Nephrol* 2002;13:10–18.
- 31 Wildman SS, Shirley DG, King BF, Unwin RJ: Indirect evidence for regulation of sodium reabsorption in sodium-restricted rats by an ATP-gated heterotrimeric P2X receptor. *J Physiol* 2004;558P:32–33.
- 32 Zhang Y, Sanchez D, Gorelik J, Klenerman D, Lab M, Edwards C, Korchev Y: Basolateral P2X₄-like receptors regulate the extracellular ATP-stimulated epithelial Na⁺ channel activity in renal epithelia. *Am J Physiol Renal Physiol* 2007;292:F1734–F1740.
- 33 Piper AS, Docherty RJ: One-way cross-desensitization between P2X purinoceptors and vanilloid receptors in adult rat dorsal root ganglion neurones. *J Physiol* 2000;523: 685–696.
- 34 King BF, Liu M, Townsend-Nicholson A, Burnstock G: Inhibitory interaction between activated TRPV1 and P2X₃ receptors. *Proc Physiol Soc* 2006;3:C35.
- 35 Bailey MA, Shirley DG, King BF, Burnstock G, Unwin RJ: Extracellular nucleotides and renal function; in Alpern RJ, Hebert SC (eds): Seldin and Geibisch's *The Kidney: Physiology and Pathophysiology*, ed 4. Amsterdam, Elsevier, 2007, pp 477–496.