

Kent Academic Repository

Full text document (pdf)

Citation for published version

Winter, Samantha L. and Burnley, Mark and Forrest, Sarah M. and Challis, John H. (2015) Complexity of isometric force production is associated with time to achieve steady state when moving to a new force level. In: 25th International Society of Biomechanics Conference, July 12-16, 2015, Glasgow, UK.

DOI

Link to record in KAR

<http://kar.kent.ac.uk/53127/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

ISB 2015

Neurological and Motor Control

ISB 2015-1036

COMPLEXITY OF ISOMETRIC FORCE PRODUCTION IS ASSOCIATED WITH TIME TO ACHIEVE STEADY STATE WHEN MOVING TO A NEW FORCE LEVEL

Samantha L. Winter¹, Mark Burnley¹, Sarah Forrest², John Challis³

¹School of Sport and Exercise Sciences, University of Kent, Chatham Maritime, ²Sport and Exercise Science, Aberystwyth University, Aberystwyth, United Kingdom, ³Department of Kinesiology, The Pennsylvania State University, State College, United States

Preferred Presentation: Oral Presentation

If your abstract is not accepted as an oral do you wish to be considered for a poster?: Yes

Clinical Biomechanics Award: No

David Winter Young Investigator Awards: No

Emerging Scientific Award sponsored by Professor J De Luca: No

Promising Scientist Award sponsored by Motion Analysis: No


Introduction and Objectives: Motor output and function has been shown to decline in later life, for example older adults have increased stimulus response times [1]; reduced strength [2] and changes in the variability of force production [2]. The loss of complexity hypothesis [3] suggests that a decrease in complexity is associated with ageing as the integration of complex physiological systems declines, and that this reduced complexity is associated with constrained behaviour and a loss of adaptability or flexibility in response. The functional consequences of this loss of complexity in the fluctuations seen in isometric force records have not been investigated. However, it is possible that the reduced complexity, and more slowly repeating and constrained fluctuations in force seen in older adults [4] would limit ability to adapt to altered force requirements. It was hypothesised that reduced complexity for isometric force records in older adults would be associated with an increase in the time taken to achieve a steady state when moving to a new isometric force level.

Methods: Young (18 to 25 years) and older (65 to 72 years) neurologically healthy adults produced isometric abduction contractions of the index finger using the First Dorsal Interosseus. Participants established a steady state force at a given effort level (a percentage of maximum voluntary contraction force - %MVC) by matching a target displayed visually. A second feedback condition used both the visual display and an audible tone that varied in pitch when subjects moved below or above the target. The target was then suddenly and unpredictably increased or decreased at a randomly decided time point and participants were instructed to match this new target as quickly as they could, and to hold the new contraction intensity as steadily as possible (Figure 1). The conditions investigated were: increasing force from 5% to 25% MVC, 25% to 50% MVC, 25% to 75% MVC, and decreasing the force from 25% to 5% MVC, 50% to 75% MVC and 75% to 25% MVC. The functional reaction time was taken to be the point at which the force exceeded 3 standard deviations (SD) from the mean force in the direction of the new force level. The time to steady state was the time from the end of the functional reaction time to the start of the 3 second window with the lowest standard deviation for force. The complexity of the force record before the target change, after the target change, and during simple continuous force production with no target change was determined for the different effort levels (5%, 25%, 50%, 75% MVC) using Approximate Entropy (ApEn) [5].

Results: Confirming previous results older adults exhibited reduced complexity indicated by reduced ApEn values at effort levels >25% MVC ($p < 0.001$) and this was the case for both the simple continuous force task ($p < 0.001$) and the section of the force record after the change in target force ($p = 0.0438$). There was no difference in the MVC abduction force for older and younger adults ($p = 0.885$). There was also no difference between older and younger adults in the magnitude of the fluctuations in force, as quantified by the standard deviation of force ($p > 0.05$). Functional reaction time was increased by around a third in older adults ($p < 0.001$) in both feedback conditions. Functional reaction time was reduced for both age groups by the addition of the audible signal. Confirming the hypothesis, the time to steady state after the functional reaction time increased from a mean of 2.83s in younger adults to 3.23s for older adults ($p = 0.025$), and

this difference was unaffected by feedback condition. Post-hoc tests showed that the time to steady state differed for old and young adults at starting effort levels >5% MVC.

Figure:



Caption: Figure 1: Display of force record showing a trial moving from an initial target of 50% MVC to a target of 25% MVC.

Conclusion: Force levels associated with reduced complexity in older adults compared to younger adults were associated with an increased time to steady state when moving from an isometric contraction at one effort level to an isometric contraction at a new effort level. This suggests that there is a change in the neuromuscular system with age evidenced by the reduced complexity that constrains response to altered force requirements. These findings have implications for the control of muscle force production in later life.

References: [1] Fozard et al., *J. Gerontol.*, 49: P179-189, 1994.

[2] Marmon et al., *Med. Sci. Sports Exerc.*, 43: 560-567, 2011.

[3] Lipsitz et al., *J. Am. Med. Assoc.*, 267: 1806-1809, 1992.

[4] Vaillancourt et al., *J. Appl. Physiol.*, 94: 903-912, 2003.

[5] Pincus, *PNAS*, 88: 2297-2301, 1991.

Disclosure of Interest: None Declared