

Kent Academic Repository

Full text document (pdf)

Citation for published version

Richardson, Katharine and Hopker, James G. (2016) One minute to assess frailty, but what should we do next? *Anaesthesia*, 71 . pp. 622-626. ISSN 0003-2409.

DOI

<https://doi.org/10.1111/anae.13321>

Link to record in KAR

<http://kar.kent.ac.uk/53184/>

Document Version

Publisher pdf

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Editorial

One minute to assess frailty, but what should we do next?

The capacity to limit age-related diseases has been proposed as one of the main mechanisms responsible for successful ageing in old individuals. Indeed, age-related decline in physiological systems has been suggested to result in an increased likelihood of frailty occurring. Fried et al. [1] suggested patients with frailty present with at least three of the following; weight loss; exhaustion; a reduction in physical activity; and reduced mobility and grip strength. Factors that are associated with frailty include: age; comorbidities (e.g. hypertension); behavioural habits (e.g. exercise, smoking, & alcohol consumption) [2]; and those patients at both extremes of body mass [3]. The majority of these factors are potentially reversible, except for age and comorbidities, the latter of which could be managed or controlled. Frailty is a debilitating condition that reduces the quality of life in elderly populations [2], which is unsurprising, as it has previously been correlated with disability [3]. However, it is important to note that disability and frailty are different entities, and being disabled is not a prerequisite for frailty [4]. Nevertheless, frail individuals are often less able to

perform the activities of daily living that promote independence.

Pre-operative frailty in patients has been linked with increased postoperative morbidity [5], including delirium [6], sepsis [7], prolonged hospital stay [5] and postoperative re-admission [8], compared with non-frail counterparts. Moreover, Fried et al. [4] suggest that frail patients may also suffer from a 'vulnerability to stressors'. Indeed, the prospect of having surgery and accompanying medical examinations and investigations will be a considerable stressor to the frail patient, which may further exacerbate their level of 'vulnerability'. Even though it may be difficult to reverse the deleterious effects of severe frailty in patients before surgery, it has been shown that moderate or borderline frailty is responsive to a targeted intervention [9]. Therefore, identifying patients that are frail, or at risk of developing frailty postoperatively, is important. Specifically, the correct diagnosis of frailty in the pre-operative setting affords the opportunity to implement interventions aimed at optimising the patient before surgery.

The study of O'Neil et al. [10] in this issue of *Anaesthesia* investigated the utility of assessing pre-operative vascular patients for frailty via 'clinical impression'.

Three hundred and ninety-two patients were assessed in pre-assessment by a healthcare professional (i.e. consultant anaesthetist or specialist nurse) for their 'clinical impression' as to whether the patient was deemed to be 'fit for the proposed operation', and thus considered as being 'non frail'. Study follow-up on incidence of mortality demonstrated that patients deemed to be 'frail' had a 2.44 increased likelihood of dying at any given postoperative time point compared with those deemed 'not frail'. Moreover, a striking feature of the study was that 20% of the 'frail' patients had died at 13 months, compared with 34 months in the 'not frail' patients, suggesting pre-assessment clinical judgement was associated with mortality outcome. Despite these interesting results, it is questionable whether a 'clinical impression' truly identifies 'frail' and 'non-frail' patients. Unfortunately, O'Neil et al's study did not provide comparison with previously validated assessment tools e.g. the timed up and go test (TUG) [11]. Therefore, further research is required to identify whether a 'clinical impression' is able to provide a valid assessment of frailty. However, frailty aside, it is interesting that the 'clinical impression' did appear to identify those at risk of a poor postoperative

This editorial accompanies an article by O'Neill et al., *Anaesthesia* 2016; 71: doi: 10.1111/anae.13404.

outcome. By assessing postoperative complications alongside 'clinical impression' it might be possible to identify specific complications that 'frail' patients are more likely to experience. In turn, targeted peri-operative strategies could be put in place for this patient population.

Geriatric patients should be assessed for their level of frailty pre-operatively, according to both the American College of Surgeons National Surgical Quality Improvement Program and the American Geriatrics Society [12], and the Association of Anaesthetists of Great Britain and Ireland [13]. However, frailty has recently been described as 'the elephant in the operating room', as it is often not diagnosed or treated peri-operatively [14]. Indeed, despite the ease by which frailty can be assessed [11], and its association with postoperative outcome [5], it is not currently explicitly included within the National Institute for Health and Care Excellent (NICE) guidelines for pre-operative tests [15]. Therefore, routine assessment of frailty is not consistently implemented across UK pre-assessment centres. If we are failing to identify patients who are frail, it is difficult to formulate pre- and postoperative strategies to enhance the potential for a successful surgical outcome.

O'Neil et al. [10] clearly demonstrate the initial subjective evaluation of a patient has the potential to be a useful tool in evaluating frailty, and therefore those at heightened risk for postoperative mortality. The clinicians used by O'Neil et al. were all experienced at assessing vascular patients, thus

whether these findings are transferable to other clinicians that do not routinely assess this subset is questionable. In this regard Hubbard et al. [14] suggest that subjective measures of frailty may be limited by their inter-rater agreement. The assessment and diagnosis of frailty should be equally accessible for healthcare professionals who occasionally encounter frail patients, and those who do so on a routine basis (i.e. geriatricians) [14]. Thus, perhaps the combination of more objective measures, such as the TUG test, in addition to a 'clinical impression', might be more amenable for those clinicians who have less experience identifying frailty within patients.

Various risk factors based on past history, clinical examination and investigations, have been shown to correlate with postoperative outcome after surgery [16]. Moreover, previous literature has demonstrated a relationship between cardiorespiratory capacity and postoperative complications as measured by cardiopulmonary exercise testing (CPET) [17], and the six-minute walk test [18]. Cardiopulmonary exercise testing measures have been shown to be able to distinguish those patients at risk of specific postoperative complications (i.e. cardiac or respiratory) [17]. Barakat et al. [17] demonstrated that CPET-derived measures of anaerobic threshold and ventilatory equivalents for CO₂ ($\dot{V}E/\dot{V}CO_2$) are predictive of postoperative cardiac and respiratory complications in vascular patients. Patient data obtained from CPET during pre-assessment can in turn be used to

estimate survival and postoperative risk. One such survival calculator has recently been presented in *Anaesthesia* [19]. This survival calculator encompasses several pre-operative measures including: age; weight; height; blood results (i.e. creatinine, haemoglobin); previous medical history; and CPET findings. Carlisle et al. [19] demonstrated significantly lower five-year mortality in those patients stratified by the risk calculator as 'medium-to-high risk' in comparison with those classified 'high risk', with a hazard ratio of 0.58. This indicates a 42% reduced likelihood of mortality in 'medium-to-high risk' vs. 'high-risk' patients. Carlisle et al's data provide encouraging evidence as to the utility of risk calculations within the peri-operative setting, but unfortunately they do not include any specific measures of frailty. Including a measure of 'frailty' and/or 'clinical impression' as a variable within the risk calculation may therefore further improve its predictive ability. However, it is currently difficult to accurately quantify risk in elderly patients for several reasons: 1, the peri-operative literature within this population is sparse; 2, interpretation of the peri-operative literature is confounded by discrepancies in the tools used to identify postoperative outcome measures, and variability regarding the postoperative day at which outcome measures (i.e. complications) were recorded [20]; 3, most of the current tools to assess postoperative outcome measures are also not specific to the elderly; 4, hospital-specific outcome measures (e.g. length of stay, mortality,

complications, re-admission) may not be in line with what patients want to know before making a surgical decision, and elderly frail patients may be more inclined to assess the likelihood of being discharged to home/residential care, or other more patient-specific outcomes [20]. Addressing the limitations of the current literature to allow for a more accurate quantification of risk in the elderly, and refinement of mortality risk estimations, must be a key priority for future research in the area.

Following the identification of frailty and a medium-high level of risk in pre-assessment, it is important that strategies are put in place to improve a patient's likelihood for a beneficial postoperative outcome. Indeed, there are studies assessing the effect of pre-operative exercise training on the frail population that have demonstrated some reversibility of the condition [21,22]. When considering surgical patients, pre-operative exercise training is often termed 'prehabilitation' and is seen as an intervention designed to enhance functional capacity in anticipation of the forthcoming surgical insult. Hoogeboom et al. [21] demonstrated that in-hospital prehabilitation training (aerobic and strength based training, twice a week for three to six weeks) afforded clinically meaningful reductions in the time to perform the TUG test in frail patients undergoing orthopaedic surgery. Hoogeboom et al's study found a 4.4 second reduction in the time taken to do the TUG test, which is 3 seconds greater than the predefined clinically bene-

ficial reduction in the TUG test [23]. Unfortunately, Hoogeboom et al. [21] did not follow patients postoperatively to assess the impact of their training intervention on outcome measurements. A more recent case report [22] demonstrated a four-week prehabilitation programme (including moderate aerobic and resistance exercises with nutritional counselling) led to progressive and 'remarkable' improvements in the patient's level of frailty prior to his radical cystectomy. The case report also documents no adverse events during or after surgery, and a timely discharge on the seventh postoperative day. Combining this kind of multimodal approach to prehabilitation, with a similarly targeted rehabilitation programme after surgery, may afford a better long-term prognosis for the frail patient. Indeed, a 12-week post-operative resistance exercise programme (three times a week of progressive knee extension and leg press exercises) increased elderly hip replacement patient's stair walking power and muscle peak torque compared with a standard rehabilitation programme (one hour per day of functional exercises to improve mobility and strength without external loading) [24]. Rehabilitation may therefore also act to prevent those patients that are borderline frail from developing postoperative frailty, although this is yet to be explored.

By combining 'clinical impression' with simple objective tests such as the TUG, it could be possible to include a time-efficient assessment of frailty as part of rou-

tine pre-assessment clinical decision-making. The inclusion of a measure of frailty within pre-assessment has the potential to enable the planning of individualised patient optimisation strategies in both the pre- and postoperative stages. However, although there is some literature-based evidence of positive outcomes following the use of multi-modal prehabilitation and rehabilitation strategies in frail patients, more research is clearly required. Performing longitudinal and multicentre intervention studies using targeted prehabilitation and rehabilitation of frail patients may improve our understanding of how to optimise these patients both pre- and postoperatively, with the aim of improving the prognosis of a positive postoperative outcome.

Acknowledgements

No conflicts of interest declared.

K. Richardson

*Clinical Exercise Physiologist
Anaesthesia and Intensive Care
Medicine Medway Maritime
Hospital
Kent, UK*

J. Hopker

*Senior Lecturer
School of Sport and Exercise Sciences
University of Kent, UK
Email: j.g.hopker@kent.ac.uk*

This editorial accompanies an article by O'Neill et al., *Anaesthesia* 2016; 71: doi: 10.1111/anae.13404.

References

1. Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: evidence for a phenotype. *Journals of Gerontology. Series A, Biological Sciences and Medical Sciences* 2001; **56**: M146–56.

2. Lin CC, Li CI, Chang CK, et al. Reduced health-related quality of life in elders with frailty: a cross-sectional study of community-dwelling elders in Taiwan. *PLoS One*, 2011; **6**: e21841.
3. Fugate Woods N, LaCroix AZ, Gray SL, et al. Frailty: emergence and consequences in women aged 65 and older in the Women's Health Initiative Observational Study. *Journal of the American Geriatrics Society* 2005; **53**: 1321–30.
4. Fried LP, Ferrucci L, Darer J, Williamson JD, Anderson G. Untangling the concepts of disability, frailty, and comorbidity: implications for improved targeting and care. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 2004; **59**: M255–M263.
5. Makary MA, Segev DL, Pronovost PJ, et al. Frailty as a predictor of surgical outcomes in older patients. *Journal of the American College of Surgeons* 2010; **210**: 901–8.
6. Leung JM, Tsai TL, Sands LP. Brief report: preoperative frailty in older surgical patients is associated with early postoperative delirium. *Anesthesia and Analgesia* 2011; **112**: 1199–201.
7. Reisinger KW, van Vugt JL, Tegels JJ, et al. Functional compromise reflected by sarcopenia, frailty, and nutritional depletion predicts adverse postoperative outcome after colorectal cancer surgery. *Annals of Surgery* 2015; **261**: 345–52.
8. Arya S, Brahmabhatt R, Gander J, et al. Preoperative frailty independently increases risk of unplanned readmission and associated complications after peripheral vascular interventions. *Journal of the American College of Surgeons* 2014; **3**: S162.
9. Gill TM, Baker DI, Gottschalk M, Peduzzi PN, Allore H, Byers A. A program to prevent functional decline in physically frail, elderly persons who live at home. *New England Journal of Medicine* 2002; **347**: 1068–74.
10. O'Neil BR, Batterham AM, Hollingsworth AC, Durrand JW, Danjoux GR. First impressions count: Frailty judged by initial clinical impression predicts medium-term mortality in vascular surgical patients. *Anaesthesia* 2015; m/s 19524. (page range to be updated later)
11. Huisman MG, van Leeuwen BL, Ugolini G, et al. "Timed Up & Go": a screening tool for predicting 30-day morbidity in onco-geriatric surgical patients? A multicenter cohort study. *PLoS One* 2014; **9**: e86863.
12. Chow WB, Rosenthal RA, Merkow RP, Ko CY, Esnaola NF. Optimal preoperative assessment of the geriatric surgical patient: a best practices guideline from the American College of Surgeons National Surgical Quality Improvement Program and the American Geriatrics Society. *Journal of the American College of Surgeons* 2012; **2154**: 453–66.
13. Griffiths R, Beech F, Brown A, et al. Peri-operative care of the elderly 2014: Association of Anaesthetists of Great Britain and Ireland. *Anaesthesia* 2014; **69**: 81–98.
14. Hubbard RE, Story DA. Patient frailty: the elephant in the operating room. *Anaesthesia* 2014; **69**: 26–34.
15. National Institute for Health and Care Excellence. Preoperative tests. The use of routine preoperative tests for elective surgery. Nice Clinical Guideline 3. Pg 1-48, 2003. <https://www.nice.org.uk/guidance/cg3/resources/guidance-preoperative-tests-pdf> (accessed 10/09/2015).
16. Leung JM, Dzankic S. Relative importance of preoperative health status versus intraoperative factors in predicting postoperative adverse outcomes in geriatric surgical patients. *Journal of the American Geriatrics Society* 2001; **49**: 1080–5.
17. Barakat HM, Shahin Y, McCollum PT, Chetter IC. Prediction of organ-specific complications following abdominal aortic aneurysm repair using cardiopulmonary exercise testing. *Anaesthesia* 2015; **70**: 679–85.
18. Lee L, Schwartzman K, Carli F, et al. The association of the distance walked in 6 min with pre-operative peak oxygen consumption and complications 1 month after colorectal resection. *Anaesthesia* 2013; **68**: 811–69.
19. Carlisle JB, Danjoux G, Kerr K, Snowden C, Swart M. Validation of long-term survival prediction for scheduled abdominal aortic aneurysm repair with an independent calculator using only pre-operative variables. *Anaesthesia* 2015; **70**: 654–65.
20. Peden CJ, Grocott MPW. National Research Strategies: what outcomes are important in peri-operative elderly care? *Anaesthesia* 2014; **69**: 61–9.
21. Hoogeboom TJ, Dronkers JJ, van den Ende CH, Oosting E, van Meeteren NL. Preoperative therapeutic exercise in frail elderly scheduled for total hip replacement: a randomized pilot trial. *Clinical Rehabilitation* 2010; **24**: 901–10.
22. Carli F, Awasthi R, Gillis C, Kassouf W. Optimizing a frail elderly patient for radical cystectomy with a prehabilitation program. *Canadian Urological Association Journal* 2014; **8**: 11–2.
23. Wright AA, Cook CE, Baxter GD, Dockerty JD, Abbott JH. A comparison of 3 methodological approaches to defining major clinically important improvement of 4 performance measures in patients with hip osteoarthritis. *Journal of Orthopaedic and Sports Physical Therapy* 2011; **41**: 319–27.
24. Suetta C, Andersen JL, Dalgas U, et al. Resistance training induces qualitative changes in muscle morphology, muscle architecture, and muscle function in elderly postoperative patients. *Journal of Applied Physiology* 2008; **105**: 180–6.

doi:10.1111/anae.13321