

# Kent Academic Repository

## Full text document (pdf)

### Citation for published version

Winter, SL and Challis, JH (2008) The force-length relationship in vivo: a simulation study.  
In: British Association of Sport and Exercise Sciences Annual Conference, 2nd-4th September 2008, Brunel University, U. K..

### DOI

### Link to record in KAR

<http://kar.kent.ac.uk/53277/>

### Document Version

Author's Accepted Manuscript

#### Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

#### Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

#### Enquiries

For any further enquiries regarding the licence status of this document, please contact:

[researchsupport@kent.ac.uk](mailto:researchsupport@kent.ac.uk)

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

## The force-length relationship in vivo: a simulation study

S.L.Winter<sup>1</sup>, & J.H.Challis<sup>2</sup>

<sup>1</sup>*Aberystwyth University* and <sup>2</sup>*The Pennsylvania State University*

The force-length relationship of muscle is a fundamental mechanical property of muscle and an important determinant of resultant joint moments. In vivo a muscle may operate over all or only part (ascending limb, descending limb or plateau region) of the force-length curve for physiologically realistic ranges of motion. There are several possible sources of this variability, for example anatomical and architectural differences and differences in muscle-tendon mechanical properties. Such features are reflected in the parameters usually included in muscle models. Reported values for these parameters typically vary between different muscles and vary between subjects for a given muscle. The purpose of this study was to determine the effect of these parameters on the section of the force-length relationship that a muscle operates over. A generalised model of a mono-articular muscle-tendon complex operating over a 90 degree joint range of motion was formulated and variables were systematically varied. The parameters investigated were: ratio of tendon resting length to muscle fibre optimum length ( $L_{TR} / Lf_{OPT}$ ) (varied from 0.5 to 11.5), ratio of muscle fibre optimum length to average moment arm ( $Lf_{OPT} / r$ ) (0.5 to 5), normalised tendon strain at maximum isometric force ( $c$ ) (0 to 8%), muscle fibre pennation angle ( $\theta_{PENN}$ ) (0 to 45 degrees). The joint angle at which the optimum muscle fibre length occurred ( $\theta_{REF}$ ) was varied throughout

the 90 degree range of motion. The values for each parameter were based on reported ranges for five mono-articular muscles with different functional roles. It was shown that  $L_{TR} / Lf_{OPT}$  was important in determining the section of the force-length relationship that a muscle operated over. The effect of this ratio was modulated by  $Lf_{OPT} / r$ . The muscle operated over only one limb at intermediate values of these two ratios ( $L_{TR} / Lf_{OPT} = 5$ ;  $Lf_{OPT} / r = 3$ ), whether this was the ascending or descending limb was determined by the precise relative values of  $c$ ,  $\theta_{PENN}$ ,  $\theta_{REF}$ , and  $L_{TR} / Lf_{OPT}$ . At higher values of the two ratios the whole force-length relationship was used, at lower values only a small section of one limb was used. It was concluded that that inter-individual variability in the expressed section of the force-length relationship is possible, particularly for muscles with intermediate values of  $Lf_{OPT} / r$  and  $L_{TR} / Lf_{OPT}$  such as the brachialis and vastus lateralis. These results have implications for understanding the training adaptations of muscles for sport.