



Kent Academic Repository

Mahoney, Patrick, Schmidt, Christopher W Schmidt, Deter, Chris, Remy, Ashley, Slavin, Philip, Johns, Sarah E., Miskiewicz, Justyna J. and Nystrom, Pia (2016) *Deciduous enamel 3D microwear texture analysis as an indicator of childhood diet in medieval Canterbury, England*. Journal of Archaeological Science, 66 . pp. 128-136. ISSN 0305-4403.

Downloaded from

<https://kar.kent.ac.uk/53641/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1016/j.jas.2016.01.007>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

CC BY-NC-ND (Attribution-NonCommercial-NoDerivatives)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

1 **Deciduous enamel 3D microwear texture analysis as an indicator of childhood**
2 **diet in medieval Canterbury, England**

3

4 Patrick Mahoney^a, Christopher W. Schmidt^b, Chris Deter^a, Ashley Remy^b, Philip Slavin^c,
5 Sarah E. Johns^a, Justyna J. Miszkiewicz^d, Pia Nystrom^e.

6

7 ^aSchool of Anthropology and Conservation, University of Kent, UK.

8 ^bDepartment of Anthropology, University of Indianapolis, USA.

9 ^cSchool of History, University of Kent, UK

10 ^dDepartment of Medicine, Imperial College London, UK.

11 ^eDepartment of Archaeology, University of Sheffield, UK.

12

13 Corresponding author: Patrick Mahoney.

14 School of Anthropology and Conservation,

15 University of Kent. Canterbury. UK. CT2 7NR.

16 Email: p.mahoney@kent.ac.uk

17

18

19

20

21

22

23

24

25

26 **Abstract**

27
28 **This study** conducted the first three dimensional microwear texture analysis of human
29 deciduous teeth to reconstruct the physical properties of medieval childhood diet (age 1-8yrs)
30 **at St Gregory's Priory and Cemetery (11th to 16th century AD) in Canterbury, England.**
31 Occlusal texture complexity surfaces of maxillary molars from juvenile skeletons ($n=44$) were
32 examined to assess dietary hardness. Anisotropy values were calculated to reconstruct dietary
33 toughness, as well as jaw movements during chewing. Evidence of weaning was sought, and
34 variation in the physical properties of food was assessed against age and socio-economic
35 status. Results indicate that weaning had already commenced in the youngest children. Diet
36 became tougher from four years of age, and harder from age six. Variation in microwear
37 texture surfaces was related to historical textual evidence that refers to lifestyle developments
38 for these age groups. Diet did not vary with socio-economic status, which differs to
39 previously reported patterns for adults. We conclude, microwear texture analyses can provide
40 a non-destructive tool for revealing subtle aspects of childhood diet in the past.

41

42 **Keywords**

43 Dental microwear; medieval childhood diet.

44

45

46

47

48

49

50

51

52 **1. Introduction.**

53 Human diet during the 11th to 16th century in medieval England is best understood for adults,
54 higher status families, or ‘closed-communities’ such as monastic settlements (Dyer, 2000: 83;
55 Slavin, 2012: 8; Woolgar, 2010). Knowledge of childhood diet during this period is generally
56 more limited because it was not a focus for medieval writers. Although limited, there is some
57 historical textual evidence that provides weaning and other childrearing advice related to food
58 consumption (Fildes, 1986: 213, 1988: 76). A few isotopic studies have also reported dietary
59 weaning age and subsequent protein consumption for medieval village and urban centres in
60 the north of England (Burt, 2013, 2015; Fuller et al., 2003; Mays et al., 2002; Richards et al.,
61 2002), but not for the south-east. Neither is anything known about the physical properties
62 (hardness, toughness) of medieval childhood diet.

63 Here, we conduct the first intra-specific three dimensional (3D) dental microwear
64 texture analysis (DMTA) of human deciduous teeth to reconstruct the physical properties of
65 childhood diet in medieval Canterbury, south-east England (Fig.1). DMTA is a non-
66 destructive methodology that provides evidence of the hardness and toughness of foods eaten
67 by an individual (Scott et al. 2005, 2006; Ungar et al., 2003) in the days and weeks preceding
68 death (Grine, 1986). For example, dietary hardness and toughness has been reconstructed
69 from DMTA of permanent tooth enamel for archaeological samples of hunter-gatherers, fossil
70 hominins, and Neanderthals (El Zaatari et al., 2011, El Zaatari and Hublin, 2014; Schmidt et
71 al., [in press](#); Ungar et al., 2008a, 2010). However, few studies have examined microwear
72 surfaces of deciduous enamel (e.g., Bullington, 1991). Our study is the first to apply the 3D
73 methodology to human deciduous teeth.

74

75

76

77 *1.1 Childhood diet in Medieval England*

78 Physicians in sixteenth century Europe advised the introduction of mixed-feeding (gradual
79 [introduction of non-milk foods leading to a relative decrease in the contribution of breast milk](#)
80 [to total diet: Humphrey, 2014](#)) between seven to nine months of age (Fildes, 1986: 245).
81 Historical records from this period indicate that a child was finally weaned (removal of breast
82 milk) between 12-18 months (Fildes, 1986: [her Table 15.3-4](#); 1995: [her Table 4.7](#)). This latter
83 age range is compatible with isotopic evidence from the medieval village of Wharram Percy
84 and the urban Fishergate House cemetery in the north of England, which suggests weaning
85 occurred in the second year after birth (Burt, 2013, 2015; Fuller et al., 2003; Mays et al.,
86 2002; Richards et al., 2002).

87 Pap (flour, milk, egg yolk) or panada (bread in broth with butter or oil) were popular
88 supplementary foods during mixed-feeding (Fildes, 1986: 213; Orme, 2003: 71). Insights into
89 early childhood diet after mixed-feeding have been gained from historical textual accounts.
90 Grain products were an important component of medieval diet (Slavin, 2012: 169; Stone,
91 2006:11), and bread with butter, porridge, and gruel, were typical early childhood foods
92 (Orme, 2003: 71-72). However, little is known about dietary variation with age. Isotopic
93 evidence from Wharram Percy indicates that children may have consumed a post-weaning
94 diet that that was lower in protein compared to older individuals (Richards et al., 2002).

95 Socio-economic status could determine the quality, variety, and type of foods consumed
96 by adults (e.g., Dyer, 2006: 201-9; Powell et al., 2001: 298; Woolgar, 2006: 196; Woolgar et
97 al., 2006: 270). Outside of periods of religious observance (primarily Advent and Lent)
98 wealthier lay households and monastic communities regularly consumed meat, but other than
99 pork, it contributed less to the peasant diet (DeWitte and Slavin, 2013; Dyer, 2000: 84-86;
100 Powell et al., 2001: 308). Higher social strata preferred white bread made from wheat, while

101 those of lower socio-economic status usually consumed coarser whole grain bread (Campbell,
102 2010; Stone, 2006: 17; Slavin, 2012: 180).

103 It is unclear if the relationship between adult status and food consumption extends to
104 children from this period (Burt, 2013). In medieval York, lower status children consumed
105 higher status and more expensive foods after weaning (Burt, 2015). Furthermore, a study of
106 gross dental wear on deciduous teeth from medieval sites in the south of England, including
107 Canterbury, reported no differences between higher and lower status burials of similarly aged
108 children (Dawson and Robson Brown, 2013). Thus, the relationship between food
109 consumption and status for children in this period might be more complex than that reported
110 for adults.

111

112 *1.2. Study Aims*

113 This study conducts the first intra-specific microwear texture analysis of human deciduous
114 teeth to reconstruct the physical properties of childhood diet in Medieval Canterbury (Fig. 1).
115 All dental samples were from human juvenile skeletons ($n=44$) aged between one to eight
116 years of age, which were recovered during excavation of St Gregory's priory and cemetery
117 (11th to 15th Century AD) in Canterbury (Hicks and Hicks, 2001). The site is unique in south-
118 east England as it contained a large number of well-preserved juveniles. It has two burial
119 areas, a priory and a cemetery, *which* correspond with higher and lower socio-economic
120 status respectively (see section three).

121 The *study aims* are, 1) to search for microwear evidence of dietary weaning in the
122 youngest children. 2) Determine if variation in the physical properties of diet correlates with
123 age. 3) Compare microwear from those buried in the higher status priory to those buried in
124 the lower status cemetery. Prior to these analyses, we conduct a preliminary experimental

125 study to explore microwear texture formation processes on human deciduous enamel
126 compared to permanent enamel.

127

128 *1.3. Dental microwear texture analysis*

129 Microscopic wear in the form of scratches and pits is laid down on the occlusal surface of
130 tooth enamel as hard particles are sheared between or compressed into opposing crowns as the
131 jaw moves through the chewing cycle (Gordon, 1982). Food contaminated by grit that is
132 harder than enamel, such as quartz inclusions, is one microwear causal agent (Lucas et al.,
133 2013; Peters, 1982; Teaford and Lytle, 1996). Two dimensional (2D) dental microwear
134 analyses have been used since the 1950's to explore jaw movements of extinct mammals and
135 modern humans (Butler, 1952; Dahlberg, 1960; Mills, 1955). Subsequent 2D studies
136 described microwear patterns by their frequency, size, and orientation in extant and fossil
137 mammals (Grine, 1981; Puech, 1979; Walker, 1976; Walker et al., 1978) leading to a range of
138 quantitative studies that sought to infer aspects of diet in past human populations (e.g.,
139 Mahoney, 2006; Pastor, 1993; Schmidt, 2001; Teaford et al., 2001). Methodological
140 developments led to DMTA, the 3D characterization of microwear surfaces (Scott et al.
141 2006). This automated quantification of microwear in three dimensions minimizes inter-
142 observer measurement error (Grine et al., 2002), and thus holds great potential for the future
143 of dietary reconstruction in an archaeological context.

144 Dental microwear texture analysis is based upon the principle that an enamel surface can
145 look different when observed at different scales. A surface may appear smooth when
146 observed at a coarse scale but can appear rough at a finer scale. The texture of an enamel
147 surface can be quantified in three dimensions by combining white-light confocal profilometry
148 with scale-sensitive fractal analysis (Scott et al. 2005, 2006; Ungar et al., 2003). In this study
149 we focus upon two texture variables that have been previously related to dietary hardness and
150 toughness in extant primates:

- 151 - Area-scale fractal **complexity** (Asfc) (Scott et al., 2005). Values for complexity measure
152 changes in surface topography across different scales. Enamel with pits and scratches of
153 different sizes superimposed onto each other, or a surface that is heavily pitted, would
154 typically have higher complexity values (Ungar et al., 2008b, 2010). Consumption of
155 hard abrasive foods, which are ‘crushed’ between opposing enamel surfaces, and
156 correlated with frequent dental pits in 2D microwear analyses (Teaford, 1985; Teaford
157 and Walker, 1984), tends to produce relatively higher Asfc values (and lower levels of
158 **anisotropy**) in some primate hard seed and hard fruit eaters (Scott et al., 2005, 2006,
159 2012). Thus, a tooth surface that is dominated by dental pits with a high Asfc value has
160 been used to infer a hard and abrasive diet (Scott et al., 2012; Ungar et al., 2010).
- 161
- 162 - Exact proportion length-scale **anisotropy** (epLsar) (Scott et al., 2005). Values for the
163 anisotropy of microwear texture surfaces measure the orientation of surface features.
164 Enamel dominated by scratches all orientated in the same direction produces a high
165 anisotropy value (Ungar et al., 2008a). Low anisotropy values indicate low similarity in
166 wear feature orientation. Tougher foods which are ‘sheared’ between opposing enamel
167 surfaces, and correlated with frequent dental scratches in 2D microwear analyses
168 (Teaford, 1993; Teaford and Walker, 1984), can produce comparatively higher epLsar
169 values (and lower levels of Asfc) in some primate species that consume leaves, stems and
170 other tough fibrous foods (Scott et al., 2005, 2006, 2012). Therefore, enamel covered
171 with scratches mainly orientated in the same direction with a high epLsar value has been
172 used to infer consumption of tough abrasive foods (Scott et al., 2012; Ungar et al., 2010).
173 Jaw movement also has been reconstructed from dental scratches (Butler, 1952; Gordon,
174 1982; Scott et al., 2006; Young and Robson, 1987), whereby high epLsar values indicate
175 more consistent rather than varied jaw movements during chewing (Ungar et al., 2010).

176 While texture values within a species are variable, and texture surfaces for harder or
177 tougher diets will often overlap (Strait et al., 2013), the key correlations between microwear
178 and the physical properties of a diet established in the 1980s (Teaford, 1985; Teaford and
179 Oyen, 1989; Teaford and Walker, 1984), have been confirmed more recently in studies of
180 texture surfaces from mammals, and in experimental studies (e.g., Schubert et al., 2010;
181 Schultz, 2013, Xia et al., 2015; Hua et al., 2015). Thus, DMTA distinguishes between extant
182 primates of known diet, and these correlations provide a base-line from which to infer diet in
183 historic and pre-historic populations.

184

185 *1.4 Potential sources of deciduous microwear texture variation in Medieval Canterbury.*

186 Breast feeding will produce no microwear and the introduction of abrasive foods should
187 produce tooth wear. After weaning, flour prepared using traditional milling methods could
188 introduce hard abrasive grit into cereal foods, which has been identified as a source of
189 microwear in 2D studies (e.g., Teaford and Lytle, 1996). In medieval Canterbury, cereal
190 foods such as these came from regional farmlands, demesnes, and local grain traders
191 (Campbell, 2010; Slavin, 2012: 52-55, 2014). During the Middle Ages, grain was ground and
192 prepared for consumption by local mills in Canterbury, one of which was owned by St.
193 Gregory's priory (Hastead, 1800; Somner, 1703). Mills in medieval Kent often used
194 limestone and sandstones querns for milling (Farmer, 1992; Keller, 1989), which can
195 introduce a residue of grit into foods (Teaford and Lytle, 1996).

196 Consumption of meat can alter microwear texture surfaces (El Zaatari, 2010). 19th
197 century Fuegian hunter-gatherers with a diet that consisted mainly of meat had a lower mean
198 Asfc but higher epLsar value, relative to other hunter-gatherer populations (El Zaatari, 2010).
199 Chewing tough meat that contained some abrasives would require repetitive shearing motions
200 of the jaw, leading to many scratches orientated in the same direction. The consumption of

201 meat in medieval England varied by status amongst adults (above). If childhood status, or
202 age, also determined access to meat, then this might contribute variation to epLsar values
203 amongst the Canterbury children.

204 Beyond hard and abrasive, and tough foods, there are several other potential microwear
205 formation processes that should be considered when interpreting deciduous enamel textures.
206 First, bite force potential will differ significantly between younger and older children, as the
207 muscles of mastication gain size and strength (Kamegai et al., 2005). As such, more force
208 exerted during chewing would provide more opportunity for hard particles to be driven into
209 enamel as microwear accumulates for the first time. Thus, variation in microwear texture
210 surfaces between children of different ages might relate in part to differences in bite force.
211 Lateral movement of the mandible will also increase with age, as the mandible increases in
212 size. Greater lateral movement, as the mandible moves through the chewing cycle, might
213 produce longer scratches, though this would not necessarily alter an anisotropy value.

214 ‘Teething’, and the use of pacifiers by young children, could contribute microwear that
215 was unrelated to diet. Dental eruption in humans typically commences around the sixth post-
216 natal month as deciduous central incisors emerge through the gum line (Hillson, 2014: [his](#)
217 [Table 4](#)). The second molar is the last deciduous tooth to erupt, usually towards the start of
218 the third post-natal year (Hillson, 2014: [his Table 4](#)). Infants biting on pacifiers might scratch
219 the enamel surface. For example, in medieval England, a child might be given a piece of
220 coral during teething (Hanawalt, 1993:52). This potential source of microwear is more likely
221 in younger infants, and more likely to accumulate on early erupting incisors. Selecting later
222 erupting teeth can reduce the potential for pacifiers to obscure a diet-microwear relationship.

223

224

225 **2. Preliminary experimental study**

226 The efficacy of DMTA as an indicator of the physical properties of diet has been
227 demonstrated numerous times using adult permanent teeth (see Section 1.3). Its value,
228 however, has not been demonstrated to the same extent on deciduous teeth. Would a single
229 microwear formation process produce similar microwear texture values for both deciduous
230 and permanent enamel? Or instead, would these two enamel types differ in an unexpected
231 way when exposed to the same formation process. For example, [deciduous enamel is more
232 porous and relatively softer than permanent enamel](#) (e.g., Wilson and Beynon, 1989). To
233 investigate microwear formation processes on deciduous [relative to permanent](#) enamel we
234 undertook an experimental study before we examined microwear texture surfaces of juveniles
235 from St Gregory's priory and cemetery.

236 Complexity and anisotropy values were calculated for thirteen deciduous incisors. The
237 deciduous teeth were experimentally abraded, complexity and anisotropy values were re-
238 calculated, and compared to those from before the experiment. We repeated the experiment
239 on six permanent premolars, and compared the results to the deciduous teeth.

240

241 *2.1. Samples*

242 Thirteen deciduous mandibular incisors were donated by former students to the Indiana
243 Prehistory Laboratory, University of Indianapolis, knowing these teeth would be used in wear
244 experiments. [These teeth were included in the experimental study because the labial surface
245 of each tooth showed no signs of gross dental wear.](#) The six permanent maxillary premolars
246 were from an early 20th century cemetery population in Indiana. [The premolars were selected
247 because the mesial surface of each tooth was unworn. There was no reason to suspect that the
248 enamel microstructure of the dental samples used in the experimental study would influence](#)

249 microwear formation processes in a way that would differ to the dental samples from
250 Canterbury.

251

252 *2.2 Experimental procedures*

253 Microwear preparation and analytical procedures are described in section four. Before each
254 tooth was experimentally abraded, a target area was identified, and the complexity and
255 anisotropy of that area was recorded. The same target area was then abraded experimentally
256 and the texture values were recorded again. Thus, we produced “before” and “after”
257 experimental data sets.

258 One tooth was fixed to a square metal block weighing 1.1 kg using industrial adhesive
259 fixing tape. The tape was folded over the block, and over the tooth cervix and root. Each
260 tooth was positioned so that a relatively flat surface would be scratched. For premolars, that
261 was on the mesial aspect of the tooth. For incisors, it was the labial surface just inferior to the
262 incisal margin. The orientation of rods relative to the enamel surface can influence enamel
263 resistance to abrasion (Rensberger, 2000: his Fig 18.7), but the rod orientation in the target
264 area for both tooth types is similar. The block, with attached tooth, was placed onto a piece of
265 abrasive paper with a 200-grit size (Buehler©) that had been taped to a flat table-top. We
266 chose a grit size of 200 rather than a finer grit size to maximise scratch formation. Only the
267 tooth surface contacted the abrasive paper. The metal block was balanced by hand and pulled
268 across the length of the paper for a distance of 20 centimetres, taking approximately three
269 seconds. A square wooden block was placed next to the abrasive paper and used as a guide to
270 ensure that the distance travelled by the metal block was in a straight line. Each tooth was
271 abraded once. This process produced wear facets that were visible to the naked eye; most
272 were approximately one to two millimetres in diameter.

273 We tested the null hypothesis that there would be no difference between the complexity
274 and anisotropy post-experimental abrasion values from deciduous enamel, when compared to
275 permanent enamel, using a Mann Whitney U test.

276

277 *2.3 Experimental results*

278 Table 1 shows the experimentally induced microwear texture values. Mean Asfc
279 increased by 5.25 for deciduous teeth, and by 4.64 for permanent teeth, from before to after
280 the experimental abrasion. Mean anisotropy increased by 0.0053 for deciduous teeth, and by
281 0.0062 for permanent teeth, from before to afterwards. The post-experimental complexity
282 and abrasion values did not differ significantly when compared between deciduous and
283 permanent teeth ($p= 0.844$, $p=0.116$, respectively). Therefore, the null hypothesis was
284 retained.

285 **Table 1**
286 Mean experimentally induced microwear texture values.

	Deciduous (n)			Permanent (n)		
	Before	After	Increase	Before	After	Increase
Complexity	1.30	6.55	5.25	1.69	6.33	4.64
Asfc	(13)	(13)		(6)	(6)	
Anisotropy	0.0031	0.0084	0.0053	0.0012	0.0074	0.0062
epLsar	(13)	(13)		(5)	(5)	

287

288 *2.4 Discussion of experimental results*

289 The experimentally created wear was statistically indistinguishable when compared
290 between deciduous and permanent teeth, indicating that microwear forms in a similar way
291 when these two enamel types are subjected to the same force applied in the same direction.
292 However, there were slight differences in the mean values from the two enamel types.
293 Deciduous enamel accumulated a slightly more complex surface with fewer similarly

294 orientated scratches during the course of the experiment, relative to permanent enamel. We
295 observed that incisor enamel surfaces touching the abrasive paper were more curved
296 compared to premolars. So, these slight differences in the degree to which the microwear
297 values changed could be an artefact of this experiment, as force would have been applied to a
298 smaller area on the incisors compared to the premolars. Future studies can explore this in
299 more detail, to determine if facet size plays a key role in microwear formation. Overall, it is
300 clear that deciduous and permanent enamel produce similar microwear texture surfaces when
301 subjected to the same force applied in the same direction.

302

303 *2.5 Limitations of experimental study*

304 Results from the experimental study underscore the efficacy of the DMTA variables
305 employed here. However, other DMTA variables commonly used in studies of dietary
306 inference were not examined. Scale of maximum complexity, textural fill volume, and
307 heterogeneity are yet to be analysed. Moreover, our study only included experimental wear
308 generated in a single direction with a single force. It may be possible that a threshold exists
309 whereby extreme force, or the direction of a force, can distinguish between adult and
310 deciduous microwear texture surfaces.

311

312

313

314

315

316

317

318

319 **3. Study samples**

320 *3.1. St Gregory Priory and Cemetery*

321 The archaeological site is located about 300 meters north of present day Canterbury Cathedral
322 just outside the medieval city wall (Fig. 1). It was in use from the mid-11th to early 16th
323 century, and was excavated between 1988-1991 (Hicks and Hicks, 2001: 1-146). The priory
324 was founded by the Archbishop of Canterbury, Lanfranc, in AD 1084 (Sparks, 2001: 371).
325 Originally it was served by priests, and subsequently by Augustine canons, who cared for the
326 sick at nearby St John's hospital and provided free burial for the poor in the cemetery (Brent,
327 1897; Duncombe, 1785; Somner, 1703; Sparks, 2001: 371).

328 All skeletons were previously excavated (Anderson and Andrews, 2001: 338-370). A
329 total of 91 burials were recovered from inside and around the priory, which included a male
330 with a chalice and a gold-embroidered monastic-like garment suggesting this was a burial
331 location for clergy (Anderson and Andrews, 2001). However, the presence of children and
332 adult females within the priory indicates that this was not a 'closed' monastic community.
333 Instead, these were members of wealthier families (Hicks and Hicks, 2001), who paid for the
334 prestigious burial location, which was a popular way of displaying socio-economic status for
335 wealthy lay people in this period (Daniell, 1997: 96-97).

336 The cemetery was established just before the priory (Sparks, 1988: 31), and a total of
337 1342 skeletons were recovered during excavation. Historical textual records indicate that the
338 cemetery served poorer families from local parishes, people who could not afford burial fees,
339 and patients from nearby St. John's hospital (Brent, 1879; Somner, 1703). It was in constant
340 use until a few years after the priory was dissolved in the 16th century (Sparks, 1988: 32,
341 2001: 376).

342

343 **4. Materials and methods**

344 *4.1 Sample selection*

345 Microwear values were produced for deciduous maxillary first and second molars from 44
346 juvenile skeletons aged one to eight years. These skeletons were selected because they
347 retained the skeletal elements needed to estimate age-at-death. We focused upon maxillary
348 molars because they have thicker enamel (Mahoney, 2013), that (usually) has relatively less
349 gross wear compared to their mandibular isomeres. This was important as dentin microwear
350 was not a focus of the present study. The molars selected were also suitable for cleaning and
351 casting for microwear. Microwear values were subdivided into age groups, which were
352 created from skeletal age-at-death (see below) and the timing of dental eruption (Table 2).

353 Two juvenile skeletons from the priory dated to the earliest Lanfranc period (11th
354 century). The microwear Asfc and epLsar values for these individuals were within the range
355 of microwear values for juveniles from the priory which dated to the 14th-16th centuries.
356 Following this, microwear values were treated as one time period for subsequent analyses.
357 The cemetery burials were not sub-divided by century during excavation.

358
359 **Table 2**
360 **Deciduous microwear samples**

<i>n</i>	Age in yrs	Tooth type ¹
7	1-2	Udm1
16	2.1-4	Udm1, Udm2
14	4.1-6	Udm1, Udm2
7	6.1-8	Udm1, Udm2

361 ¹Udm1 = maxillary first molar. Udm2 = maxillary second molar

362

363

364 *4.2 Preparation and microwear texture data*

365 All teeth were prepared in the Human Osteology Research Lab, University of Kent, using
366 standard methods (e.g., Mahoney, 2006; Nystrom et al., 2004; Schmidt, 2001). The occlusal
367 surface of each tooth was cleaned using 95% ethanol and cotton wool. Impressions of phase
368 II facets were taken using a rubber-based addition-curing silicone (Coltène-Whaledent
369 Lightbody President Jet[®]). The first impression was discarded and a second impression was
370 taken and used to create the cast. The dental impression was set into dental putty (Coltène-
371 Whaledent, President Putty[®]). An epoxy resin and hardner (Buehler EpoxiCure[®]) was poured
372 into the impression to produce a cast of the occlusal surface.

373 Microwear texture data were produced in the Indiana Prehistory Lab, University of
374 Indianapolis. Resin dental casts were examined using a Sensofar[®] White Light Confocal
375 Profiler at a magnification of 100x. The microscope collected data from four contiguous
376 areas totalling 276 x 204 μm^2 . After digitally stitching the original four areas together, the
377 final study area was 242 x 181 μm^2 . Data came from Phase II wear facets (usually facet 9).
378 Data cloud manipulation was undertaken using Sensoscan[®] software, where the data were
379 levelled and non-microwear entities (primarily any remaining dirt) were removed. Analysis
380 of the data cloud required the use of Sfrax[®] and Toothfrax[®], which are scale-sensitive fractal
381 analysis programs customized for dental microwear texture analysis. Microwear variables
382 Asfc and epLsar were recorded as scale-dependent relative values (Scott et al., 2006).

383

384 *4.3. Estimating age-at-death*

385 For the one child aged 1 year, we estimated age-at-death using enamel formation times
386 (Mahoney, 2011). For the rest of the children we used a combination of enamel formation
387 times (Moorrees et al., 1963a,b), timing of dental eruption (Schour and Massler, 1941; Al-

388 [Qahtani et al., 2010](#)), long bone length (Hoppa, 1992; Scheuer et. al., 1980), and fusion of
389 cervical vertebra (Scheuer and Black, 2000).

390

391 *4.4. Statistical analyses*

392 The distribution of each microwear variable for each childhood age group (1-2, 2.1-4, 4.1-6,
393 6.1-8yrs) was checked with a one sample Kolmogorov–Smirnov test and did not differ
394 significantly from a normal curve. However, sample sizes were unequal. Thus, microwear
395 was compared between the four age groups using a non-parametric Kruskal-Wallis test.
396 Multiple post-hoc pair-wise comparisons of the age groups were undertaken using a
397 Tamhane-2 test. Microwear was compared between the two status groups using a Mann
398 Whitney U test.

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427 **5. Results**

428 Microwear descriptive statistics are summarized in Table 3. Figure 2 illustrates microwear
 429 texture surfaces. A Kruskal-Wallis test revealed that the complexity of microwear surfaces
 430 differed significantly between the four childhood age groups ($H=9.037$, $p=0.029$), but
 431 anisotropy did not ($H=6.572$, $p=0.087$). Post-hoc tests of pair-wise mean differences using
 432 the T2 statistic indicates that children aged 4.1-6 years of age had a significantly lower mean
 433 complexity value compared to younger (aged 2.1-4yrs; $p=0.017$) or older children (6.1-8yrs;
 434 $p=0.011$). Microwear did not differ significantly between higher and lower status children
 435 within the age group 2.1-4yrs, or within the age group 4.1-6 yrs.

436

437 **Table 3**
 438 Deciduous microwear mean values

Group	Age	n	Asfc		epLsar	
			mean	sd	mean	sd
a	1-2	7	1.81	0.57	0.0039	0.0019
b	2.1-4	16	2.14	0.61	0.0029	0.0016
c	4.1-6	14	1.63	0.43	0.0033	0.0014
d	6.1-8	7	2.25	0.41	0.0019	0.0010
e	1-8	44	1.95	0.57	0.0030	0.0016
Group	Status ¹					
b	lower	12	2.09	0.58	0.0028	0.0017
b	higher	4	2.28	0.80	0.0036	0.0007
c	lower	9	1.74	0.54	0.0030	0.0015
c	higher	5	1.53	0.31	0.0040	0.0007

439 ¹Lower = cemetery; higher = priory.

440 **6. Discussion**

441 *6.1. Weaning amongst one to two year olds.*

442 Microwear was present on molars from seven children in this age group (Table 3). The
443 presence of microwear suggests that mixed-feeding, at least, had commenced. On average,
444 their molar surfaces had a low Asfc and higher epLsar value, which is usually associated with
445 the consumption of tougher foods amongst extant primates (section 1.3). Therefore, at first
446 glance, there appears to be discrepancy between the microwear and the type of diet consumed
447 by the youngest children, as textual accounts indicate that a soft and limited range of foods,
448 such as pap or panda (Orme, 2003:71) would have been consumed. However, a high epLsar
449 can also be indicator of jaw movements during chewing (Ungar et al., 2010). On average, the
450 one to year olds had the most anisotropic texture surfaces compared to all other childhood
451 groups. Thus, the orientation of their microwear was the most organized, reflecting the
452 fewest changes in jaw direction during chewing. This makes sense, when viewed alongside
453 the limited range of foods consumed by this age group. The low mean complexity value for
454 the youngest children, relative to the 2.1-4yr olds, more likely represents the consumption of
455 soft foods (Scott et al., 2012). Flour in pap or panda, contaminated during the milling process
456 is one potential source of the abrasive particles that caused microwear for this age group.

457 The youngest infant with microwear was aged 1 year, which implies that **mixed-feeding**
458 might have **commenced** slightly earlier for some children in Canterbury, compared to children
459 from the contemporary Fishergate House cemetery in the north of England **where breast milk**
460 **continued to be a significant part of the diet until age 18 months** (Burt, 2013, 2015).
461 However, there is no change in microwear throughout the course of the year, which might
462 have indicated a transition from mixed-feeding to fully weaned. Instead, a child aged 1.25
463 years had a similar complexity value compared to another aged 1.75 years (Asfc= 1.71 and
464 1.69 respectively). This may simply reflect a gradual change in feeding practices that is not

465 detectable from microwear. Alternatively, breast-feeding might have been completely
466 removed from the infant diet at or around the start of the second year after birth. If this was
467 the case for the Canterbury children then their weaning age would lie within the lower end of
468 the age-range recommended for weaning in texts from the period (Fildes, 1995: 115). It
469 would also lie within the lowermost end of the weaning age-range indicated by isotopic
470 studies at contemporary Wharram Percy in the north of England, where breast-feeding ceased
471 between one to two years of age (Mays et al., 2002).

472

473 6.2. Variation in diet with age

474 Dental microwear texture analysis results suggest that the physical properties of diet for
475 children in medieval Canterbury varied from one age group to the next.

476

477 6.2.1. *Two to four years of age.*

478 Children aged two to four display an increased mean complexity of enamel surfaces
479 combined with a lower mean anisotropy, relative to one to two year olds. When this
480 combination of microwear features are compared with the base-line texture surfaces from
481 extant primates (section 1.3), it implies that the Canterbury children in this age group
482 consumed a range of foods that included relatively harder and more abrasive items. These
483 texture surfaces might be expected, as their diet was probably no longer focused upon just soft
484 infant foods like pap and panda. A more varied diet is also suggested by the lowered mean
485 anisotropy value, indicating that jaw movements were more disorganized during chewing.
486 Increased bite force relative to the infants (Kamegai et al., 2005) might be a factor here as
487 well, driving hard particles deeper into the enamel surface leading to a higher complexity
488 value.

489

490

491 6.2.2. *Four to six years of age.*

492 There was a significant change in the physical properties of diet amongst children in this
493 age group. The four to six year olds had significantly less texture complexity than either
494 younger (2.1-4yrs) or older (6.1-8yrs) children. The lowered complexity was matched by a
495 higher mean anisotropy value, which approached significance when compared to the less
496 anisotropic enamel from the 6.1-8 year olds. This combination of microwear features, lower
497 Asfc and higher epLsar (section 1.3), implies that the diet of children in medieval Canterbury
498 had altered, and now included tougher foods.

499 A change in diet between age four to six could relate in part to a period in which
500 childhood routines started to change (Bailey et al., 2008; Hanawalt, 1977: 64). Greater
501 mobility allowed children to accompany adults outside of their home and into the work place,
502 paradoxically leading to more time spent in adult company (Flemming, 2001; Hanawalt 1977,
503 1988:158). More time in adult company may have given more access to adult dietary staples,
504 such as a meat or vegetable pottage (e.g., Brears, 2008). A greater component of meat in the
505 diet of the Canterbury children might explain the change in microwear (e.g., El-Zaatari,
506 2010), especially if this was a permanent supplement to early childhood foods.

507 Support for the idea that children in this age group accessed ‘tougher’ adult dietary
508 staples, rather than returning to a soft diet similar to the infants, is provided by examining
509 their bite force potential. Children in this age group would have exerted significantly more
510 force during chewing compared to the one to two year olds (Kamegai et al., 2005). If the
511 change in the microwear pattern of the four to six year olds occurred because they re-accessed
512 a soft infant diet, whilst for example caring for a younger sibling (Hanawalt, 1988: 157), then
513 you would expect the enamel of the older children to have a higher complexity value, as
514 abrasive particles from the shared foods would have been driven deeper into their enamel.
515 This idea is not supported by the mean microwear texture values, which show that the older

516 children had a lower, not a higher mean Asfc value, relative to the infants. Neither does
517 ‘teething’ nor a ‘sick-bed’ diet seem likely causal agents. All deciduous teeth would have
518 erupted by around the age of 2.5 years, so pacifiers would not have contributed to the
519 microwear of this age group, or to the preceding age group. A sick-bed diet would not
520 necessarily contribute to the microwear of only this age group.

521

522 *6.2.3. Six to eight years of age.*

523 Children in this age group had the roughest texture surfaces with many pits and scratches
524 of different sizes overlying each other. The scratches were the least orientated compared to
525 all other childhood age groups, leading to the lowest epLsar value. The reduced range of
526 complexity and anisotropy values for this group indicates that fewer children deviated away
527 from the rough and disorganized wear features. If the tougher diet of the preceding age
528 group marks the introduction of ‘adult foods’, then the increase in food hardness in the eldest
529 children might indicate the addition of hard ‘adult’ foods. This idea is supported by
530 historical textual accounts. From around the age of seven onwards children were treated
531 increasingly like young adults and were given independent tasks outside of their home
532 (Hanawalt, 1977, 1988: 158; Fleming, 2001: 64), including apprenticeships or employment as
533 household servants (Bailey et al., 2008; Dunlop, 1912). It might be expected therefore, that
534 this change in a child’s social network would provide reduced opportunity for a distinct
535 childhood diet as they entered a new environment.

536

537 *6.3. Childhood status and diet*

538 Mean complexity and anisotropy values for children aged two to four years, or four to six
539 years of age, did not vary consistently with status (Table 3). This finding lends support to the

540 idea that the relationship between status and food consumption for medieval children might
541 be more complex compared to adults (Burt, 2013, 2015; Dawson and Robson Brown, 2013).

542

543 **7. Conclusion**

544 This study conducted the first 3D intra-specific dental microwear texture analysis of
545 childhood diet. We searched for evidence of dietary weaning, evaluated variation in the
546 physical properties of diet against age, and compared higher with lower status children.
547 Results indicate that **mixed-feeding** in Canterbury could commence by the end of a child's
548 first year. After weaning, and until the age of eight, there was no simple trajectory in the
549 physical properties of the foods that were consumed in the weeks before death. Diet
550 contained abrasives for all age groups. Texture surfaces indicated that, on average, the four to
551 six year olds consumed a diet that included tough foods whilst the eldest children consumed
552 the hardest diet. We related these changes in microwear texture surfaces to medieval textual
553 records that refer to lifestyle developments for these age groups. Our study also lends support
554 to the idea that the relationship between socio-economic status and diet for children in
555 medieval England might not be as clear as it is for adults. We conclude that deciduous dental
556 microwear texture analyses hold great potential for revealing very subtle changes to childhood
557 diet in the past.

558

559 **Acknowledgement**

560 Research funded by a British Academy-Leverhulme Trust Research Grant (SG-121921) to
561 PM and CWS. We thank two anonymous reviewers and an Associate Editor for comments
562 that improved the manuscript.

563

564

565 **References**

- 566 Al-Qahtani, S.J., Hector, M.P., Liversidge, H.M., 2010. Brief Communication: The London
567 atlas of human tooth development and eruption. *Am. J. Phys. Anth.* 142: 481–490.
- 568 Anderson, T., Andrews, J., 2001. The human remains. In: St Gregory's Priory Northgate
569 Canterbury Excavations 1988-1991. Canterbury Archaeological Trust LTD, Canterbury.
570 pp. 338-370
- 571 Bailey, B.G., Bernard, M.E., Carrier, G., Cherise Elliott, L., Langdon, J., Leishman, N,
572 Mlynarz, M., Mykhed, O. and Sidders, L.C., 2008. Coming of age and the family in medieval.
573 England. *J. Fam. Hist.* 33: 41-60.
- 574 Brears, P., 2008. *Cooking and dining in medieval England.* Prospect Books., London.
- 575 Brent, J., 1879. *Canterbury in the olden time.* Simpkin, Marshall and Co., London.
- 576 Bullington, J., 1991. Deciduous dental microwear of prehistoric juveniles from the lower
577 Illinois valley. *Am. J. Phys. Anth.* 84: 59-73.
- 578 Burt, N.M., 2013. Stable isotope ratio analysis of breastfeeding and weaning practices of
579 children from Medieval Fishergate House York. *Am. J. Phys. Anth.* 152: 407-16.
- 580 Burt, N.M., 2015. Individual dietary patterns during childhood: an archaeological application
581 of a stable isotope micro-sampling method for tooth dentin. *J. Arch. Sci.* 53: 277-290.
- 582 Butler, P.M., 1952. The milk-molars of *Perissodactyla* with remarks on molar occlusion.
583 *Proc. Zoo. Soc. Lond.* 121: 777-812.
- 584 Campbell, B.M.S., 2010. Agriculture in Kent in the high middle ages. In: Sweetinburgh S,
585 (Ed.), *Later medieval Kent 1220-1540.* Boydell Press, Woodbridge. pp. 25-53.
- 586 Dahlberg, A. 1960. Microscopic studies of abrasion and attrition on tooth surfaces. *J. Dent.*
587 *Res.* 39(4): 713-4.
- 588 Daniell, C., 1997. *Death and burial in Medieval England 1066–1550.* Routledge, London.

- 589 Dawson, H., Robson Brown, K., 2013. Exploring the relationship between dental wear and
590 status in late medieval subadults from England. *Am. J. Phys. Anth.* 150: 433–441.
- 591 DeWitte, S., Slavin, P., 2013. Between famine and death. Physiological stress and dairy
592 deficiency in England on the eve of the Black Death (1315-50): New evidence from
593 paleoepidemiology and manorial accounts. *J. Interdis. Hist.* 44: 37-60.
- 594 Duncombe, J., 1785. The history and antiquities of the three archiepiscopal hospitals and
595 other charitable foundations at and near Canterbury. *Bibliotheca Topographica Britannica* No
596 XXX, London.
- 597 Dunlop, O.J., 1912. English apprenticeship and child labour: a history. T Fisher Unwin,
598 London.
- 599 Dyer, C., 2000. *Everyday life in Medieval England*. Hambledon and London, London.
- 600 Dyer, C., 2006. Seasonal patterns in food consumption in the latter Middle Ages. In:
601 Woolgar, C.M., Serjeantson, D., Waldron, T., (Eds), *Food in medieval England*. Oxford
602 University Press, Oxford. pp. 201-14.
- 603 El-Zaatari, S., 2010. Occlusal microwear texture analysis and the diets of
604 historical/prehistoric hunter-gatherers. *Int. J. Osteoarchaeol.* 20: 67–87.
- 605 El-Zaatari, S., Grine, F.E., Ungar, P.U., Hublin, J-J., 2011. Ecogeographic variation in
606 Neanderthal dietary habits: Evidence from occlusal molar microwear texture analysis. *J.*
607 *Hum. Evol.* 61:411-424
- 608 El-Zaatari, S., Hublin, J-J., 2014. Diet of upper Paleolithic modern humans: evidence from
609 microwear texture analysis. *Am. J. Phys. Anth.* 153, 4: 570-81.
- 610 Farmer, D., 1992. Millstones for medieval manors. *Agr. Hist. Rev.* 40: 97-111.
- 611 Fildes, V.A., 1986. *Breasts, bottles and babies: a history of infant feeding*. Edinburgh
612 University Press, Edinburgh.

- 613 Fildes, V.A., 1988. *Wet nursing: a history from antiquity to the present*. Basil Blackwell,
614 New York.
- 615 Fildes, V.A., 1995. Chapter 4: The culture and biology of breastfeeding: an historical review
616 of Western Europe. In: Stuart-Macadam, P., Dettwyler, K., (Eds), *Breastfeeding: biocultural*
617 *perspectives*. Aldine De Gruyter, New York. pp. 101-126.
- 618 Flemming, P., 2001. *Family and household in Medieval England*. Palgrave Macmillan,
619 Basingstoke.
- 620 Fuller, B.T., Richards, M.P., May, S.A., 2003. Carbon and Nitrogen isotope variations in
621 tooth dentine serial sections from Wharren Percy. *J. Arch. Sci.* 30: 1673–1684.
- 622 Goldberg, G., 1992. *Women, work, and life cycle in a medieval economy: Women in work*
623 *and Yorkshire c. 1300-1520*. Oxford University Press, Oxford.
- 624 Gordon, K.D., 1982. A study of microwear on chimpanzee molars: implications for dental
625 microwear analysis. *Am. J. Phys. Anth.* 59: 195-215.
- 626 Grine, F.E., 1981. Trophic differences between gracile and robust australopithecines: a
627 scanning electron microscope analysis of occlusal events. *South. Af. J. Sci.* 77: 203-30.
- 628 Grine, F.E., 1986. Dental evidence for dietary differences in *Australopithecus* and
629 *Paranthropus*: a quantitative analysis of permanent molar microwear. *J. Hum. Evol.* 15: 783-
630 822.
- 631 Grine, F.E., Ungar, P.S., Teaford, M.F., 2002. Error rates in dental microwear quantification
632 using scanning electron microscopy. *Scanning* 24: 144-53
- 633 Hanawalt, B., 1977. Child rearing among the lower classes of late Medieval England. *J. Inter.*
634 *Hist.* VIII: 1–23.
- 635 Hanawalt, B., 1988. *The ties that bound: Peasant families in Medieval England*. Oxford
636 University Press, Oxford.

- 637 Hanawalt, B., 1993. Growing up in Medieval London. The experience of childhood in
638 history. Oxford University Press.
- 639 Hasted, E., 1800. The History and Topographical Survey of the County of Kent: Volume 11
640 Hicks, M., Hicks, A., 2001. St Gregory's Priory Northgate Canterbury Excavations 1988-
641 1991. Canterbury Archaeological Trust LTD, Canterbury.
- 642 Hillson, S., 2014. Tooth development in human evolution and bioarchaeology. Cambridge
643 University Press: Cambridge.
- 644 Hoppa, R.D., 1992. Evaluating human skeletal growth an Anglo-Saxon example. *Inter. J.*
645 *Osteoarch.* 2: 275–288.
- 646 Hua, L.C., Brandt, E.T., Meullenet J.F., Zhou, Z.R., Ungar, P.S., 2015. Technical note: an in
647 vitro study of dental microwear formation using the BITE Master II chewing machine. *Am. J.*
648 *Phys. Anth.* 158: 769–775.
- 649 Humphrey L., 2014. Isotopic and trace element evidence of dietary transitions in early life.
650 *Ann. Hum. Biol.* 41: 348–357.
- 651 Kamegai, T., Tatsuki, T., Nagano, H., Mitsunashi, H., Kumeta, J., Tatsuki, Y., Kamegai, T.,
652 Ina, D., 2005. A determination of bite force in northern Japanese children. *Eur. J. Orth.* 27:
653 53–57.
- 654 Keller, P.T., 1989. Quern Production at Folkstone, South-East Kent. *Britannia* 20: 193-200.
- 655 Lucas, P.W., Omar, R., Al-Fadhalah, K., Almusallam, A.S., Henry, A.G., Michael, S., Thai,
656 L.A., Watzke, J., Strait, D.S., Atkins, A.G., 2013. Mechanisms and causes of wear in tooth
657 enamel: implications for hominin diets. *J. R. Soc. Interface.* 10: 2012-0923.
- 658 Mahoney, P., 2006. Dental microwear from Natufian hunter-gatherers and early Neolithic
659 farmers: comparisons within and between samples. *Am. J. Phys. Anth.* 129: 39–44.
- 660 Mahoney, P., 2011. Human deciduous mandibular molar incremental enamel development.
661 *Am. J. Phys. Anth.* 144: 204–214.

- 662 Mahoney, P., 2013. Testing functional and morphological interpretations of enamel thickness
663 along the deciduous tooth row in human children. *Am. J. Phys. Anthropol.* 151: 518–525.
- 664 Mays, S., Richards, M., Fuller, B., 2002. Bone stable isotope evidence for infant feeding in
665 mediaeval England. *Antiquity* 76: 654–656.
- 666 Mills, J.R.. 1955. Ideal dental occlusion in the primates. *The Dental Practitioner.* 6(2): 47-63.
- 667 Moorrees, C.F.A., Fanning, E.A., Hunt, E.E., 1963a. Formation and resorption of three
668 deciduous teeth in children. *Am. J. Phys. Anth.* 19:99–108.
- 669 Moorrees, C.F.A., Fanning, E.A., Hunt, E.E., 1963b. Age variation of formation stages for ten
670 permanent teeth. *J. Dent. Res.* 42: 1490-1502.
- 671 Nystrom, P., Phillips-Conroy, J.E., Jolly, C.J., 2004. Dental microwear in anubis and hybrid
672 baboons (*Papio hamadryas*, sensu lato) living in Awash National Park, Ethiopia. *Am. J.*
673 *Phys. Anth.* 125: 279-291
- 674 Orme, N., 2003. *Medieval Children.* Yale University Press, New Haven.
- 675 Pastor, R., 1993 Dental microwear among prehistoric inhabitants of the Indian subcontinent:
676 a quantitative and comparative analysis. Ph.D. Dissertation, University of Oregon, Eugene.
- 677 Peters, C., 1982. Electron-optical microscope study of incipient dental microdamage from
678 experimental seed and bone crushing. *Am. J. Phys. Anth.* 57: 283-301.
- 679 Powell, A., Serjeantson, D., Smith, P., 2001. Food consumption and disposal: the animal
680 remains. In: Hicks, M., Hicks, A., (Eds), *St Gregory's Priory Northgate Canterbury*
681 *Excavations 1988-1991.* Canterbury Archaeological Trust LTD, Canterbury. pp. 289-333.
- 682 Puech, P.F. 1979. Diet of early man – evidence from abrasion of teeth and tools. *Cur. Anth.*
683 20: 590-2.
- 684 Rensberger, J.M., 2000. Pathways to functional differentiation in mammalian enamel. In:
685 Teaford MF, Smith MF, Ferguson WJ. (Eds), *Development, function, and evolution of teeth.*
686 Cambridge University Press, Cambridge. pp. 252-268.

- 687 Richards, M.P., May, S.A., Fuller, B.T., 2002. Stable carbon and nitrogen isotope values of
688 bone and teeth reflect weaning age at the Medieval Wharram Percy site, Yorkshire, UK. *Am.*
689 *J. Phys. Anth.* 119: 205–10.
- 690 Scheuer, L., Black, S., 2000. *Developmental juvenile osteology*. Elsevier Academic Press,
691 New York.
- 692 Scheuer, J.L., Musgrave, J.H., Evens, S.P., 1980. The estimation of late fetal and perinatal age
693 from limb bone length by linear and logarithmic regression. *Ann. Hum. Biol.* 7: 257–265.
- 694 Schmidt, C.W., 2001. Dental microwear evidence for a dietary shift between two nonmaize-
695 reliant prehistoric human populations from Indiana. *Am. J. Phys. Anth.* 114:139–145.
- 696 Schmidt, C.W., Beach, J., McKinley, J., Eng, J., In press. Distinguishing dietary indicators
697 of pastoralists and agriculturists via dental microwear texture analysis. *Surface Topography:*
698 *Metrology and Properties*.
- 699 Schubert, B.W., Ungar, P.S., DeSantis, L.R.G., 2010. Carnassial microwear and dietary
700 behaviour in large carnivorans. *J. Zoo.* 280: 257–263.
- 701 Schulz, E., Piotrowski, V., Clauss, M., Mau, M., Merceron, G., 2013. Dietary abrasiveness is
702 associated with variability of microwear and dental surface texture in rabbits. *PLoS ONE*
703 8(2): e56167.
- 704 Schour, I., Massler, M., 1941. The development of the human dentition. *J. Am. Dent. Ass.*
705 28: 1153-1160.
- 706 Scott, R.S., Teaford, M.F., Ungar PS. 2012. Dental microwear texture and anthropoid diets.
707 *Am. J. Phys. Anth.* 147: 551–579.
- 708 Scott, R.S., Ungar, P.S., Bergstrom, T.S., Brown, C.A., Childs, B.E., Teaford, M.F., Walker,
709 A., 2006. Dental microwear texture analysis: technical considerations. *J. Hum. Evol.* 339–
710 349.

- 711 Scott, R.S., Ungar, P.S., Bergstrom, T.S., Brown, C.A., Grine, F.E., Teaford, M.F., Walker,
712 A., 2005. Dental microwear texture analysis shows within-species diet variability in fossil
713 hominins. *Nature* 436: 693-695.
- 714 Slavin, P., 2012. *Bread and ale for the brethren: The provisioning of Norwich cathedral*
715 *Priory, c.1260-1536*. University of Hertfordshire Press, Hatfield.
- 716 Slavin, P., 2014. Market failure during the Great Famine in England and Wales (1315-7):
717 towards the re-assessment of the institutional side of the crisis. *Past and Present* 222: 9-49
- 718 Somner, W., 1703. *The Antiquities of Canterbury*. EP Publishing Limited.
- 719 Sparks, M., 1988. High Street St Gregory's and Nos 90-91: Documentary Evidence. In:
720 Bennett, P., (Ed), *Canterbury Archaeological Trust LTD Annual Reports 1987-1988*.
721 *Canterbury Archaeological Trust LTD, Canterbury*. pp 31–32.
- 722 Sparks, M., 2001. The documentary evidence: In: Hicks, M., Hicks, A., (Eds), *St Gregory's*
723 *Priory Northgate Canterbury Excavations 1988-1991*. *Canterbury Archaeological Trust*,
724 *Canterbury*. pp. 371-76.
- 725 Stone, D.J., 2006. The consumption of field crops during in late medieval England. In:
726 Woolgar, C.M., Serjeantson, D., Waldron, T., (Eds), *Food in Medieval England*. Oxford
727 *University Press, Oxford*. pp. 11-26.
- 728 Strait, D.S., Constantino, P., Lucas, P.W., Richmond, B.G., 2013. Viewpoints: Diet and
729 dietary adaptations in early hominins: The hard food perspective. *Am. J. Phys. Anth.* 151:
730 339–355.
- 731 Teaford, M.F., 1985. Molar microwear and diet in the genus *Cebus*. *Am. J. Phys. Anth.* 66:
732 363-70.
- 733 Teaford, M.F., 1993. Dental microwear and diet in extant and extinct *Theropithecus*:
734 preliminary analyses. In: Jablonski, N.G., (Ed), *Theropithecus: The Life and Death of a*
735 *Primate Genus*. Cambridge University Press, Cambridge. pp. 331-349.

- 736 Teaford, .F., Larsen, C.S., Pastor, R.F., Noble, V.E. 2001. Dental microwear and diet in La
737 Florida. In: Larsen, C.S., (Ed), Bioarchaeology of La Florida. pp. 82-112.
- 738 Teaford, M.F., Lytle, J.D., 1996. Brief communication: diet-induced changes in rates of
739 human tooth microwear: a case study involving stone-ground maize. *Am. J. Phys. Anth.*
740 100:143-7.
- 741 Teaford, M.F., Oyen, O.J., 1989a. Differences in the rate of molar wear between monkeys
742 raised on different diets. *J. Dent. Res.* 68:1513-1518.
- 743 Teaford, M.F., Walker, A., 1984. Quantitative differences in dental microwear between
744 primate species with different diets and a comment on the presumed diet of *Sivapithecus*. *Am.*
745 *J. Phys. Anth.* 64: 191-200.
- 746 Ungar, P.S., Brown, C.A., Bergstrom, T.S., Walker, A., 2003 Quantification of dental
747 microwear by tandem scanning confocal microscopy and scale-sensitive fractal analyses. *Scan*
748 25: 185–193.
- 749 Ungar, P.S., Scott, R.S., Scott, J.E., Teaford, M., 2008b. Dental microwear analysis:
750 historical perspectives and new approaches. In: Irish, J.D., Nelson, G.C., (Eds), *Technique*
751 *and application in dental anthropology*. Cambridge University Press, Cambridge. pp. 389-425.
- 752 Ungar, P.S., Grine, F.E., Teaford, M.F., 2008a. Dental microwear and diet of the Plio-
753 Pleistocene hominin *Paranthropus boisei*. *PLoS ONE* 3, e2044.
- 754 Ungar, P.S., Scott, R.S., Grine, F.E., Teaford, M.F., 2010. Molar microwear textures and the
755 diets of *Australopithecus anamensis* and *Australopithecus afarensis*. *Phil. Trans. R. Soc. B.*
756 365: 3345–3354.
- 757 Walker, P.L. 1976 Wear striations on the incisors of ceropithecoid monkeys as an index of diet
758 and habitat preference. *Am.J. Phys. Anthropol.*45:299-308.
- 759 Walker, A., Hoeck, H. N., Perez, L. 1978. Microwear of mammalian teeth as an indicator of
760 diet. *Science.* 201(8):908-910

761 Wilson, P.R., Beynon, A.D., 1998. Mineralization differences between human deciduous and
762 permanent enamel measured by quantitative microradiography. *Arch. Oral. Biol.* 34:85-88.

763 Woolgar, C.M., 2006. Group diets in Late Medieval England. In: Woolgar CM, Serjeantson
764 D, Waldron T, editors. *Food in Medieval England*. Oxford University Press: Oxford. p 191-
765 200.

766 Woolgar, C.M., 2010. Food and middle ages. *J. Med. Hist.* 36: 1-19.

767 Woolgar, C.M., Serjeantson D, Waldron T. 2006. Conclusion. In: Woolgar CM, Serjeantson
768 D, Waldron T, editors. *Food in Medieval England*. Oxford University Press: Oxford. p 267-
769 80.

770 Xia J, Zheng J, Huang D, Tian ZR, Chen L, Zhongrong Z, Ungar PS, Qian L. 2015. New
771 model to explain tooth wear with implications for microwear formation and diet
772 reconstruction. *PNAS*. 112 (34): 10669-10672.

773 Young, W.G., Robson, S.K., 1987. Jaw movement from microwear on molar teeth of the
774 koala *Phascolarctos cinereus*. *J. Zool. Lond.* 213: 51-61.

775

776

777

778

779

780

781

782


783

784

785


786 **Fig.1.** Map of United Kingdom showing Medieval Canterbury in AD1703 (after Somner,
787 AD1703). Dental samples were from juvenile skeletons recovered during excavation of St
788 Gregory's priory and cemetery. See Section 3.1.

789
790



791
792
793
794
795
796
797
798
799
800
801
802
803
804

805 **Fig.2.** Bivariate box plot (A) subdividing each age group in Table 3 into quartiles, with dental
806 microwear texture images showing 3D representations of molar enamel surfaces from two
807 children in the cemetery. Each image represents a field of view measuring 242 x 181 μm^2 .
808 Changes in colour indicate changes in depth. (B) When many pits and scratches are present
809 together, or overlying each other, they produce a 'rougher' surface and a higher complexity
810 value. The more complex surface of the 6.1-8 year olds, combined with a relatively low
811 anisotropy value (C), implies that they had a harder diet compared to the 4.1-6 year olds.
812 Their anisotropy value is low because scratches (lower right to upper left corner; lower
813 surface to upper right corner) are not orientated in the same direction.



814

815

816

817

818

819

820