

# Kent Academic Repository

## Full text document (pdf)

### Citation for published version

Cruse, Damian and Chennu, Srivas and Chatelle, Camille and Bekinschtein, Tristan A. and Fernández-Espejo, Davinia and Pickard, John D. and Laureys, Steven and Owen, Adrian M. (2012) Bedside detection of awareness in the vegetative state? - Authors' reply. *Lancet*, 379 (9827). p. 1702. ISSN 0140-6736.

### DOI

[https://doi.org/10.1016/S0140-6736\(12\)60715-6](https://doi.org/10.1016/S0140-6736(12)60715-6)

### Link to record in KAR

<http://kar.kent.ac.uk/54645/>

### Document Version

Publisher pdf

#### Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

#### Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

#### Enquiries

For any further enquiries regarding the licence status of this document, please contact:

[researchsupport@kent.ac.uk](mailto:researchsupport@kent.ac.uk)

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Since identification of command-following is so crucial for diagnosis and care of patients in vegetative and minimally conscious states, the importance of thoroughly vetted, transparent methods cannot be overstated.

We receive grant support from the James S McDonnell Foundation.

**Andrew M Goldfine, Jonathan D Victor, Mary M Conte, Jonathan C Bardin, \*Nicholas D Schiff**  
nds2001@med.cornell.edu

Department of Neurology and Neuroscience, LC-803, Weill Cornell Medical College, New York, NY 10065, USA (AMG, JDV, MMC, JCB, NDS); and Burke Medical Research Institute, White Plains, NY, USA (AMG)

- 1 Cruse D, Chennu S, Chatelle C, et al. Bedside detection of awareness in the vegetative state: a cohort study. *Lancet* 2011; **378**: 2088–94.
- 2 Pfurtscheller G, Lopes da Silva FH. Event-related EEG/MEG synchronization and desynchronization: basic principles. *Clin Neurophysiol* 1999; **110**: 1842–57.
- 3 Bai O, Lin P, Huang D, Fei D-Y, Floeter MK. Towards a user-friendly brain-computer interface: initial tests in ALS and PLS patients. *Clin Neurophysiol* 2010; **121**: 1293–303.
- 4 Goldfine AM, Victor JD, Conte MM, Bardin JC, Schiff ND. Determination of awareness in patients with severe brain injury using EEG power spectral analysis. *Clin Neurophysiol* 2011; **122**: 2157–68.
- 5 Jennett B, Plum F. Persistent vegetative state after brain damage: a syndrome in search of a name. *Lancet* 1972; **299**: 734–37.

### Authors' reply

Andrew Goldfine and colleagues make three comments about our paper. First, they imply that we do not use “power” as features in our analyses. In the Methods section<sup>1</sup> we clearly stated that we used power values as features, in line with many published studies on motor imagery and the contemporary brain-computer interface.

Second, they argue that an unspecified factor could have covaried in time with our randomly varying task structure, leading to reliable, yet erroneous, classification. In fact, randomisation is used in task design specifically to preclude conditions under which such covariation could occur. Indeed, in our paper we report data from healthy controls who were asked simply to listen to the same task instructions and “mind-

wander”, rather than complete the imagery tasks. As we reported, no positive EEG outcomes occurred under these conditions. By contrast, 75% of these participants were able to return significantly classifiable EEG data when actually following the commands. In short, the data show that random EEG fluctuations (by any known or unknown variable) were not sufficient to return a single, significant classification.

Third, Goldfine and colleagues argue, reasonably, that a 500 ms baseline might be insufficient for comparison with an action period of 3 s. Accordingly, we have rerun our analyses, comparing the 500-ms-wide “baseline” time window before the tone (–500 ms to 0 ms) with an equal-sized time window beginning 1 s after the tone (ie, far beyond any possible “stimulus-linked startle”, of the sort described by Goldfine and colleagues<sup>2</sup>). That is to say, exactly the same number of band-power values are entered into the analyses at pre-tone and post-tone time-points. The results were unchanged; as we reported in the paper, the same three patients in the vegetative state returned significant classification values in the post-tone window, but non-significant classification values in the pre-tone window—confirming that they were, indeed, following commands.

We declare that we have no conflicts of interest.

**\*Damian Cruse, Srivas Chennu, Camille Chatelle, Tristan A Bekinschtein, Davinia Fernández-Espejo, John D Pickard, Steven Laureys, Adrian M Owen**  
dcruse@uwo.ca

Brain and Mind Institute, University of Western Ontario, London, ON N6A 5B7, Canada (DC, DF-E, AMO); Medical Research Council Cognition and Brain Sciences Unit, Cambridge, UK (DC, TAB, AMO); Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK (SC); Coma Science Group, Cyclotron Research Centre & Neurology Department, University and University Hospital of Liège, Liège, Belgium (CC, SL); and Division of Academic Neurosurgery, Addenbrooke's Hospital, Cambridge, UK (JDP)

- 1 Cruse D, Chennu S, Chatelle C, et al. Bedside detection of awareness in the vegetative state: a cohort study. *Lancet* 2011; **378**: 2088–94.
- 2 Goldfine AM, Victor JD, Conte MM, Bardin JC, Schiff ND. Determination of awareness in patients with severe brain injury using EEG power spectral analysis. *Clin Neurophysiol* 2011; **122**: 2157–68.

Disorders of consciousness are at a historic epistemological junction: those who think like Damian Cruse and colleagues<sup>1</sup> continue to search within the vegetative state for evidence of misdiagnosis. However, we believe that, when a pathological state presents itself, we must research its treatment and prognosis rather than striving to search for a different diagnosis. To try to disprove the diagnosis of the vegetative state is like appearing during a liver transplant with evidence that the diseased liver in reality still works, albeit only partly. Cruse and colleagues persist in the assertion that many diagnoses are wrong and that the incidence of vegetative state is overestimated compared with minimally conscious state. For us, this is the obvious natural consequence of the fact that differences between vegetative and minimally conscious states have not yet been categorised in many of the protocols for diagnosis and treatment used by health systems worldwide.

Finding unexpected signs of consciousness simply means changing the diagnosis from vegetative state to locked-in syndrome. It means that consciousness was not evident on clinical examination, but has come to light through paraclinical testing. In our view, the most sensible thing to do is to approach disorders of consciousness as neurologists rather than instrumentalists and to develop reliable prognostic methods, validated on cohorts of an ever-increasing number of patients.<sup>2</sup> We must also acknowledge the overlap between locked-in syndrome and vegetative state: a necessity which we have emphasised on many occasions.<sup>3,4</sup>

We declare that we have no conflicts of interest.