

Kent Academic Repository

Full text document (pdf)

Citation for published version

Ciesielski, Jakub and Janczewska, Joanna and Waterstraat, Nils (2017) On the existence of homoclinic type solutions of inhomogenous Lagrangian systems. *Differential and Integral Equations*, 30 (3-4). pp. 259-272. ISSN 0893-4983.

DOI

Link to record in KAR

<http://kar.kent.ac.uk/55882/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

On the existence of homoclinic type solutions of inhomogenous Lagrangian systems

Jakub Ciesielski, Joanna Janczewska

Faculty of Applied Physics and Mathematics

Gdańsk University of Technology

Narutowicza 11/12, 80-233 Gdańsk, Poland

jciesielski@mif.pg.gda.pl, janczewska@mif.pg.gda.pl

Nils Waterstraat

School of Mathematics, Statistics and Actuarial Science

University of Kent, Canterbury

Kent CT2 7NF, England

N.Waterstraat@kent.ac.uk

February 4, 2017

Abstract

We study the existence of homoclinic type solutions for second order Lagrangian systems of the type $\ddot{q}(t) - q(t) + a(t)\nabla G(q(t)) = f(t)$, where $t \in \mathbb{R}$, $q \in \mathbb{R}^n$, $a: \mathbb{R} \rightarrow \mathbb{R}$ is a continuous positive bounded function, $G: \mathbb{R}^n \rightarrow \mathbb{R}$ is a C^1 -smooth potential satisfying the Ambrosetti-Rabinowitz superquadratic growth condition and $f: \mathbb{R} \rightarrow \mathbb{R}^n$ is a continuous bounded square integrable forcing term. A homoclinic type solution is obtained as limit of $2k$ -periodic solutions of an approximative sequence of second order differential equations.

key words: Homoclinic type solution; Lagrangian system; Critical point

AMS Subject Classification: 37J45; 58E05; 34C37; 70H05

running head: On the existence of homoclinic type solutions

1 Introduction

The aim of this paper is to prove the existence of a solution for the second order Lagrangian system

$$\begin{cases} \ddot{q}(t) - q(t) + a(t)\nabla G(q(t)) = f(t), \\ \lim_{t \rightarrow \pm\infty} q(t) = \lim_{t \rightarrow \pm\infty} \dot{q}(t) = 0, \end{cases} \quad (1)$$

where $a: \mathbb{R} \rightarrow \mathbb{R}$ is a continuous positive bounded function, $G: \mathbb{R}^n \rightarrow \mathbb{R}$, $n \geq 1$, is a C^1 -smooth potential satisfying the Ambrosetti-Rabinowitz superquadratic growth condition and $f: \mathbb{R} \rightarrow \mathbb{R}^n$ is a continuous bounded square integrable forcing term.

Our intention is to generalise the following result by E. Serra, M. Tarallo and S. Terracini from [16] to the inhomogeneous systems (1).

Theorem 1.1 *Assume that*

($\tilde{C}1$) $G \in C^2(\mathbb{R}^n, \mathbb{R})$,

($C2$) *there exists $\mu > 2$ such that for all $x \in \mathbb{R}^n \setminus \{0\}$,*

$$0 < \mu G(x) \leq (\nabla G(x), x),$$

($\tilde{C}3$) $a \in C(\mathbb{R}, \mathbb{R})$ *is almost periodic in the sense of Bohr and*

$$\inf_{t \in \mathbb{R}} a(t) > 0.$$

Then the problem

$$\begin{cases} \ddot{q}(t) - q(t) + a(t)\nabla G(q(t)) = 0, \\ \lim_{t \rightarrow \pm\infty} q(t) = \lim_{t \rightarrow \pm\infty} \dot{q}(t) = 0 \end{cases} \quad (2)$$

has at least one nonzero solution.

Here and subsequently, $(\cdot, \cdot): \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ denotes the standard inner product in \mathbb{R}^n , and $|\cdot|: \mathbb{R}^n \rightarrow [0, \infty)$ is the induced norm. Let us recall that a function a is almost periodic in the sense of Bohr if for every $\varepsilon > 0$ there is a finite linear combination of sine and cosine functions that is of distance less than ε from a with respect to the supremum norm.

The proof of Theorem 1.1 in [16] is of variational nature, i.e. a solution is found as a critical point of a suitable functional. The lack of a group of symmetries for which the functional is invariant, which exists in the case of periodic potentials, is faced by a property of Palais-Smale sequences introduced by E. Séré (see [15]) and Bochner's criterion of almost periodicity (see [4]).

Let us now consider the inhomogeneous Lagrangian systems (1). Intuitively, if the forcing term $f(t)$ in (1) is sufficiently small, then a homoclinic type solution should exist simply because of the existence in the homogenous case.

Our main result affirms this and it also deals with the question how large the forcing term f in (1) can be:

Theorem 1.2 *Assume that*

($C1$) $G \in C^1(\mathbb{R}^n, \mathbb{R})$ *and $|\nabla G(q)| = o(|q|)$ as $|q| \rightarrow 0$,*

($C2$) *there exists $\mu > 2$ such that for all $x \in \mathbb{R}^n \setminus \{0\}$,*

$$0 < \mu G(x) \leq (\nabla G(x), x),$$

($C3$) $a \in C(\mathbb{R}, \mathbb{R})$ *and $\inf_{t \in \mathbb{R}} a(t) > 0$,*

($C4$) $M := \sup\{a(t)G(x) : t \in \mathbb{R}, |x| = 1\} < \frac{1}{2}$,

($C5$) $(\int_{\mathbb{R}} |f(t)|^2 dt)^{\frac{1}{2}} < \frac{1}{2\sqrt{2}}(1 - 2M)$.

Then the inhomogenous Lagrangian system (1) has at least one homoclinic type solution.

Let us briefly discuss our assumptions in Theorem 1.2. Condition (C2) is the superquadratic growth condition due to A. Ambrosetti and P. Rabinowitz [1]. Since G has a global minimum at 0 by (C2), (C1) is more general than (C1). Moreover, it is readily seen by (C2) that for every $q \neq 0$ the map

$$(0, \infty) \ni \xi \mapsto G(\xi^{-1}q)|\xi|^\mu \in \mathbb{R}$$

is non-increasing, which yields the following inequalities:

$$G(q) \leq G\left(\frac{q}{|q|}\right)|q|^\mu, \text{ if } 0 < |q| \leq 1 \quad (3)$$

and

$$G(q) \geq G\left(\frac{q}{|q|}\right)|q|^\mu, \text{ if } |q| \geq 1. \quad (4)$$

As $\mu > 2$, the inequality (4) implies that G grows faster than $|\cdot|^2$ at infinity.

Clearly, (C3) is more general than (C3). Note that (C3) and (C4) imply that a is bounded, which, however, is also true for every almost periodic function in the sense of Bohr.

The last two conditions (C4) and (C5) are closely related. Namely, the forcing term f needs to be sufficiently small in $L^2(\mathbb{R}, \mathbb{R}^n)$, but the upper bound on the norm of f depends on the restriction of the space variable of the potential $a(t)G(x)$ to the unit sphere in \mathbb{R}^n .

The study of homoclinic solutions for Lagrangian systems has received much attention in recent years, especially when the potential is periodic in time. The existence problem of homoclinics has been widely investigated by variational methods, see for example in [2, 3, 5, 11–13, 15]. Existence results for perturbed systems were given in [6–10, 14].

Our proof of Theorem 1.2 is also of variational nature. Let us point out, however, that it is quite different from Serra, Tarallo and Terracini's proof of Theorem 1.1 in [16]. Here, we find a solution of (1) as a limit in $C_{loc}^2(\mathbb{R}, \mathbb{R}^n)$ of a sequence $\{q_k\}_{k \in \mathbb{N}}$, $q_k \in W_{2k}^{1,2}(\mathbb{R}, \mathbb{R}^n)$, obtained by an approximation scheme introduced by Krawczyk in [10], where every q_k is a critical point of a suitable functional I_k that we introduce below.

We now prove Theorem 1.2 in the following section and we nicely round off the paper by two numerical examples in a final section.

2 Proof of Theorem 1.2

In what follows, we let E be the Sobolev space $W^{1,2}(\mathbb{R}, \mathbb{R}^n)$ of $W^{1,2}$ -functions on \mathbb{R} with values in \mathbb{R}^n equipped with the norm

$$\|q\|_E = \left(\int_{-\infty}^{\infty} (|q(t)|^2 + |\dot{q}(t)|^2) dt \right)^{\frac{1}{2}}.$$

For each $k \in \mathbb{N}$, we denote by $E_k = W_{2k}^{1,2}(\mathbb{R}, \mathbb{R}^n)$ the Sobolev space of $2k$ -periodic $W^{1,2}$ -functions with the norm

$$\|q\|_{E_k} = \left(\int_{-k}^k (|q(t)|^2 + |\dot{q}(t)|^2) dt \right)^{\frac{1}{2}}.$$

Then let $L_{2k}^\infty(\mathbb{R}, \mathbb{R}^n)$ be the space of $2k$ -periodic, essentially bounded and measurable functions from \mathbb{R} into \mathbb{R}^n with the norm

$$\|q\|_{L_{2k}^\infty} = \text{ess sup}\{|q(t)|: t \in [-k, k]\}.$$

We note for later reference that

$$\|q\|_{L_{2k}^\infty} \leq \sqrt{2}\|q\|_{E_k} \quad (5)$$

for all $k \in \mathbb{N}$ and $q \in E_k$ (cf. [6, Fact 2.8]). Finally, let $C_{loc}^2(\mathbb{R}, \mathbb{R}^n)$ denote the space of C^2 -functions with the topology of almost uniform convergence of functions and all their derivatives up to second order.

The following result can be found in [10, Thm. 1.3].

Theorem 2.1 *Let $f: \mathbb{R} \rightarrow \mathbb{R}^n$ be a non-trivial, bounded, continuous and square integrable map and $V: \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ a C^1 -smooth potential such that $\nabla_q V: \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is bounded in the time variable. Assume that for each $k \in \mathbb{N}$ the boundary value problem*

$$\begin{cases} \ddot{q}(t) + \nabla_q V_k(t, q(t)) = f_k(t), \\ q(-k) - q(k) = \dot{q}(-k) - \dot{q}(k) = 0, \end{cases} \quad (6)$$

where $f_k: \mathbb{R} \rightarrow \mathbb{R}^n$ is a $2k$ -periodic extension of $f|_{[-k, k]}$ and $V_k: \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a $2k$ -periodic extension of $V|_{[-k, k] \times \mathbb{R}^n}$, has a periodic solution $q_k \in E_k$ and $\{\|q_k\|_{E_k}\}_{k \in \mathbb{N}}$ is a bounded sequence in \mathbb{R} . Then there exists a subsequence $\{q_{k_j}\}_{j \in \mathbb{N}}$ converging in the topology of $C_{loc}^2(\mathbb{R}, \mathbb{R}^n)$ to a function $q \in E$ which is a homoclinic type solution of the Newtonian system

$$\ddot{q}(t) + \nabla_q V(t, q(t)) = f(t), \quad t \in \mathbb{R}. \quad (7)$$

Our aim is to obtain a homoclinic type solution of (1) by Theorem 2.1 as a limit in $C_{loc}^2(\mathbb{R}, \mathbb{R}^n)$ of a sequence $\{q_k\}_{k \in \mathbb{N}}$ such that for each $k \in \mathbb{N}$, $q_k \in E_k$ is a $2k$ -periodic solution of the boundary value problem

$$\begin{cases} \ddot{q}(t) - q(t) + a_k(t)\nabla G(q(t)) = f_k(t), \\ q(-k) - q(k) = \dot{q}(-k) - \dot{q}(k) = 0, \end{cases} \quad (8)$$

where f_k is as above and $a_k: \mathbb{R} \rightarrow \mathbb{R}$ is a $2k$ -periodic extension of $a|_{[-k, k]}$.

To this purpose, we now define for $k \in \mathbb{N}$ a functional $I_k: E_k \rightarrow \mathbb{R}$ by

$$I_k(q) = \frac{1}{2}\|q\|_{E_k}^2 - \int_{-k}^k a_k(t)G(q(t))dt + \int_{-k}^k (f_k(t), q(t))dt. \quad (9)$$

Then $I_k \in C^1(E_k, \mathbb{R})$ and, moreover,

$$\begin{aligned} I_k'(q)v &= \int_{-k}^k (\dot{q}(t), \dot{v}(t))dt + \int_{-k}^k (q(t), v(t))dt \\ &\quad - \int_{-k}^k (a_k(t)\nabla G(q(t)), v(t))dt + \int_{-k}^k (f_k(t), v(t))dt. \end{aligned} \quad (10)$$

Hence

$$I'_k(q)q = \|q\|_{E_k}^2 - \int_{-k}^k (a_k(t)\nabla G(q(t)), q(t))dt + \int_{-k}^k (f_k(t), q(t))dt. \quad (11)$$

Let us note for later reference that by [6, Fact 2.2], if

$$m := \inf\{a(t)G(x) : t \in \mathbb{R}, |x| = 1\}$$

and $\mu > 2$ is defined as in Theorem 1.2, then for all $\zeta \in \mathbb{R} \setminus \{0\}$ and $q \in E_k \setminus \{0\}$

$$\int_{-k}^k a_k(t)G(\zeta q(t))dt \geq m|\zeta|^\mu \int_{-k}^k |q(t)|^\mu dt - 2km. \quad (12)$$

Clearly, critical points of the functional I_k are classical $2k$ -periodic solutions of (8). We will now obtain a critical point of I_k by using the Mountain Pass Theorem from [1]. This theorem provides the minimax characterisation for a critical value which is important for our argument. Let us recall its statement for the convenience of the reader.

Theorem 2.2 *Let E be a real Banach space and $I: E \rightarrow \mathbb{R}$ a C^1 -smooth functional. If I satisfies the following conditions:*

- (i) $I(0) = 0$,
- (ii) every sequence $\{u_j\}_{j \in \mathbb{N}} \subset E$ such that $\{I(u_j)\}_{j \in \mathbb{N}}$ is bounded in \mathbb{R} and $I'(u_j) \rightarrow 0$ in E^* as $j \rightarrow \infty$ contains a convergent subsequence (Palais-Smale condition),
- (iii) there exist constants $\rho, \alpha > 0$ such that $I|_{\partial B_\rho(0)} \geq \alpha$,
- (iv) there is some $e \in E \setminus \overline{B}_\rho(0)$ such that $I(e) \leq 0$,

where $B_\rho(0)$ denotes the open ball in E of radius ρ about 0, then I has a critical value $c \geq \alpha$ given by

$$c = \inf_{g \in \Gamma} \max_{s \in [0,1]} I(g(s)),$$

where

$$\Gamma := \{g \in C([0,1], E) : g(0) = 0, g(1) = e\}.$$

The following lemma, in combination with Theorem 2.1, is the keystone of our proof of Theorem 1.2.

Lemma 2.3 *For each $k \in \mathbb{N}$, the functional I_k given by (9) has the mountain pass geometry, i.e. it satisfies all assumptions of Theorem 2.2.*

Proof. We fix $k \in \mathbb{N}$ and we now show the assumptions (i)-(iv) in Theorem 2.2 for I_k . It is clear that $I_k(0) = 0$, which is (i). In order to show the Palais-Smale condition (ii), we consider a sequence $\{u_j\}_{j \in \mathbb{N}} \subset E_k$ such that $\{I_k(u_j)\}_{j \in \mathbb{N}} \subset \mathbb{R}$ is bounded and $I'_k(u_j) \rightarrow 0$ in E_k^* as $j \rightarrow +\infty$. Consequently, there exists a constant $C_k > 0$ such that for all $j \in \mathbb{N}$ we have

$$|I_k(u_j)| \leq C_k, \quad \|I'_k(u_j)\|_{E_k^*} \leq C_k. \quad (13)$$

By (9) and (C2) we get

$$\begin{aligned} \|u_j\|_{E_k}^2 &\leq 2I_k(u_j) - 2 \int_{-k}^k (f_k(t), u_j(t)) dt \\ &\quad + \frac{2}{\mu} \int_{-k}^k (a_k(t) \nabla G(u_j(t)), u_j(t)) dt. \end{aligned} \quad (14)$$

As

$$\int_{-k}^k (a_k(t) \nabla G(u_j(t)), u_j(t)) dt = \|u_j\|_{E_k}^2 - I'_k(u_j)u_j + \int_{-k}^k (f_k(t), u_j(t)) dt$$

by (11), we obtain

$$\begin{aligned} \left(1 - \frac{2}{\mu}\right) \|u_j\|_{E_k}^2 &\leq 2I_k(u_j) - \frac{2}{\mu} I'_k(u_j)(u_j) - 2 \int_{-k}^k (f_k(t), u_j(t)) dt \\ &\quad + \frac{2}{\mu} \int_{-k}^k (f_k(t), u_j(t)) dt \end{aligned}$$

and so

$$\begin{aligned} \left(1 - \frac{2}{\mu}\right) \|u_j\|_{E_k}^2 &\leq 2I_k(u_j) \\ &\quad + \left(\frac{2}{\mu} \|I'_k(u_j)\|_{E_k^*} + \left(2 - \frac{2}{\mu}\right) \|f_k\|_{L^2_{2k}}\right) \|u_j\|_{E_k}, \end{aligned} \quad (15)$$

where we denote by

$$\|q\|_{L^2_{2k}} = \left(\int_{-k}^k |q(t)|^2 dt \right)^{\frac{1}{2}}$$

the norm of the space $L^2_{2k}(\mathbb{R}, \mathbb{R}^n)$ of all $2k$ -periodic L^2 -functions. Combining (15) with (C5) and (13) we get

$$\left(1 - \frac{2}{\mu}\right) \|u_j\|_{E_k}^2 - \left(\frac{2C_k}{\mu} + \frac{1}{\sqrt{2}} \left(1 - \frac{1}{\mu}\right) (1 - 2M)\right) \|u_j\|_{E_k} - 2C_k \leq 0,$$

which yields the boundedness of $\{u_j\}_{j \in \mathbb{N}}$ in E_k as $\mu > 2$ by (C2). Going to a subsequence if necessary, we can assume that there exists a function $u \in E_k$ such that $u_j \rightharpoonup u$ weakly in E_k as $j \rightarrow +\infty$, and hence $\{u_j\}_{j \in \mathbb{N}}$ also converges to u uniformly, as E_k is compactly embedded in $C([-k, k], \mathbb{R}^n)$. This shows in particular that $\|u_j - u\|_{L^2_{2k}} \rightarrow 0$ as $j \rightarrow \infty$.

Applying (10) we have

$$\begin{aligned} I'_k(u_j)(u_j - u) &= \int_{-k}^k (\dot{u}_j(t), \dot{u}_j(t) - \dot{u}(t)) dt + \int_{-k}^k (u_j(t), u_j(t) - u(t)) dt \\ &\quad - \int_{-k}^k (a_k(t) \nabla G(u_j(t)), u_j(t) - u(t)) dt + \int_{-k}^k (f_k(t), u_j(t) - u(t)) dt \end{aligned}$$

and

$$\begin{aligned} I'_k(u)(u_j - u) &= \int_{-k}^k (\dot{u}(t), \dot{u}_j(t) - \dot{u}(t))dt + \int_{-k}^k (u(t), u_j(t) - u(t))dt \\ &\quad - \int_{-k}^k (a_k(t)\nabla G(u(t)), u_j(t) - u(t))dt + \int_{-k}^k (f_k(t), u_j(t) - u(t))dt, \end{aligned}$$

which yields

$$\begin{aligned} \|\dot{u}_j - \dot{u}\|_{L^2_{2k}}^2 &= (I'_k(u_j) - I'_k(u))(u_j - u) - \|u_j - u\|_{L^2_{2k}}^2 \\ &\quad + \int_{-k}^k a_k(t)(\nabla G(u_j(t)) - \nabla G(u(t)), u_j(t) - u(t))dt. \end{aligned}$$

As $I'_k(u_j)$ is bounded by (13), ∇G is continuous and $u_j \rightarrow u$ uniformly, we see that $\|\dot{u}_j - \dot{u}\|_{L^2_{2k}}^2 \rightarrow 0$. Hence $\|\dot{u}_j - \dot{u}\|_{E_k} \rightarrow 0$ and the Palais-Smale condition is shown.

In the next step we will prove that there exist constants $\rho > 0$ and $\alpha > 0$ independent of $k \in \mathbb{N}$ such that $I_k|_{\partial B_\rho(0)} \geq \alpha$, which is (iii). Assume that $0 < \|q\|_{L^\infty_{2k}} \leq 1$. By (3) and (C4) we obtain

$$\int_{-k}^k a_k(t)G(q(t))dt \leq \int_{-k}^k a_k(t)G\left(\frac{q(t)}{|q(t)|}\right)|q(t)|^\mu dt \leq M \int_{-k}^k |q(t)|^2 dt \leq M\|q\|_{E_k}^2.$$

Combining this with (9) we get

$$I_k(q) \geq \frac{1}{2}(1 - 2M)\|q\|_{E_k}^2 - \|f\|_{L^2}\|q\|_{E_k}.$$

Let $\rho = \frac{1}{\sqrt{2}}$ and $\alpha = \frac{1}{\sqrt{2}}\left(\frac{1}{2\sqrt{2}}(1 - 2M) - \|f\|_{L^2}\right)$. From (C5) it follows that $\alpha > 0$. Using (5), if $\|q\|_{E_k} = \rho$, then $0 < \|q\|_{L^\infty_{2k}} \leq 1$, which implies $I_k(q) \geq \alpha$.

It remains to prove (iv), i.e. that for all $k \in \mathbb{N}$ there is $e_k \in E_k$ such that $\|e_k\|_{E_k} > \rho$ and $I_k(e_k) \leq 0$. Applying (9) and (12), we have

$$I_k(\zeta q) \leq \frac{\zeta^2}{2}\|q\|_{E_k}^2 - m|\zeta|^\mu \int_{-k}^k |q(t)|^\mu dt + |\zeta|\|f_k\|_{L^2_{2k}}\|q\|_{E_k} + 2km$$

for all $\zeta \in \mathbb{R} \setminus \{0\}$ and $q \in E_k \setminus \{0\}$. Let us choose $0 \neq Q \in E_1$ such that $Q(\pm 1) = 0$. As $\mu > 2$ and $m > 0$, there exists $\zeta \in \mathbb{R} \setminus \{0\}$ such that $\|\zeta Q\|_{E_1} > \rho$ and $I_1(\zeta Q) < 0$.

We set $e_1(t) = \zeta Q(t)$ and define for each positive integer $k > 0$,

$$e_k(t) = \begin{cases} e_1(t) & \text{if } |t| \leq 1 \\ 0 & \text{if } 1 < |t| \leq k. \end{cases}$$

Then $e_k \in E_k$, $\|e_k\|_{E_k} = \|e_1\|_{E_1} > \rho$ and $I_k(e_k) = I_1(e_1) < 0$ for all $k \in \mathbb{N}$.

Consequently, by Theorem 2.2, the action I_k has a critical value $c_k \geq \alpha$ given by

$$c_k = \inf_{g \in \Gamma_k} \max_{s \in [0,1]} I_k(g(s)), \quad (16)$$

where $\Gamma_k = \{g \in C([0,1], E_k) : g(0) = 0, g(1) = e_k\}$.

□

From Lemma 2.3 we conclude that for all $k \in \mathbb{N}$ there exists $q_k \in E_k$ such that

$$I_k(q_k) = c_k, \quad I'_k(q_k) = 0,$$

where the values c_k are given by (16). By Theorem 2.1, in order to finish the proof of Theorem 1.2, it suffices to show that the sequence of real numbers $\{\|q_k\|_{E_k}\}_{k \in \mathbb{N}}$ is bounded. For this purpose, set

$$M_0 = \max_{s \in [0,1]} I_1(se_1).$$

As $I_k(se_k) = I_1(se_1)$ for all $s \in [0,1]$, $k \in \mathbb{N}$, and $se_k \in \Gamma_k$, we have by (16)

$$c_k = \inf_{g \in \Gamma_k} \max_{s \in [0,1]} I_k(g(s)) \leq M_0. \quad (17)$$

Using (9) and (11) we obtain

$$\begin{aligned} c_k = I_k(q_k) - \frac{1}{2} I'_k(q_k) q_k &= \frac{1}{2} \int_{-k}^k (a_k(t) \nabla G(q_k(t)), q_k(t)) dt \\ &\quad - \int_{-k}^k a_k(t) G(q_k(t)) dt + \frac{1}{2} \int_{-k}^k (f_k(t), q_k(t)) dt \end{aligned}$$

for all $k \in \mathbb{N}$, and by (C2),

$$c_k \geq \left(\frac{\mu}{2} - 1\right) \int_{-k}^k a_k(t) G(q_k(t)) dt - \frac{1}{2} \|f_k\|_{L^2_{2k}} \|q_k\|_{E_k}.$$

Hence

$$\int_{-k}^k a_k(t) G(q_k(t)) dt \leq \frac{1}{\mu - 2} \left(2c_k + \|f_k\|_{L^2_{2k}} \|q_k\|_{E_k}\right).$$

Combining this with (9), (17) and (C5), for each $k \in \mathbb{N}$, we have

$$\begin{aligned} c_k = I_k(q_k) &= \frac{1}{2} \|q_k\|_{E_k}^2 - \int_{-k}^k a(t) G(q_k(t)) dt + \int_{-k}^k (f_k(t), q_k(t)) dt \\ &\geq \frac{1}{2} \|q_k\|_{E_k}^2 - \frac{1}{\mu - 2} \left(2c_k + \|f_k\|_{L^2_{2k}} \|q_k\|_{E_k}\right) - \|f_k\|_{L^2_{2k}} \|q_k\|_{E_k}, \end{aligned}$$

and so

$$\|q_k\|_{E_k}^2 - \frac{1}{\sqrt{2}} \frac{\mu - 1}{\mu - 2} (1 - 2M) \|q_k\|_{E_k} - \frac{2\mu M_0}{\mu - 2} \leq 0,$$

which completes the proof of Theorem 1.2.




Figure 1: Example 3.1, an approximative solution of (8) for $k = 10$

3 One-dimensional Examples

In this section we present two simple one-dimensional examples, i.e. we consider the case $n = 1$.

Example 3.1 Let a function $a: \mathbb{R} \rightarrow \mathbb{R}$, a forcing term $f: \mathbb{R} \rightarrow \mathbb{R}$ and a potential $G: \mathbb{R} \rightarrow \mathbb{R}$ be given as follows:

$$\begin{aligned} a(t) &= \frac{1}{5}e^{-t^2} + \frac{1}{10}, \quad t \in \mathbb{R}, \\ f(t) &= \frac{2}{5}e^{-\frac{t^2}{2}}, \quad t \in \mathbb{R}, \\ G(q) &= q^4, \quad q \in \mathbb{R}. \end{aligned}$$

It is easy to check that a , f and G satisfy the assumptions (C1) – (C5).

The figures 1-5 show the graphs of approximative solutions q_k of (8) for $k = 10, 16, 90, 140, 200$.

Example 3.2 Let us define functions $a, f, G: \mathbb{R} \rightarrow \mathbb{R}$ as follows:

$$\begin{aligned} a(t) &= \frac{1}{\pi} \arctan(t) + \frac{1}{2}, \quad t \in \mathbb{R}, \\ f(t) &= \frac{1}{2}e^{-\frac{t^2}{2}}, \quad t \in \mathbb{R}, \\ G(q) &= q^4, \quad q \in \mathbb{R}. \end{aligned}$$

Again, it is readily seen that the assumptions (C1) – (C5) are satisfied.

The figures 6-9 show the graphs of approximative solutions q_k of (8) for $k = 10, 16, 90, 140$.




Figure 2: Example 3.1, an approximative solution of (8) for $k = 16$




Figure 3: Example 3.1, an approximative solution of (8) for $k = 90$




Figure 4: Example 3.1, an approximative solution of (8) for $k = 140$




Figure 5: Example 3.1, an approximative solution of (8) for $k = 200$




Figure 6: Example 3.2, an approximative solution of (8) for $k = 10$




Figure 7: Example 3.2, an approximative solution of (8) for $k = 16$




Figure 8: Example 3.2, an approximative solution of (8) for $k = 90$




Figure 9: Example 3.2, an approximative solution of (8) for $k = 140$

Acknowledgments

The second and the third author are supported by Grant PPP-PL no. 57217076 of DAAD and MNiSW.

References

- [1] A. Ambrosetti, P.H. Rabinowitz, Dual variational methods in critical point theory and applications, *J. Funct. Anal.* 14 (1973), 349–381.
- [2] G. Arioli, A. Szulkin, Homoclinic solutions for a class of systems of second order differential equations, *Topol. Methods Nonlinear Anal.* 6 (1995), no. 1, 189–197.
- [3] M.L. Bertotti, P. Montecchiari, Connecting orbits for some classes of almost periodic Lagrangian systems, *J. Differential Equations* 145 (1998), 453–468.
- [4] A.S. Besicovitch, *Almost Periodic Functions*, Dover Publications, Inc., 1954.
- [5] V. Coti Zelati, Multibump homoclinic solutions to periodic orbits in a center manifold, *Ricerche Mat.* 54 (2005), no. 2, 433–446.
- [6] M. Izydorek, J. Janczewska, Homoclinic solutions for a class of the second order Hamiltonian systems, *J. Differential Equations* 219 (2005), 375–389.
- [7] M. Izydorek, J. Janczewska, Homoclinic solutions for nonautonomous second order Hamiltonian systems with a coercive potential, *J. Math. Anal. Appl.* 335 (2007), no. 2, 1119–1127.
- [8] J. Janczewska, An approximative scheme of finding almost homoclinic solutions for a class of Newtonian systems, *Topol. Methods Nonlinear Anal.* 33 (2009), no. 1, 169–177.
- [9] J. Janczewska, Two almost homoclinic solutions for second-order perturbed Hamiltonian systems, *Commun. Contemp. Math.* 14 (2012), no. 4, 1250025, 9 pp.
- [10] R. Krawczyk, A note on an approximative scheme of finding almost homoclinic solutions for Newtonian systems, *Calculus of variations and PDEs*, 107–113, Banach Center Publ., 101, Polish Acad. Sci. Inst. Math., Warsaw, 2014.
- [11] P.H. Rabinowitz, Homoclinic orbits for a class of Hamiltonian systems, *Proc. Roy. Soc. Edinburgh Sect. A* 114 (1990), no. 1-2, 33–38.
- [12] P.H. Rabinowitz, Variational methods for Hamiltonian systems, *Handbook of dynamical systems*, vol. 1A, 1091–1127, North-Holland, Amsterdam, 2002.
- [13] P.H. Rabinowitz, K. Tanaka, Some results on connecting orbits for a class of Hamiltonian systems, *Math. Z.* 206 (1991), no. 3, 473–499.

- [14] A. Salvatore, Multiple homoclinic orbits for a class of second order perturbed Hamiltonian systems, *Dynamical systems and differential equations* (Wilmington, NC, 2002), *Discrete Contin. Dyn. Syst.* 2003, suppl., 778–787.
- [15] E. Séré, Existence of infinitely many homoclinic orbits in Hamiltonian systems, *Math. Zeit.* 209 (1991), 27–42.
- [16] E. Serra, M. Tarallo, S. Terracini, On the existence of homoclinic solutions for almost periodic second order systems, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 13 (1996), no. 6, 783–812.