

Kent Academic Repository

Full text document (pdf)

Citation for published version

Judge, Edmund and Naboko, Serguei and Wood, Ian (2016) Eigenvalues for perturbed periodic Jacobi matrices by the Wigner-von Neumann approach. *Integral Equations and Operator Theory*, 85 (3). pp. 427-450. ISSN 0378-620X.

DOI

<https://doi.org/10.1007/s00020-016-2302-5>

Link to record in KAR

<http://kar.kent.ac.uk/55954/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Eigenvalues for perturbed periodic Jacobi matrices by the Wigner-von Neumann approach

Edmund Judge, Sergey Naboko and Ian Wood

Abstract. The Wigner-von Neumann method, which has previously been used for perturbing continuous Schrödinger operators, is here applied to their discrete counterparts. In particular, we consider perturbations of arbitrary T -periodic Jacobi matrices. The asymptotic behaviour of the subordinate solutions is investigated, as too are their initial components, together giving a general technique for embedding eigenvalues, λ , into the operator's absolutely continuous spectrum. Introducing a new rational function, $C(\lambda; T)$, related to the periodic Jacobi matrices, we describe the elements of the a.c. spectrum for which this construction does not work (zeros of $C(\lambda; T)$); in particular showing that there are only finitely many of them.

Keywords. Spectral theory, Periodic Jacobi operators, Wigner-von Neumann potential, Subordinate solutions.

1. Introduction

First published in 1929, the Wigner-von Neumann method provides a way of embedding eigenvalues into the absolutely continuous spectrum of a one-dimensional Schrödinger operator [16]. Specifically, the operator is perturbed by a potential of the form

$$\frac{c \sin(2\omega x + \varphi)}{x}$$

causing the eigenvalue $E = \omega^2$ to become embedded in the interval $[0, \infty)$ of a.c. spectrum. Since then the method has been adapted to embed multiple

Edmund Judge was supported by the Engineering and Physical Sciences Research Council (grant EP/M506540/1). Sergey Naboko was supported by the Russian Science Foundation (grant 15-11-30007), NCN 2013/09/BST1/04319 and Marie Curie grant (2013-2014 yy) PIFI-GA-2011-299919.

with complex entries and non-zero diagonal, but we stick to the simple case to make our constructions more transparent.

Our goal is to make an ansatz for a possible eigenvector (introduced in Section 5) and an ansatz for the potential (Section 6). We then establish the asymptotics of the potential needed to realize our ansatz. Additionally, we must confirm that the subordinate solution constructed in this way also satisfies the initial equations encoded within the Jacobi matrix (Section 7), thus giving an embedded eigenvalue. Section 2 contains some general results on T -periodic Jacobi operators, while Sections 3 contains some preliminary results for the construction. In Section 4 we introduce the aforementioned function $C(\lambda; T)$ and analyze its properties, in particular that it is a rational function of λ .

2. Preliminary results on the spectrum of a period- T Jacobi operator

Before we consider perturbations, it is best to state some results describing the structure of the spectrum of unperturbed T -periodic Jacobi operators.

It is not hard to see that the spectrum of a period- T Jacobi matrix, J_T , is such that

$$\sigma(J_T) \subseteq [-\max\{a_1 + a_2, a_2 + a_3, \dots, a_T + a_1\} - \min\{b_1, \dots, b_T\}, \max\{a_1 + a_2, a_2 + a_3, \dots, a_T + a_1\} + \max\{b_1, \dots, b_T\}].$$

The inclusion is sharp for period-1 and period-2 Jacobi operators, which can easily be proved. Moreover, the operator is self-adjoint and in the case of the vanishing diagonal its spectrum is symmetric w.r.t. zero.

The next elementary lemma gives information which will be useful in determining the entries of the monodromy matrix associated to J_T .

Lemma 2.1. *Let*

$$A_s = \begin{pmatrix} 0 & 1 \\ c_s & \frac{\lambda - b_s}{a_s} \end{pmatrix}$$

with $a_s, c_s \neq 0$. Let $m \in \mathbb{N}$ and

$$A(\lambda) = \begin{pmatrix} a_{11}(\lambda) & a_{12}(\lambda) \\ a_{21}(\lambda) & a_{22}(\lambda) \end{pmatrix} = \prod_{s=1}^m A_s.$$

Then, for $m \geq 2$,

$$\begin{aligned} a_{11}(\lambda) &= c_1 \frac{\lambda^{m-2}}{m-1} + P_{m-3}(\lambda), & a_{12}(\lambda) &= \frac{\lambda^{m-1}}{m-1} + P_{m-2}(\lambda), \\ & \prod_{s=2} a_s & \prod_{s=1} a_s & \\ a_{21}(\lambda) &= c_1 \frac{\lambda^{m-1}}{m} + \tilde{P}_{m-2}(\lambda), & \text{and } a_{22}(\lambda) &= \frac{\lambda^m}{m} + P_{m-1}(\lambda), \\ & \prod_{s=2} a_s & \prod_{s=1} a_s & \end{aligned}$$

where $P_{m-1}(\lambda), P_{m-2}(\lambda), \tilde{P}_{m-2}(\lambda)$ and $P_{m-3}(\lambda)$ are real polynomials in λ of degree less than or equal to $m-1, m-2, m-2$ and $m-3$, respectively, and $P_k(\lambda) = 0$ for $k < 0$.

Proof. We use induction on m , $m \geq 2$. For the base case of $m = 2$ we have $a_{11} = c_1, a_{12} = \frac{\lambda-b_1}{a_1}, a_{21} = c_1 \frac{\lambda-b_2}{a_2}$ and $a_{22} = c_2 + \frac{(\lambda-b_2)(\lambda-b_1)}{a_2 a_1}$. Immediately, they satisfy the hypothesis. Now assume the result holds up to $m = k$. To prove the result for $m = k+1$, observe that by induction,

$$\begin{aligned} \prod_{s=1}^{k+1} \begin{pmatrix} 0 & 1 \\ c_s & \frac{\lambda-b_s}{a_s} \end{pmatrix} &= \begin{pmatrix} 0 & 1 \\ c_{k+1} & \frac{\lambda-b_{k+1}}{a_{k+1}} \end{pmatrix} \prod_{s=1}^k \begin{pmatrix} 0 & 1 \\ c_s & \frac{\lambda}{a_s} \end{pmatrix} \\ &= \begin{pmatrix} 0 & 1 \\ c_{k+1} & \frac{\lambda-b_{k+1}}{a_{k+1}} \end{pmatrix} \begin{pmatrix} c_1 \frac{\lambda^{k-2}}{\prod_{s=2}^{k-1} a_s} + P_{k-3} & \frac{\lambda^{k-1}}{\prod_{s=1}^{k-1} a_s} + P_{k-2} \\ c_1 \frac{\lambda^{k-1}}{\prod_{s=2}^k a_s} + \tilde{P}_{k-2} & \frac{\lambda^k}{\prod_{s=1}^k a_s} + P_{k-1} \end{pmatrix} \\ &= \begin{pmatrix} c_1 \frac{\lambda^{k-1}}{\prod_{s=2}^k a_s} + \tilde{P}_{k-2} & \frac{\lambda^k}{\prod_{s=1}^k a_s} + P_{k-1} \\ c_1 \frac{\lambda^k}{\prod_{s=2}^{k+1} a_s} + \tilde{P}_{k-1} & \frac{\lambda^{k+1}}{\prod_{s=1}^{k+1} a_s} + P_k \end{pmatrix}. \quad \square \end{aligned}$$

Corollary 2.2. *Let M be the monodromy matrix for an arbitrary period- T operator, i.e.*

$$M(\lambda) = \begin{pmatrix} m_{11}(\lambda) & m_{12}(\lambda) \\ m_{21}(\lambda) & m_{22}(\lambda) \end{pmatrix} := B_T(\lambda) B_{T-1}(\lambda) \dots B_1(\lambda)$$

and $B_i(\lambda)$ are the transfer matrices given by

$$B_i(\lambda) := \begin{pmatrix} 0 & 1 \\ \frac{-a_{i-1}}{a_i} & \frac{\lambda-b_i}{a_i} \end{pmatrix}, \lambda \in \mathbb{C},$$

where $i = 1, 2, \dots, T$, with $a_0 := a_T$. Then, $\det(M(\lambda)) = 1$ and for all $T \geq 1$ we have

$$\begin{aligned} m_{11}(\lambda) &= -a_T \frac{\lambda^{T-2}}{\prod_{s=1}^{T-1} a_s} + P_{T-3}(\lambda), & m_{12}(\lambda) &= \frac{\lambda^{T-1}}{\prod_{s=1}^{T-1} a_s} + P_{T-2}(\lambda), \\ m_{21}(\lambda) &= -\frac{\lambda^{T-1}}{\prod_{s=1}^{T-1} a_s} + \tilde{P}_{T-2}(\lambda), & m_{22}(\lambda) &= \frac{\lambda^T}{\prod_{s=1}^T a_s} + P_{T-1}(\lambda), \end{aligned}$$

where $P_{T-1}(\lambda), P_{T-2}(\lambda), \tilde{P}_{T-2}(\lambda)$ and $P_{T-3}(\lambda)$ are real polynomials in λ of degree less than or equal to $T-1, T-2, T-2$ and $T-3$, respectively.

Proof. We will refer to the situation that $\sigma_{ess}(J_{\epsilon,\eta})$ consists of T distinct real intervals as the non-degenerate case. The proof will consist of two parts: We show that non-degeneracy is stable under small perturbations, while on the other hand the degenerate case is not stable. We initially introduce some notation.

Define the transfer matrices for $J_{\epsilon,\eta}$ as

$$B_i(\lambda) := \begin{pmatrix} 0 & 1 \\ \frac{-a_{i-1}}{a_i} & \frac{\lambda-b_i}{a_i} \end{pmatrix},$$

for $i = 3, 4, \dots, T$, and

$$B_{1,\epsilon,\eta}(\lambda) := \begin{pmatrix} 0 & 1 \\ \frac{-a_T}{a_1+\eta} & \frac{\lambda-b_1-\epsilon}{a_1+\eta} \end{pmatrix}, \quad B_{2,\epsilon,\eta}(\lambda) := \begin{pmatrix} 0 & 1 \\ -\frac{a_1+\eta}{a_2} & \frac{\lambda-b_2}{a_2} \end{pmatrix}.$$

Let M_0 be the monodromy matrix for $J_{0,0}$, i.e. $M_0 := B_T B_{T-1} \dots B_{2,0,0} B_{1,0,0}$, and $M_{\epsilon,\eta}$ the monodromy matrix for $J_{\epsilon,\eta}$, i.e.

$$M_{\epsilon,\eta} := B_{T,0,0} B_{T-1,0,0} \dots B_{3,0,0} B_{2,\epsilon,\eta} B_{1,\epsilon,\eta} = M_0 B_{1,0,0}^{-1} B_{2,0,0}^{-1} B_{2,\epsilon,\eta} B_{1,\epsilon,\eta}. \quad (3)$$

By Corollary 2.2 we have

$$M_0(\lambda) = \begin{pmatrix} p_1(\lambda) & p_2(\lambda) \\ p_3(\lambda) & p_4(\lambda) \end{pmatrix},$$

where $p_1(\lambda), p_2(\lambda), p_3(\lambda)$ and $p_4(\lambda)$ are real polynomials in λ of order $T-2, T-1, T-1$ and T respectively. Then, as $\lambda \mapsto (\text{Tr}(M_0(\lambda)) \pm 2)$ are two polynomials each of degree T in λ , there are at most $2T$ real zeros of these functions, providing at most T intervals of a.c. spectrum. Recall from Lemma 2.5 that all of the parabolic points for J_T are real.

(Step One) It needs to be shown that if $J_{0,0}$ is non-degenerate, then adding sufficiently small ϵ, η to the operator does not cause two previously distinct parabolic points to overlap. The argument is simple: For each pair of distinct parabolic points (λ_j, λ_k) of $J_{0,0}$ there exists $\delta_{j,k} > 0$ such that for the corresponding parabolic points of $J_{\epsilon,\eta}$ we have $\lambda_j(\epsilon, \eta) \neq \lambda_k(\epsilon, \eta)$ for $|\epsilon|, |\eta| < \delta_{j,k}$, using the fact that the roots of a polynomial depend continuously on its coefficients. (See, for example, Appendix A in [17].) Then, since there are at most $2T$ parabolic points in total, we can define

$$\delta := \min_{j,k} \delta_{j,k} > 0$$

which implies

$$\lambda_m(\epsilon, \eta) \neq \lambda_n(\epsilon, \eta)$$

for all $|\epsilon|, |\eta| < \delta, m, n \in \{1, \dots, 2T\}, m \neq n$. This shows that the non-degenerate case is stable.

(Step Two) We now show that the case where two of the intervals of essential spectrum of $J_{0,0}$ overlap is unstable. Let λ_0 be a parabolic point for $J_{0,0}$. We will only consider the case when $\text{Tr}(M_0(\lambda_0)) = 2$, the case $\text{Tr}(M_0(\lambda_0)) = -2$ can be dealt with similarly. Assume that $\lambda_0 \notin \partial\sigma_{ess}(J_{0,0})$. Then $\frac{d}{d\lambda} \text{Tr}(M_0(\lambda_0)) = 0$, otherwise $\text{Tr}(M_0(\lambda)) - 2$ would change sign at λ_0

and λ_0 would separate the elliptic and hyperbolic regions, implying $\lambda_0 \in \partial\sigma_{ess}(J_{0,0})$.

We now show that in most cases a diagonal perturbation is sufficient to split the overlapping intervals. Assume that

$$|p_3(\lambda_0)| + |p'_3(\lambda_0)| \neq 0. \quad (4)$$

Let λ (depending on ϵ) be a degenerate parabolic point for some $J_{\epsilon,0}$, i.e.

$$\text{Tr}(M_{\epsilon,0}(\lambda)) = 2 \quad \text{and} \quad \frac{d}{d\lambda}\text{Tr}(M_{\epsilon,0}(\lambda)) = 0 \quad (5)$$

Our objective is to show that this cannot happen for sufficiently small $\lambda - \lambda_0$ and ϵ . Due to continuous dependence of the roots on the small parameter ϵ , there is no need to consider the case $\text{Tr}(M_{\epsilon,0}(\lambda)) = -2$. Also, (4) will hold (for the same polynomial p_3 from M_0) with λ_0 replaced by λ in a sufficiently small neighbourhood of λ_0 . Noting that $B_{2,\epsilon,0}$ is independent of ϵ we get that

$$\begin{aligned} M_{\epsilon,0} &= B_T B_{T-1} \dots B_{2,0,0} \left(B_{1,0,0} - \frac{\epsilon}{a_1} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right) \\ &= M_0 \left(I - \frac{\epsilon}{a_1} B_{1,0,0}^{-1} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right). \end{aligned}$$

As

$$B_{1,0,0}^{-1}(\lambda) = \begin{pmatrix} \frac{\lambda - b_1}{a_T} & -\frac{a_1}{a_T} \\ 1 & 0 \end{pmatrix},$$

we get

$$\text{Tr}(M_{\epsilon,0}(\lambda)) = p_1(\lambda) + p_4(\lambda) + \frac{\epsilon p_3(\lambda)}{a_T}. \quad (6)$$

Now, Equations (6) and (5) combine to give the new conditions

$$p_1(\lambda) + p_4(\lambda) + \frac{\epsilon p_3(\lambda)}{a_T} = 2 \quad (7)$$

and

$$p'_1(\lambda) + p'_4(\lambda) + \frac{\epsilon p'_3(\lambda)}{a_T} = 0. \quad (8)$$

If Equations (7) and (8) are both satisfied then we obtain

$$(2 - p_1(\lambda) - p_4(\lambda))p'_3(\lambda) + (p'_1(\lambda) + p'_4(\lambda))p_3(\lambda) = 0. \quad (9)$$

By invoking Corollary 2.2 we observe that the product of polynomials on the left hand side equals

$$2p'_3(\lambda) - \left(\frac{p_1(\lambda) + p_4(\lambda)}{p_3(\lambda)} \right)' p_3^2(\lambda). \quad (10)$$

Note that the term $2p'_3(\lambda)$ is a polynomial of degree not greater than $(T-2)$, the rational function $\left(\frac{p_1(\lambda) + p_4(\lambda)}{p_3(\lambda)} \right)'$ is of order 0, and the term $p_3^2(\lambda)$ is a polynomial of degree $2(T-1)$. Combining these observations we have that the entire last term of the expression is a polynomial of degree $2(T-1)$. Since $2(T-1)$ is greater than $(T-2)$ the whole expression is of degree $2(T-1)$.

Clearly, this is not identically zero. Furthermore, this means there are at most $2T - 2$ roots, say μ_1, \dots, μ_{2T-2} which are independent of ϵ . Then, since under our assumptions $|p_3(\lambda)| + |p'_3(\lambda)| \neq 0$, we calculate the valid values for ϵ by substituting $\lambda := \mu_i$ into either Equation (7) or (8), and, so, there are at most $2T - 2$ valid values for ϵ . In particular, for any sufficiently small $\epsilon \neq 0$, the value λ cannot be a degenerate parabolic point for $J_{\epsilon,0}$. Therefore, all degenerate parabolic points satisfying (4) will be split into non-degenerate points for $|\epsilon| \neq 0$ sufficiently small.

It remains to deal with the exceptional case $p_3(\lambda_0) = p'_3(\lambda_0) = 0$. In this case, we use a perturbation with $\epsilon = 0, \eta \neq 0$. Note that since $\text{Tr}(M_0(\lambda_0)) = 2$ and $\det M_0(\lambda_0) = 1$, we have that $p_3(\lambda_0) = 0$ implies $p_1(\lambda_0) = p_4(\lambda_0) = 1$. Then

$$B_{1,0,0}^{-1} B_{2,0,0}^{-1} B_{2,0,\eta} B_{1,0,\eta} = \begin{pmatrix} \frac{a_1}{a_1+\eta} & \frac{\lambda-b_1}{a_T} \left(\frac{a_1+\eta}{a_1} - \frac{a_1}{a_1+\eta} \right) \\ 0 & \frac{a_1+\eta}{a_1} \end{pmatrix}$$

and using (3) we get

$$\text{Tr} M_{0,\eta}(\lambda) = p_1(\lambda) \frac{a_1}{a_1+\eta} + p_3(\lambda) \frac{\lambda-b_1}{a_T} \left(\frac{a_1+\eta}{a_1} - \frac{a_1}{a_1+\eta} \right) + p_4(\lambda) \frac{a_1+\eta}{a_1}.$$

Evaluating at λ_0 , we get

$$\text{Tr} M_{0,\eta}(\lambda_0) = \frac{a_1}{a_1+\eta} + \frac{a_1+\eta}{a_1} = \frac{a_1^2 + (a_1+\eta)^2}{a_1(a_1+\eta)} = 2 + \frac{\eta^2}{a_1(a_1+\eta)} > 2,$$

for all $|\eta| \neq 0$, so λ_0 is a hyperbolic point for $J_{0,\eta}$ for $\eta \neq 0$. Choosing $|\eta|$ sufficiently small such that no non-degenerate parabolic points can degenerate (see Step One), this implies that the total degeneracy of the roots at must have decreased by at least one.

Repeating the procedure finitely many times, we can ensure that all roots of $\text{Tr} M_{\epsilon,\eta}(\lambda) - 2$ are simple for sufficiently small non-zero (ϵ, η) . Note that in each step, ϵ or η may be chosen arbitrarily small. Since the set of non-degenerate points is open by Step 1, the set of points (ϵ, η) is an open dense set. \square

As a consequence of the above, we obtain the following theorem.

Theorem 2.7. *For almost all choices of parameters $(a_1, \dots, a_T, b_1, \dots, b_T) \in (\mathbb{R}^+)^T \times \mathbb{R}^T$ the essential spectrum (which equals the absolutely continuous spectrum) of the associated Hermitian T -periodic Jacobi matrix consists of T distinct real intervals.*

3. Solutions to period-T difference equations

In this section, and the next, the subsidiary functions our eigenvector will depend upon are defined. From now on until the end of the paper we will assume for simplicity that $b_i = 0$ for all i . This restriction is simple to remove.

Firstly, define $B_j(\lambda) := \begin{pmatrix} 0 & 1 \\ -\frac{a_{j-1}}{a_j} & \frac{\lambda}{a_j} \end{pmatrix}$, where $j \in \{1, \dots, T\}$, and

$$M(\lambda) := B_T(\lambda)B_{T-1}(\lambda) \dots B_1(\lambda).$$

Then, if $\lambda \in \sigma_{ell}(J_T)$ we have that $\sigma(M(\lambda)) = \{e^{i\theta(\lambda)}, e^{-i\theta(\lambda)}\}$ for some real-valued function $\theta(\lambda)$ (the quasi-momentum). Therefore there exists an invertible matrix V such that $M = V^{-1} \begin{pmatrix} \mu & 0 \\ 0 & \bar{\mu} \end{pmatrix} V$, where

$$\mu(\lambda) = e^{i\theta(\lambda)}. \quad (11)$$

Lemma 3.1. *Let $\lambda \in \sigma_{ell}(J_T)$. Then, for any non-zero solution, $(\psi_n)_{n \geq 1}$, to the period- T difference equation, $a_{n-1}u_{n-1} + a_n u_{n+1} = \lambda u_n$, $n > 1$, we have the expression*

$$(Im(\psi_n))^2 = \eta_s(\lambda) \sin(2(k-1)\theta(\lambda) + \phi_s(\lambda)) + \gamma_s(\lambda),$$

where $n = T(k-1) + s$, $s \in \{0, \dots, T-1\}$ and η_s, γ_s are real functions which, along with ϕ_s , are independent of k and $\theta(\lambda)$ is given by Equation (11).

Proof. Since ψ_n satisfies the difference equation, and $n = T(k-1) + s$ with $s \in \{0, \dots, T-1\}$ we have for any $\begin{pmatrix} \psi_0 \\ \psi_1 \end{pmatrix} \in \mathbb{C}^2 \setminus \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

$$\begin{aligned} \begin{pmatrix} \psi_n \\ \psi_{n+1} \end{pmatrix} &= B_s \dots B_1 M^{(k-1)} \begin{pmatrix} \psi_0 \\ \psi_1 \end{pmatrix} \\ &= B_s \dots B_1 V \begin{pmatrix} \mu^{(k-1)}(\lambda) & 0 \\ 0 & \bar{\mu}^{(k-1)}(\lambda) \end{pmatrix} V^{-1} \begin{pmatrix} \psi_0 \\ \psi_1 \end{pmatrix} \\ &= \begin{pmatrix} \alpha_s(\lambda)e^{i(k-1)\theta(\lambda)} + \beta_s(\lambda)e^{-i(k-1)\theta(\lambda)} \\ \kappa_s(\lambda)e^{i(k-1)\theta(\lambda)} + \chi_s(\lambda)e^{-i(k-1)\theta(\lambda)} \end{pmatrix}, \end{aligned}$$

for some functions $\alpha_s, \beta_s, \kappa_s, \chi_s$ of $\lambda \in \sigma_{ell}(J_T)$ and s . In the case of $s = 0$ we interpret $B_0 \dots B_1$ to equal the identity, and $B_1 \dots B_1 = B_1$. Consequently,

$$\psi_n = \psi_{T(k-1)+s} = \alpha_s(\lambda)e^{i(k-1)\theta(\lambda)} + \beta_s(\lambda)e^{-i(k-1)\theta(\lambda)}.$$

Thus,

$$Im(\psi_n) = \tilde{\alpha}_s(\lambda) \sin((k-1)\theta) + \tilde{\beta}_s(\lambda) \cos((k-1)\theta), \quad (12)$$

where $\tilde{\alpha}_s(\lambda) := Re(\alpha_s(\lambda)) - Re(\beta_s(\lambda))$, $\tilde{\beta}_s(\lambda) := Im(\alpha_s(\lambda)) + Im(\beta_s(\lambda))$ are real-valued functions of λ .

Furthermore, using the double-angle formulae, $\sin(2x) = 2 \sin(x) \cos(x)$ and $\cos(2x) = \cos^2(x) - \sin^2(x)$, we have

$$\begin{aligned} (Im(\psi_n))^2 &= \tilde{\alpha}_s^2 \sin^2((k-1)\theta) + \tilde{\beta}_s^2 \cos^2((k-1)\theta) \\ &\quad + \tilde{\alpha}_s \tilde{\beta}_s (2 \sin((k-1)\theta) \cos((k-1)\theta)) \\ &= \eta_s \sin(2(k-1)\theta + \phi_s) + \gamma_s, \end{aligned}$$

where η_s, ϕ_s and γ_s are real-valued. \square

Remark 3.2. Clearly, by suitably choosing ψ_0, ψ_1 the vector $(\eta_0, \dots, \eta_{T-1})$ can be arranged to be non-trivial.

Lemma 3.3. *Let $\lambda \in \sigma_{ell}(J_T)$. Then, there exists a particular non-zero solution, $(\varphi_n)_{n \geq 1}$, to the period- T difference equation $a_{n-1}u_{n-1} + a_n u_{n+1} = \lambda u_n, n > 1$ which has the property*

$$\varphi_n(\lambda) = \varphi_s(\lambda) e^{i(k-1)\theta(\lambda)} \quad (13)$$

for some non-trivial set of functions $(\varphi_s)_{s=0}^{T-1}$, where $n = T(k-1) + s, s \in \{0, \dots, T-1\}$.

Proof. Recall from above that $M(\lambda)$ has eigenvalues $e^{\pm i\theta(\lambda)}$, i.e.

$$M(\lambda) \begin{pmatrix} \varphi_0 \\ \varphi_1 \end{pmatrix} = e^{i\theta(\lambda)} \begin{pmatrix} \varphi_0 \\ \varphi_1 \end{pmatrix},$$

for some φ_0, φ_1 . Define $\varphi_2, \dots, \varphi_{T-1}$ by

$$\begin{pmatrix} \varphi_s \\ \varphi_{s+1} \end{pmatrix} := B_s B_{s-1} \dots B_1 \begin{pmatrix} \varphi_0 \\ \varphi_1 \end{pmatrix}.$$

Then using the notation $n = T(k-1) + s$,

$$\begin{aligned} \begin{pmatrix} \varphi_n \\ \varphi_{n+1} \end{pmatrix} &= B_s \dots B_1 M^{(k-1)} \begin{pmatrix} \varphi_0 \\ \varphi_1 \end{pmatrix} \\ &= B_s \dots B_1 e^{(k-1)i\theta(\lambda)} \begin{pmatrix} \varphi_0 \\ \varphi_1 \end{pmatrix} \\ &= e^{(k-1)i\theta(\lambda)} \begin{pmatrix} \varphi_s \\ \varphi_{s+1} \end{pmatrix}. \end{aligned}$$

Consequently,

$$\varphi_n = \varphi_s e^{(k-1)i\theta(\lambda)}. \quad \square$$

Remark 3.4. Henceforth, the eigenvector of the monodromy matrix will be normalized with $\varphi_0 = 1$. Subsequent calculations in Lemma 4.3 will confirm the validity of this choice for almost every λ .

4. Properties of the function $C(\lambda; T)$

In this section we introduce a new, analytic function of λ , $C(\lambda; T)$. This will play an important role in the asymptotic expansion of our eigenvector, (u_n) . Its zeros will give values of λ where our construction fails. Here we explore its properties and structure.

Definition 4.1. For $\lambda \in \sigma_{ell}(J_T)$, let $C(\lambda; T) := \operatorname{Re} \left(\sum_{s=1}^T \varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda) \right)$, where φ_s are as in Lemma 3.3.

Note that Definition 4.1 is invariant w.r.t. the choice of branches μ and $\bar{\mu}$ on $\sigma_{ell}(J_T)$. Indeed, since $\lambda \in \sigma_{ell}(J_T) \subset \mathbb{R}$, all matrix elements of $B_s(\lambda), s = 1, 2, \dots, T$ and $M(\lambda)$ are real polynomials, $\varphi_0 = 1$ and

$$\varphi_1(\lambda) = (\mu - m_{11}(\lambda)) m_{12}^{-1}(\lambda)$$

changes under the transformation $\mu \mapsto \bar{\mu}$ to the complex conjugate function $\varphi_1(\lambda) \mapsto \bar{\varphi}_1(\lambda)$, $\lambda \in \sigma_{ell}(J_T)$. For the last fact the inclusion $\lambda \in \sigma_{ell}(J_T)$ is essential. Hence for all $s = 1, \dots, T$ $\varphi_s(\lambda) \mapsto \bar{\varphi}_s(\lambda)$ and the expression for $C(\lambda; T)$, $\lambda \in \sigma_{ell}(J_T)$, transforms into

$$Re \left(\sum_{i=1}^T \bar{\varphi}_s(\lambda) \varphi_{s-1}(\lambda) \right) = Re \left(\overline{\sum_{i=1}^T \varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda)} \right) = C(\lambda; T).$$

Definition 4.2. Consider a rational function in the variable x of the form $\frac{P(x)}{Q(x)}$ where $P(x), Q(x)$ are polynomials. The order of the rational function is defined to be the difference in degree of the polynomials $P(x)$ and $Q(x)$: degree $P - \text{degree } Q$.

Lemma 4.3. *The function $C(\lambda; T)$ is a rational function on $\sigma_{ell}(J_T)$ and can be extended, uniquely, as a rational function to \mathbb{C} .*

Proof. Let $\lambda \in \sigma_{ell}(J_T)$. First, the special cases of $T = 1$ and $T = 2$ must be considered separately. For $T = 1$, $\varphi_0 = 1$ and $\varphi_1 = \mu$, so

$$C(\lambda; 1) = Re(\varphi_1 \bar{\varphi}_0) = Re(\mu) = \frac{\text{Tr}(M(\lambda))}{2} = \frac{\lambda}{2a_1}.$$

For $T = 2$ we have

$$M(\lambda) = \begin{pmatrix} -\frac{a_2}{a_1} & \frac{\lambda}{a_1} \\ -\frac{\lambda}{a_1} & \frac{\lambda^2}{a_1 a_2} - \frac{a_1}{a_2} \end{pmatrix}.$$

By defining $\varphi_0 := 1$, φ_1 is such that

$$M(\lambda) \begin{pmatrix} \varphi_0 \\ \varphi_1 \end{pmatrix} = e^{i\theta(\lambda)} \begin{pmatrix} \varphi_0 \\ \varphi_1 \end{pmatrix}.$$

Consequently,

$$\varphi_1 = \frac{a_1 \mu + a_2}{\lambda}, \varphi_2 = \mu.$$

Then, using $\mu \bar{\mu} = 1$,

$$\begin{aligned} C(\lambda; 2) &= Re(\varphi_2 \bar{\varphi}_1 + \varphi_1 \bar{\varphi}_0) \\ &= Re \left(\mu \frac{a_1 \bar{\mu} + a_2}{\lambda} + \frac{a_1 \mu + a_2}{\lambda} \right) \\ &= \frac{(a_1 + a_2)}{\lambda} (1 + Re(\mu)) = \frac{(a_1 + a_2)}{\lambda} \left(1 + \frac{\text{Tr}(M(\lambda))}{2} \right) \\ &= \frac{(a_1 + a_2)}{2\lambda a_1 a_2} (\lambda^2 - (a_1^2 + a_2^2) + 2a_1 a_2) = \frac{(a_1 + a_2)}{2\lambda a_1 a_2} (\lambda^2 - |a_1 - a_2|^2). \end{aligned} \tag{14}$$

Thus, the assertion holds for both of these cases.

For $T \geq 3$ we define $\varphi_0 := 1$ and follow a similar technique to the case for $T = 2$. Here we see that the normalisation $\varphi_0 = 1$ is valid unless $m_{12}(\lambda) = 0$. Consequently, for $m_{12}(\lambda) \neq 0$,

$$\varphi_1 = \frac{\mu - m_{11}}{m_{12}},$$

where m_{11}, m_{12} are as described in Corollary 2.2. Throughout the proof P_k will denote a polynomial of at most degree k , while R_k, \tilde{R}_k will denote rational functions of order at most k . Using Lemma 2.2, again, and a similar calculation as in the case of Lemma 3.3, for $s = 2, \dots, T$ we obtain

$$\varphi_s = \left(\frac{\lambda^{s-1}}{\prod_{j=1}^{s-1} a_j} + P_{s-2} \right) \frac{\mu - m_{11}}{m_{12}} + \left(-\frac{a_T \lambda^{s-2}}{\prod_{j=1}^{s-1} a_j} + P_{s-3} \right),$$

where P_{-1} and \tilde{P}_{-1} are both identically zero. Note that $Im(\mu)$ is an algebraic but not rational function of λ . Indeed, $\text{Tr}(M(\lambda))$ is a polynomial in λ and $\det(M(\lambda)) = 1$, therefore $Im(\mu)$ is the root of $\left(\frac{\text{Tr}(M(\lambda))}{2}\right)^2 + 1$, which would be the square of a rational function iff $\text{Tr}(M(\lambda))$ were equal to a constant. However, since $\mu\bar{\mu} = 1$ and

$$Re(\mu) = Re(\bar{\mu}) = \frac{\text{Tr}(M(\lambda))}{2},$$

$Re(\varphi_s(\lambda)\bar{\varphi}_{s-1}(\lambda))$ is clearly a rational function of λ , $C(\lambda; T)$ is also a rational function of λ , $\lambda \in \sigma_{ell}(J_T)$. Now we see that $C(\lambda; T)$ is well-defined as an analytic function not only on $\sigma_{ell}(J_T)$, but everywhere on \mathbb{C} except at the roots of $m_{12}(\lambda)$. \square

Remark 4.4. The function $C(\lambda; T)$ only fails to be defined when the polynomial $m_{12}(\lambda)$, defined in Corollary 2.2, is equal to 0. For $\lambda \in \sigma_{ell}(J_T)$ we have $m_{12}(\lambda) \neq 0$ since if $m_{12}(\lambda) = 0$ then the eigenvalues of the monodromy matrix for real λ are real, and as usual their product is 1. Indeed, since $m_{12}(\lambda) = 0$, we have that the monodromy matrix is lower-triangular and therefore $m_{11}(\lambda)$ and $m_{22}(\lambda)$ are the (real) eigenvalues. Thus, λ is either in the hyperbolic or parabolic case, contradicting that $\lambda \in \sigma_{ell}(J_T)$, and so the denominator has no roots in $\sigma_{ell}(J_T)$.

Our technique for embedding eigenvalues fails for values λ when the function $C(\lambda; T) = 0$. It is important to understand when this situation arises.

Remark 4.5. For the case $T = 1$, the function $C(\lambda; 1)$ has only one root at $\lambda = 0$. From Equation (14) we know that for the case $T = 2$ the function $C(\lambda; 2)$ has no zeros for $\lambda \in \sigma_{ell}(J_2)$ as its two roots, $\lambda_{\pm} = \pm|a_1 - a_2|$, are parabolic points. For the case $T = 3$ the function

$$C(\lambda; 3) = \frac{\lambda \left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} \right)}{2(\lambda^2 - a_1^2)} \left(\lambda^2 - (a_1^2 + a_2^2 + a_3^2) + \frac{2(a_1 + a_2 + a_3)}{\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3}} \right) \quad (15)$$

has a zero at $\lambda = 0$. In order to preclude any other roots in the generalised interior of the a.c. spectrum (see Definition 2.3) it is sufficient to establish that $|\text{Tr}(M(\lambda))| \geq 2$ whenever $C(\lambda; 3) = 0$. A simple calculation shows that

this is equivalent to

$$g(a_1, a_2) := (a_1^3 + a_1^3 a_2 + a_2^3 + a_1 a_2^3 + a_2 + a_1 - a_1^2 a_2 - a_1 a_2^2 - a_1 a_2)(a_1 + a_2 + 1)^2 - (a_1 + a_2 + a_1 a_2)^3 \geq 0,$$

where, by homogeneity, w.l.o.g $a_3 = 1$. Numerical calculations of the roots of g suggest that this function is non-negative for $a_1, a_2 > 0$. More generally, we believe that for even T the function $C(\lambda; T)$ has no zeros in the generalized interior of the a.c. spectrum, and for odd T there is a single solution at $\lambda = 0$.

Remark 4.6. Now consider a different formula for $C(\lambda; T)$ having a ‘‘symplectic character’’. Using it one can easily deduce, in a slightly different way, the rationality of $C(\lambda; T)$. Introducing the indefinite matrix $\hat{\mathcal{J}} := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ in \mathbb{C}^2 one can rewrite the expression for $C(\lambda; T)$ in the following form (we assume below that $\lambda \in \sigma_{ell}(J_T)$):

$$\begin{aligned} C(\lambda; T) &= \frac{1}{2} \sum_{s=1}^T (\varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda) + \varphi_{s-1}(\lambda) \bar{\varphi}_s(\lambda)) \\ &= \frac{1}{2} \sum_{s=1}^T \left\langle \hat{\mathcal{J}} \left(\prod_{k=1}^{s-1} B_k(\lambda) \right) \begin{pmatrix} \varphi_0 \\ \varphi_1(\lambda) \end{pmatrix}, \left(\prod_{k=1}^{s-1} B_k(\lambda) \right) \begin{pmatrix} \varphi_0 \\ \varphi_1(\lambda) \end{pmatrix} \right\rangle_{\mathbb{C}^2} \\ &= \frac{1}{2} \sum_{s=1}^T \left\langle \left(\prod_{k=1}^{s-1} B_k(\bar{\lambda}) \right)^* \hat{\mathcal{J}} \left(\prod_{k=1}^{s-1} B_k(\lambda) \right) \begin{pmatrix} \varphi_0 \\ \varphi_1(\lambda) \end{pmatrix}, \begin{pmatrix} \varphi_0 \\ \varphi_1(\bar{\lambda}) \end{pmatrix} \right\rangle_{\mathbb{C}^2} \\ &= \sum_{s=1}^T \left\langle F_s(\lambda) \left[\begin{pmatrix} 1 \\ -m_{11}(\lambda) m_{12}^{-1}(\lambda) \end{pmatrix} + \mu(\lambda) \begin{pmatrix} 0 \\ m_{12}^{-1}(\lambda) \end{pmatrix} \right], \right. \\ &\quad \left. \left[\begin{pmatrix} 1 \\ -m_{11}(\bar{\lambda}) m_{12}^{-1}(\bar{\lambda}) \end{pmatrix} + \mu(\bar{\lambda}) \begin{pmatrix} 0 \\ m_{12}^{-1}(\bar{\lambda}) \end{pmatrix} \right] \right\rangle_{\mathbb{C}^2} \end{aligned}$$

where we denoted the real matrix polynomials

$$\frac{1}{2} \left(\prod_{k=1}^{s-1} B_k(\bar{\lambda}) \right)^* \hat{\mathcal{J}} \left(\prod_{k=1}^{s-1} B_k(\lambda) \right)$$

by $F_s(\lambda)$, $s > 1$ and $F_1(\lambda) := \frac{\hat{\mathcal{J}}}{2}$. Therefore

$$\begin{aligned} C(\lambda; T) &= \sum_{s=1}^T \left\{ \left\langle F_s(\lambda) \begin{pmatrix} 1 \\ -m_{11}(\lambda) m_{12}^{-1}(\lambda) \end{pmatrix}, \begin{pmatrix} 1 \\ -m_{11}(\bar{\lambda}) m_{12}^{-1}(\bar{\lambda}) \end{pmatrix} \right\rangle_{\mathbb{C}^2} \right. \\ &\quad + \text{Tr}(M(\lambda)) \left\langle F_s(\lambda) \begin{pmatrix} 1 \\ -m_{11}(\lambda) m_{12}^{-1}(\lambda) \end{pmatrix}, \begin{pmatrix} 1 \\ m_{12}^{-1}(\bar{\lambda}) \end{pmatrix} \right\rangle_{\mathbb{C}^2} \\ &\quad \left. + \left\langle F_s(\lambda) \begin{pmatrix} 1 \\ m_{12}^{-1}(\lambda) \end{pmatrix}, \begin{pmatrix} 1 \\ m_{12}^{-1}(\bar{\lambda}) \end{pmatrix} \right\rangle_{\mathbb{C}^2} \right\} \quad (16) \end{aligned}$$

where we used that $\text{Tr}(M(\lambda)) = \mu(\lambda) + \bar{\mu}(\lambda)$, $\lambda \in \sigma_{ell}(J_T)$. From the last expression, taking into consideration that $F_s(\lambda)$, $m_{11}(\lambda)$, $m_{12}(\lambda)$, $\text{Tr}(M(\lambda))$

are polynomials in λ , we see immediately that $C(\lambda; T)$ is a rational function of λ on $\sigma_{ell}(J_T)$ and therefore admits unique analytic continuation as a rational function to the whole of \mathbb{C} , given by Formula (16). Moreover, using the last formula one can give an upper bound for the order of $C(\lambda; T)$ as a rational function, but in the next theorem we will present an explicit calculation of the order.

Theorem 4.7. *The function $C(\lambda; T)$ is a rational function of λ of order 1. Moreover, its asymptotic expansion is given by*

$$C(\lambda; T) \sim \frac{1}{2} (a_1^{-1} + \cdots + a_T^{-1}) \lambda, \lambda \rightarrow \infty.$$

Proof. (Step One) For $\lambda \in \sigma_{ell}(J_T)$ we have

$$\begin{aligned} C(\lambda; T) &= \sum_{s=1}^T \varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda) - i \operatorname{Im} \left(\sum_{s=1}^T \varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda) \right) \\ &= \sum_{s=1}^T \varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda) - \frac{1}{2} \sum_{s=1}^T (\varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda) - \varphi_{s-1}(\lambda) \bar{\varphi}_s(\lambda)) \end{aligned}$$

Note that by the constancy of the “discrete Wronskian” we know that (using $a_0 := a_T$)

$$a_s (\varphi_{s+1}(\lambda) \bar{\varphi}_s(\lambda) - \varphi_s(\lambda) \bar{\varphi}_{s+1}(\lambda)) = a_{s-1} (\varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda) - \varphi_{s-1}(\lambda) \bar{\varphi}_s(\lambda)), \quad (17)$$

$s = 1, 2, \dots, T-1$. The last identity can be easily proved using the fact that both $\varphi_s(\lambda)$ and $\bar{\varphi}_s(\lambda)$ are solutions to the recurrence relations. Applying Equation (17) one obtains

$$\begin{aligned} C(\lambda; T) &= \left(\sum_{s=1}^T \varphi_s(\lambda) \bar{\varphi}_{s-1}(\lambda) \right) \\ &\quad - \frac{1}{2} (a_1^{-1} + \cdots + a_T^{-1}) a_T (\varphi_1(\lambda) \bar{\varphi}_0(\lambda) - \varphi_0(\lambda) \bar{\varphi}_1(\lambda)) \quad (18) \end{aligned}$$

since

$$\begin{aligned} \varphi_1(\lambda) \bar{\varphi}_0(\lambda) - \varphi_0(\lambda) \bar{\varphi}_1(\lambda) &= \varphi_1(\lambda) - \bar{\varphi}_1(\lambda) \\ &= (\mu(\lambda) - m_{11}(\lambda) m_{12}^{-1}(\lambda)) - (\bar{\mu}(\lambda) - m_{11}(\lambda) m_{12}^{-1}(\lambda)) \\ &= (\mu(\lambda) - \bar{\mu}(\lambda)) m_{12}^{-1}(\lambda) \end{aligned}$$

admits analytic continuation from $\sigma_{ell}(J_T)$ to $\mathbb{C} \setminus \sigma_{ess}(J_T)$ as an analytic (algebraic, but not rational) function $(\mu(\lambda) - \mu^{-1}(\lambda)) m_{12}^{-1}(\lambda)$. It asymptotically behaves like

$$-\operatorname{Tr}(M(\lambda)) m_{12}^{-1}(\lambda) \sim -\lambda^T \left(\prod_{s=1}^T a_s \right)^{-1} \left(\lambda^{T-1} \left(\prod_{s=1}^{T-1} a_s \right)^{-1} \right)^{-1} = -\frac{\lambda}{a_T}$$

assuming that the branch of the analytic function $\mu(\lambda)$ has been chosen so that $\mu(\lambda) \rightarrow 0$, as $\lambda \rightarrow \infty$. Note that in $\mathbb{C}^+ \cup \mathbb{C}^-$ we are in the hyperbolic

situation (Lemma 2.5) so the eigenvalues of $M(\lambda)$ (which are $\mu(\lambda), \mu^{-1}(\lambda)$ as $\det(M(\lambda)) \equiv 1$) with one of them (at our choice) behaving at infinity like $\mu(\lambda) \sim \left(\lambda^T / \left(\prod_{s=1}^T a_s \right) \right)^{-1}$ and the other one like $\mu^{-1}(\lambda) \sim \left(\lambda^T / \prod_{s=1}^T a_s \right)$. Note that the correct choice of the branch of $\mu(\lambda)$ (despite invariance of the definition of $C(\lambda; T)$ under that choice) is crucial for our proof.

Therefore the second term in Equation (18) admits the asymptotics

$$\frac{1}{2} (a_1^{-1} + \dots + a_T^{-1}) \lambda$$

as $\lambda \rightarrow \infty$ according to our choice of the branch $\mu(\lambda)$. The opposite choice of the branch changes the sign in the above mentioned asymptotics of the second term and therefore leads to a sophisticated calculation of the first term, which we are not able to produce here.

(Step Two) We next analyse the asymptotics at infinity of the first term in Formula (18). Since $\varphi_0 = 1$,

$$\varphi_1(\lambda) = (\mu(\lambda) - m_{11}(\lambda))m_{12}^{-1}(\lambda) = O(\lambda^{-1}),$$

as $\mu(\lambda) = O(\lambda^{-T}), \lambda \rightarrow \infty$. For the function $\varphi_s(\lambda)$ we have

$$\begin{aligned} \begin{pmatrix} \varphi_s(\lambda) \\ \varphi_{s+1}(\lambda) \end{pmatrix} &= B_s(\lambda) \dots B_1(\lambda) \begin{pmatrix} \varphi_0 \\ \varphi_1(\lambda) \end{pmatrix} \\ &= (B_{s+1}^{-1}(\lambda) \dots B_T^{-1}(\lambda)) M(\lambda) \begin{pmatrix} \varphi_0 \\ \varphi_1(\lambda) \end{pmatrix} \\ &= \mu(B_{s+1}^{-1}(\lambda) \dots B_T^{-1}(\lambda)) \begin{pmatrix} \varphi_0 \\ \varphi_1(\lambda) \end{pmatrix} \\ &= [\mu(B_{s+1}^{-1}(\lambda) \dots B_T^{-1}(\lambda))] \begin{pmatrix} 1 \\ O(\frac{1}{\lambda}) \end{pmatrix} \\ &= \begin{pmatrix} O(\lambda^{(T-s)-T}) \\ O(\lambda^{(T-s)-T}) \end{pmatrix} = O(\lambda^{-s}) \end{aligned}$$

since obviously the matrix function

$$\mu(\lambda) B_{s+1}^{-1}(\lambda) \dots B_T^{-1}(\lambda) = O(\lambda^{(T-s)-T}),$$

as

$$B_s^{-1}(\lambda) = \frac{a_s}{a_{s-1}} \begin{pmatrix} \lambda & -1 \\ \frac{a_s}{a_{s-1}} & 0 \end{pmatrix} = O(\lambda)$$

and $\mu(\lambda) = O(\lambda^{-T}), \lambda \rightarrow \infty$. Hence

$$\varphi_s(\lambda) = O(\lambda^{-s}), \lambda \rightarrow \infty,$$

$s = 1, 2, \dots, T$.

(Step Three) We now analyse the asymptotics of the analytic continuation of $\overline{\varphi_{s-1}(\lambda)} \equiv \overline{\varphi_{s-1}(\overline{\lambda})}$. Taking the complex conjugate for $\lambda \in \sigma_{ell}(J_T)$

we get

$$\begin{pmatrix} \overline{\varphi}_s(\overline{\lambda}) \\ \overline{\varphi}_{s+1}(\overline{\lambda}) \end{pmatrix} = B_s(\lambda) \dots B_1(\lambda) \begin{pmatrix} 1 \\ (\mu^{-1}(\lambda) - m_{11}(\lambda)) m_{12}^{-1}(\lambda) \end{pmatrix}.$$

Since $\mu^{-1}(\lambda) = O(\lambda^T)$, $\lambda \rightarrow \infty$, for the analytic continuation to \mathbb{C} this gives

$$\begin{pmatrix} \overline{\varphi}_s(\overline{\lambda}) \\ \overline{\varphi}_{s+1}(\overline{\lambda}) \end{pmatrix} = B_s(\lambda) \dots B_1(\lambda) \begin{pmatrix} 1 \\ O(\lambda) \end{pmatrix},$$

where the matrix polynomial $B_s(\lambda) \dots B_1(\lambda) = O(\lambda^s)$, $\lambda \rightarrow \infty$. So,

$$\overline{\varphi}_{s+1}(\overline{\lambda}) = O(\lambda^{s+1}),$$

$\lambda \rightarrow \infty$, $s = 0, 1, \dots, T-1$.

(Step Four) Combining both asymptotic formulas for $\varphi_s(\lambda)$ and $\overline{\varphi}_s(\overline{\lambda})$ we finally obtain

$$\begin{aligned} \sum_{s=1}^T \varphi_s(\lambda) \overline{\varphi}_{s-1}(\overline{\lambda}) &= \sum_{s=1}^T O(\lambda^{-s}) \cdot O(\lambda^{s-1}) \\ &= \sum_{s=1}^T O(\lambda^{-1}) = O(\lambda^{-1}) \end{aligned}$$

as $\lambda \rightarrow \infty$, which leads to the formula

$$C(\lambda; T) = \frac{1}{2} (a_1^{-1} + \dots + a_T^{-1}) \lambda + O(1), \lambda \rightarrow \infty.$$

As a corollary we obtain that the function $C(\lambda; T)$ is always of order exactly 1 and is therefore never identically zero. \square

5. The ansatz for the eigenvector and its asymptotics

In this section we plan to elaborate on the explicit construction of the eigenvector associated with the eigenvalue embedded in the a.c. spectrum of the Jacobi matrix with a diagonal perturbation of Coulomb-type decay.

The following classical result will be used in the next lemma.

Proposition 5.1. (see [22]). *Assume $\alpha, \gamma \in \mathbb{R}$, $\gamma > 0$, then the following estimate holds:*

$$\sum_{k=n}^{\infty} \frac{e^{ik\alpha}}{k^\gamma} = O(1/n^\gamma), n \rightarrow \infty, \iff \frac{\alpha}{2\pi} \notin \mathbb{Z}.$$

We will now introduce the function ω_n which is an important part of the eigenvector of the embedded eigenvalue.

Lemma 5.2. *Let $\lambda \in \sigma_{ell}(J_T)$, $\alpha > 1$ and*

$$\omega_n(\lambda) := \sum_{m=n+1}^{\infty} m^{-\alpha} \text{Im}(\varphi_m(\lambda)) \text{Im}(\varphi_{m-1}(\lambda)), \quad (19)$$

where (φ_n) is defined as in (13). Then

$$\omega_n = \frac{C(\lambda; T)}{2(\alpha - 1)Tn^{\alpha-1}} + O\left(\frac{1}{n^\alpha}\right), n \rightarrow \infty. \quad (20)$$

Moreover, $\omega_n \in l^2$ for $\alpha > \frac{3}{2}$.

Remark 5.3. Formula (20) shows that at zeros of $C(\lambda; T)$ the asymptotics for the function ω_n change drastically. This proves the importance of our analysis in Section 4.

Proof. The proof is divided into two cases.

Case 1 If $n = T(k - 1)$ then by Lemma 3.3 we obtain the relation

$$\begin{aligned} \omega_n &= \sum_{j=k-1}^{\infty} \sum_{s=1}^T (Tj + s)^{-\alpha} \operatorname{Im}(\varphi_{Tj+s}) \operatorname{Im}(\varphi_{Tj+s-1}) \\ &= \sum_{j=k-1}^{\infty} (Tj)^{-\alpha} \left(\sum_{s=1}^{T-1} [\operatorname{Im}(e^{ij\theta}\varphi_s) \operatorname{Im}(e^{ij\theta}\varphi_{s-1})] \right. \\ &\quad \left. + \operatorname{Im}(e^{i(j+1)\theta}\varphi_0) \operatorname{Im}(e^{ij\theta}\varphi_{T-1}) \right) + O(k^{-\alpha}) \\ &= \sum_{j=k-1}^{\infty} (Tj)^{-\alpha} \left(\left[\sum_{s=1}^{T-1} \frac{-1}{4} (e^{ij\theta}\varphi_s - e^{-ij\theta}\bar{\varphi}_s) (e^{ij\theta}\varphi_{s-1} - e^{-ij\theta}\bar{\varphi}_{s-1}) \right] \right. \\ &\quad \left. - \frac{1}{4} (e^{i(j+1)\theta}\varphi_0 - e^{-i(j+1)\theta}\bar{\varphi}_0) (e^{ij\theta}\varphi_{T-1} - e^{-ij\theta}\bar{\varphi}_{T-1}) \right) + O(k^{-\alpha}). \end{aligned}$$

Then, $\theta(\lambda) \notin \pi\mathbb{Z}$ as $\lambda \in \sigma_{\text{ell}}(J_T)$, so by Proposition 5.1

$$\begin{aligned} \omega_n &= \frac{T^{-\alpha}}{4} \sum_{j=k-1}^{\infty} \frac{1}{j^\alpha} \left(\sum_{s=1}^{T-1} (\varphi_s \bar{\varphi}_{s-1} + \bar{\varphi}_s \varphi_{s-1}) + e^{i\theta}\varphi_0 \bar{\varphi}_{T-1} + e^{-i\theta}\bar{\varphi}_0 \varphi_{T-1} \right) \\ &\quad + O(k^{-\alpha}) \\ &= T^{-\alpha} \sum_{j=k-1}^{\infty} \frac{j^{-\alpha}}{2} C(\lambda; T) + O(k^{-\alpha}). \end{aligned}$$

Thus we can apply the Integral Test and obtain

$$\omega_n = \frac{C(\lambda; T)}{2(\alpha - 1)Tn^{\alpha-1}} + O\left(\frac{1}{n^\alpha}\right).$$

Finally, if $C(\lambda; T) \neq 0$,

$$\omega_n \asymp n^{1-\alpha} \in l^2 \iff \alpha > \frac{3}{2}.$$

This proves the result for Case 1.

Case 2 If $n = T(k - 1) + s_n$ with $s_n \in \{1, \dots, T - 1\}$. Then

$$\omega_n = \sum_{j=k}^{\infty} \sum_{s=1}^T (Tj+s)^{-\alpha} \operatorname{Im}(\varphi_{Tj+s}) \operatorname{Im}(\varphi_{Tj+s-1}) + F(n)$$

where, noting that $s_n + 1 \geq 2$,

$$\begin{aligned} F(n) &:= \sum_{s=s_n+1}^T (T(k-1)+s)^{-\alpha} \operatorname{Im}(\varphi_{T(k-1)+s}) \operatorname{Im}(\varphi_{T(k-1)+s-1}) \\ &= \sum_{s=s_n+1}^T (T(k-1)+s)^{-\alpha} \operatorname{Im}\left(e^{i(k-1)\theta} \tilde{\varphi}_s\right) \operatorname{Im}\left(e^{i(k-1)\theta} \tilde{\varphi}_{s-1}\right) \\ &= O(k^{-\alpha}) = O(n^{-\alpha}). \end{aligned}$$

Thus, the remainder can be absorbed in the error term. \square

We now make an ansatz for the eigenvector of the embedded eigenvalue, $\lambda \in \sigma_{\text{ell}}(J_T)$, in the form

$$u_n = \operatorname{Im}(\varphi_n) \omega_n.$$

Theorem 5.4. *The sequence, (u_n) , has the asymptotic form*

$$u_n = \frac{\tilde{\eta}_s \sin(n\theta/T + \tilde{\zeta}_s)}{n^{\alpha-1}} + O\left(\frac{1}{n^\alpha}\right),$$

where $\tilde{\eta}_s$ and $\tilde{\zeta}_s$ are real functions, $\alpha > 1$, $n = T(k-1) + s$ with $s \in \{0, \dots, T-1\}$ and $\theta(\lambda)$ as in Equation (11). Moreover, the vector $(\tilde{\eta}_s)_{s=0}^{T-1}$ is equal to the product of $C(\lambda; T)$ with some non-zero vector. Therefore the only source of vanishing leading terms in the function u_n is the vanishing of $C(\lambda; T)$.

Proof. By Equation (12)

$$\begin{aligned} \operatorname{Im}(\varphi_n) &= \tilde{\alpha}_s \sin((k-1)\theta) + \tilde{\beta}_s \cos((k-1)\theta) \\ &= \sqrt{\tilde{\alpha}_s^2 + \tilde{\beta}_s^2} \left(\frac{\tilde{\alpha}_s}{\sqrt{\tilde{\alpha}_s^2 + \tilde{\beta}_s^2}} \sin((k-1)\theta) + \frac{\tilde{\beta}_s}{\sqrt{\tilde{\alpha}_s^2 + \tilde{\beta}_s^2}} \cos((k-1)\theta) \right) \\ &= \eta'_s \sin((k-1)\theta + \phi'_s) \end{aligned}$$

where $\eta'_s := \sqrt{\tilde{\alpha}_s^2 + \tilde{\beta}_s^2}$ and ϕ'_s are real functions of λ . Then, using Lemma 5.2, we obtain

$$\begin{aligned} u_n &= \operatorname{Im}(\varphi_n) \omega_n \\ &= (\eta'_s \sin((k-1)\theta + \phi'_s)) \left(\frac{2C(\lambda; T)}{(\alpha-1)Tn^{\alpha-1}} + O\left(\frac{1}{n^\alpha}\right) \right) \\ &= \frac{\tilde{\eta}_s \sin((k-1)\theta + \phi'_s)}{n^{\alpha-1}} + O\left(\frac{1}{n^\alpha}\right), \end{aligned}$$

where $\tilde{\eta}_s := \frac{C(\lambda; T)\eta'_s}{2(\alpha-1)T}$. Finally, we wish to express our eigenvector in terms of n . Thus,

$$\begin{aligned} u_n &= \frac{\tilde{\eta}_s \sin((k-1)\theta + \phi'_s)}{n^{\alpha-1}} + O\left(\frac{1}{n^\alpha}\right) \\ &= \frac{\tilde{\eta}_s \sin(n\theta/T + \tilde{\zeta}_s)}{n^{\alpha-1}} + O\left(\frac{1}{n^\alpha}\right), \end{aligned}$$

where $\tilde{\zeta}_s := \phi'_s - s\theta/T$. □

6. The structure of the potential and its asymptotics

The following theorem gives an explicit formula for the potential, and the eigenvector, in terms of the solutions φ_n of the periodic problem, λ and the parameter α .

Theorem 6.1. *Let $\lambda \in \sigma_{ell}(J_T)$ with $C(\lambda; T) \neq 0$. Define $\omega_n(\lambda)$ as in (19) and $\varphi_n(\lambda)$ as in (13) and let $\alpha > \frac{3}{2}$, $n = T(k-1) + s$, $s \in \{0, \dots, T-1\}$,*

$$q_n = -a_{n-1}(Im(\varphi_{n-1}))^2 \left(\frac{n^{-\alpha}}{\omega_n}\right) + a_n(Im(\varphi_{n+1}))^2 \left(\frac{(n+1)^{-\alpha}}{\omega_n}\right). \quad (21)$$

Then

$$u_n(\lambda) = \omega_n(\lambda) Im(\varphi_n(\lambda)) \quad (22)$$

satisfies

$$a_{n-1}u_{n-1} + a_nu_{n+1} + (q_n - \lambda)u_n = 0 \quad (23)$$

for $n \geq 2$. Moreover, q_n has the following asymptotic behaviour:

$$q_n = \frac{1}{n} (\rho_s(\lambda) \sin(2n\theta(\lambda)/T + \zeta_s(\lambda)) + \delta_s(\lambda)) + O\left(\frac{1}{n^2}\right), \quad (24)$$

where ρ_s, ζ_s and δ_s are real functions.

Remark 6.2. In Formula (21) we assume without loss of generality that $\omega_n \neq 0 \forall n = 1, 2, \dots$. Indeed, due to the condition that $C(\lambda; T) \neq 0$ and Formula (20) we see that $\omega_n \neq 0 \forall n \geq L$, where L is sufficiently large. If ω_n vanishes for some $n < L$, then one can change the ansatz for ω_n , (19), by introducing into the sum over m an extra multiple, c_m , where $c_m = 1 \forall m \geq L$. The values c_1, c_2, \dots, c_{L-1} can be chosen in a suitable way such that $\omega_1, \omega_2, \dots, \omega_{L-1}$ are not equal to zero.

Remark 6.3. It can be shown by a lengthy calculation that if $C(\lambda; T) \neq 0$ then

$$\sum_{s=0}^{T-1} |\rho_s(\lambda)|^2 > 0,$$

showing that the potential is genuinely of the form $\frac{1}{n}$ times an oscillating term. This also follows, without any calculation, from the fact that no eigenvalues can be embedded in $\sigma_{ell}(J_T)$ by a potential (q_n) with $q_n = O(\frac{1}{n^2})$. The

proof of this fact for $T = 1$ (the discrete Schrödinger case) is well-known, see, e.g. [15].

Proof. (Step One) Check $u_n(\lambda)$ in (22) satisfies (23). Then, for $n \geq 2$,

$$\begin{aligned}
& a_{n-1}u_{n-1} + a_n u_{n+1} - \lambda u_n = -q_n u_n \\
& \iff a_{n-1}\omega_{n-1}Im(\varphi_{n-1}) + a_n\omega_{n+1}Im(\varphi_{n+1}) - \lambda\omega_n Im(\varphi_n) \\
& \quad = -q_n\omega_n Im(\varphi_n) \\
& \iff Im(a_{n-1}\varphi_{n-1} + a_n\varphi_{n+1} - \lambda\varphi_n)\omega_n + Im(\varphi_{n-1})a_{n-1}(\omega_{n-1} - \omega_n) \\
& \quad + a_n Im(\varphi_{n+1})(\omega_{n+1} - \omega_n) = -q_n Im(\varphi_n)\omega_n \\
& \iff Im(\varphi_{n-1})a_{n-1}(\omega_{n-1} - \omega_n) \\
& \quad + a_n Im(\varphi_{n+1})(\omega_{n+1} - \omega_n) = -q_n Im(\varphi_n)\omega_n \tag{25}
\end{aligned}$$

where we have used that

$$\omega_{n-1} - \omega_n = n^{-\alpha} Im(\varphi_n) Im(\varphi_{n-1}), \tag{26}$$

and that φ_n satisfies the three-term recurrence relation (23). Choosing q_n as in (21) guarantees the equality (25).

Since $\lambda \in \sigma_{ell}(J_T)$, $|\varphi_n| = |\varphi_s|$, which means that $a_{n-1}(Im(\varphi_{n-1}))^2$ and $a_n(Im(\varphi_{n+1}))^2$ are oscillating factors in the variable k . Then one can expect growth or decay in q_n to come from the components $\left(\frac{n-\alpha}{\omega_n}\right)$ and $\left(\frac{(n+1)-\alpha}{\omega_n}\right)$. By Lemma 5.2 we have the relation $\omega_n \asymp n^{1-\alpha}$, and so we obtain

$$\frac{(n+1)^{-\alpha}}{\omega_n} \asymp n^{-1},$$

which gives a Coulomb-type decay for q_n .

(Step Two) We now prove (24). Using Lemmas 3.1 and 5.2 for $n = T(k-1) + s$, $s \in \{1, \dots, T-1\}$, we immediately obtain

$$\begin{aligned}
q_n &= -a_{s-1}(\eta_{s-1} \sin(2(k-1)\theta + \phi_{s-1}) + \gamma_{s-1}) \left(\frac{2(\alpha-1)T}{nC(\lambda; T)} + O\left(\frac{1}{n^2}\right) \right) \\
& \quad + a_s(\eta_{s+1} \sin(2(k-1)\theta + \phi_{s+1}) + \gamma_{s+1}) \left(\frac{2(\alpha-1)T}{nC(\lambda; T)} + O\left(\frac{1}{n^2}\right) \right) \tag{27} \\
&= \frac{2(\alpha-1)T}{nC(\lambda; T)} \left((-\eta_{s-1}a_{s-1} \cos \phi_{s-1} + \eta_{s+1}a_s \cos \phi_{s+1}) \sin(2(k-1)\theta) \right. \\
& \quad \left. + (\eta_{s+1}a_s \sin \phi_{s+1} - \eta_{s-1}a_{s-1} \sin \phi_{s-1}) \cos(2(k-1)\theta) \right. \\
& \quad \left. - \gamma_{s-1}a_{s-1} + \gamma_{s+1}a_s \right) + O\left(\frac{1}{n^2}\right) \\
&= \frac{1}{n} ([\rho_s(\lambda) \sin(2(k-1)\theta(\lambda) + \zeta'_s(\lambda))] + \delta_s) + O\left(\frac{1}{n^2}\right),
\end{aligned}$$

for real functions ζ'_s of λ ,

$$\rho_s^2 := \frac{4T^2(\alpha-1)^2}{C(\lambda; T)^2} \left((\eta_{s+1}a_s \cos \phi_{s+1} - \eta_{s-1}a_{s-1} \cos \phi_{s-1})^2 + (\eta_{s+1}a_s \sin \phi_{s+1} - \eta_{s-1}a_{s-1} \sin \phi_{s-1})^2 \right),$$

$$\text{and } \delta_s := \frac{2T(\alpha-1)}{C(\lambda; T)} (-\gamma_{s-1}a_{s-1} + \gamma_{s+1}a_s).$$

For the special cases of $s \in \{0, T-1\}$ we must be careful because $n-1$ and $n+1$ will produce different values in the parameter k to those contained in the n -th element. When $s=0$ (i.e. $n=(k-1)T$):

$$\begin{aligned} q_n &= -a_{T-1} (\eta_{T-1} \sin(2(k-2)\theta + \phi_{T-1}) + \gamma_{T-1}) \left(\frac{2(\alpha-1)T}{nC(\lambda; T)} + O\left(\frac{1}{n^2}\right) \right) \\ &\quad + a_T (\eta_1 \sin(2(k-1)\theta + \phi_1) + \gamma_1) \left(\frac{2(\alpha-1)T}{nC(\lambda; T)} + O\left(\frac{1}{n^2}\right) \right) \\ &= O\left(\frac{1}{n^2}\right) + \frac{2(\alpha-1)T}{nC(\lambda; T)} \left(-\eta_{T-1}a_{T-1} \sin(2(k-1)\theta + \phi_{T-1} - 2\theta) \right. \\ &\quad \left. + \eta_1 a_0 \sin(2(k-1)\theta + \phi_1) - a_{T-1}\gamma_{T-1} + a_T\gamma_1 \right). \end{aligned}$$

This is of the same form as (27). Consequently,

$$q_n = O\left(\frac{1}{n^2}\right) + \frac{1}{n} [\rho_0(\lambda) \sin(2(k-1)\theta(\lambda) + \zeta'_0) + \delta_0(\lambda)],$$

for functions $\zeta'_0, \delta_0 := \frac{2T(\alpha-1)}{C(\lambda; T)} (-\gamma_{T-1}a_{T-1} + \gamma_1 a_T)$ and

$$\begin{aligned} \rho_0^2 &:= \frac{4T^2(\alpha-1)^2}{C(\lambda; T)^2} \left((\eta_1 a_T \cos \phi_1 - \eta_{T-1} a_{T-1} \cos(\phi_{T-1} - 2\theta))^2 \right. \\ &\quad \left. + (\eta_1 a_T \sin \phi_1 - \eta_{T-1} a_{T-1} \sin(\phi_{T-1} - 2\theta))^2 \right). \end{aligned}$$

Similarly, when $s=T-1$ (i.e. $n=kT-1$):

$$\begin{aligned} q_n &= -a_{T-2} (\eta_{T-2} \sin(2(k-1)\theta + \phi_{T-2}) + \gamma_{T-2}) \left(\frac{2(\alpha-1)T}{nC(\lambda; T)} + O\left(\frac{1}{n^2}\right) \right) \\ &\quad + a_T (\eta_0 \sin(2k\theta + \phi_0) + \gamma_0) \left(\frac{2(\alpha-1)T}{nC(\lambda; T)} + O\left(\frac{1}{n^2}\right) \right) \\ &= O\left(\frac{1}{n^2}\right) + \frac{1}{n} (\rho_{T-1}(\lambda) \sin(2(k-1)\theta(\lambda) + \zeta'_{T-1}) + \delta_{T-1}(\lambda)), \end{aligned}$$

for functions $\zeta'_{T-1}, \delta_{T-1} := \frac{2T(\alpha-1)}{C(\lambda; T)} (-\gamma_{T-2}a_{T-2} + \gamma_0 a_{T-1})$ and

$$\begin{aligned} \rho_{T-1} &:= \frac{4T^2(\alpha-1)^2}{C(\lambda; T)^2} \left((\eta_0 a_{T-1} \cos(\phi_0 + 2\theta - \eta_{T-2} a_{T-2} \cos \phi_{T-2}))^2 \right. \\ &\quad \left. + (\eta_0 a_{T-1} \sin(\phi_0 + 2\theta) - \eta_{T-2} a_{T-2} \sin \phi_{T-2})^2 \right). \end{aligned}$$

Thus, for all $s \in \{0, \dots, T-1\}$, we have the result:

$$q_n = O\left(\frac{1}{n^2}\right) + \frac{1}{n} [\rho_s(\lambda) \sin(2(k-1)\theta(\lambda) + \zeta'_s) + \delta_s(\lambda)].$$

However, we still wish to express our potential in terms of the variable n . This follows simply from defining the new function ζ_s , where $\zeta_s := \zeta'_s - s\theta/T$. \square

Remark 6.4. Concerning the roots of $C(\lambda; T)$ for $\lambda \in \sigma_{ell}(J_T)$, we may say the following. As has been stated in the theorem, using a sufficiently slowly decaying potential, $q_n = O(\frac{1}{n})$, it is possible to introduce a subordinate l^2 -solution for any fixed $\lambda \in \sigma_{ell}(J_T)$, except at roots of $C(\lambda; T)$. However we believe that at any root of $C(\lambda; T)$, $\lambda \in \sigma_{ell}(J_T)$, the existence of the subordinate l^2 -solution can still be obtained by using a potential, $q_n = O(\frac{1}{n})$, $n \rightarrow \infty$.

Remark 6.5. The last statement in the previous remark is true for the case of the discrete Schrödinger operator. To see this choose a candidate eigenvector of the form $u_n = \frac{(-1)^{\lfloor \frac{n}{2} \rfloor}}{n}$ for $n \geq 1$ and a potential defined by $q_n = \frac{2n(-1)^n}{n^2-1}$, for $n \geq 2$ and $q_1 = \frac{1}{2}$. Then clearly $u_2 + q_1 u_1 = 0$ and the recurrence equations

$$\frac{(-1)^{\lfloor \frac{n+1}{2} \rfloor}}{n+1} + \frac{(-1)^{\lfloor \frac{n-1}{2} \rfloor}}{n-1} + \frac{2n(-1)^n}{n^2-1} \frac{(-1)^{\lfloor \frac{n}{2} \rfloor}}{n} = 0,$$

are also satisfied for $n \geq 2$. Thus the eigenvalue $\lambda = 0$ becomes embedded in the a.c. spectrum of the operator.

7. Embedded eigenvalues

Theorem 6.1 guarantees a subordinate solution of the recurrence relation (23), but does not guarantee an embedded eigenvalue since it still remains to be seen if the first-row equation of the Jacobi matrix is satisfied, i.e.

$$q_1 u_1 + a_1 u_2 = \lambda u_1.$$

The next result shows that it is always possible to make λ an eigenvalue by suitably modifying the potential, slightly.

Theorem 7.1. *Assume $\lambda \in \sigma_{ell}(J_T)$, $C(\lambda; T) \neq 0$, $\alpha > \frac{3}{2}$ and let u_n be given by (22) for $n \geq 2$ and q_n by (21) for $n \geq 3$. Then it is possible to choose $u_1, q_1, q_2 \in \mathbb{R}$ such that $\lambda \in \sigma_p(J_T + Q)$, where Q is an infinite diagonal matrix with entries (q_n) .*

Remark 7.2. Note that

$$\sigma_{ess}(J_T) = \sigma_{a.c.}(J_T) = \sigma_{a.c.}(J_T + Q) = \sigma_{ess}(J_T + Q)$$

and

$$\sigma_{ell}(J_T) = \sigma_{ell}(J_T + Q).$$

The coinciding of the essential spectrum of J_T and $J_T + Q$ follows from the classical Weyl Theorem [5]. The preservation of the a.c. spectrum under the perturbation q_n follows from the combination of subordinancy theory [3]

and asymptotic Levinson-type theory [4]. A similar result for the continuous Schrödinger case was proven by Behncke [2].

Proof. By Theorem 6.1 we have

$$a_{n-1}u_{n-1} + a_n u_{n+1} + (q_n - \lambda)u_n = 0$$

for $n \geq 3$. However, we also need to satisfy

$$q_1 u_1 + a_1 u_2 = \lambda u_1 \tag{28}$$

and

$$a_1 u_1 + (q_2 - \lambda)u_2 + a_2 u_3 = 0. \tag{29}$$

We have two cases:

1. If $u_2 \neq 0$ then defining $q_2 := \frac{-\lambda u_2 - a_2 u_3 - a_1 u_1}{u_2}$ with $u_1 := -\frac{a_1 u_2}{q_1 - \lambda}$, with q_1 as a free parameter and not equal to λ , ensures all conditions are satisfied.
2. If $u_2 = 0$ then defining $u_1 := -\frac{a_2 u_3}{a_1}$ and $q_1 := \lambda$, with q_2 as a free parameter, ensures all conditions are satisfied. \square

We are grateful to the unknown referee for his/her very useful remarks.

References

- [1] S. Albeverio, “On bound states in the continuum of N -body systems and the virial theorem”, *Ann. Physics* **71** (1972), no. 1, 167-276.
- [2] H. Behncke, “Absolute continuity of Hamiltonians with von Neumann Wigner potentials”, *Proc. Amer. Math. Soc.* **111** (1991), no. 2, 373-384.
- [3] D. J. Gilbert, D. B. Pearson, “On Subordinacy and analysis of the spectrum of one-dimensional Schrödinger operators”, *J. Math. Anal. Appl.* **128** (1987) no. 1, 30-56.
- [4] J. Janas, M. Moszynski, “New discrete Levinson type asymptotics of solutions of linear systems”, *J. Difference. Equ. Appl.* **12** (2006), no. 2, 133-163.
- [5] T. Kato, “Perturbation theory for linear operators”, Springer Verlag (1966).
- [6] S. Khan, D.B. Pearson, “Subordinacy and spectral theory for infinite matrices”, *Helv. Phys. Acta.* **65** (1992), no. 4, 505-527.
- [7] A. Kiselev, C. Remling, B. Simon, “Effective perturbation methods for one-dimensional Schrödinger operators”, *J. Diff. Equa.* **151** (1999), no. 2, 290-312.
- [8] H. Krüger, “On the existence of embedded eigenvalues”, *J. Math. Anal. App.* **395** (2012), no. 2, 776-787.
- [9] D. Krutikov, “On eigenvalues of discrete Schrödinger operators with potentials of Coulomb-type decay”, *Lett. Math. Phys.* **62** (2002), no. 3, 185-192 .
- [10] P. Kurasov, S. Naboko, “Wigner-von Neumann perturbations of a periodic potential: spectral singularities in bands”, *Proc. Cam. Phil. Soc.*, **142** (2007), no. 1, 161-183.
- [11] M. Lukic, D. C. Ong, “Generalized Prüfer variables for perturbations of Jacobi and CMV matrices”, arXiv:1409.7116.

- [12] M. Lukic, D.C. Ong, “Wigner-von Neumann type perturbations of periodic Schrödinger operators”, *Trans. Amer. Math. Soc.* **367** (2014), no. 1, 707-724.
- [13] S. Naboko, “On the dense point spectrum of the Schrödinger and Dirac operators”, *Teoret. Mat. Fiz.* **68** (1986), 18-28.
- [14] S. Naboko, S. Simonov, “Zeroes of the spectral density of the periodic Schrödinger operator with Wigner-von Neumann potential”, *Proc. Cam. Phil. Soc.* **153** (2012), no. 1, 33-58.
- [15] S. N. Naboko, S. I. Yakovlev, “The discrete Schrödinger operator: the point spectrum lying on the continuous spectrum”, *St. Petersburg Math. J.* **4** (1993), no. 3, 559-568.
- [16] J. von Neumann, E. P. Wigner, “Über merk würdige diskrete Eigenwerte”, *Z. Phys.* **30** (1929), 465-467.
- [17] A. Ostrowski, “Solution of equations in Euclidean and Banach spaces”, Academic Press (1973).
- [18] M. Reed, B. Simon, “Methods of Modern Mathematical Physics, IV: Analysis of Operators”, New York, Academic Press (1978).
- [19] B. Simon, “Orthogonal polynomials on the unit circle, part 1: classical theory”, American Mathematical Society, Providence (2005).
- [20] B. Simon, “Some Jacobi matrices with decaying potential and dense point spectrum”, *Comm. Math. Phys.* **87** (1982/1983), 253-258.
- [21] B. Simon, “Some Schrödinger operators with dense point spectrum”, *Proc. Amer. Math. Soc.* **125** (1997), no. 1, 203-208.
- [22] A. Zygmund, “Trigonometric Series, vol 1”, Cambridge University Press (1959).

Edmund Judge

SMSAS, Cornwallis Building, University of Kent, Canterbury,
Kent CT2 7NF, United Kingdom
e-mail: ej75@kent.ac.uk

Sergey Naboko

Depart. of Math. Physics, Institute of Physics, St.Petersburg University,
St.Petergoff, St.Petersburg, 198904, Russia
e-mail: sergey.naboko@gmail.com

Ian Wood

SMSAS, Cornwallis Building, University of Kent, Canterbury,
Kent CT2 7NF, United Kingdom
e-mail: i.wood@kent.ac.uk