



# Kent Academic Repository

Simon, R., Csengeri, T., Bontemps, S., Wyrowski, F., Menten, K.M., Leurini, S., Urquhart, J.S., Motte, F., Schuller, F., Testi, L. and others (2016) *The first Galaxy scale hunt for the youngest high-mass protostars*. EAS Publications Series, 75-76 . pp. 255-258. ISSN 1633-4760.

## Downloaded from

<https://kar.kent.ac.uk/59983/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1051/eas/1575050>

## This document version

Author's Accepted Manuscript

## DOI for this version

## Licence for this version

CC BY-NC-ND (Attribution-NonCommercial-NoDerivatives)

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal* , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

## Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

*Title : will be set by the publisher*  
*Editors : will be set by the publisher*  
*EAS Publications Series, Vol. ?, 2018*

## THE FIRST GALAXY SCALE HUNT FOR THE YOUNGEST HIGH-MASS PROTOSTARS

T. Csengeri<sup>1</sup>, S. Bontemps<sup>2</sup>, F. Wyrowski<sup>1</sup>, K. M. Menten<sup>1</sup>, S. Leurini<sup>1</sup>, J. S. Urquhart<sup>1</sup>, F. Motte<sup>3</sup>, F. Schuller<sup>4</sup>, L. Testi<sup>4</sup>, L. Bronfman<sup>5</sup>, H. Beuther<sup>6</sup>, S. Longmore<sup>7</sup>, B. Commerçon<sup>8</sup>, Th. Henning<sup>6</sup>, A. Palau<sup>9</sup>, J. C. Tan<sup>10</sup>, G. Fuller<sup>11</sup>, N. Peretto<sup>12</sup>, A. Duarte-Cabral<sup>13</sup> and A. Traficante<sup>11</sup>

**Abstract.** The origin of massive stars is a fundamental open issue in modern astrophysics. Pre-ALMA interferometric studies reveal precursors to early B to late O type stars with collapsing envelopes of 15–20  $M_{\odot}$  on 1000–3000 AU size-scales. To search for more massive envelopes we selected the most massive nearby young clumps from the ATLASGAL survey to study their protostellar content with ALMA. Our first results using the intermediate scales revealed by the ALMA ACA array providing 3–5'' angular resolution, corresponding to  $\sim 0.05$ –0.1 pc size-scales, reveals a sample of compact objects. These massive, dense cores are on average two-times more massive than previous studies of similar types of objects. We expect that once the full survey is completed, it will provide a comprehensive view on the origin of the most massive stars.

---

<sup>1</sup> Max Planck Institute for Radioastronomy, [csengeri@mpifr-bonn.mpg.de](mailto:csengeri@mpifr-bonn.mpg.de)

<sup>2</sup> OASU/LAB, Université Bordeaux

<sup>3</sup> Laboratoire AIM Paris Saclay

<sup>4</sup> European Southern Observatory

<sup>5</sup> Departamento de Astronomía, Universidad de Chile

<sup>6</sup> Max Planck Institute for Astronomy

<sup>7</sup> Astrophysics Research Institute, Liverpool John Moores University

<sup>8</sup> Ecole Normale Supérieure de Lyon

<sup>9</sup> Centro de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México

<sup>10</sup> Departments of Astronomy and Physics, University of Florida

<sup>11</sup> Jodrell Bank Centre for Astrophysics, University of Manchester

<sup>12</sup> School of Physics & Astronomy, Cardiff University

<sup>13</sup> School of Physics and Astronomy, University of Exeter

## 1 Introduction


The origin of stellar masses, in particular of massive stars, is one of the most fundamental open issues of modern astrophysics. In spite of recent progress in our understanding of high-mass star-formation, a coherent picture is still lacking of the earliest evolutionary stages, i.e. , the onset of collapse and the initial fragmentation of massive, dense cores (MDCs) (e.g. Tan *et al.* 2014). Potential precursors of early B to possibly late O stars with  $M_{\text{env}} \sim 10 - 20 M_{\odot}$  have been frequently revealed (e.g. Bontemps *et al.* 2010, Zhang *et al.* 2009, Rathborne *et al.* 2011, Wang *et al.* 2011, Longmore *et al.* 2011, Palau *et al.* 2013, Beuther *et al.* 2013). Their typical sizes of 1000–3000 AU and low bolometric luminosities make them excellent analogs for the low-mass Class 0 stage (Duarte-Cabral *et al.* 2013).

The larger effective Jeans masses of these protostars can be explained by a combination of turbulence and magnetic fields at small scale (McKee & Tan 2003), or by collapse from larger scales at which Jeans masses are larger due to lower average densities (e.g. Hennebelle & Chabrier 2009). Recent numerical models reproduce up to  $10 M_{\odot}$  fragments from a  $100 M_{\odot}$  collapsing core (Commerçon *et al.* 2011), further suggesting that the combined effect of magnetic fields and radiative feedback determines the early fragmentation of massive cores (see also Krumholz *et al.* 2009). To test formation scenarios the precursors of the more massive objects need to be revealed, which becomes now feasible by performing statistical studies using the capabilities of the Atacama Large Millimeter/submillimeter Array (ALMA).

## 2 A sample of the brightest ATLASGAL sources

The ATLASGAL survey (Schuller *et al.* 2009, Csengeri *et al.* 2014) is the most sensitive and extensive ground-based unbiased survey of the inner Galaxy at submillimeter wavelengths, and provides an unprecedented view on all stages of massive star formation. Over 10 000 compact sources have been identified (Csengeri *et al.* 2014), and we have made substantial progress in characterising various evolutionary stages of massive clumps by using ancillary radio and mid-infrared data (e.g. Urquhart *et al.* 2014), and assigned distances to a large number of sources (Wienen *et al.* 2015). Given its higher angular resolution ( $19''.2$ ) compared to *Herschel* at submillimeter wavelengths, the ATLASGAL survey is better suited to disentangle the more evolved from the pristine cold sources.

Selected from ATLASGAL, we identified a complete sample of 45 objects which are massive ( $>650 M_{\odot}$ ) and dense, with surface density exceeding the theoretical  $1 \text{ g cm}^{-2}$  threshold (Krumholz *et al.* (2009) for forming high-mass stars. In addition they lack bright mid-infrared embedded objects, suggesting that they are in their earliest evolutionary phase. Our selection of these mid-infrared quiet massive clumps is complete within 4.5 kpc, and to date represents the best potential sites to host the next generation of the most massive stars currently forming in our Galaxy.



**Fig. 1.** Line-free continuum observations at  $870\ \mu\text{m}$  of 40 of the targeted sources with ALMA ACA at  $3\text{--}5''$  angular resolution.

### 3 ALMA observations of precursors of the youngest massive protoclusters

We used ALMA to perform the first systematic survey at high angular-resolution to look for high-mass protostars in this sample of massive clumps. As a first step we present here the results of the Atacama Compact Array (ACA) observations, which provides a  $3\text{--}5''$  angular resolution corresponding to  $\sim 0.05\text{--}0.1$  pc physical scales (Fig. 1). This size-scale shows the fragmentation from clump to core scales, and reveals a sample of MDCs.

The first result of this survey is to confirm the presence of compact embedded sources towards our selection of mid-infrared quiet ATLASGAL clumps. Only one source out of the entire survey is found to be resolved out lacking embedded compact objects, while the rest all shows at least one centrally concentrated fragment.

Interestingly, the sample shows limited fragmentation, with only 2–3 MDCs revealed per clump, which is similar to earlier findings by Motte *et al.* (2007) and Bontemps *et al.* (2010) in the systematic study of the Cygnus-X region. A first mass estimate shows, however that they are on average twice as massive as the MDCs in Cygnus-X. Among them we also find several candidates to be the most massive MDCs known to date (Csengeri *et al. in prep*).

## 4 Conclusions and Outlook

A first analysis of fragmentation on intermediate scales reveals a large number of MDCs within the sample of mid-infrared quiet ATLASGAL sources targeted with ALMA. They are extreme in terms of their mass and surface density, potentially hosting the precursors of the most massive stars in our Galaxy. Complementary molecular line data will be used to further investigate the nature of their embedded sources. The full capacities of ALMA requested for this survey will provide 0.6" angular resolution, corresponding to individual protostars. This dataset will ultimately provide a comprehensive view of the earliest phase of the formation of high-mass protostars.

## References

- Beuther, H.; Linz, H.; Tackenberg, J. et al. 2013, *A&A*, 553, 115  
Bontemps, S.; Motte, F.; Csengeri, T.; Schneider N. 2010, *A&A*, 524, 18  
Commerçon, B.; Hennebelle, P.; Henning, Th., 2011, *ApJ*, 742, 9  
Csengeri, T., Urquhart, J. 2014, *A&A*, 565, 75  
Duarte-Cabral, A.; Bontemps, S.; Motte, F et al. *A&A*, 558, 125  
Hennebelle & Chabrier 2008, *ApJ*, 684, 395  
Krumholz, M. R.; McKee, Ch. F. 2008, *Nature*, 451, 1082  
Krumholz, M. R.; Klein, R. I.; McKee, Ch. F. 2012, *ApJ*, 754, 71  
Longmore, S. N.; Pillai, T.; Keto, E. et al. 2011, *ApJ*, 726, 97  
McKee & Tan 2003, *ApJ*, 585, 850  
Motte, F.; Bontemps, S.; Schilke, P.; 2007, *A&A*, 476, 1243  
Palau, A.; Fuente, A.; Girart, J. M. et al. , 2013, *ApJ*, 762, 120  
Rathborne, J. M.; Garay, G.; Jackson, J. M. et al. 2011, *ApJ*, 741, 120  
Schuller, F.; Menten, K. M.; Contreras, Y.; 2009, *A&A*, 504, 415  
Tan, J. C.; Beltrán, M. T.; Caselli, P. et al. 2014, *PPVI*, University of Arizona Press, Tucson, 49-172  
Urquhart, J., Moore, T., Csengeri, T., 2014, *MNRAS*, 443, 1555  
Wienen, M., Wyrowski, F. 2015, *A&A*, in press, ArXiv:150300007W  
Zhang, Q.; Wang, Y.; Pillai, T.; Rathborne, J. 2009, *ApJ*, 696, 268  
Wang, Y.; Beuther, H.; Bik, A. et al. 2011, *A&A*, 527, 32