



Kent Academic Repository

Salim, K, Fenton, TR, Bacha, J, Urien-Rodriguez, H, Bonnert, T, Skynner, HA, Watts, E, Kerby, J, Heald, A, Beer, M and others (2002) *Oligomerization of G-protein-coupled receptors shown by selective co-immunoprecipitation*. *Journal of Biological Chemistry*, 277 (18). pp. 15482-15485. ISSN 0021-9258.

Downloaded from

<https://kar.kent.ac.uk/61530/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1074/jbc.M201539200>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Oligomerization of G-protein-coupled Receptors Shown by Selective Co-immunoprecipitation*

Received for publication, February 14, 2002, and in revised form, February 15, 2002
Published, JBC Papers in Press, February 19, 2002, DOI 10.1074/jbc.M201539200

Kamran Salim‡, Tim Fenton, Jamil Bacha, Hector Urien-Rodriguez, Tim Bonnert,
Heather A. Skynner, Emma Watts, Julie Kerby, Anne Heald, Margaret Beer, George McAllister,
and Paul C. Guest

From the Neuroscience Research Centre, Merck Sharp & Dohme Research Laboratories, Harlow,
Essex CM20 2QR, United Kingdom

Recent studies have shown that G-protein-coupled receptors (GPCRs) can assemble as high molecular weight homo- and hetero-oligomeric complexes. This can result in altered receptor-ligand binding, signaling, or intracellular trafficking. We have co-transfected HEK-293 cells with differentially epitope-tagged GPCRs from different subfamilies and determined whether oligomeric complexes were formed by co-immunoprecipitation and immunoblot analysis. This gave the surprising result that the 5HT_{1A} receptor was capable of forming hetero-oligomers with all GPCRs tested including the 5HT_{1B}, 5HT_{1D}, EDG₁, EDG₃, GPR₂₆, and GABA_{B2} receptors. The testing of other GPCR combinations showed similar results with hetero-oligomer formation occurring for the 5HT_{1D} with the 5HT_{1B} and EDG₁ receptor. Control studies showed that these complexes were present in co-transfected cells before the time of lysis and that the hetero-oligomers were comprised of GPCRs at discrete stoichiometries. These findings suggest that GPCRs have a natural tendency to form oligomers when co-transfected into cells. Future studies should therefore investigate the presence and physiological role of GPCR hetero-oligomers in cells in which they are endogenously expressed.

Recent studies have shown that G-protein-coupled receptors (GPCRs)¹ may form dimers or higher order oligomers (1–7). This has led to some re-evaluation of the mechanisms thought to be involved in GPCR function. Co-expression studies with chimeric α_{2C} -adrenergic/M3 muscarinic receptor constructs have shown that intermolecular interactions can occur between different GPCR subtypes (1, 2). More direct studies have shown that the GABA_BR1 and GABA_BR2 receptors are not functional as separate units and can only form a functional receptor complex with the correct pharmacological properties and plasma membrane expression when they are co-expressed (8–12). The finding that co-transfection of the κ and δ opioid receptors results in the formation of heterodimers with ligand binding and functional properties that are distinct from singly transfected receptors suggests that heterodimerization may be involved in modulation of GPCR function (13). Also, Western

blot studies have shown that the 5HT_{1D} and 5HT_{1B} receptors form homodimers when expressed alone although they preferentially form heterodimers when co-expressed (14). In the present study, we have directly examined whether 5HT₁ receptors are capable of forming oligomeric complexes with a variety of GPCRs by specific immunoprecipitation of each receptor followed by identification of the co-precipitated proteins using immunoblot analysis. The main objective was to determine the specificity of the interactions across different GPCR subfamilies to gain further insight into the mechanism of GPCR hetero-oligomerization.

EXPERIMENTAL PROCEDURES

Materials—Restriction endonucleases and other DNA-modifying enzymes were from New England Biolabs (Beverly, MA) or Amersham Biosciences. Dulbecco's modified Eagle's medium and fetal calf serum were from Invitrogen. All other biochemicals were obtained from Sigma unless specified otherwise.

Construction of Epitope-tagged GPCRs and Transient Transfections—The c-Myc epitope (EQKLISEEDL) was inserted at the amino termini of 5HT_{1A} and 5HT_{1D} receptors by PCR mutagenesis using a Stratagene Robocycler with Hot-Top assembly (Amsterdam, Holland) as described by Nelson and Long (15). The FLAG epitope (DYKDDDDK) was inserted by a similar procedure into the amino termini of 5HT_{1A}, 5HT_{1B}, 5HT_{1D}, EDG₁, and EDG₂ receptors. FLAG-tagged EDG₃, GPR₂₆, and GABA_{B2} receptors were a gift from Dr. Kevin Lynch (University of Virginia, Charlottesville, VA). PCR products were cloned into the eucaryotic expression vector pCDNA3.1⁺ (Invitrogen) using standard techniques (16). Inserts were sequenced with an ABI Prism dye terminator cycle sequencing kit and analyzed on an Applied Biosystems 373A stretch DNA sequencer. Transient transfection of HEK-293 cells with the epitope-tagged constructs was performed by calcium phosphate-mediated gene transfer as described previously (17).

Immunoprecipitation—Forty-eight hours after transfection cells were washed three times in phosphate-buffered saline, harvested by scraping, and centrifuged for 5 min at 500 × g. The pelleted cells were homogenized at 4 °C by drawing up and down through a 20-gauge syringe needle in 1 ml of cell lysis buffer (Sigma). The homogenates were centrifuged for 20 min at 14,000 × g at 4 °C, and the supernatants were combined with 12.5 μ l (packed gel) of either anti-c-Myc or anti-FLAG M2 affinity agarose (Sigma) and mixed overnight at 4 °C. The immunoadsorbents were recovered by centrifugation for 5 min at 700 × g and washed three times by resuspension and centrifugation (5 min at 700 × g) in cell lysis buffer and two times in 50 mM Tris (pH 7.5) containing 0.1% (w/v) SDS and 150 mM NaCl. The samples were eluted into 60 μ l of SDS loading buffer (Sigma).

Immunoblot Analysis—Eluted samples were heated for 3 min at 100 °C and subjected to SDS-PAGE on slab gels (180 × 160 × 1.5 mm) polymerized from 10% (w/v) acrylamide and 0.1% N,N'-methylenebisacrylamide in Tris/glycine/SDS buffer (Novex, San Diego, CA) using the discontinuous buffer system of Laemmli (18). The electrophoresed proteins were subjected to semi-dry electrophoretic transfer onto polyvinylidene difluoride membranes as described previously (19). The membranes were incubated with either mouse c-Myc antibodies (1:1000) or mouse FLAG antibodies (1:1000) and then with peroxidase-conjugated sheep anti-mouse serum (1:1000) for detection of immunoreactive bands

* The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

‡ To whom correspondence should be addressed: Dept. of Biochemistry and Molecular Biology, Merck Sharp & Dohme, Terlings Park, Harlow, Essex CM20 2QR, UK; Tel.: 01279-440-495; Fax: 01279-440-390; E-mail: kamran_salim@merck.com.

¹ The abbreviation used is: GPCR, G-protein-coupled receptor.




FIG. 1. Immunoblot analysis of transfected HEK-293 cells showing homo-oligomerization of 5HT_{1A}, 5HT_{1B}, and 5HT_{1D} receptors. **A**, HEK-293 cells (5×10^7) expressing the FLAG-tagged 5HT_{1A} (lane 1), 5HT_{1B} (lane 2), or 5HT_{1D} receptor (lane 3) were immunoprecipitated and subjected to immunoblot analysis using FLAG antibodies as described under "Experimental Procedures." **B**, the same immunoprecipitated samples were subjected to immunoblot analysis under non-reducing (minus dithiothreitol) conditions. The migration of monomers is indicated by open arrows (approximate molecular sizes are: 5HT_{1A}, 46 kDa; 5HT_{1B}, 44 kDa; 5HT_{1D}, 42 kDa), the putative dimers are indicated by solid arrows, and immunoglobulin heavy chains are indicated by an arrowhead. The molecular size markers are shown in kDa. The immunoblots shown are representative of at least three independent experiments.

by enhanced chemiluminescence (Amersham Biosciences). Molecular size calibration was achieved with the MultiMark standards (Novex).

RESULTS

Homo-oligomerization of 5HT_{1A}, 5HT_{1B}, and 5HT_{1D} Receptors—HEK-293 cells transfected with either FLAG-tagged 5HT_{1A}, 5HT_{1B}, or 5HT_{1D} receptor constructs were immunoprecipitated with anti-FLAG-agarose, and immunoblot analysis of the precipitates revealed the presence of 46, 44, and 42-kDa FLAG-immunoreactive bands, respectively (Fig. 1A). Higher molecular weight immunoreactive forms were also present, which might represent oligomeric forms of the 5HT₁ receptors. An immunoreactive band that migrated at a size predicted for a FLAG-tagged 5HT_{1A} receptor dimer (92 kDa) showed a strong increase in immunostaining with the reciprocal decrease of the 46-kDa monomer band when the disulfide bond-reducing agent dithiothreitol was omitted from the sample loading buffer (Fig. 1B). An immunoreactive band that migrated at a size predicted for a FLAG-tagged 5HT_{1B} receptor dimer (88 kDa) showed increased immunostaining with the loss of the 44-kDa monomer band when the samples were electrophoresed in the absence of dithiothreitol (Fig. 1B). Similarly, a possible FLAG-tagged 5HT_{1D} receptor dimer (82 kDa) showed an increase in immunostaining on non-reducing gels with the reciprocal disappearance of the 41-kDa monomer band (Fig. 1B). Similar results have been shown previously for the 5HT_{1B} and 5HT_{1D} receptors, which led to the suggestion that they form homodimers (14). The mechanism of 5HT_{1B} receptor homodimerization appeared to differ from that of the 5HT_{1A} and 5HT_{1D} receptors because a high proportion 5HT_{1B} receptor homodimer was observed even when gels were run under disulfide bond-reducing conditions (Fig. 1A).

Hetero-oligomerization of 5HT_{1A} and 5HT_{1D} Receptors with Other GPCRs—To directly test for hetero-oligomerization of different GPCR combinations we used sequential co-immunoprecipitation and immunoblot analyses of cells co-transfected with differentially epitope-tagged receptors. This showed that the c-Myc-tagged 5HT_{1A} receptor co-precipitated with FLAG-tagged versions of all receptors tested including the 5HT_{1A}, 5HT_{1B}, 5HT_{1D}, EDG₁, EDG₃, GPR₂₆, and GABA_{B2} receptors (Fig. 2, A and B). The same finding was observed for the c-Myc-tagged 5HT_{1D} receptor, which was co-precipitated with FLAG-tagged versions of the 5HT_{1B} and EDG₁ receptors (Fig. 2, A and B). Previous studies have also shown that the 5HT_{1B} and 5HT_{1D} receptors form heterodimers (14). The finding that the FLAG- and c-Myc-tagged versions of the 5HT_{1A} receptor were co-precipitated provides direct evidence that this GPCR




FIG. 2. Sequential immunoprecipitation/immunoblot analysis of transfected HEK-293 cells showing oligomerization of 5HT_{1A} and 5HT_{1D} receptors with other GPCRs. HEK-293 cells (5×10^7) either co-expressing the c-Myc-tagged 5HT_{1A} receptor with FLAG-tagged versions of the 5HT_{1A} (lane 1), 5HT_{1B} (lane 2), 5HT_{1D} (lane 3), EDG₁ (lane 4), EDG₃ (lane 5), GPR₂₆ (lane 6), and GABA_{B2} (lane 7) receptors or co-expressing the c-Myc-tagged 5HT_{1D} receptor with FLAG-tagged versions of the 5HT_{1B} (lane 8) and EDG₁ (lane 9) receptors were immunoprecipitated using either anti-c-Myc (**A**) or anti-FLAG-agarose (**B**), and the precipitates were subjected to immunoblot analysis using FLAG (**A**) or c-Myc (**B**) antibodies as described under "Experimental Procedures." The molecular size markers are shown in kDa (approximate molecular sizes are: 5HT_{1A}, 46 kDa; 5HT_{1B}, 44 kDa; 5HT_{1D}, 42 kDa; EDG₁, 43 kDa; EDG₃, 42 kDa; GPR₂₆, 38 kDa; GABA_{B2}, 106 kDa). The immunoblots shown are representative of four independent experiments. IP, immunoprecipitate.

forms a homo-oligomer. This was also found for the 5HT_{1D} receptor using the c-Myc- and FLAG-tagged versions of this GPCR (data not shown). Co-precipitation of high molecular weight FLAG-immunoreactive 5HT_{1B}, 5HT_{1D}, EDG₁, and EDG₃ species with the c-Myc-5HT_{1A} receptor (Fig. 2A) suggests that these GPCRs might be present as homodimers and trimers in higher order hetero-oligomeric complexes comprised of at least one c-Myc-5HT_{1A} receptor and two or more of the FLAG-tagged constructs.

Sequential Immunoprecipitation/Immunoblot Analysis of Mixed Singly Transfected HEK-293 Cells—To rule out the possibility that the observed hetero-oligomerizations were caused by nonspecific aggregation of receptors, extracts from cells separately expressing the c-Myc-tagged 5HT_{1A} and FLAG-tagged 5HT_{1B} receptors were mixed, immunoprecipitated with anti-c-




FIG. 3. Sequential immunoprecipitation/immunoblot analysis of mixed singly transfected HEK-293 cells. HEK-293 cells (5×10^7) expressing either the c-Myc-tagged 5HT_{1A} receptor (lane 1), the FLAG-tagged 5HT_{1B} receptor (lane 2), or a mixture of both membranes (lane 3) were immunoprecipitated (IP) with either anti-c-Myc agarose (A and D) or anti-FLAG agarose (B and C), and the precipitates were subjected to immunoblot (Blot) analysis with either c-Myc (A and B) or FLAG (C and D) antibodies as described under "Experimental Procedures." The immunoblots shown are representative of at least three independent experiments.

Myc agarose, and subjected to immunoblot analysis with FLAG antibodies. In this case, the FLAG-5HT_{1B} receptor was not detected in the precipitated material (Fig. 3A). Similarly, immunoprecipitation of the mixed extract with anti-FLAG agarose followed by immunoblot analysis with c-Myc antibodies did not detect the c-Myc-5HT_{1A} receptor (Fig. 3B). This confirmed that successful immunoprecipitation of 5HT_{1A}/5HT_{1B} hetero-oligomers requires co-expression within the same cells before the time of lysis. Thus, the oligomerization of the c-Myc-tagged 5HT_{1A} and FLAG-tagged 5HT_{1B} receptors in the present study is not likely to be caused by nonspecific aggregation. The same results were found for all of the other GPCR/GPCR hetero-oligomers tested in this study (data not shown).

Sequential Immunoprecipitation/Immunoblot Analysis of HEK-293 Cells Transfected with Varying Amounts of 5HT_{1A} and EDG₁ Receptors—To determine whether the hetero-oligomers observed in this study are comprised of GPCRs at discrete protein/protein ratios, HEK-293 cells were co-transfected with a fixed amount (25 μ g) of the c-Myc-tagged 5HT_{1A} receptor and varying amounts (25, 5, 1, and 0.2 μ g) of the FLAG-tagged EDG₁ receptor. The relative amounts of the c-Myc-5HT_{1A} receptor that were associated with the decreasing levels of the FLAG-EDG₁ receptor were determined by sequential immunoprecipitation and immunoblot analysis. Immunoprecipitation of the FLAG-EDG₁ receptor with anti-FLAG-agarose resulted in co-precipitation of similar decreasing levels of c-Myc-5HT_{1A} receptor (Fig. 4B) as shown by immunoblot analysis with c-Myc antibodies. Similarly, immunoprecipitation of the c-Myc-5HT_{1A} receptor resulted in co-precipitation of decreasing levels of the FLAG-EDG₁ receptor (Fig. 4D). Sequential immunoprecipitation/immunoblot analysis of the lysates with c-Myc/c-Myc or FLAG/FLAG confirmed that the c-Myc-5HT_{1A} receptor was actually expressed in the cells at relatively constant levels and that expression of the FLAG-EDG₁ receptor decreased accordingly with the amounts of cDNA transfected into the cells (Fig. 4, A and C). Similar results were obtained for all of the other receptors under study (data not shown).

DISCUSSION

We have used a co-immunoprecipitation approach to provide direct evidence that GPCRs from different subfamilies are capable of forming hetero-oligomers when co-expressed in HEK-293 cells. Cells were co-transfected with GPCR constructs containing different epitope tags followed by immunoprecipitation of one receptor and detection of co-precipitated receptors by immunoblot analysis. The 5HT_{1A} receptor formed a homo-oligomer when expressed alone and formed hetero-oligomers when co-expressed with the closely related 5HT_{1B} and 5HT_{1D}




FIG. 4. Sequential immunoprecipitation/immunoblot analysis of HEK-293 cells transfected with varying amounts of 5HT_{1A} and EDG₁ receptors. HEK-293 cells (5×10^7) were co-transfected with 25 μ g of the c-Myc-tagged 5HT_{1A} receptor and either 25 (lane 1), 5 (lane 2), 1 (lane 3), or 0.2 (lane 4) μ g of the FLAG-tagged EDG₁ receptor followed by immunoprecipitation (IP) using either anti-c-Myc agarose (A and D) or anti-FLAG agarose (B and C), and the precipitates were subjected to immunoblot (Blot) analysis using c-Myc (A and B) or FLAG (C and D) antibodies as described under "Experimental Procedures." The molecular sizes of the c-Myc-5HT_{1A} (45 kDa) and FLAG-EDG₁ (42 kDa) receptor constructs are shown. The immunoblots shown are representative of three independent experiments.

receptors (~36% amino acid homology) and also with the more distantly related EDG₁, EDG₃, GPR₂₆, and GABA_{B2} receptors (4–16% amino acid homology). Interestingly, hetero-oligomer formation was shown for all other GPCR combinations tested including the 5HT_{1D} receptor with the 5HT_{1B} and EDG₁ receptors. The heterodimerization of 5HT_{1B} and 5HT_{1D} receptors has been shown previously in insect SF9 cells using a viral expression system followed by Western blot analysis (14).

The results of this study suggest that the receptor complexes were pre-existing in cells at the time of lysis and were not caused by nonspecific aggregation. Oligomerization was observed only in lysates from co-transfected cells and not in mixed lysates from singly transfected cells. Also, the observed receptor/receptor interactions appeared to occur at defined stoichiometries in co-transfected cells because decreased expression of one receptor resulted in a similar incremental decrease in the amount of the second receptor immunoprecipitated. The appearance of high molecular weight immunoreactive bands corresponding in size to 5HT_{1A} and 5HT_{1D} receptor homooligomers appeared to require intact disulfide bonds because these were only detected when gels were run under non-reducing conditions. Little or no 5HT_{1A} and 5HT_{1D} receptor homooligomers could be detected when gels were run under reducing conditions. In contrast, a significant proportion of the 5HT_{1B} receptor was detected as high molecular weight oligomers when gels were run under reducing conditions with increased levels of the oligomer and reciprocal loss of the monomer under non-reducing conditions. This indicates that the mechanism of 5HT_{1B} receptor homo-oligomer formation differs from that of the 5HT_{1A} and 5HT_{1D} receptors. Also, the observed immunoprecipitation of the putative 5HT_{1B} receptor dimer by immunoprecipitation of the 5HT_{1A} receptor indicates that these receptors may be present as higher order hetero-oligomers such as trimers or tetramers.

A major prerequisite for the physiological assembly of hetero-oligomeric GPCRs is co-expression in the same cells. To demonstrate this would require detailed immunological or specific binding experiments that localize the receptors under study to the same cells combined with co-precipitation studies using receptor-specific antibodies or ligands. The GPCRs tested in this study show a wide range of tissue distributions in the brain. Within the 5HT receptor subfamily 5HT_{1A}, 5HT_{1B}, and 5HT_{1D} receptors all show pre- and postsynaptic localizations and can be found together in some of the same brain regions including the cortex and dorsal raphe nucleus (20–22). The 5HT_{1B} receptor shows co-expression with the 5HT_{1A} receptor in

the hippocampus and with the 5HT_{1D} receptor in the olfactory tubercle although the 5HT_{1B} receptor is expressed separately in the anterior caudate putamen, hypothalamus, and thalamus, and the 5HT_{1D} receptor is localized separately in the trigeminal nucleus and in parts of the cerebellum (21). Hetero-oligomerization of the 5HT_{1A} with the GABA_{B2} and GPR26 receptors may be of interest because all of these receptors are widely distributed throughout the cortex and may therefore share some regions of overlap (22–24). The physiological relevance of the hetero-oligomerization of 5HT_{1A} receptor with the EDG₁ and EDG₃ receptors is not clear because no detailed studies have been carried out on the distribution of the latter receptors within the brain. However, the interactions observed between the 5HT_{1A} and EDG₃ receptors are not likely to occur *in vivo* because the relative expression levels of the latter are relatively low in the brain with the highest levels observed in peripheral tissues such as lung and heart (25, 26).

Supporting evidence for oligomerization of endogenously expressed GPCRs comes from Western blot studies of brain tissues revealing the presence of high molecular weight immunoreactive forms of the dopamine D2 receptor (27) and the A1 adenosine receptor (28). More recently an immunoelectron microscopy study showed that CCR5 and CXCR4 receptors are clustered and closely apposed on microvilli and in trans-Golgi vesicles of human macrophages and T cells (29). The strongest evidence comes from studies showing that GABA_BR1 and GABA_BR2 receptors could be co-immunoprecipitated from rat cortex (9).

This is the first study in which the phenomenon of GPCR oligomerization has been investigated using a large number of diverse receptor subtypes. Interestingly, all of the receptors tested were capable of forming hetero-oligomers when co-expressed in tissue culture cells suggesting that this might be a general characteristic of these receptors. These findings highlight the importance for further studies to investigate the occurrence of GPCR hetero-oligomerization in tissues and cells where they are endogenously expressed. Such studies will help to establish the physiological role of such complexes in GPCR function.

REFERENCES

- Maggio, R., Vogel, Z., and Wess, J. (1993) *Proc. Natl. Acad. Sci. U. S. A.* **90**, 3103–3107
- Maggio, R., Vogel, Z., and Wess, J. (1993) *FEBS Lett.* **319**, 195–200
- Hebert, T. E., Moffett, S., Morello, J. P., Loisel, T. P., Bichet, D. G., Barret, C., and Bouvier, M. (1996) *J. Biol. Chem.* **271**, 16384–16392
- Romano, C., Yang, W. L., and O'Malley, K. L. (1996) *J. Biol. Chem.* **271**, 28612–28616
- Fukushima, Y., Asano, T., Saitoh, T., Anai, M., Funaki, M., Ogihara, T., Katagiri, H., Matsuhashi, N., Yazaki, Y., and Sugano, K. (1997) *FEBS Lett.* **409**, 283–286
- Cvejić, S., and Devi, L. A. (1997) *J. Biol. Chem.* **272**, 26959–26964
- Nimchinsky, E. A., Hof, P. R., Janssen, W. G. M., Morrison, J. H., and Schmauss, C. (1997) *J. Biol. Chem.* **272**, 29229–29237
- Jones, K. A., Borowsky, B., Tamm, J. A., Craig, D. A., Durkin, M. M., Dai, M., Yao, W. J., Johnson, M., Gunwaldsen, C., Huang, L. Y., Tang, C., Shen, Q., Salon, J. A., Morse, K., Laz, T., Smith, K. E., Nagarathnam, D., Noble, S. A., Branchek, T. A., and Gerald, C. (1998) *Nature* **396**, 674–679
- Kaupmann, K., Malitschek, B., Schuler, V., Heid, J., Froestl, W., Beck, P., Mosbacher, J., Bischoff, S., Kulik, A., Shigemoto, R., Karschin, A., and Bettler, B. (1998) *Nature* **396**, 683–687
- White, J. H., Wise, A., Main, M. J., Green, A., Fraser, N. J., Disney, G. H., Barnes, A. A., Emson, P., Foord, S., and Marshall, F. H. (1998) *Nature* **396**, 679–682
- Ng, G. Y., Clark, J., Coulombe, N., Ethier, N., Hebert, T. E., Sullivan, R., Kargman, S., Chateaufneuf, A., Tsukamoto, N., McDonald, T., Whiting, P., Mezey, E., Johnson, M. P., Liu, Q., Kolakowski, L. F., Jr., Evans, J. F., Bonner, T. I., O'Neill, G. P. (1999) *J. Biol. Chem.* **274**, 7607–7610
- Kuner, R., Kohr, G., Grunewald, S., Eisenhardt, G., Bach, A., and Kornau, H. C. (1999) *Science* **283**, 74–77
- Jordan, B. A., and Devi, L. A. (1999) *Nature* **399**, 697–700
- Xie, Z., Lee, S. P., O'Dowd, B. F., and George, S. R. (1999) *FEBS Lett.* **456**, 63–67
- Nelson, R. M., and Long, G. L. (1989) *Anal. Biochem.* **180**, 147–151
- Davis, L. G., Kuehl, W. M., and Battey, J. F. (1994) *Basic Methods in Molecular Biology*, Appleton & Lange, Norwalk, CT
- Chen, C. A., and Okayama, H. (1988) *BioTechniques* **6**, 632–638
- Laemmli, U. K. (1970) *Nature* **227**, 680–685
- Guest, P. C., Ravazzola, M., Davidson, H. W., Orci, L., and Hutton, J. C. (1991) *Endocrinology* **129**, 734–740
- Fink, K., Zentner, J., and Gothert, M. (1995) *Naunyn-Schmiedeberg's Arch. Pharmacol.* **352**, 451–454
- Bonaventure, P., Voorn, P., Luyten, W. H., Jurzak, M., Schotte, A., and Leysen, J. E. (1998) *Neuroscience* **82**, 469–484
- Barnes, N. M., and Sharp, T. (1999) *Neuropharmacology* **38**, 1083–1152
- Durkin, M. M., Gunwaldsen, C. A., Borowsky, B., Jones, K. A., and Branchek, T. A. (1999) *Mol. Brain Res.* **71**, 185–200
- Lee, D. K., Lynch, K. R., Nguyen, T., Im, D. S., Cheng, R., Saldivia, V. R., Liu, Y., Liu, I. S., Heng, H. H., Seeman, P., George, S. R., O'Dowd, B. F., and Marchese, A. (2000) *Biochim. Biophys. Acta* **149**, 311–323
- Yamaguchi, F., Tokuda, M., Hatase, O., and Brenner, S. (1996) *Biochem. Biophys. Res. Commun.* **227**, 608–614
- Zhang, G., Contos, J. J., Weiner, J. A., Fukushima, N., and Chun, J. (1999) *Gene (AMST.)* **227**, 89–99
- Ng, G. Y., O'Dowd, B. F., Lee, S. P., Chung, H. T., Brann, M. R., Seeman, P., and George, S. R. (1996) *Biochem. Biophys. Res. Commun.* **227**, 200–204
- Ciruela, F., Casado, V., Mallol, J., Canela, E. I., Lluis, C., and Franco, R. (1995) *J. Neurosci. Res.* **42**, 818–828
- Singer, I. I., Scott, S., Kawka, D. W., Chin, J., Daugherty, B. L., DeMartino, J. A., DiSalvo, J., Gould, S. L., Lineberger, J. E., Malkowitz, L., Miller, M. D., Mitnaul, L., Siciliano, S. J., Staruch, M. J., Williams, H. R., Zweerink, H. J., and Springer, M. S. (2001) *J. Virol.* **75**, 3779–3790

Oligomerization of G-protein-coupled Receptors Shown by Selective Co-immunoprecipitation

Kamran Salim, Tim Fenton, Jamil Bacha, Hector Urien-Rodriguez, Tim Bonnert, Heather A. Skynner, Emma Watts, Julie Kerby, Anne Heald, Margaret Beer, George McAllister and Paul C. Guest

J. Biol. Chem. 2002, 277:15482-15485.

doi: 10.1074/jbc.M201539200 originally published online February 19, 2002

Access the most updated version of this article at doi: [10.1074/jbc.M201539200](https://doi.org/10.1074/jbc.M201539200)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 28 references, 8 of which can be accessed free at <http://www.jbc.org/content/277/18/15482.full.html#ref-list-1>