

Kent Academic Repository

Full text document (pdf)

Citation for published version

Aziz, M. A. and Tollington, Simon and Barlow, Adam and Greenwood, Christina and Goodrich, John M. and Smith, Olutolani and Shamsuddoha, Mohammad and Islam, M. Anwarul and Groombridge, Jim J. (2017) Using non-invasively collected genetic data to estimate density and population size of tigers in the Bangladesh Sundarbans. *Global Ecology and Conservation* . ISSN 2351-9894.

DOI

<https://doi.org/10.1016/j.gecco.2017.09.002>

Link to record in KAR

<http://kar.kent.ac.uk/63702/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

1 Title

2 Using non-invasively collected genetic data to estimate density and
3 population size of tigers in the Bangladesh Sundarbans.

4

5 Author names and affiliations

6

7 M. Abdul Aziz ^{a,1,*}, Simon Tollington ^{a,g}, Adam Barlow ^b, Christina Greenwood ^b, John M.
8 Goodrich ^c, Olutolani Smith ^{c,d}, Mohammad Shamsuddoha ^e, M. Anwarul Islam ^f, Jim J.
9 Groombridge ^a

10

11

12 ^a Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation,
13 University of Kent, Canterbury CT2 7NZ, UK

14

15 ^b WildTeam, Surfside, St Merryn, Padstow PL28 8NU, Cornwall, UK

16

17 ^c Panthera Foundation, 8 West 40th Street, NY 10018, USA

18

19 ^d Department of Genetics, Evolution & Environment, University College London, London,
20 UK

21

22 ^e Department of Zoology, Jahangirnagar University, Dhaka 1342, Bangladesh

23

24 ^f WildTeam, Bangladesh and Department of Zoology, University of Dhaka, Bangladesh

25

26 ^g The North of England Zoological Society, Chester Zoo, Caughall Road, Chester, CH2 1LH,
27 UK

28

29 * Corresponding author: Department of Zoology, Jahangirnagar University, Savar, Dhaka
30 1342, Bangladesh; Email: maaaziz78@gmail.com; Phone: +88 0176256193

31

32

33

34

35

36

37

¹ Present address: Department of Zoology, Jahangirnagar University, Dhaka 1342, Bangladesh

38

39 **Abstract**

40 Population density is a key parameter to monitor endangered carnivores in the wild. The
41 photographic capture-recapture method has been widely used for decades to monitor tigers,
42 *Panthera tigris*, however the application of this method in the Sundarbans tiger landscape is
43 challenging due to logistical difficulties. Therefore, we carried out molecular analyses of
44 DNA contained in non-invasively collected genetic samples to assess the tiger population in
45 the Bangladesh Sundarbans within a spatially explicit capture-recapture (SECR) framework.
46 By surveying four representative sample areas totalling 1,994 km² of the Bangladesh
47 Sundarbans, we collected 440 suspected tiger scat and hair samples. Genetic screening of
48 these samples provided 233 authenticated tiger samples, which we attempted to amplify at 10
49 highly polymorphic microsatellite loci. Of these, 105 samples were successfully amplified,
50 representing 45 unique genotype profiles. The capture-recapture analyses of these unique
51 genotypes within the SECR model provided a density estimate of $2.85 \pm \text{SE } 0.44$ tigers/100
52 km² (95% CI: 1.99-3.71 tigers/100 km²) for the area sampled, and an estimate of 121 tigers
53 (95% CI: 84-158 tigers) for the total area of the Bangladesh Sundarbans. We demonstrate that
54 this non-invasive genetic surveillance can be an additional approach for monitoring tiger
55 populations in a landscape where camera-trapping is challenging.

56

57 **Keywords:** Bangladesh; Bengal tiger; genetic sampling; population density; Sundarbans.

58

59 **1. Introduction**

60 Wild tigers now survive within 76 Tiger Conservation Landscapes, representing only seven
61 percent of their ancestral range (Dinerstein et al., 2007). Monitoring changes in the tiger
62 population at each of these landscapes is fundamental in assessing the level of threats and the
63 effectiveness of management actions (Walston et al., 2010). The Indian and Bangladesh
64 Sundarbans, representing 10,236 km² of mangrove forest, has been identified as one of 11
65 Tiger Conservation Landscapes of global priority for long-term conservation in the region
66 (Sanderson et al., 2006). A reliable monitoring approach is, therefore, critical to guide the
67 management of this landscape (Ahmad et al., 2009).

68

69 The first tiger population survey in the Sundarbans Reserved Forest of Bangladesh utilised
70 the “pug-mark” method, but this approach was subsequently abandoned due to its

71 methodological shortcomings (Karanth, 2005). Since 2007, a secondary sign survey has been
72 regularly used to monitor changes in relative tiger abundance across the whole forest (Barlow
73 et al., 2008). This type of survey provides reasonable power to detect changes in the relative
74 abundance of tigers over time, but the numerical relationship between the relative abundance
75 index and the actual tiger population size population is not known (Barlow et al., 2008;
76 Hayward et al., 2002). A rough population estimate has also been generated using the
77 estimated home range size of adult female tigers living in the Sundarbans Reserved Forest,
78 but the limitations of this study included a small sample size of only two female tigers
79 (Barlow, 2009). Camera trap surveys carried out in both the Indian and Bangladesh
80 Sundarbans have met with various difficulties, mainly associated with the enormous, daily
81 tidal transformation of this landscape. For example, researchers in India concentrated their
82 efforts around watering holes, because of the difficulty in identifying tiger travel routes in
83 this densely vegetated and muddy habitat, this resulted in very low capture-recapture rates
84 and corresponding density estimates compared to other tiger sites (Karanth and Nichols,
85 2000). Further camera-trap surveys in both the Indian and Bangladesh Sundarbans attempted
86 to improve capture-recapture rates by using various forms of lures to encourage tigers to
87 come to the camera trap locations (Dey et al., 2015; Jhala et al., 2011). It is not clear,
88 however, if the use of lures in this way has any meaningful effect on the resulting density and
89 population estimates generated from this approach (Dey et al., 2015; Jhala et al., 2011).
90 Moreover, a recent study suffered from the loss of more than half of their camera traps due to
91 suspected theft in the field (Hossain et al., 2016).

92

93 Advances in DNA technology have enabled researchers to incorporate non-invasive genetic
94 techniques to survey tiger populations at some sites (Bhagavatula and Singh, 2006; Mondol et
95 al., 2009a), but this approach has never been used before in the Sundarbans. The objective of
96 this study was, therefore, to investigate if this non-invasive genetic approach could be used to
97 provide a reliable density and population estimate for the tiger population in the Sundarbans
98 landscape.

99

100 **2. Methods**

101

102 **2.1. Study site**

103 The Sundarbans, the largest contiguous mangrove forest of the world, is located on the
104 Ganges-Brahmaputra delta (Giri et al., 2007). The part of the Sundarbans in Bangladesh
105 (21°30'–22°30' N, 89°00'–89°55' E) covers 6,017 km², of which 4,267 km² is forest and the
106 remaining area is comprised of water bodies (Iftekhar and Islam, 2004a). The Sundarbans is
107 bordered on the south by the Bay of Bengal and on the north and east sides by a landmass
108 dominated by human settlements (Hussain and Acharya, 1994). Two rivers, the Raimangal
109 and the Hariabhangra, mark the international boundary between Bangladesh and India, and
110 separate the Indian part of the Sundarbans (Fig. 1).

111

112 The Bangladesh Sundarbans is managed as a reserve forest, except three demarcated areas
113 within the forest that were declared wildlife sanctuaries (IUCN Category III and VI) in 1996
114 for the higher protection of wildlife and their habitat (BFD, 2012). The sanctuaries comprise
115 the Sundarbans West (715 km²), Sundarbans South (370 km²), and Sundarbans East (312
116 km²), and were collectively declared a UNESCO World Heritage Site in 1997 (BFD, 2012;
117 Iftekhar and Islam, 2004a) (Fig. 1).

118

119 The Sundarbans is one of the most biologically diverse mangrove forests in the world,
120 supporting 330 species of plants, more than 400 species of fishes, 35 species of reptiles, over
121 300 species of birds, and 42 species of mammals (Islam and Wahab, 2005; IUCN–
122 Bangladesh, 2001). The tiger is the only large terrestrial carnivore in the Sundarbans; their
123 major prey species include spotted deer (*Axis axis*), wild boar (*Sus scrofa*) and barking deer
124 (*Muntiacus muntjak*) (Khan, 2008). Several small carnivores found in the forest include
125 fishing cat (*Prionailurus viverrinus*), jungle cat (*Felis chaus*) and leopard cat (*Prionailurus*
126 *bengalensis*).

127

128 The Sundarbans Reserved Forest is mostly comprised of two tree species; Sundri (*Heritiera*
129 *fomes*; 39%) and Gewa (*Excoecaria agalloch*; 39%), with other species constituting only
130 16% of the forest cover (Iftekhar and Saenger, 2008). The Sundarbans is characterized by a
131 maritime, humid climate with very seasonal weather patterns (Iftekhar and Islam, 2004b).
132 Most of this area is less than one meter above the sea level (Canonizado and Hossain, 1998),
133 and consists of vegetated islands that are inundated regularly by two high and low tides each
134 day with a mean amplitude of 3-4 m (Chaffey et al., 1985; Gopal and Chauhan, 2006).

135

136 **2.2. Sampling approach**

137 To collect non-invasive genetic samples (scat or hair), four sampling areas (totalling 1,994
138 km²) were selected within the Bangladesh Sundarbans: Satkhira block (SB, 554 km²), West
139 Wildlife Sanctuary (WS, 715 km²), East Wildlife Sanctuary with additional areas (ES, 383
140 km²) and Chandpai block (CB, 342 km²) (Fig. 1). Location, protection status and level of
141 human use were considered in selecting these sample areas. The ES and WS areas have
142 higher protection status and are situated away from human settlements, whereas the CB and
143 SB areas have lower protection status and are located close to local villages. The Forest
144 Department issues permission to local people for collecting forest and aquatic resources (e.g.,
145 nypa palms, honey, fish and crabs) from SB and CB sample areas, but not from the ES and
146 WS (Aziz et al., 2017).

147

148 Following standard capture-recapture approaches (Karanth, 1995; Karanth and Nichols,
149 1998) to select sampling points, each sampling area was divided into 2×2 km grid cells
150 creating a total of 373 grid cells for potential sampling. A survey team of four trained field
151 staff searched each grid cell with three separate transects (using one transect each time).
152 Starting points for each transect were selected by where the grid cell could be easily accessed
153 by boat. From the start point the field team walked each transect roughly in the direction of
154 the opposite side of the grid square. Each transect was walked for a length of 1 km, or until
155 the observers could not continue further because of particularly dense habitat or a large water
156 body obstructing their way. The field team walked in parallel along the line of each transect,
157 with the distance between the first and last observer being maintained at approximately 15 m
158 (5 m between each observer). Five survey field teams, each with four observers, were used to
159 simultaneously survey a sample area over a short (13-22 days) period of time for sample
160 collection.

161

162 Field teams managed to survey 10 grid cells with four transects, 297 grid cells with three
163 transects, 7 grid cells with two transects, and 32 grid cells with one transect. A total of 27
164 (11%) grid cells were not surveyed due to inaccessibility and security issues.

165

166 Winter months were chosen for sampling to avoid extreme weather conditions, and to
167 maximise the chance of collecting dry samples. We sampled SB areas from 20 November to
168 11 December 2014, WS areas from 17 to 30 December in 2014, and areas ES and CB from 4

169 to 26 February 2015. Survey teams recorded location data for each sample using handheld
170 Global Positioning System (GPS) Garmin GPSMAP 64.

171

172 Suspected tiger scat samples, identified by size and associated signs (Johnsingh, 1983;
173 Karanth et al., 1995), were collected in 100ml polypropylene tubes (ThermoFisher Scientific,
174 UK) using twigs to avoid contamination. All scat samples were air-dried before being
175 preserved with silica gel desiccant, and stored at -20 °C until extraction within a month of
176 collection. Suspected tiger hairs, identified by being associated with territorial scratch marks
177 on trees, were also collected (Sharma et al., 2012). High quality tiger blood and tissue
178 samples were also collected from captive tigers or confiscated tiger products that originated
179 from the Bangladesh Sundarbans to provide reference genotypes for comparison with our
180 field collected samples: one blood sample (from a rescued tiger), five tissue samples (from
181 confiscated skins) and four hair samples (from confiscated and rescued tigers) were collected.

182

183 **2.3. DNA extraction**

184 All biological samples were transported to the Durrell Institute of Conservation and Ecology,
185 University of Kent, for analyses under permits (Permit No. BD 9118404) from the
186 Convention on International Trade in Endangered Species (CITES), and Department for
187 Environment, Food and Rural Affairs, United Kingdom (Authorization no. AHVLA:
188 TARP/2015/111).

189

190 Genomic DNA from scat samples was extracted using QIAamp DNA Stool mini kits
191 (QIAGEN Inc.) following the manufacturer's instructions. Approximately 200 mg of scat
192 material was scraped from the outer surface of each scat sample with a sterilized razor blade
193 and then incubated overnight with 1.5 ml ASL buffer on a mechanical rotator at 56 °C. The
194 DNA supernatant from the sample was lysed with 300 µl AL buffer plus 25 µl proteinase K
195 and incubated at 70 °C for 15 min. Four microlitres of carrier RNA (ThermoFisher Scientific,
196 UK) was added to AL buffer to increase DNA yield from scat samples. To extract DNA from
197 blood, tissue, and hair samples, we used DNeasy™ Blood and Tissue Kits (QIAGEN Inc.);
198 approximately 10 hairs of each sample was added to 300 µl AL buffer incorporating 20 µl of
199 proteinase K and 20 µl of DTT (Dithiothreitol, Biotech), and then incubated at 56 °C
200 overnight or until the sample was completely digested. The elution of DNA was carried out in
201 75 µl buffer solution. Strict protocols was observed to reduce the chances of contamination
202 including using aerosol barrier pipette tips, separate pre and post PCR rooms and UV PCR

203 hoods for sample preparation. A negative control (with no biological material) was included
204 with each batch of extractions to monitor for possible contamination during the DNA
205 extraction procedure.

206

207 **2.4. Species authentication**

208 Scat morphology and associated secondary signs have been used to identify scat samples of
209 the study species (Bagchi et al., 2003; Karanth et al., 1995). However, non-target scats can
210 potentially be misidentified and collected when such field protocols are used in isolation
211 (Farrell et al., 2000), and therefore more reliable DNA-based identification of non-invasive
212 scat samples is necessary to avoid inadvertent sampling of scat from non-target species
213 (Bhagavatula and Singh, 2006). A PCR-based assay was used to reliably identify tiger
214 samples (Bhagavatula and Singh, 2006; Davison et al., 2002; Mondol et al., 2009a), so that
215 only genetically authenticated samples were included in further downstream analyses
216 (Mondol et al., 2009a). All field-collected samples were screened using tiger-specific NADH₅
217 gene fragment of 225 base pairs (fwd TTACTAGGACTCCTCCTAGCC; rev
218 GAATAGGGTTGTGATGGCCCC) that has been successfully used in other non-invasive tiger
219 studies (Mukherjee et al., 2007). In this screening process, PCR reaction volumes (total 27
220 µl) contained 3 µl of template DNA, 12.5 µl MyTaq Redmix (containing dNTPs and MgCl₂;
221 Bioline, UK), 5 µM of each forward and reverse primer, 4 µM BSA (Bovine Serum Albumin,
222 New England Biolabs Inc.) and 8.5 µl dH₂O. PCR cycling conditions for this screening
223 process consisted of an initial hot start of 95 °C for 1 min followed by 45 cycles of 95 °C for
224 15 s, 55 °C for 15 s and 72 °C for 15 s, and a final incubation period of 10 min at 72 °C using
225 a G-Storm Thermal Cycler (Labtech France). PCR products were then purified and
226 sequenced using a 3730XL analyser (Macrogen, Amsterdam, Netherlands). Mitochondrial
227 DNA (mtDNA) sequences were edited using Jalview v2 (Waterhouse et al., 2009), and then
228 cross-checked with sequences from the Genbank database (National Center for
229 Biotechnology Information, NCBI) to ensure that positive PCR samples were in fact tiger.
230 This level of rigorous screening process using a tiger-specific gene fragment and the resultant
231 sequence blasting ensured that authenticated samples were not contaminated with prey DNA,
232 and/or not sourced from other wild cats that might have been eaten by tigers. DNA samples
233 that showed poor quality, or no bands in the species-specific PCRs after three independent
234 extraction attempts were removed before microsatellite amplification (Kohn et al., 1999).

235

236 **2.5. Microsatellite amplification and sex determination**

237 A range of microsatellite primers have been developed in the domestic cat (Menotti-
238 Raymond et al., 1999), and successfully applied in investigating population abundance
239 (Mondol et al., 2009a), genetic structure (Mondol et al., 2009b; Reddy et al., 2012), spatial
240 genetics (Sharma et al., 2012), and connectivity of tiger populations across India (Joshi et al.,
241 2013). Considering the high number of alleles observed in these studies (Bhagavatula and
242 Singh, 2006; Menotti-Raymond et al., 1999; Mondol et al., 2009a), a preliminary set of 14
243 loci were selected for this study (Table A.1). These loci were then optimised using a subset (n
244 = 10) of field-collected scat samples and reference samples ($n = 10$). Based on levels of PCR
245 amplification success, allelic richness, and extent of genotyping errors, a set of 10 loci were
246 chosen to genotype all field-collected samples that had been genetically authenticated as
247 being from tiger (Table A.2). A felid specific zinc-finger (Zfx and Zfy) locus was also
248 optimised using samples from known male ($n = 1$) and female tigers ($n = 2$) for sex
249 determination (Pilgrim et al., 2005).

250

251 Four multiplexes were designed to include the full set of loci. All forward primers were
252 fluorescently labelled for gene-scanning (Table A.2). Each microsatellite PCR reaction
253 volume (10 μ l) contained 5 μ l Qiagen multiplex PCR buffer mix (Qiagen Inc.), 0.2 μ M
254 labelled forward primer (Eurofins Genomics), 0.2 μ M unlabelled reverse primer, 2 μ M BSA,
255 and 3 μ l of DNA template. For all multiplex reactions, the PCR temperature regime included
256 an initial denaturation step of 95 °C for 15 min, 45 cycles of denaturation (94 °C for 30 s),
257 annealing (T_a ranges from 52 °C to 57 °C for 90 s for four multiplexes; Table A.2), extension
258 (72 °C for 90 s), and a final extension of 10 min at 72 °C, using a G-Storm Thermal Cycler.
259 All PCR products were genotyped using an Applied Biosystems 3730 DNA Analyser and
260 ROX 500 ROX™ as the size-standard.

261

262 **2.6. Genotype data validation**

263 To reduce the possibility of genotyping errors, we discarded any DNA samples that
264 amplified at fewer than three loci at the first PCR attempt; these were re-extracted from
265 source and included in subsequent PCRs thus ensuring that poor quality samples were
266 immediately eliminated (Creel et al., 2003). Furthermore, we employed the comparative
267 genotyping approach (Frantz et al., 2003; Hansen et al., 2008) by ensuring that equivalent
268 heterozygote genotype profiles were scored at least twice and corresponding homozygote
269 genotypes at least three times (up to a maximum of five). This approach ensured a level of

270 rigour in resolving the true genotype of each scat sample and was less laborious and more
271 cost-effective than the multiple tubes approach (Taberlet et al., 1997). A consensus genotype
272 was achieved if genotypes matched 100% at all loci in at least two repeats. If genotype
273 consensus was not reached in five independent scoring attempts the samples were removed
274 from the analysis (Jackson et al., 2016). A negative control was included with each batch of
275 PCR reaction to monitor for possible contamination. Genotyping errors due to stuttering were
276 checked using the program MICROCHECKER v2.2.3 (van Oosterhout et al., 2004). Allele
277 frequencies, observed (H_o) and expected (H_e) heterozygosity, allelic dropout, false alleles
278 and tests for adherence to the Hardy-Weinberg equilibrium were quantified using GIMLET
279 v1.3.3 (Valiere, 2002). Alleles were identified and scored using GENEMAPPER v3.7
280 (Applied Biosystems, MA, USA).

281

282 **2.7. Individual identification**

283 The set of 10 polymorphic loci were used to create consensus genotype profiles for all
284 samples. To distinguish between closely related individuals and to avoid an overestimation of
285 population size (Kohn et al., 1999; Waits et al., 2001), we determined the required number of
286 loci using the probability of identify for siblings, PID(sibs), based on polymorphic
287 information content (PIC) of the loci (Bhagavatula and Singh, 2006; Mondol et al., 2009a;
288 Waits et al., 2001). In addition, three reference samples were sourced from confirmed
289 siblings, which we used to estimate the PID(sibs) in order to determine the required number
290 of loci that could sufficiently distinguish between them. By combining this result with PIC
291 values for the microsatellite loci, we determined a set of five polymorphic loci that were
292 sufficient to distinguish siblings within the population. The program GIMLET v1.3.3 was
293 used to estimate PID(sibs) for the microsatellite loci (Valière, 2002). We then compared
294 consensus genotype profiles in the program CERVUS v3.0 (Marshall et al., 1998) to identify
295 matched genotypes with a minimum of loci criteria. The identity module of CERVUS
296 produced a matrix of pair-wise comparisons that allowed us to separate matched and
297 unmatched individuals based on the criteria of a minimum five loci. While examining the
298 pair-wise matrix, we carefully checked genotypes that differed by fewer than three loci,
299 where we allowed up to two mismatches considering genotyping errors in the dataset (Creel
300 et al., 2003). Matching genotypes based on five or more loci were considered to be sourced
301 from the same individual and classified as a recapture (Budowle, 2004; Mondol et al., 2009a).
302 Incomplete or partial genotype profiles, genotyped at 5 – 10 loci, were also used following
303 approaches used in studies involving tigers (Bhagavatula and Singh, 2006; Mondol et al.,

2009a), and badgers (Frantz et al., 2003). When partial profiles were used, we carefully considered samples that had amplified the most informative loci, namely Fca279, Fca232, Fca090, Fca672, and D15. Although it is possible that an incomplete genotype might actually have originated from a new individual (Mondol et al., 2009a) using incomplete genotype profiles in this way provides a conservative population estimate (Bhagavatula and Singh, 2006; Frantz et al., 2003) by minimising the possibility of creating non-existent individuals through genotyping error (Mondol et al., 2009a).

311

312 **2.8. Density estimation**

313 To estimate tiger population density, we used a likelihood-based spatially explicit capture-
314 recapture (SECR) approach that has become widely used for estimating densities of large
315 carnivores, including tigers (Kalle et al., 2011), leopards, *Panthera pardus* (Kalle et al.,
316 2011), jaguars, *Panthera onca* (Sollmann et al., 2013), and European wildcat, *Felis silvestris*
317 *silvestris* (Kéry et al., 2010). SECR uses detection locations to fit a spatial likelihood-based
318 model, avoiding the need to estimate ad hoc effective sample area. Moreover, the ‘area
319 search polygon’ (sample area) approach in SECR allows an analysis of all detections
320 (capture-recaptures) of all individuals by pooling them together as a ‘single session’ (Efford,
321 2011), avoiding the difficulty of assigning non-invasive samples to predefined sample
322 occasions.

323

324 The SECR model assumes that no animal activity centres can occur in non-habitat beyond the
325 animal’s range (Efford, 2011; Efford et al., 2009). Therefore, density estimates can
326 potentially be biased if non-habitat is included in the ‘sample area polygon’ (Efford, 2011;
327 Gerber et al., 2012). Tigers in the Sundarbans are known to navigate water bodies up to but
328 rarely exceeding 1.5 km wide (Barlow, 2009). Therefore, ‘non-habitat’ of tigers (e.g., water
329 bodies more than 1.5 km wide and human settlements on the northern boundary of the area
330 sampled) were removed from the buffer area; defined as the adjoining area of the sample area
331 polygon ($3 \times \sigma$) where activity centres of sampled tigers can occur (Efford, 2011; Gerber et
332 al., 2012; Mace et al., 1994). For the SB and CB sample areas, tiger movement is restricted
333 on the north side by densely populated human settlements separated by rivers. The WS
334 sample area is bounded on the south side by the Bay of Bengal and on the west side and most
335 of the east sides by rivers >3 km wide. Similarly, tiger movement is restricted on the south
336 side of the ES sample area by the Bay of Bengal and on the east side by rivers >3 km wide

337 (Fig. 1). Consequently, these areas were also excluded from the overall sample area in the
338 SECR analysis.

339

340 Two matrices of spatiotemporal detection history and spatiotemporal search area polygons
341 were used in the SECR analysis for estimating density parameters. The spatiotemporal
342 detection history included capture-recapture locations for each individual tiger and the
343 spatiotemporal search area polygon contained geographic coordinates defining the area
344 sampled. Using these two input datasets, a detection model was fitted by maximum
345 likelihood, with parameters, g_0 (detection probability at the activity centre of the animal's
346 home range), and σ (the spatial movement parameter away from the centre of the animal's
347 home range). Using the detection function as half-normal, g_0 and σ were modelled as
348 constant to estimate overall and sample area-wise tiger density (Borchers and Efford, 2008;
349 Efford, 2011). The SECR analysis was carried out in the R package SECR v2.10.3, and
350 ArcGIS v10.3 was used for creating polygons of areas sampled.

351

352 **3. Results**

353

354 **3.1. Species and individual identification**

355 A total of 440 samples of putative tiger faeces and hair were collected. Molecular
356 identification using tiger-specific NADH₅ primers confirmed the existence of 233 (53%) tiger
357 samples after replicate extraction and amplification procedures. The remaining samples were
358 discarded from further analysis because they failed to produce quality, identifiable tiger
359 DNA. A final set of 105 separate scat and hair samples were genotyped at 5-10 loci (see
360 Table 1 for full sample information).

361

362 A higher level of amplification success was obtained for the reference samples (13 loci
363 showed 100% amplification) than the field collected samples (78 - 100%) (Table A.1). Using
364 the set of 10 microsatellite loci, we were able to derive consensus genotypes, based on a
365 minimum of five loci, for 105 scat and hair samples (45% of the tiger-positive samples). A
366 higher genotyping success rate was obtained for samples from the CB sample area (58%)
367 compared to the SB area (39%).

368

369 The CERVUS analysis yielded high proportions of pairwise matrix with zero difference
370 (ranging from 49% to 74% for sample areas) as well as pairwise matrix that differed by more

371 than 7 loci (ranging from 15% to 26% for sample areas) of the final sample genotypes. Using
372 a minimum of five loci criteria, a total of 45 individual tigers comprising six from SB, 15
373 from WS, 14 from CB and 10 from the ES sample area was identified from 105 (capture and
374 recaptures) genotype profiles (Table 2). Sexing of individuals was attempted for these 45
375 individuals resulting in a total of 11 males and 24 females. The sex of the remaining 10
376 individuals could not be determined due to inconclusive genotypes.

377 All loci were polymorphic with a mean number of alleles of $5.50 \pm \text{SD } 1.65$ per locus. The
378 marker set revealed a level of polymorphism sufficient to distinguish between individuals,
379 with a mean PIC of 0.58. Several loci showed allelic dropout and false alleles in the dataset.
380 Deviations from Hardy-Weinberg equilibrium were also detected for loci FCA304, FCA279
381 in ES; D15 in SB; and FCA230, FCA279 for samples from the CB area (Table 3).

382

383 **3.2. Estimating tiger density**

384 The estimated probabilities of detections of 45 tigers ranged from 0.02 to 0.04 across the four
385 sample areas, with the highest in the ES area and lowest in the SB area (Table 2). The null
386 model, $D(\cdot)g0(\cdot)\sigma(\cdot)$, yielded an overall tiger density of $2.85 \pm 0.44 \text{ SE tigers}/100 \text{ km}^2$ (95%
387 CI: 1.99-3.71). The highest density of tigers was estimated for the CB area ($3.18 \pm \text{SE } 0.90$)
388 followed by the ES ($3.17 \pm \text{SE } 1.04$), WS ($2.99 \pm \text{SE } 0.80$) and SB ($1.86 \pm \text{SE } 0.81$) (Table
389 2). By extrapolating the overall tiger density of $2.85 \pm \text{SE } 0.44 \text{ tigers}/100 \text{ km}^2$ to the total of
390 $4,247 \text{ km}^2$ occupied by tigers (Dey et al., 2015), we estimate that the Bangladesh Sundarbans
391 may currently support approximately 121 tigers (95% CI: 84-158).

392

393 **4. Discussion**

394

395 **4.1. Identifying species and individual identity of tigers**

396 Although there are no large carnivores in the Sundarbans except tigers, DNA-based screening
397 to genetically confirm species ensures that samples from non-target species are removed prior
398 to downstream analysis (Mondol et al., 2009a; Mukherjee et al., 2007). The low PCR
399 amplification rate (53%) in this study compared to higher success rates reported from drier
400 areas in India (e.g. 93% in Bandipur National Park, India; Mondol et al., 2009a), may be a
401 consequence of inferior sample quality due to the humid and wet mangrove habitat in the
402 Sundarbans.

403

404 Each microsatellite locus used in this study amplified a region less than 160 base pairs, so
405 they were appropriately-sized to amplify low quality, potentially highly fragmented faecal
406 DNA (Bhagavatula and Singh, 2006; Frantzen et al., 1998). The overall genotyping success
407 rate of all samples (46%) was relatively low because of the rigorous screening process
408 undertaken to reduce genotyping errors. Although no genotyping errors were detected in the
409 reference samples, field samples produced 5-26% genotyping errors for five loci (Table A.1;
410 Table 3). These error rates, however, are reasonably low when compared to other non-
411 invasive genetic studies of tigers (2-65%) (Bhagavatula and Singh, 2006), and other carnivore
412 species such as wolves (3-33%) (Lucchini et al., 2002).

413

414 The five most informative loci with a PID(sibs) value of 0.0186 (0.0003 for full microsatellite
415 panel) and mean PIC value of 0.58 demonstrated that together these loci could successfully
416 distinguish even siblings with 99% certainty. We also note that the PID(sibs) for the five least
417 informative loci accounted to be 0.0193, which is close to the value of the five most
418 informative loci, might be due to the fact that the number of alleles for our loci set ranged
419 from 4 to 7, mostly with 5 alleles (see the Table 3 for details). These PID(sibs) estimates
420 closely aligned with the suggested value of approximately 0.01 for studies intended to
421 estimating population density following mark-recapture approach (Waits et al., 2001). With
422 this level of statistical rigour, we determined that a minimum of five loci of the set used in
423 this study were sufficient to distinguish unique genotype profile from the pool of pair-wise
424 genotype matrix. Using the conservative cumulative PID (sibs) attained for the five loci, and
425 with suggestions made in similar tiger studies (Mondol et al., 2009a), we therefore avoided
426 an overestimation of population size by reducing the incidence of false individuals due to
427 genotyping error (Creel et al., 2003; Bhagavatula and Singh, 2006). There is a general
428 consensus that genotyping error might not be completely eliminated from the dataset
429 demonstrated in non-invasive genetic studies (Bhagavatula and Singh, 2006; Creel et al.,
430 2003; Mondol et al., 2009a), it is therefore reasonable to allow genotypes with one or more
431 mismatches to be scored as identical to avoid an overestimate (Creel et al., 2003). Moreover,
432 it is plausible that our scored individuals represent an underestimate of the true population
433 abundance, however on the management perspective this would be an impetus for protected
434 area managers to intensify monitoring and law enforcement for this important tiger
435 landscape.

436

437 **4.2. Estimating tiger density**

438 Our estimates of tiger density and population size most likely accounts for adult and sub-
439 adult tigers, which gives a sex ratio of 2.18 females : 1 male, similar to sex ratios recorded in
440 other sites for adult tigers (Barlow et al., 2009). It is also important to note that the strict
441 methodological procedures followed in our study might have excluded an unknown number
442 of tigers. It is possible that our sampling approach was not representative of the entire
443 Bangladesh Sundarbans population or that it did not account for tigers from all demographic
444 groups. For example, tiger scats from juveniles may not have been detected because this
445 group tends to have more limited, clumped movement patterns, (Smith, 1978), or they may
446 not have been collected due to the similarity in size to the scats of other species (e.g. fishing
447 cat or leopard cat). Moreover, survey teams did not collect suspected tiger scats that were
448 degraded due to being submerged by tidal waters in areas of low elevation. Therefore the
449 overall lower probability of detections in our study may be due to the sampling approach
450 and/or subsequent screening of samples.

451

452 However, despite these methodological limitations, our density estimates (2.85 tigers/100
453 km²) are comparable to the 2.17 tigers/100 km² estimated in a recent camera-trap survey in
454 the Bangladesh Sundarbans (Dey et al., 2015). Our extrapolated population size estimates
455 (95% CI: 84-158 tigers) are also in line with the camera trap results (SE interval: 84-130
456 tigers) produced by Dey et al. (2015). It is relevant to mention that, due to the overlapping
457 sampling period between these studies (Dey et al. sampled between 2013-2015; this study
458 sampled between 2014-2015), we extrapolated our density estimates for the entire
459 Sundarbans in order for meaningful comparisons to be made. Both studies used SECR
460 modelling in density estimates. Furthermore, our density estimates for SB ($1.86 \pm SE 0.81$
461 tigers/100 km²) and ES sample areas ($3.17 \pm SE 1.04$ tigers/100 km²) were also similar to
462 estimates from camera-trap data in Block III ($2.77 \pm SE 0.78$ tigers/100 km²), and Block I
463 ($3.70 \pm SE 0.91$ tigers/100 km²) (Table 4). These two areas overlap with our study area and
464 that of the camera-trap study by Dey et al. (2015).

465

466 In contrast, a study by Barlow (2009) estimated much higher tiger density (9.33 tigers/100
467 km²) but similar population size (133-200 adult and sub-adult individuals) in the Bangladesh
468 Sundarbans using telemetry. The difference in the estimate of tiger density is most likely due
469 to differences in sampling method (telemetry versus DNA sampling) or changes in the tiger
470 population in the time that separated the two surveys. Although current tiger densities in the

471 Bangladesh Sundarbans may also be lower than densities estimated in Nepal and Bhutan
472 (e.g., Karki et al., 2015; Thinley and Curtis, 2015), combining our study's estimates with the
473 estimated $4.3 \pm \text{SE } 0.3$ tigers/100 km² for the Indian Sundarbans (Jhala et al., 2015, 2011),
474 supports previous assertions (Barlow, 2009) that the entire Sundarbans has the capacity to
475 support one of the largest tiger populations, up to 197 tigers (95% CI: 146-254), in the world.

476

477 **4.3. Conservation implications**

478 We have demonstrated the utility of noninvasive genetic sampling to assess the tiger
479 population of the Bangladesh Sundarbans, complementing camera trap and secondary sign
480 surveys already employed in this landscape (Barlow et al., 2008; Dey et al., 2015). For
481 assessing population parameters in conventional camera trap studies, it is critically important
482 to place camera trap on routes regularly travelled by tigers (Karanth and Nichols, 1998) in
483 order to obtain improved detections for precise estimates. The topography of the Sundarbans
484 mangrove habitat is only few meters above the sea level, therefore, most of the forest land is
485 regularly washed by tidal waters twice daily, leaving few recognizable tiger signs that could
486 be used for camera placements. As a result, previous camera trap studies were able to obtain
487 limited detections both in Bangladesh (Khan, 2012) and the Indian Sundarbans (Karanth and
488 Nichols, 2000), except the one that used lures and baits (Dey et al., 2015). Moreover, a recent
489 study could not recover more than half of their camera traps from the Bangladesh Sundarbans
490 due to suspected theft (Hossain et al., 2016). Given these challenges with camera trapping,
491 we have demonstrated that non-invasive genetic sampling approach could overcome these
492 constraints with considerable success. Additionally, potential statistical biases related to using
493 various types of lures to bring tigers to camera-traps sites (Kéry et al., 2010; Mowat and
494 Strobeck, 2000; Noyce et al., 2001) or disease transmission (Thiry et al., 1988) using some
495 forms of lures can be overcome by using the non-invasive genetic technique.

496

497 The limitations of camera trap studies include the requirement to follow typical field designs
498 for installation, they need to be maintained in often challenging conditions in which they are
499 prone to failure and even theft, and the logistics for vast survey areas such as the Sundarbans
500 are considerable. Conversely, a non-invasive genetic sampling approach is much easier to
501 implement where all the genetic samples collected over a short period of time could be
502 pooled together without assigning them into different sampling sessions, and can be analysed
503 adopting the 'area search' SECR approach (Efford, 2011).

504

505 Estimates of population density and size of our study are almost similar to camera trap
506 surveys. One of the major challenges of non-invasive genetic technique is to ensure good
507 quality DNA extraction for successful amplification and individual detection. However, we
508 note that genetic sampling can provide additional demographic and population-level
509 information which can be useful for detailed monitoring of these populations. For example,
510 genetic status, sex ratios, family size, effective population size, patterns of dispersal etc.

511

512 Finally, non-invasive genetic sampling can be advantageous over camera trapping for other
513 low density and secretive carnivores (e.g., fishing cat) which cannot be detected and
514 identified by camera trapping using their natural markings.

515

516 We conclude that non-invasive genetic sampling is an appropriate method for assessing tiger
517 population in the Sundarbans mangrove habitat where camera trapping techniques face a
518 range of constraints in relation to limited detections due to unsuitable habitat condition.
519 Therefore, the future monitoring of tigers to determine long-term patterns of population
520 demography and genetic health in this habitat could be carried out by non-invasive genetic
521 sampling.

522

523 **Acknowledgements**

524 We are thankful to the Chief Conservator of Forest (CCF), Ishtiaque Ahmad and Md. Yunus
525 Ali, for providing research and CITES permits. Divisional Forest Officers of the Sundarbans
526 along with field staff extended their support during data collection. This study would not
527 have been possible without the collective effort of our survey teams comprising local
528 fishermen and honey hunters. M. Abdul Aziz was awarded with a Commonwealth
529 Scholarship by the Commonwealth Scholarship Commission, UK and the fieldwork was
530 supported by the Panthera, and WildTeam's Bagh Conservation Activity project financed by
531 USAID Bangladesh. The Society for Conservation Biology provided a part of cost for M.
532 Abdul Aziz for attending training at the University of Nottingham Malaysia Campus during
533 this research. Joe Smith and Wai-Ming Wong of Panthera provided support during this study.
534 We are grateful to the staff of WildTeam, especially Iqbal Hussain, Rezvin Akter, Mahbulul
535 Alam, Nasir Uddin, Abdullah Al Mamun, Alam Howlader, Rubyat Ahmed, Rezu Azam,
536 Sohel Ahmed and Amit Mondol for their support during data collection. Thanks are due to
537 the staff of Dulahazara Safari Park, Bangladesh for cordial assistance in obtaining tiger
538 samples from captive facility. Andrew Kitchener of the National Museums, Scotland and Jo

539 Cook of the Zoological Society of London helped obtain tiger tissue samples for initial
540 optimization of the microsatellite loci. Mark Darmaraj, WWF Malaysia and Mathew Linkie,
541 Fauna and Flora International provided useful suggestions on the initial study design. The
542 SECR modelling analyses were enriched with helpful discussions of Professor David
543 Borchers and Dr Ben Stevenson of the Centre for Research into Ecological and
544 Environmental Modelling of the University of St Andrews, Scotland; Mike Meredith of the
545 Biodiversity Conservation Society Sarawak, Malaysia, and Professor Murray Efford of the
546 Department of Mathematics & Statistics of the University of Otago, New Zealand. We thank
547 three anonymous reviewers for their constructive comments and helpful suggestions, which
548 greatly improved this manuscript.

549

550 **References**

- 551 Ahmad, M.I.U., Greendwood, C.J., Barlow, A.C.D., Islam, M.A., Hossain, A.N.M., Khan,
552 M.M.H., Smith, J.L.D., 2009. Bangladesh Tiger Action Plan 2009-2017. Ministry of
553 Environment and Forests, Bangladesh Forest Department, Dhaka.
- 554 Aziz, M.A., Tollington, S., Barlow, A., Goodrich, J., Shamsuddoha, M., Islam, M.A.,
555 Groombridge, J.J., 2017. Investigating patterns of tiger and prey poaching in the
556 Bangladesh Sundarbans: Implications for improved management. *Glob. Ecol. Conserv.*
557 9, 70–81. doi:10.1016/j.gecco.2016.12.001
- 558 Bagchi, S., Goyal, S.P., Sankar, K., 2003. Prey abundance and prey selection by tigers
559 (*Panthera tigris*) in a semi-arid, dry deciduous forest in western India. *J. Zool.* 260, 285–
560 290.
- 561 Barlow, A., Ahmed, M., Rahman, M., Howlader, A., Smith, A., Smith, J., 2008. Linking
562 monitoring and intervention for improved management of tigers in the Sundarbans of
563 Bangladesh. *Biol. Conserv.* 141, 2032–2040.
- 564 Barlow, A.C.D., 2009. The Sundarbans tiger: evolution, population status and conflict
565 management. University of Minnesota.
- 566 Barlow, A.C.D., McDougal, C., Smith, J.L.D., Gurung, B., Bhatta, S.R., Kumal, S., Mahato,
567 B., Tamang, D.B., 2009. Temporal variation in tiger (*Panthera tigris*) populations and its
568 implications for monitoring. *J. Mammal.* 90, 472–478.
- 569 BFD, 2012. Protected Areas of Bangladesh. Bangladesh Forest Department, Dhaka.
- 570 Bhagavatula, J., Singh, L., 2006. Genotyping faecal samples of Bengal tiger *Panthera tigris*
571 *tigris* for population estimation: a pilot study. *BMC Genet.* 7, 48. doi:10.1186/1471-
572 2156-7-48

573 Borchers, D.L., Efford, M.G., 2008. Spatially explicit maximum likelihood methods for
574 capture-recapture studies. *Biometrics* 64, 377–385.

575 Budowle, B., 2004. SNP typing strategies. *Forensic Sci. Int.* 146.

576 Canonizado, J.A., Hossain, M.A., 1998. Integrated forest management plan for the
577 Sundarbans reserved forest. Dhaka.

578 Chaffey, D.R., Miller, F.R., Sandom, J.H., 1985. A forestry inventory of the Sundarbans,
579 Bangladesh. Surrey, England.

580 Creel, S., Spong, G., Sands, J.L., Rotella, J., Zeigle, J., Joe, L., Murphy, K.M., Smith, D.,
581 2003. Population size estimation in Yellowstone wolves with error-prone noninvasive
582 microsatellite genotypes. *Mol. Ecol.* 12, 2003–2009. doi:10.1046/j.1365-
583 294X.2003.01868.x

584 Davison, A., Birks, J.D.S., Brookes, R.C., Braithwaite, T.C., Messenger, J.E., 2002. On the
585 origin of faeces: Morphological versus molecular methods for surveying rare carnivores
586 from their scats. *J. Zool.* 257, 141–143. doi:DOI:10.1017/S0952836902000730

587 Dey, T.K., Kabir, M.J., Ahsan, M.M., Islam, M.M., Chowdhury, M.M.R., Hassan, S., Roy,
588 M., Qureshi, Q., Naha, D., Kumar, U., Jhala, Y.V., 2015. First Phase Tiger Status Report
589 of Bangladesh Sundarbans. Bangladesh Forest Department, Ministry of Environment
590 and Forests, Government of Bangladesh.

591 Dinerstein, E., Loucks, C., Wikramanayake, E., Ginsberg, J., Sanderson, E., Seidensticker, J.,
592 Forrest, J., Bryja, G., Heydlauff, A., Klenzendorf, S., Leimgruber, P., Mills, J., O'Brien,
593 T.G., Shrestha, M., Simons, R., Songer, M., 2007. The Fate of Wild Tigers. *Bioscience*
594 57, 508. doi:10.1641/B570608

595 Efford, M.G., 2011. Estimation of population density by spatially explicit capture-recapture
596 analysis of data from area searches. *Ecology* 92, 2202–2207.

597 Efford, M.G., Borchers, D.L., Byrom, A.E., 2009. Density Estimation by Spatially Explicit
598 Capture–Recapture: Likelihood-Based Methods, in: Thomson, D.L., Cooch, E.G.,
599 Conroy, M.J. (Eds.), *Modeling Demographic Processes In Marked Populations*. Springer
600 US, Boston, MA, pp. 255–269. doi:10.1007/978-0-387-78151-8_11

601 Farrell, L.E., Roman, J., Sunquist, M.E., 2000. Dietary separation of sympatric carnivores
602 identified by molecular analysis of scats. *Mol. Ecol.* 9, 1583–1590.

603 Frantz, A.C., Pope, L.C., Carpenter, P.J., Roper, T.J., Wilson, G.J., Delahay, R.J., Burke, T.,
604 2003. Reliable microsatellite genotyping of the Eurasian badger (*Meles meles*) using
605 faecal DNA. *Mol. Ecol.* 12, 1649–1661.

606 Frantzen, M. a, Silk, J.B., Ferguson, J.W., Wayne, R.K., Kohn, M.H., 1998. Empirical

607 evaluation of preservation methods for faecal DNA. *Mol. Ecol.* 7, 1423–8.

608 Gerber, B.D., Karpanty, S.M., Kelly, M.J., 2012. Evaluating the potential biases in carnivore
609 capture-recapture studies associated with the use of lure and varying density estimation
610 techniques using photographic-sampling data of the Malagasy civet. *Popul. Ecol.* 54,
611 43–54.

612 Giri, C., Pengra, B., Zhu, Z., Singh, A., Tieszen, L.L., 2007. Monitoring mangrove forest
613 dynamics of the Sundarbans in Bangladesh and India using multi-temporal satellite data
614 from 1973 to 2000. *Estuar. Coast. Shelf Sci.* 73, 91–100. doi:10.1016/j.ecss.2006.12.019

615 Gopal, B., Chauhan, M., 2006. Biodiversity and its conservation in the Sundarban Mangrove
616 Ecosystem. *Aquat. Sci.* 68, 338–354. doi:10.1007/s00027-006-0868-8

617 Hansen, H., Ben-David, M., McDonald, D.B., 2008. Effects of genotyping protocols on
618 success and errors in identifying individual river otters (*Lontra canadensis*) from their
619 faeces. *Mol. Ecol. Notes* 8, 282–289.

620 Hayward, G.D., Miquelle, D.G., Smirnov, E.N., Nations, C., 2002. Monitoring Amur tiger
621 populations: Characteristics of track surveys in snow. *Wildl. Soc. Bull.* 30, 1150–1159.

622 Hossain, A.N.M., Barlow, A., Barlow, C.G., Lynam, A.J., Chakma, S., Savini, T., 2016.
623 Assessing the efficacy of camera trapping as a tool for increasing detection rates of
624 wildlife crime in tropical protected areas. *Biol. Conserv.* 201, 314–319.
625 doi:10.1016/j.biocon.2016.07.023

626 Hussain, Z., Acharya, G., 1994. Mangroves of the Sundarbans, vol. II: Bangladesh. IUCN -
627 The International Conservation Union, Bangkok.

628 Iftekhar, M.S., Islam, M.R., 2004a. Managing mangroves in Bangladesh: A strategy analysis.
629 *J. Coast. Conserv.* 10, 139–146. doi:10.1652/1400-
630 0350(2004)010[0139:MMIBAS]2.0.CO;2

631 Iftekhar, M.S., Islam, M.R., 2004b. Degeneration of Bangladesh's Sundarbans mangroves: a
632 management issue. *Int. For. Rev.* 6, 123–135. doi:10.1505/ifor.6.2.123.38390

633 Iftekhar, M.S., Saenger, P., 2008. Vegetation dynamics in the Bangladesh Sundarbans
634 mangroves: a review of forest inventories. *Wetl. Ecol. Manag.* 16, 291–312.
635 doi:10.1007/s11273-007-9063-5

636 Islam, M., Wahab, M., 2005. A review on the present status and management of mangrove
637 wetland habitat resources in Bangladesh with emphasis on mangrove fisheries and
638 aquaculture. *Hydrobiologia* 542, 165–190. doi:10.1007/1-4020-4111-X_19

639 IUCN–Bangladesh, 2001. The Bangladesh Sundarbans: a photoreal sojourn. IUCN - The
640 International Conservation Union Bangladesh country office, Dhaka, Bangladesh.

641 Jackson, H.A., Bunbury, N., Przelomska, N., Groombridge, J.J., 2016. Evolutionary
642 distinctiveness and historical decline in genetic diversity in the Seychelles Black Parrot
643 *Coracopsis nigra barklyi*. *Ibis* (Lond. 1859). 158, 380–394. doi:10.1111/ibi.12343

644 Jhala, Y. V., Qureshi, Q., Gopal, R., 2015. Status of Tigers in India 2015.

645 Jhala, Y. V., Qureshi, Q., Gopal, R., Sinha, P.R., 2011. Status of tigers, co-predators and prey
646 in India, 2010. New Delhi and Dehradun.

647 Johnsingh, A.J.T., 1983. Large mammalian prey and predators in Bandipur, India. *J. Bombay*
648 *Nat. Hist. Soc.* 80, 1–57.

649 Joshi, A., Vaidyanathan, S., Mondol, S., Edgaonkar, A., Ramakrishnan, U., 2013.
650 Connectivity of tiger (*Panthera tigris*) populations in the human-influenced forest mosaic
651 of Central India. *PLoS One* 8, e77980. doi:10.1371/journal.pone.0077980

652 Kalle, R., Ramesh, T., Qureshi, Q., Sankar, K., 2011. Density of tiger and leopard in a
653 tropical deciduous forest of Mudumalai Tiger Reserve, southern India, as estimated
654 using photographic capture-recapture sampling. *Acta Theriol. (Warsz)*. 56, 335–342.

655 Karanth, K.U., 2005. Joining the dots but missing the cats? *Cat News* 43, 8–11.

656 Karanth, K.U., 1995. Estimating tiger *Panthera tigris* populations from camera-trap data using
657 capture—recapture models. *Biol. Conserv.* 71, 333–338. doi:10.1016/0006-
658 3207(94)00057-W

659 Karanth, U.K., Nichols, J.D., 2000. Ecological status and conservation of tigers in India.
660 Bangalore, India. Bangalore.

661 Karanth, U.K., Nichols, J.D., 1998. Estimation of Tiger Densities in India Using
662 Photographic Captures and Recaptures. *Ecology* 79, 2852–2862.

663 Karanth, U.K., Sunquist, M.E., Sunquist, M.E., 1995. Prey Selection by Tiger, Leopard and
664 Dhole in Tropical Forests. *J. Anim. Ecol.* 64, 439–450. doi:10.2307/5647

665 Karki, J.B., Pandav, B., Jnawali, S.R., Shrestha, R., Pradhan, N.M.B., Lamichane, B.R.,
666 Khanal, P., Subedi, N., Jhala, Y. V., 2015. Estimating the abundance of Nepal’s largest
667 population of tigers *Panthera tigris*. *Oryx* 49, 150–156.
668 doi:10.1017/S0030605313000471

669 Kéry, M., Gardner, B., Stoeckle, T., Weber, D., Royle, J.A., 2010. Use of spatial capture-
670 recapture modeling and DNA data to estimate densities of elusive animals. *Conserv.*
671 *Biol.* 25, 356–64. doi:10.1111/j.1523-1739.2010.01616.x

672 Khan, M.M.H., 2012. Population and prey of the Bengal Tiger *Panthera tigris tigris*
673 (Linnaeus, 1758) (Carnivora: Felidae) in the Sundarbans, Bangladesh. *J. Threat. Taxa* 4,
674 2370–2380.

675 Khan, M.M.H., 2008. Prey selection by tigers *Panthera tigris* (Linnaeus 1758) in the
676 Sundarbans East Wildlife Sanctuary of Bangladesh. *Journal Bombay Nat. Hist. Soc.*
677 105, 255–263.

678 Kohn, M.H., York, E.C., Kamradt, D. a, Haught, G., Sauvajot, R.M., Wayne, R.K., 1999.
679 Estimating population size by genotyping faeces. *Proc. Biol. Sci.* 266, 657–663.
680 doi:10.1098/rspb.1999.0686

681 Lucchini, V., Fabbri, E., Marucco, F., Ricci, S., Boitani, L., Randi, E., 2002. Noninvasive
682 molecular tracking of colonizing wolf (*Canis lupus*) packs in the western Italian Alps.
683 *Mol. Ecol.* 11, 857–868.

684 Mace, R.D., Minta, S.C., Manley, T.L., Aune, K.E., 1994. Estimating grizzly bear population
685 size using camera sightings. *Wildl. Soc. Bull.* 22, 74–82.

686 Marshall, T.C., Slate, J., Kruuk, L.E.B., Pemberton, J.M., 1998. Statistical confidence for
687 likelihood-based paternity inference in natural populations. *Mol. Ecol.* 7, 639–655.
688 doi:10.1046/j.1365-294x.1998.00374.x

689 Menotti-Raymond, M., David, V.A., Lyons, L.A., Schaffer, A.A., Tomlin, J.F., Hutton,
690 M.K., O’Brien, S.J., 1999. A Genetic Linkage Map of Microsatellites in the Domestic
691 Cat (*Felis catus*). *Genomics* 57, 9–23.

692 Mondol, S., Karanth, K.U., Kumar, N.S., Gopaldaswamy, A.M., Andheria, A., Ramakrishnan,
693 U., 2009a. Evaluation of non-invasive genetic sampling methods for estimating tiger
694 population size. *Biol. Conserv.* 142, 2350–2360. doi:10.1016/j.biocon.2009.05.014

695 Mondol, S., Karanth, K.U., Ramakrishnan, U., 2009b. Why the Indian subcontinent holds the
696 key to global tiger recovery. *PLoS Genet.* 5, e1000585.
697 doi:10.1371/journal.pgen.1000585

698 Mowat, G., Strobeck, C., 2000. Estimating population size of grizzly bears using hair capture,
699 DNA profiling, and mark-recapture analysis. *J. Wildl. Manage.* 64, 183–193.

700 Mukherjee, N., Mondol, S., Andheria, A., Ramakrishnan, U., 2007. Rapid multiplex PCR
701 based species identification of wild tigers using non-invasive samples. *Conserv. Genet.*
702 8, 1465–1470. doi:10.1007/s10592-007-9289-z

703 Noyce, K. V, Garshelis, D.L., Coy, P.L., 2001. Differential vulnerability of black bears to
704 trap and camera sampling and resulting biases in mark-recapture estimates. *Ursus* 12,
705 211–226.

706 Pilgrim, K.L., Mckelvey, K.S., Riddle, A.E., Schwartz, M.K., 2005. Felid sex identification
707 based on noninvasive genetic samples. *Mol. Ecol. Notes* 5, 60–61. doi:10.1111/j.1471-
708 8286.2004.00831.x

709 Reddy, P.A., Gour, D.S., Bhavanishankar, M., Jaggi, K., Hussain, S.M., Harika, K., Shivaji,
710 S., 2012. Genetic evidence of tiger population structure and migration within an isolated
711 and fragmented landscape in Northwest India. PLoS One 7, e29827.
712 doi:10.1371/journal.pone.0029827

713 Sanderson, E., Forrest, J., Loucks, C., Ginsberg, J., Dinerstein, E., Seidensticker, J.,
714 Leimgruber, P., Songer, M., Heydlauff, A., O'Brien, T., Bryja, G., Klenzendorf, S.,
715 Wikramanayake, E., 2006. Setting Priorities for the Conservation and Recovery of Wild
716 Tigers: 2005-2015, The Technical Assessment. New York - Washington D.C.

717 Sharma, S., Dutta, T., Maldonado, J.E., Wood, T.C., Panwar, H.S., Seidensticker, J., 2012.
718 Spatial genetic analysis reveals high connectivity of tiger (*Panthera tigris*) populations in
719 the Satpura-Maikal landscape of Central India. Ecol. Evol. 3, 48–60.
720 doi:10.1002/ece3.432

721 Smith, J.L.D., 1978. Smithsonian Tiger Ecology Project Report 13. Washington D.C.

722 Sollmann, R., Tôrres, N.M., Furtado, M.M., de Almeida Jácomo, A.T., Palomares, F.,
723 Roques, S., Silveira, L., 2013. Combining camera-trapping and noninvasive genetic data
724 in a spatial capture–recapture framework improves density estimates for the jaguar. Biol.
725 Conserv. 167, 242–247. doi:10.1016/j.biocon.2013.08.003

726 Taberlet, P., Griffin, S., Uhres, E., Waits, L.P., Dubois Paganon, C., Bouvet, J., Camarra, J.J.,
727 Hanotte, O., Burke, T., 1997. Noninvasive genetic tracking of the endangered Pyrenean
728 brown bear population. Mol. Ecol. 6, 869–876.

729 Thinley, P., Curtis, P.D., 2015. Estimating Wild Tiger (*Panthera tigris* Linnaeus) Abundance
730 and Density using a Spatially-explicit Capture-recapture Model in Jigme Dorji National
731 Park, Bhutan. Bhutan J. Nat. Resour. Dev. doi:10.17102/cnr.2015.01

732 Thiry, E., Brochier, B., Schwers, a, Thomas, I., Dubuisson, J., 1988. Diseases of wild
733 animals transmissible to domestic animals. Rev. Sci. Tech. l'OIE 7, 705–736.

734 Valiere, N., 2002. GIMLET: A computer program for analysing genetic individual
735 identification data. Mol. Ecol. Notes.

736 Valière, N., 2002. a computer program for analysing genetic GIMLET. Mol. Ecol. Notes 2,
737 377–379. doi:10.1046/j.1471-8278

738 van Oosterhout, C., Hutchinson, W.F., Wills, D.P.M., Shipley, P., 2004. micro-checker:
739 software for identifying and correcting genotyping errors in microsatellite data. Mol.
740 Ecol. Notes 4, 535–538. doi:10.1111/j.1471-8286.2004.00684.x

741 Waits, L.P., Luikart, G., Taberlet, P., 2001. Estimating the probability of identity among
742 genotypes in natural populations: cautions and guidelines. Mol. Ecol. 10, 249–256.

743 doi:10.1046/j.1365-294X.2001.01185.x
744 Walston, J., Robinson, J.G., Bennett, E.L., Breitenmoser, U., da Fonseca, G.A.B., Goodrich,
745 J., Gumal, M., Hunter, L., Johnson, A., Ullas Karanth, K., Leader-Williams, N.,
746 MacKinnon, K., Miquelle, D., Pattanavibool, A., Poole, C., Rabinowitz, A., Smith,
747 J.L.D., Stokes, E.J., Stuart, S.N., Vongkhamheng, C., Wibisono, H., 2010. Bringing the
748 tiger back from the brink-the six percent solution. PLoS Biol. 8, 6–9.
749 doi:10.1371/journal.pbio.1000485
750 Waterhouse, A.M., Procter, J.B., Martin, D.M.A., Clamp, M., Barton, G.J., 2009. Jalview
751 Version 2--a multiple sequence alignment editor and analysis workbench.
752 Bioinformatics 25, 1189–1191. doi:10.1093/bioinformatics/btp033
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769

770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790

Table 1. Summary of scat samples (plus number of hair samples in parentheses) collected, screened and genotyped from each of the sample areas of the Bangladesh Sundarbans between November 2014 and February 2015.

Sampling area	Area* (km ²)	Protection status	Sampling duration (days)	Samples collected from field	Samples screened as tiger	Samples genotyped for at least 5 loci	Samples sexed
Satkhira Block (SB)	342	Reserve forest	21	62(15)	23(10)	10(5)	6(5)
West Wildlife Sanctuary (WS)	715	Protected area	13	124(28)	61(21)	21(12)	16(12)
Chandpai Block (CB)	554	Reserve forest	21	91(36)	57(17)	27(6)	19(6)
East Wildlife Sanctuary (ES)	383	Protected area	21	62(22)	29(15)	18(6)	17(6)
Totals	1,994	--	76	440	233	105	87

* Area included forest land and waterbodies

791

792

793

794

795

796

797

798 **Table 2.** Sample area (forest land only), capture-recaptures and density parameter estimates with spatially explicit capture-recapture (SECR)
 799 model for area-wise and overall estimates of tigers using non-invasively collected DNA data from the Bangladesh Sundarbans between
 800 November 2014 and February 2015.
 801

Name of sample area	Area* (km ²)	No. of individuals detected	No. of total detections	Tiger density (D ± SE per 100 km ²)	Probability of detection (g0 ± SE)	Spatial distance moved (σ ± SE km)
Satkhira Block (SB)	275	6	15	1.86 ± 0.81	0.0226 ± 0.0098	3.989 ± 0.825
West Wildlife Sanctuary (WS)	414	15	33	2.99 ± 0.85	0.0185 ± 0.0057	3.920 ± 0.506
Chandpai Block (CB)	418	14	33	3.18 ± 0.90	0.0224 ± 0.0071	3.088 ± 0.438
East Wildlife Sanctuary (ES)	290	10	24	3.17 ± 1.04	0.0361 ± 0.0128	2.918 ± 0.416
Overall (all sampled areas)	1,397	45	105	2.85 ± 0.44	0.0223 ± 0.0038	3.478 ± 0.262

802 * Area estimated excluding waterbodies.
 803
 804
 805
 806
 807
 808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818

819 **Table 3.** Genetic variability at 10 microsatellite loci for field samples (n = 105*) collected
 820 from the Bangladesh Sundarbans between November 2014 and February 2015.
 821

Locus	Allele size range (bp)	No. of alleles/locus	Dropout rate	False allele rate	H _E	H _O	P _{ID(sibs)}
FCA279	97-107	7	0	0.19	0.78	0.5	8.14E-02
FCA232	99-113	5	0	0	0.78	0.42	6.79E-03
FCA090	107-117	5	0	0	0.77	0.38	6.61E-04
FCA672	93-105	6	0	0	0.67	0.24	1.45E-05
D15	119-139	5	0	0.12	0.68	0.39	9.61E-05
FCA304	121-129	4	0.26	0	0.67	0.34	2.44E-06
FCA126	138-144	4	0	0	0.68	0.15	4.17E-07
F41	111-135	6	0.05	0	0.63	0.59	7.61E-08
FCA230	103-115	7	0	0	0.54	0.14	1.19E-09
E7	137-151	5	0	0	0.56	0.28	4.61E-09

822 *Sample area-wise amplified samples: SB (n= 15), WS (n=33), CB (n=33), ES (n=24)

823 He: Expected heterozygosity, Ho: Observed heterozygosity.

824
 825
 826
 827

828
829
830
831
832

Table 4. Sample area-wise comparison of tiger density estimates between this study and a camera-trap survey by Dey et al. (2015) in the Bangladesh Sundarbans.

Sampling area	Study method	No. of individuals detected	SECR model	Tiger density ($D \pm SE$ per 100 km ²)	Probability of detection ($g0 \pm SE$)	Spatial distance moved ($\sigma \pm SE$ km)
Area-wise						
Satkhira Block ^a	DNA study	6	$D(.)g0(.)\sigma(.)$	1.86 ± 0.81	0.0226 ± 0.0098	3.989 ± 0.825
Block III (Satkhira) ^a	Camera traps	13	$D(.)g0(bk)\sigma(.)$	2.77 ± 0.78	0.0100 ± 0.0020	4.270 ± 0.050
East Wildlife Sanctuary ^b	DNA study	10	$D(.)g0(.)\sigma(.)$	3.17 ± 1.04	0.0361 ± 0.0127	2.918 ± 0.416
Block I (Sarankhola) ^b	Camera traps	18	$D(.)g0(bk)\sigma(.)$	3.70 ± 0.91	0.0100 ± 0.0030	3.370 ± 0.350
Overall						
Sampling area (1,397 km ²)	DNA study	48	$D(.)g0(.)\sigma(.)$	2.85 ± 0.44	0.0231 ± 0.0038	3.478 ± 0.262
Sampling area (1,265 km ²)	Camera traps	38	$D(.)g0(bk)\sigma(.)$	2.17 (1.73-2.68)	Not available	Not available

833 Note: ^aSatkhira Block completely overlapped with Block III (Satkhira), and ^bEast Wildlife Sanctuary with Block I (Sarankhola) of camera-trap
834 study (Dey et al., 2015).

835
836
837
838
839
840
841
842
843

844
845
846

Table A.1. Characteristics of 14 microsatellite loci optimized for reference samples (RS, n = 10) and field-collected samples (FS, n = 10).

Locus*	Allele size range (bp)	Amplification success (%)		Allelic dropout		False alleles		No. of alleles/locus		Expected heterozygosity, H_E		Observed heterozygosity, H_O		Probability of identity, $P_{ID(sibs)}$	
		RS	FS	RS	FS	RS	FS	RS	FS	RS	FS	RS	FS	RS	FS
FCA090	111-113	90	78	0	0	0	0	4	5	0.69	0.81	0.63	0.63	1.86E-02	1.64E-01
FCA672	93-105	100	100	0	0	0	0	4	5	0.76	0.81	0.5	0.8	4.23E-01	3.99E-01
FCA232	99-103	100	89	0	0	0	0	5	5	0.73	0.71	0.3	0.33	1.87E-01	7.62E-02
D15	119-139	100	89	0	0	0	0	4	4	0.71	0.71	0.67	0.9	8.54E-02	3.71E-02
FCA279	99-107	100	100	0	0	0	0	3	3	0.62	0.66	0.44	0.6	9.15E-03	1.86E-02
FCA304	121-129	100	89	0	0	0	0	4	3	0.5	0.66	0.3	0.44	8.49E-04	4.88E-03
F41	111-133	100	89	0	0	0	0.14	5	5	0.55	0.66	0.33	0.4	1.44E-03	9.44E-03
FCA126	140-144	100	89	0	0.11	0	0	3	3	0.71	0.49	0.22	0.1	3.96E-02	1.53E-03
FCA309	98-100	100	89	0	0.11	0	0	2	2	0.5	0.48	0.11	0.1	3.07E-04	2.58E-03
FCA230	105-115	100	100	0	0	0	0	6	3	0.57	0.29	0.5	0.2	2.58E-03	2.74E-04
E7	138-151	100	100	0	0	0	0	3	3	0.43	0.27	0.3	0.1	1.23E-04	2.74E-04
FCA043	120-130	100	89	0	0	0	0	2	2	0.48	0.52	0.3	0.44	1.91E-04	5.40E-04
FCA052	108-114	100	100	0	0	0	0	3	2	0.61	0.39	0.4	0.3	4.81E-03	3.35E-04
FCA164	80-90	100	100	0	0	0	0	2	2	0.53	0.53	0.78	0.78	5.04E-04	9.09E-04

*All loci optimized from Menotti-Raymond et al. (1999), except E7 and D15 (Bhagavatula and Singh, 2006).


847
848
849
850
851
852
853
854

855 **Table A.2.** Locus name, primer sequences, annealing temperature (AT), forward primer fluorescent dye (FD), and PCR multiplexes (PM) used
 856 in this study.
 857

Locus name	Forward sequence	Reverse sequence	AT (°C)	FD	PM
F41	GTCTGCATCTTCAAATAGGA	GTACCTGAGTTGGCTGTTGA	56	FAM	Set 1
D15	TGTGACCTTTCTCTAGTTTC	GCACAAAACATTCAGTCTCC	55	FAM	Set 1
Fca232	ATGACCATCTCAAACCTCATGG	AGCTGAGTTTGCCTTTATCATG	56	HEX	Set 1
Fca304	TCATTGGCTACCACAAAGTAGG	CTGCATGCCATTGGGTAAC	56	FAM	Set 2
E7	GCCCCAAAGCCCTAAAATAA	GCATGTTCGGACAGTAAAGCA	55	NED	Set 2
ZN (ZF _x /Zfy)	AAGTTTACACAACCACCTGG	CACAGAATTTACACTTGTGCA	55	NED	Set 2
Fca126	GCCCCTGATACCCTGAATG	CTATCCTTGCTGGCTGAAGG	56	HEX	Set 3
Fca672	AAGTTGCTTGACACACTGC	TCCAAGAGCCTTTTCAGTTAGG	56	HEX	Set 3
Fca090	ATCAAAAGTCTTGAAGAGCATGG	TGTTAGCTCATGTTTCATGTGTCC	52	HEX	Set 4
Fca230	AAGAATGGACTTGGGAAATGG	AAACCACAACAGGCAAAAGG	52	NED	Set 4
Fca279	AGCCAAGTAATATTCCTCTGTG	GTCCATCCGCAGATGAATG	52	FAM	Set 4

858 All loci optimized from Menotti-Raymond et al. (1999), except D15, E7 (Bhagavatula and Singh, 2006), and ZN (Pilgrim et al., 2005).

859
 860
 861



862

863 **Fig. 1.** Location of Bangladesh, and Sundarbans with wildlife sanctuaries, sample areas and
 864 sample grids. Sample area: SB – Sathkira Block, CB – Chandpai Block, WS – West Wildlife
 865 Sanctuary, ES – East Wildlife Sanctuary.

