

# Kent Academic Repository

## Full text document (pdf)

### Citation for published version

Green, Reuben Mark and Paget, Rowena E. (2017) Iterated Inflations of Cellular Algebras. *Journal of Algebra*, 493 . pp. 341-345. ISSN 0021-8693.

### DOI

<https://doi.org/10.1016/j.jalgebra.2017.09.030>

### Link to record in KAR

<http://kar.kent.ac.uk/63860/>

### Document Version

Publisher pdf

#### Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

#### Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

#### Enquiries

For any further enquiries regarding the licence status of this document, please contact:

[researchsupport@kent.ac.uk](mailto:researchsupport@kent.ac.uk)

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>



Contents lists available at ScienceDirect

Journal of Algebra

[www.elsevier.com/locate/jalgebra](http://www.elsevier.com/locate/jalgebra)



## Iterated inflations of cellular algebras



Reuben Green <sup>\*,1</sup>, Rowena Paget

*School of Mathematics, Statistics and Actuarial Science, Sibson Building,  
University of Kent, CT2 7FS, UK*

### ARTICLE INFO

*Article history:*

Received 7 April 2017

Available online 5 October 2017

Communicated by Volodymyr Mazorchuk

*Keywords:*

Cellular algebras

Iterated inflations

### ABSTRACT

We present a result characterising iterated inflations of cellular algebras, derived from the work of König and Xi. This result is intended to replace an incorrect proposition in the literature, and gives explicit and readily checked conditions which establish that an algebra is an iterated inflation of cellular algebras, and hence is cellular, with cellular data directly related to the cellular data of the constituent cellular algebras.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

The concept of a *cellular algebra* was introduced by Graham and Lehrer in [2], and has found wide application in the representation theory of finite groups and associative algebras; it has been especially useful in the study of *diagram algebras*. We refer the reader to the existing literature for the definition and basic properties of cellular algebras; we use the notation of [3]. We work over a field  $k$ , and all our  $k$ -algebras are finite-dimensional; since all tensor products are taken over  $k$ , we abbreviate  $\otimes_k$  to  $\otimes$ .

In [4], König and Xi introduced the notion of an *iterated inflation* of cellular algebras. Informally, iterated inflation allows us to prove that an algebra is cellular by showing that it may be constructed using some family of existing cellular algebras in a certain

\* Corresponding author.

*E-mail addresses:* [rmg29@kent.ac.uk](mailto:rmg29@kent.ac.uk) (R. Green), [r.e.paget@kent.ac.uk](mailto:r.e.paget@kent.ac.uk) (R. Paget).

<sup>1</sup> Supported by EPSRC grant [EP/M508068/1].

way. There are, however, various technical details and conditions which must be satisfied in order for such a construction to yield a cellular algebra as desired. In [4] and [5], these conditions are not presented in a single definition or result, but rather are developed in the course of the text. In [7], Xi gave the following lemma to provide a precise characterisation of iterated inflations of cellular algebras:

**Lemma.** (Xi — Lemma 3.3 in [7]) *Let  $A$  be an algebra with an anti-involution  $\sigma$ . Suppose there is a vector space decomposition  $A = \bigoplus_{j=1}^m V_j \otimes B_j \otimes V_j$  where  $V_j$  is a vector space and  $B_j$  is a cellular algebra with respect to an anti-involution  $\sigma_j$  and a cell chain  $J_1^{(j)} \subseteq \dots \subseteq J_{s_j}^{(j)} = B_j$  for each  $j$ . Define  $J_t = \bigoplus_{j=1}^t V_j \otimes B_j \otimes V_j$ . Assume that*

- (i) *the restriction of  $\sigma$  on  $V_j \otimes B_j \otimes V_j$  is given by  $w \otimes b \otimes v \mapsto v \otimes \sigma_j(b) \otimes w$*
- (ii) *for each  $j$ , there is a bilinear form  $\phi_j : V_j \times V_j \rightarrow B_j$  such that  $\sigma_j(\phi_j(w, v)) = \phi_j(v, w)$  for all  $v, w \in V_j$*
- (iii) *for  $x, y, u, v \in V_j$  and  $b, c \in B_j$ , we have*

$$(x \otimes b \otimes y)(u \otimes c \otimes v) = x \otimes b\phi_j(y, u)c \otimes v \pmod{J_{j-1}}$$

- (iv)  *$(V_j \otimes J_l^{(j)} \otimes V_j) + J_{j-1}$  is an ideal in  $A$  for all  $l$  and  $j$ .*

*Then  $A$  is a cellular algebra.*

This lemma was used, for example, to prove the cellularity of the partition algebra in [7], and of the BMW-algebra in [6]; both of these papers have become standard references for these facts. The lemma is, however, incorrect, as the following example shows. Recall that the matrix algebra  $M_n(k)$  is cellular with respect to the anti-involution which takes a matrix to its transpose. In particular, the field  $k$  is cellular with respect to the identity map. Let  $k$  be any field, and define  $A$  to be the  $k$ -algebra obtained by equipping  $M_2(k) \oplus k \oplus k$  with the (commutative) multiplication

$$\left( \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \alpha, \beta \right) \left( \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}, \gamma, \delta \right) = \left( \begin{pmatrix} \gamma a_{11} & \delta a_{12} \\ \delta a_{21} & \gamma a_{22} \end{pmatrix} + \begin{pmatrix} \alpha b_{11} & \beta b_{12} \\ \beta b_{21} & \alpha b_{22} \end{pmatrix}, \alpha\gamma, \beta\delta \right),$$

which has identity

$$\left( \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, 1, 1 \right).$$

Let  $\sigma$  be the anti-involution  $\sigma(M, \alpha, \beta) = (M^T, \alpha, \beta)$  on  $A$ . Let  $V_1 = V_2 = V_3 = k$  and define cellular algebras  $B_1 = M_2(k)$  and  $B_2 = B_3 = k$ . Thus we obtain an isomorphism of vector spaces

$$A \cong (V_1 \otimes B_1 \otimes V_1) \oplus (V_2 \otimes B_2 \otimes V_2) \oplus (V_3 \otimes B_3 \otimes V_3)$$

from the mapping  $(M, \alpha, \beta) \mapsto (1 \otimes M \otimes 1, 1 \otimes \alpha \otimes 1, 1 \otimes \beta \otimes 1)$ . Define  $k$ -bilinear forms  $\phi_1, \phi_2, \phi_3$ , where  $\phi_i$  maps  $V_i \times V_i$  to  $B_i$ , by taking  $\phi_1$  to be zero and  $\phi_2(1, 1) = \phi_3(1, 1) = 1$ . One may check that these definitions satisfy the hypotheses of Xi’s lemma (note that  $B_j$  has cell chain  $\{0\} \subseteq B_j$  for each  $j$ ). Suppose for a contradiction that  $A$  is cellular, and let

$$e = \left( \left( \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, 0, 1 \right) \right).$$

Then  $e$  is an idempotent of  $A$ , and one may see by direct calculation that  $eAe$  is a 3-dimensional subspace of  $A$  upon which  $\sigma$  does not act as the identity map. Further, we have  $\sigma(e) = e$  and so by Proposition 4.3 of [3], the algebra  $eAe$  must be cellular with respect to the restriction of  $\sigma$ . But it follows easily from the definition of a cellular algebra (and the fact that the only expression of 3 as a sum of squares is  $1 + 1 + 1$ ) that the anti-involution on a 3-dimensional cellular algebra is the identity map; thus  $A$  is not cellular.<sup>2</sup>

We now present a result which can serve as a convenient replacement for the incorrect lemma above. This result characterises iterated inflations via two readily checked conditions; these two conditions were noted for the Brauer algebra by König and Xi in [5] (Lemmas 5.4 and 5.5). The proof of this theorem is a straightforward verification of the axioms for a cellular algebra given in [2], and thus is omitted.

**Theorem 1.** *Let  $A$  be a  $k$ -algebra, with an anti-involution  $\sigma$ . Suppose that we have, up to isomorphism of  $k$ -vector spaces, a decomposition*

$$A \cong \bigoplus_{i \in I} V_i \otimes B_i \otimes V_i$$

of  $A$ , where  $I$  is a finite partially ordered set, each  $V_i$  is a  $k$ -vector space, and each  $B_i$  is a cellular algebra over  $k$  with respect to an anti-involution  $\sigma_i$  and cellular data  $(\Lambda_i, M_i, C)$ . We shall henceforth consider  $A$  to be identified with this direct sum of tensor products.

Suppose that for each  $i \in I$ , we have a basis  $\mathcal{V}_i$  for  $V_i$  and a basis  $\mathcal{B}_i$  for  $B_i$ , such that:

1. For each  $i \in I$ , we have for any  $u, v \in \mathcal{V}_i$  and any  $b \in \mathcal{B}_i$  that

$$\sigma(u \otimes b \otimes v) = v \otimes \sigma_i(b) \otimes u. \tag{1}$$

2. Let  $\mathcal{A}$  be the basis of  $A$  consisting of all elements  $u \otimes b \otimes v$  for all  $u, v \in \mathcal{V}_i$  and all  $b \in \mathcal{B}_i$ , as  $i$  ranges over  $I$ . Then for any  $i \in I$  we have maps  $\phi_i : \mathcal{A} \times \mathcal{V}_i \rightarrow \mathcal{V}_i$  and

---

<sup>2</sup> If  $k$  has characteristic 2, then  $A$  and  $\sigma$  may be shown to satisfy the alternative definition of cellularity given by Goodman and Graber in [1], demonstrating that this definition is indeed different from that given by Graham and Lehrer over fields of characteristic 2.

$\theta_i : \mathcal{A} \times \mathcal{V}_i \rightarrow B_i$  such that for any  $u, v \in \mathcal{V}_i$  and any  $b \in \mathcal{B}_i$ , we have for any  $a \in \mathcal{A}$  that

$$a \cdot (u \otimes b \otimes v) \equiv \phi_i(a, u) \otimes \theta_i(a, u)b \otimes v \pmod{J(< i)} \tag{2}$$

where  $J(< i) = \bigoplus_{l < i} V_l \otimes B_l \otimes V_l$ .

Then  $A$  is cellular with respect to  $\sigma$  and the cellular data  $(\Lambda, M, C)$ , where  $\Lambda$  is the set  $\{(i, \lambda) : i \in I \text{ and } \lambda \in \Lambda_i\}$  with the lexicographic order,  $M(i, \lambda)$  is  $\mathcal{V}_i \times M_i(\lambda)$ , and  $C_{(x,X),(y,Y)}^{(i,\lambda)} = x \otimes C_{X,Y}^\lambda \otimes y$ .

Note that we may use *any* bases of the cellular algebras  $B_i$  to check the conditions of **Theorem 1**: we need not use the cellular bases of the  $B_i$ . This is convenient as cellular bases are often awkward to work with in practice.

**Proposition 2.** *Let  $A$  be an algebra satisfying the hypotheses of **Theorem 1**. Then the multiplication in each “layer” of  $A$  is “governed” by a bilinear form as in Xi’s lemma: for each  $i \in I$  there is a unique  $B_i$ -valued  $k$ -bilinear form  $\psi_i$  on  $V_i$  such that for any  $u, v, x, y \in V_i$  and  $b, c \in B_i$ , we have  $\psi_i(y, u) = \sigma_i(\psi_i(u, y))$  and*

$$(x \otimes c \otimes y)(u \otimes b \otimes v) \equiv x \otimes c \psi_i(y, u)b \otimes v \pmod{J(< i)}. \tag{3}$$

**Proof (outline).** We shall show that there is a bilinear form  $\psi_i$  satisfying (3) by proving that we have a map  $\psi_i : \mathcal{V}_i \times \mathcal{V}_i \rightarrow B_i$  satisfying (3); the rest is then easy. First note that (1) holds even when  $u, v, b$  are not required to be basis elements, and likewise for  $v, b$  in (2); these equations are used extensively in what follows. By applying  $\sigma^2 = \text{id}_A$  we may show that

$$(x \otimes c \otimes y)(u \otimes \sigma_i(b) \otimes u) \equiv x \otimes c \sigma_i(\theta_i(u \otimes b \otimes u, y)) \otimes \phi_i(u \otimes b \otimes u, y) \tag{4}$$

modulo  $J(< i)$ , for  $u, x, y \in \mathcal{V}_i$  and  $b, c \in \mathcal{B}_i$ . Now for  $u, x, y \in \mathcal{V}_i$  and  $c \in \mathcal{B}_i$ , we may on the one hand apply (2) to  $(x \otimes c \otimes y)(u \otimes 1_{B_i} \otimes u)$ , and on the other hand we may expand  $1_{B_i}$  in terms of the basis  $\sigma(\mathcal{B}_i)$  and apply (4). By carefully comparing the results of these calculations, we may deduce that if  $\theta_i(x \otimes c \otimes y, u) \neq 0$  then  $\phi_i(x \otimes c \otimes y, u)$  is some scalar multiple of  $x$ , say  $\alpha(x, c, y, u)x$ , and further that

$$\phi_i(x \otimes c \otimes y, u) \otimes \theta_i(x \otimes c \otimes y, u) \otimes u = x \otimes c \psi_i(y, u) \otimes u$$

for a value  $\psi_i(y, u) \in B_i$  depending only on  $y$  and  $u$ ; hence we may deduce that  $\alpha(x, c, y, u)\theta_i(x \otimes c \otimes y, u)$  is equal to  $c \psi_i(y, u)$ . Then for  $u, v, x, y \in \mathcal{V}_i$  and  $b, c \in \mathcal{B}_i$ , we apply (2) to the left-hand side of (3) and then apply the above results, and hence obtain the right-hand side of (3).  $\square$

The first part of our final result is implicit in [4] and [5], and a more explicit formulation was given in [7]; we believe that the second part has not previously appeared in the literature. The proof is easy and is omitted.

**Proposition 3.** *Let  $A$  be as in Theorem 1, let  $(i, \lambda) \in \Lambda$ , and let  $\Delta^\lambda$  be the cell module of  $B_i$  corresponding to  $\lambda$ . The cell module  $\Delta^{(i, \lambda)}$  of  $A$  may be obtained by equipping  $V_i \otimes \Delta^\lambda$  with the action given, for  $a \in \mathcal{A}$ ,  $x \in \mathcal{V}_i$  and  $z \in \Delta^\lambda$ , by*

$$a(x \otimes z) = \phi_i(a, x) \otimes \theta_i(a, x)z.$$

*Recall that the simple modules of a cellular algebra may be obtained as quotients of the cell modules by the radicals of certain bilinear forms; with the above construction of the cell module, if  $\langle \cdot, \cdot \rangle$  is the bilinear form on  $V_i \otimes \Delta^\lambda$  and  $\langle \cdot, \cdot \rangle_\lambda$  is the bilinear form on  $\Delta^\lambda$ , then for any  $x, y \in V_i$  and any  $z, w \in \Delta^\lambda$ , we have*

$$\langle x \otimes z, y \otimes w \rangle = \langle z, \psi_i(x, y)w \rangle_\lambda = \langle \psi_i(y, x)z, w \rangle_\lambda.$$

## Acknowledgments

The authors are grateful to the referee for a number of suggestions which have improved this paper; in particular, the referee suggested a significant simplification of the proof that the algebra used in the example is not cellular.

## References

- [1] F. Goodman, J. Graber, Cellularity and the Jones basic construction, *Adv. in Appl. Math.* 46 (2011) 312–362.
- [2] J.J. Graham, G.I. Lehrer, Cellular Algebras, *Invent. Math.* 123 (1) (1996) 1–34.
- [3] S. König, C.C. Xi, On the structure of cellular algebras, *Canad. Math. Soc. Conf. Proc.*, vol. 24, Amer. Math. Soc., Providence, RI, 1999, pp. 365–386.
- [4] S. König, C.C. Xi, Cellular algebras: inflations and Morita equivalences, *J. Lond. Math. Soc.* (2) 60 (1999) 700–722.
- [5] S. König, C.C. Xi, A characteristic free approach to Brauer algebras, *Trans. Amer. Math. Soc.* 353 (2001) 1489–1505.
- [6] C.C. Xi, On the quasi-heredity of Birman–Wenzl algebras, *Adv. Math.* 154 (2000) 280–298.
- [7] C.C. Xi, Partition Algebras are Cellular, *Compos. Math.* 119 (1999) 99–109.