

Kent Academic Repository

Full text document (pdf)

Citation for published version

Scholz, Aleks and Llama, Joe and Muzic, Koraljka and Faller, Sarah and Froebrich, Dirk and Stelzer, Beate (2018) Discovery of a Magnetic White Dwarf with Unusual Short-period Variability. *Research Notes of the American Astronomical Society*, 2 (27). ISSN 2515-5172.

DOI

<https://doi.org/10.3847/2515-5172/aaadbf>

Link to record in KAR

<http://kar.kent.ac.uk/66560/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Discovery of a magnetic white dwarf with unusual short-period variability

ALEKS SCHOLZ,¹ JOE LLAMA,² KORALJKA MUZIC,³ SARAH FALLER,¹ DIRK FROEBRICH,⁴ AND BEATE STELZER^{5,6}

¹*SUPA, School of Physics & Astronomy, University of St Andrews, North Haugh, St Andrews, KY16 9SS, United Kingdom*

²*Lowell Observatory, 1400 W. Mars Hill Rd. Flagstaff, AZ 86001, USA*

³*SIM/CENTRA, Faculdade de Ciencias de Universidade de Lisboa, Ed. C8, Campo Grande, P-1749-016 Lisboa, Portugal*

⁴*Centre for Astrophysics and Planetary Science, University of Kent, Canterbury, CT2 7NH, UK*

⁵*Institut für Astronomie und Astrophysik, Eberhard Karls Universität, Sand 1, D-72076 Tübingen, Germany*

⁶*INAF-Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, 90134, Palermo, Italy*

(Accepted for publication in RNAAS)

Keywords: (stars:) white dwarfs — (stars:) starspots — stars: magnetic field – stars: peculiar

We report the discovery of a magnetic white dwarf which shows periodic variability with $P = 110$ min, color-dependent amplitudes and a transient phase shift in the blue compared to the red lightcurve – a previously unknown type of variability for this type of object. We attribute the variations either to a close ultracool (thus far undetected) companion or, more likely, to magnetic spots with unusual temperature structure.

The object SDSS 160357.93+140929.97 (hereafter SDSS 16+14) was identified as a magnetic white dwarf in Kepler et al. (2013), with an effective temperature of 10123 K and a magnetic field strength of 43 MG. We monitored this target on May 3rd and June 5th 2015, each time over a time period of 3 h, using the optical camera EFOSC2 at the ESO/NTT (program 095.D-0245). The g-band lightcurves shows a photometric period of ~ 100 min, with a min-max amplitude of $\sim 3\%$ (compared to a photometric noise of 1-2%).¹

We obtained high signal-to-noise lightcurves for SDSS 16+14 using Lowell Observatory’s Discovery Channel Telescope (DCT) and the Large Monolithic Imager (LMI), on June 15th and July 4th 2017. Here we observed in the three filters g, r, i quasi-simultaneously, i.e. switching filters every few minutes. The resulting multi-band lightcurves are shown in Figure 1. This dataset confirms the presence of a photometric period with $P = 110 \pm 3$ min, averaged over the six lightcurves. The amplitudes in g, r, and i are 0.027, 0.021, 0.021 in the first run and 0.030, 0.021, 0.019 in the second run, respectively. Interestingly, the first set of observations shows a clear offset in phase of ~ 0.2 between the r/i-band lightcurves and the one in the g-band. This offset was not observed (or is significantly smaller) in the second run. Both times, the g-band lightcurve is deviating from the sinusoidal variations seen in r- and i-band.

There are two ways to explain this type of variability. The first option is to postulate the presence of a cool, unresolved companion, with the period being the orbital period. The observed period and amplitude fit into the small group of white plus brown dwarf systems in the literature (Casewell et al. 2018). The variability could be produced by reflected light, thermal re-radiation, accretion, or beaming (Maoz et al. 2015). The phaseshift may be caused by more than one of these mechanisms being involved. To validate this hypothesis, we searched for the IR excess caused by the companion, using deep H- and K-band imaging from Subaru/IRCS, in an observing run on July 1-2 2017. The object is clearly detected as a single point source, and we measure magnitudes of $H = 18.1 \pm 0.1$ and $K = 18.3 \pm 0.1$. Within the errorbars, this is consistent with a 10000 K blackbody scaled to the Sloan g-band magnitude of SDSS 16+14 and rules out L-type companions with $T > 1000$ K. Radial velocity monitoring is needed to exclude even cooler companions.

In absence of an IR excess, the favoured explanation is the presence of magnetic spots and a rotational modulation of the flux. In terms of period, amplitude, temperature and magnetic field strength, SDSS 16+14 fits into the sample of magnetic white dwarfs with measured rotation periods by Brinkworth et al. (2013). Simple blackbody spot models,

Corresponding author: Aleks Scholz
as110@st-andrews.ac.uk

¹ As part of the same run we also confirm in broadband light the periods for SDSS 151415.66+074446.50 and SDSS 125044.43+154957.36, both known to have short-period RV companions (Breedt et al. 2012).




Figure 1. DCT/LMI lightcurves in multiple bands for SDSS 16+14.

such as used by [Scholz et al. \(2009\)](#), match the color dependence of the amplitudes, with spot temperature around 5000 K and a filling factor of 2-3%. If this interpretation is correct, SDSS 16+14 is one of the fastest known rotating white dwarfs ([Kawaler 2015](#)) and among the spotted white dwarfs with the strongest magnetic field. The phaseshift observed in one of the DCT runs requires an unusual temperature structure in the spots (e.g., a hot spot and a cool spot offset in longitude). This finding encourages multi-band observations of variable magnetic white dwarfs to elucidate whether or not such spot configurations are common.

We thank Sarah Casewell, Elme Breedt, Philip Lucas, Hiro Takami, and Brendan McDonald for supporting this project. These results made use of Lowell Observatory’s Discovery Channel Telescope, and The Large Monolithic Imager (NSF/AST-1005313).

Facilities: DCT,NTT,Subaru

REFERENCES

- | | |
|---|---|
| <p>Brinkworth, C. S., Burleigh, M. R., Lawrie, K., Marsh, T. R., & Knigge, C. 2013, <i>ApJ</i>, 773, 47</p> <p>Breedt, E., Gänsicke, B. T., Girven, J., et al. 2012, <i>MNRAS</i>, 423, 1437</p> <p>Casewell, S. L., Braker, I. P., Parsons, S. G., et al. 2018, arXiv:1801.07773</p> <p>Kawaler, S. D. 2015, 19th European Workshop on White Dwarfs, 493, 65</p> | <p>Kepler, S. O., Pelisoli, I., Jordan, S., et al. 2013, <i>MNRAS</i>, 429, 2934</p> <p>Maoz, D., Mazeh, T., & McQuillan, A. 2015, <i>MNRAS</i>, 447, 1749</p> <p>Scholz, A., Xu, X., Jayawardhana, R., et al. 2009, <i>MNRAS</i>, 398, 873</p> |
|---|---|