

Kent Academic Repository

Full text document (pdf)

Citation for published version

Fahy, Geraldine and Mahoney, Patrick and Curto, Ana and Maurer, Anne-France and Barrocas-Dias, Christina and Fernandes, Teresa (2018) Did military orders influence the general population diet? Stable isotopes analysis from Medieval Tomar, Portugal. *Archaeological and Anthropological Sciences*. ISSN 1866-9557.

DOI

<https://doi.org/10.1007/s12520-018-0637-3>

Link to record in KAR

<http://kar.kent.ac.uk/66731/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

**Did military orders influence the general population diet?
Stable isotopes analysis from Medieval Tomar, Portugal**

Ana Curto*^{1,2}, Anne-France Maurer³, Cristina Barrocas-Dias^{3,4}, Patrick Mahoney¹, Teresa Fernandes^{2,5}, Geraldine E. Fahy¹

Email: *arqc3@kent.ac.uk

Phone: +351 965805316

ORCID: 0000-0003-3568-9322

¹Human Osteology Lab, Skeletal Biological Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, UK

²Research Centre for Anthropology and Health, Department of Life Sciences, University of Coimbra, Coimbra, Portugal

³HERCULES Laboratory, Évora University, Largo Marquês de Marialva 8, 7000-809 Évora, Portugal

⁴School of Sciences and Technology, Chemistry Department, University of Évora, Rua Romão Ramalho 59, Évora, Portugal

⁵School of Sciences and Technology, Biology Department, University of Évora, Pólo da Mitra, Apartado 94, Évora, Portugal

Abstract

This study integrates bone collagen stable isotope data (carbon, nitrogen and sulphur) from 33 human adult tibiae (15 females; 18 males) and 13 faunal remains from Tomar, while it was under the Military Orders domain (11th – 17th centuries). Historical literature indicates that the amount of meat consumption among Templars was lower than in individuals with similar social status. In medieval times these Military Orders had total control of towns and angling and fishing rights, but their influence on the general population diet remains unknown. While no statistically significant differences ($p>0.05$) were found between sexes, social status, or for bone collagen $\delta^{13}\text{C}$ and $\delta^{34}\text{S}$ between age groups, $\delta^{15}\text{N}$ did differ significantly with age, which may be related to tooth loss in old individuals. Additionally, the human samples have higher stable isotope differences, in comparison to faunal samples, than would be expected within the food web, particularly for $\delta^{13}\text{C}$. This human bone collagen $\delta^{13}\text{C}$ enrichment may reflect a diet rich in aquatic protein intake, which is also supported by $\delta^{34}\text{S}$ archived in human and faunal samples, and the presence of oysters and cockles shells at the excavation. The religious diet restrictions might have led to a higher intake of aquatic protein when meat consumption was not allowed.

Keywords: Europe, Iberian Peninsula, paleodiet, carbon, nitrogen, sulphur

1. Introduction

This study investigates diet in a town ruled by religious military orders and how the general population diet may have adapted to religious dietary restrictions. This study is the first of its kind to analyse carbon, nitrogen and sulphur stable isotope data from skeletons of the 11th – 17th centuries in Portugal.

1.1. Stable isotopic analysis

Analysis of stable isotope ratios from mineralized tissue has been widely used for dietary reconstruction. This technique is based on the assumption that “you are what you eat (plus a few ‰)” (DeNiro and Epstein 1976), as a consumer’s tissues reflect the isotopic array of the ingested foods. Food webs have an impact on carbon isotope values due to the correlation between animal tissues carbon values ($\delta^{13}\text{C}$) and their diet (DeNiro and Epstein 1978; Teeri and Schoeller 1979). There is enrichment in $\delta^{13}\text{C}$ in an animal’s body tissues relative to its diet due to the fractionation that occurs during the formation of tissues (van der Merwe and Vogel 1978). Primary consumers have a fractionation factor (enrichment in $\delta^{13}\text{C}$) of approximately 5‰ in their bone collagen relative to their diet (van der Merwe and Vogel 1978; Ambrose and Norr 1993) and an enrichment of 1‰ between trophic levels (DeNiro and Epstein 1978; Tieszen et al. 1983). In marine plants the main carbon source is dissolved carbonate (0‰), instead of atmospheric CO_2 (-7‰), therefore, this difference is reflected in the $\delta^{13}\text{C}$ in tissues of mammals feeding from these two different ecosystems (Tauber 1981; Chisholm et al. 1982, 1983). $\delta^{13}\text{C}$ in bone collagen can also help identifying freshwater resources. Katzenberg and Weber (1999) observed a range of – 14.2 to – 24.6‰ in fish bones in Siberia with higher $\delta^{13}\text{C}$ in species inhabiting shallow waters and lower $\delta^{13}\text{C}$ for fish inhabiting deeper open waters on the lake. Freshwater fish exhibit variation in $\delta^{13}\text{C}$ depending on the ecosystem as freshwater plants have numerous sources of carbon, unlike terrestrial plants (Zohary et al. 1994; Dufour et al. 1999).

In terrestrial ecosystems there is an increment of 3‰ to 5‰ in $\delta^{15}\text{N}$ between trophic levels when compared with consumer diet (Schoeninger et al. 1983; Minagawa and Wada 1984; Schoeninger and DeNiro 1984; Bocherens and Drucker 2003). This fractionation enables the use of stable nitrogen isotopes ($\delta^{15}\text{N}$) to infer trophic level and high $\delta^{15}\text{N}$ recorded in bone collagen usually indicates high-protein diets (Sponheimer et al. 2003). Nitrogen isotope values can also be used to differentiate between terrestrial and marine food sources (DeNiro and Epstein 1981; Schoeninger et al. 1983; Walker and DeNiro 1986; Richards and Hedges 1999), especially when combined with carbon isotope data. Bone collagen $\delta^{15}\text{N}$ can also be used to analyse access to fresh water resources, as organisms in these ecosystems exhibit higher $\delta^{15}\text{N}$ than those in terrestrial ecosystems (van Klinken et al. 2000).

Advances in mass spectrometry and methodology development, following the work of Leach et al. (1996) allow an easier and more frequent analysis of sulphur isotope data ($\delta^{34}\text{S}$). Sulphur isotope analysis can shed some light on the use of freshwater or marine resources (Nehlich and Richards 2009; Nehlich et al. 2010; Nehlich 2015), especially when combined with the analysis of carbon and nitrogen stable isotopes. A freshwater ecosystem, which is highly depending on the geological conditions and source of water sulphates (Nehlich 2015), has an impact on terrestrial $\delta^{34}\text{S}$, especially if the fauna fed on the floodplains of the river (Fry 2002; Nehlich et al. 2011). $\delta^{34}\text{S}$ at riverine ecosystems fall between -5‰ and +15‰, but the values can be outside this range in relatively small geographical scale due to specific environmental conditions (Nehlich 2015).

1.2. Historic background

The city of Tomar had a very important military role consolidating the Kingdom of Portugal by resisting the advances of the last Moroccan king of Hispania, Iacub ben Iuquf Almançor (França 1994). The construction of the Convent of Christ, a Templar stronghold, began in

1160 and was also likely around that time that the Church of Santa Maria do Olival was constructed (Conde 1996). In 1317 Pope Clement issued the Papal Bull *Pastoralis Praeeminentiae*, which instructed all Christian monarchs in Europe to arrest all Templars and seize their assets (Barber 2012). Portugal successfully lobbied the papacy and the Templars did not face a trial, instead the Order's assets and personnel were transferred to the newly established Order of Christ, a continuation of the Templars in Portugal (Valente 1998). Tomar then became a centre of Portuguese overseas expansion under Henry the Navigator, the Grand Master of the Order of Christ (Conde 1996).

Trade in Europe began to increase in the 11th century (Malgosa 2011), since Tomar was located at the main Portuguese road connecting the North of the country to the limits of the Reconquista (Conde 1996). Given Tomar's location it would have frequent movement of goods but also people and one of its functions was to receive and protect refugees in case of invasion (Conde 1996).

According to historical data, the staple medieval diet in Portugal was bread accompanied by wine, olives and olive oil (Vicente 2013). A significant part of agriculture was focused on cereals but a large percentage of the harvest was inaccessible to peasants after paying tributes to lords and the church (Vicente 2013). Chestnuts and sweet acorns could sometimes substitute the bread (Vicente 2013) and some legumes could be reduced to flour when there was a lack of cereals (Gonçalves 2004). The acorns were frequently used to feed the livestock, especially swine that also fed from various roots and mushrooms (Vicente 2013).

Cattle were not abundant, compared to sheep and goats, and only the pigs were purposely raised for meat production (Gonçalves 2004). Other sources of meat were chicken, duck and goose as well as a variety of game (Gonçalves 2004). For the peasants, hunting could represent the only access to meat; however, in Tomar angling and warren rights were

reserved for the military orders. Among medieval Iberian faunal assemblages the domestic animals predominate (Grau-Sologestoa 2017), which can be a result of hunt restrictions. Fish was an expensive food, with the exception of sardines which were more abundant and easy to preserve salted or smoked (Gonçalves 2004). Fish was indispensable during the numerous fast days that the medieval religious calendar imposed (Vicente 2013) but it was consumed more in the littoral despite the availability of Portuguese rivers (Gonçalves 2004). Molluscs and crustaceans were also part of the diet of all social status but were considered a “food of the poor” due to their abundance (Gonçalves 2004).

Various studies suggest dietary differences between sex, age groups and social status in medieval times (e.g. Adamson 2004, Kjellström et al. 2009, Linderholm et al. 2008, Polet and Katzenberg 2003, Schutkowski et al. 1999, Reitsema et al. 2010, Reitsema and Vercellotti 2012). Since fish was expensive but necessary for religious fasts and the military orders had angling and warren rights, diet in Tomar may also reflect social status. The historical literature (Barber and Bate 2002) implies that the amount of meat consumption among Templars was lower than in individuals with similar social status, and the intake of vegetables was higher. In Tomar, merchants, crafters and farmers participated actively at the local army alongside with knights, raising their status (Conde, 1996) and probably having access to similar food resources to the Templars.

2. Materials and sampling

This study analyses bone collagen stable isotope data from 33 human adult tibiae (15 females; 18 males) and 13 fauna remains (2 wild Sus; 2 domestic Sus; 1 juvenile Sus; 1 Canidae; 3 Bos; 1 Equus; 3 Ovicapridae) from Santa Maria do Olival graveyard (11th – 17th centuries), in Tomar, Portugal. Only individuals from areas 13 to 20 (2nd phase of the excavation; Annex, Fig. A1) were analysed. Areas 13, 15, 18 and 19 were considered to be a place of burial for individuals with higher social status, not only due to the proximity to the

church (Binski 1996; Daniell 1998; Graves 1989; Ottaway1992; Platt 1981; Swanson 1989) but also because of the higher frequency of structured graves. Faunal remains were collected from areas 14,17 and 20 (Annex, Fig. A1). The faunal remains from area 20 were mixed with human remains in an ossuary with at least 14 human adults. The faunal remains recovered from areas 14 and 17 were in grave fill material.

The skeletons (all ages and both sexes) distribution within the necropolis suggest that Santa Maria do Olival collection represents the general population of Tomar and not, or at least not only, the individuals from the military orders. The uniform spatial distribution between sexes within the graveyard and the use of structured graves for both males and females suggest that, at least at death, social status was not dependent on sex. However, social status seems to increase with age as older individuals were more frequently buried in structured graves.

Only individuals without signs of physiological stress were sampled in an attempt to estimate the diet of the general population and avoid isotopic data that may represent differing metabolism during disease and/or malnutrition (Steele and Daniel 1978; Hobson and Clark 1992; Hobson et al. 1993; Gaye-Siessegger et al. 2004; Fuller et al. 2005; D'Ortenzio et al. 2015). To avoid sampling individuals with physiological stress only individuals without skeletal markers of stress, such as cribra orbitalia or obvious enamel hypoplasias, were selected. Since low stature can also be associated with physiological stress (e.g. Haviland 1967; Morris and McAlpin 1979; Allen and Uauy 1994; Roberts and Manchester 2007; Moore and Ross 2013), only individuals with maximum length of the skeleton (measurement was taken during the excavation while the skeleton was still articulated, in situ, in extended supine position and used as a proxy for stature) equal or above the mean for this population ($151.8\pm 6.1\text{cm}$ for the females, $n=256$; $163.4\pm 7.5\text{cm}$ for the males, $n=287$) were sampled.

3. Methods

Sex was estimated based on pelvic (Phenice 1969; Buikstra and Ubelaker 1994) and cranial features (Buikstra and Ubelaker 1994). Adult age at death estimates employed a combination of skeleton maturation (Scheuer and Black 2000), pubic symphysis degeneration (Brooks and Suchey 1990; Buikstra and Ubelaker 1994) and auricular surface degeneration (Lovejoy et al. 1985). The skeletons analysed were classified as young (18 to 29 years), mature (30 to 60 years) and old (more than 60 years) adults.

Collagen extraction was done following Longin (1971), Brown et al. (1988) and Richards and Hedges (1999). The collagen samples were weighed into tin capsules and combusted into CO₂ and N₂ using an Elemental analyzer (Flash/EA) coupled to a Thermo Finnigan Delta^{Plus} XL isotope ratio mass spectrometer via a ConFlo III interface at NERC Isotope Geosciences Facility (Nottingham, UK). Sulphur stable isotopes were analysed at the Faculdade de Ciências da Universidade de Lisboa (Lisbon, Portugal). $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were calibrated using an in-house reference material M1360p (powdered gelatine from British Drug Houses) with expected δ values of -20.32‰ (calibrated against CH₄, IAEA) and $+8.12\text{‰}$ (calibrated against N-1 and N-2, IAEA) for carbon and nitrogen respectively. Samples were run in duplicate and the 1σ reproducibility for mass spectrometry controls for these analyses were $\delta^{15}\text{N} = \pm 0.08\text{‰}$ and $\delta^{13}\text{C} = \pm 0.07\text{‰}$. The sulphur isotope analysis was done at SIIAF (University of Lisbon), using an IsoPrime mass spectrometer. The collagen was combusted with additional V₂O₅ and a pulse of oxygen. $\delta^{34}\text{S}$ was calibrated using the inorganic international standards NBS127 ($+20.3\text{‰}$), IAEA S1 (-0.3‰) and casein protein ($+4.0\text{‰}$). Mass spectrometry control for these analyses was $\delta^{34}\text{S} = \pm 0.08\text{‰}$.

Mann-Whitney U non-parametric tests were used for pair-wise comparisons and Kruskal-Wallis non-parametric tests were used to compare more than two groups. All

statistics were computed in SPSS 24 for Windows and $p \leq 0.05$ were considered statistically significant.

4. Results

The bones from all individuals in the present study had acceptable C:N ratios (2.9 to 3.6; DeNiro 1985) and S% (0.15% to 0.35%, Nehlich and Richards 2009). Herbivores, with the exception of *Equus*, have similar values for bone collagen $\delta^{13}\text{C}$ (-21.2‰ to -20.9‰), while bone collagen $\delta^{15}\text{N}$ (4.8‰ to 7.8‰) and $\delta^{34}\text{S}$ (13.1‰ to 18.5‰) values are more variable (Figures 1 and 2). The domestic *Sus* and the only carnivore analysed (*Canidae*) have similar $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values to the herbivores (Figure 1). The faunal remains have higher bone collagen $\delta^{34}\text{S}$ than the human remains (Figure 2), only the *Equus* displays bone collagen $\delta^{34}\text{S}$ expected for an exclusive terrestrial diet.

Among the humans sampled there is an outlier, a male young adult. While his bone collagen $\delta^{15}\text{N}$ (12.3‰) are amongst the highest, his bone collagen $\delta^{13}\text{C}$ (-15.4‰) is highly enriched (Figure 1) and he displays bone collagen $\delta^{34}\text{S}$ depleted (9.3‰) compared to the other individuals (Figure 2). Overall, bone collagen $\delta^{15}\text{N}$ and $\delta^{34}\text{S}$ recorded in the humans are more variable than their bone collagen $\delta^{13}\text{C}$ (Table 1). The females show higher variance in their bone collagen $\delta^{15}\text{N}$, while the males display a higher variance in their bone collagen $\delta^{34}\text{S}$ (Table 1). There are no statistically significant differences ($p > 0.05$) in stable isotope values between sexes, social status, or for $\delta^{13}\text{C}$ and $\delta^{34}\text{S}$ between age groups. However, bone collagen $\delta^{15}\text{N}$ recorded in the human skeletons display significant differences ($p = 0.05$) with age groups (Table 2).

The individuals for which it was possible to estimate sex and age are represented in figure 3 illustrating $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ differences in bone collagen between age groups. While the young adults are above or close to the mean values for $\delta^{13}\text{C}$ (-18.6‰) and $\delta^{15}\text{N}$ (10.8‰), the old adults are all under the mean values for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ (with the exception of

one), but most fall within two standard deviations from the mean. There are no differences in bone collagen $\delta^{34}\text{S}$ between the age groups (Figure 4).

5. Discussion

5.1. General diet at Tomar

There might be dietary differences between different chronologies, especially after the 16th century with the introduction of new food sources, like C₄ plants from America, but unfortunately it was not possible to decrease the chronological interval estimated for Tomar (11th – 17th centuries). The high density of burials and the fact that Christian burials usually do not have associated artefacts did not allow reliable dating.

Herbivores' $\delta^{13}\text{C}$ (-21.3‰ to -20.1‰, Figure 1, Annex: Table A1) suggest a diet based on C₃ plants (Vogel 1978; Schoeninger and DeNiro 1984; Chisholm 1989). Despite the wide range of the estimated chronology for Tomar's necropolis (11th to 17th centuries) and the possibility that the analysed fauna represents different times (areas 14, 17 and 20), the herbivores' $\delta^{13}\text{C}$ are similar, arguing against the introduction of new food sources like maize (C₄ plants). In contrast, bone collagen $\delta^{15}\text{N}$ recorded in herbivores are more variable (4.8‰ to 7.9‰, Figure 1) and with some enrichment, particularly observed for the Ovicapridae. Enrichment in faunal bone collagen $\delta^{15}\text{N}$ may be related to variable animal husbandry practices and land management. Manured soils raise $\delta^{15}\text{N}$ in soil and plants (van Klinken et al. 2000; Bogaard et al. 2007), having an impact on the local food web. $\delta^{15}\text{N}$ enrichment is particularly evident between the Ovicapridae, which may be related to different food sources for sheep (grass; hay) and goats (bushes; tree leaves/bark). The Ovicapridae have higher $\delta^{15}\text{N}$ than Bos. Higher $\delta^{15}\text{N}$ in Ovicapridae compared to Bos are also observed in faunal remains (Annex, FigA2) from Koksijde (Polet and Katzenberg 2003) but not from Benipeixcar (Alexander et al. 2015). The domestic and wild Sus have similar $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ to the herbivores, as well as the only carnivore analysed (Canidae), suggesting a diet poor in animal

protein. While pigs are frequently kept as herbivores (e.g. Quirós Castillo 2013), dog isotopic ratios usually cluster with the humans (e.g. Haffman and Velemínský 2015; Quirós Castillo 2013; Lubritto et al. 2013). As dogs frequently eat food scraps, their isotope values can be indicative of a human diet poor in animal protein.

The mean increase from faunal (except the Equus and Canidae) to human remains is 2.3‰ (in some individuals more than 3‰, Figure 1) for $\delta^{13}\text{C}$ and 4.9‰ for the $\delta^{15}\text{N}$. Some individuals have higher enrichment than would be expected within the food web: up to 1‰ for $\delta^{13}\text{C}$ (Schoeninger et al. 1983; Minagawa and Wada 1984; Schoeninger and DeNiro 1984) and 3‰ to 5.7‰ for $\delta^{15}\text{N}$ (van der Merwe and Vogel 1978; Ambrose and Norr 1993). The trophic level increase expected based on the faunal isotope values would be between 7.8‰ and 13.6‰ for $\delta^{15}\text{N}$ and from -21.3‰ to -19.9‰ for $\delta^{13}\text{C}$ (grey area at Figure 1). While human $\delta^{15}\text{N}$ at Tomar range between 9.0‰ and 12.5‰ and can be explained by the trophic increment, their $\delta^{13}\text{C}$ vary between -19.4‰ and -17.3‰, being clearly enriched compared to the faunal remains recovered in Tomar. This $\delta^{13}\text{C}$ enrichment observed in the human remains can reflect a diet complemented by some aquatic (Tauber 1981; Chisholm et al. 1982, 1983) or C_4 plants (Vogel et al. 1978; Chisholm 1989) intake. Due to the presence of oysters and cockles shells at the excavation (areas 14 and 17) and that the analysed fauna were feeding on C_3 plants it is more probable that the $\delta^{13}\text{C}$ enrichment in the human bone collagen is related to aquatic protein than C_4 plants intake.

To better understand diet at Tomar, $\delta^{34}\text{S}$ were also analysed for some faunal and human remains. Surprisingly, the fauna $\delta^{34}\text{S}$ are higher than would be expected for terrestrial animals ($\delta^{34}\text{S} > 12\text{‰}$; Nehlich 2015) and correspond to values expected from coastal fauna influenced by marine sea spray (Nehlich 2015). However, Tomar is located at approximately 70km from the coast and sea spray sulphates only reach up to 30km inland (Wakshal and Nielsen 1982). Tide floods from the Atlantic Ocean increase the Tagus River flow and its

salinity, reaching the floodplains near Santarém (Figure 5) and increasing the sea spray reach but not enough to justify the $\delta^{34}\text{S}$ registered for the fauna on its own.

Riverine sulphates can also be found on the riverbanks and floodplains, influencing the isotopic composition of the surrounding landscape (Fry 2002; Nehlich et al. 2011) and the values observed for Tomar's fauna may be related to the livestock feeding on the floodplains. However, floodplains tend to have lower $\delta^{34}\text{S}$ than areas further away from freshwater ecosystems (Nehlich et al. 2011). Therefore it is possible that the use of algae as a fertilizer may have increased the $\delta^{34}\text{S}$ in the food web as fresh seaweed can also be used to feed livestock, mostly ruminants and pigs (Chapman and Chapman 1980). In Portugal, algae has been used in agriculture previously to the 14th century (Veiga de Oliveira et al. 1975; Vieira and Santos 1995) but it would likely be restricted to coastal areas as algae are heavy and usually not carried very far inland (McHugh 2003). Even though the seaweed could be sundried and stored to be used as winter feedstuff for sheep and cattle (Evans and Critchley 2014) it would probably not be taken so far inland.

$\delta^{34}\text{S}$ vary not only by dietary behaviour (Richards et al. 2001) but also by location (Hobson, 1999), ranging from -40‰ to +40‰ in terrestrial rocks (Nielsen et al. 1991) and between -20‰ and +20‰ in terrestrial organic matter (Peterson and Fry 1987). The oxidation of sulphides and organic sulphur by microorganisms in the soils can also result in high $\delta^{34}\text{S}$ and therefore influence the food web (Böttcher et al. 1998; Nehlich et al. 2011). Therefore it is possible that the higher $\delta^{34}\text{S}$ observed in the terrestrial fauna from Tomar may be related to the geochemistry of that area and not with agricultural or husbandry practices. Tomar is located at an area with evaporites, gypsum and marl (yellow area at Figure 5) that would increase the $\delta^{34}\text{S}$ in the food webs of this region. The Equus, with the lowest $\delta^{34}\text{S}$ (10.2‰, Figure 2), supports this hypothesis, as it was more mobile than the other domestic animals. The wild Sus also have lower $\delta^{34}\text{S}$ (14.6‰ and 14.8‰), which can also be related

with a higher mobility. Interestingly, the human collagen from Tomar has lower $\delta^{34}\text{S}$ (9.3‰ to 15.6‰) than the faunal remains (13.1‰ to 18.5‰), suggesting that those terrestrial animals were not frequently consumed by the local population, who could have relied on other food sources from another geographical area with lower $\delta^{34}\text{S}$ in its geo-ecosystem.

$\delta^{34}\text{S}$ at riverine ecosystems usually fall between -5‰ and +15‰ (Nehlich 2015), but values can fall outside this range, depending on the geological surroundings of the river basin, ultimately influencing the $\delta^{34}\text{S}$ of the river fauna (Nehlich 2010, 2011). Unfortunately, fish bones were not recovered from Tomar's excavation to confirm the values of the fish consumed by this population. If fresh water protein intake was important and with low $\delta^{34}\text{S}$, it could decrease the high $\delta^{34}\text{S}$ within the surroundings of Tomar, related to its particular geological context. However, Nabão River, that crosses Tomar, is also located within the same geological substrate and thus, $\delta^{34}\text{S}$ within its food webs are probably high. Zêzere River is located at approximately 10km from Tomar rising at Serra da Estrela (a granitic and metamorphic mountain range) and meeting the Tagus River (an international river) at about 15km from Tomar (Figure 5). Since Zêzere and Tagus rivers do not pass through an area with evaporites and gypsum, $\delta^{34}\text{S}$ of their food webs are probably lower than those at Nabão River. Human bone collagen $\delta^{34}\text{S}$ suggest that if they were eating fresh water protein it was probably coming from Zêzere and/or Tagus. Besides, their larger dimensions could offer more food sources than Nabão River. The presence of shells at the excavation suggests also some marine protein intake. Nazaré is the closest coastal town where today fish and octopus are still sundried at the beach, this way of preserving the fish might have allowed its consumption further inland, in towns like Tomar, alongside with fish from the surrounding rivers.

The lower $\delta^{34}\text{S}$ registered in human bone collagen can also be related with terrestrial intake from a geographical location with lower $\delta^{34}\text{S}$ in its food webs. Since the staple

medieval diet in Portugal was bread (Vicente 2013) it is possible that it was being made with flour from cereals grown in a location different from Tomar's surroundings. If bread, made with cereals with low $\delta^{34}\text{S}$, was consumed in high quantities, it could also have lowered the $\delta^{34}\text{S}$ of individuals, independently of the geological substrate in the surroundings of Tomar. The possibility of C_4 plants being consumed only by humans cannot be excluded. It could have been entering their diet in the form of maize flour, for example, and if the maize was not cultivated in Tomar, it could explain both the lower $\delta^{34}\text{S}$ and the higher $\delta^{13}\text{C}$ recorded in human bone collagen. However, the negative relation between $\delta^{34}\text{S}$ and $\delta^{15}\text{N}$ (Figure 4) likely indicates that the higher $\delta^{15}\text{N}$ are related with protein with lower $\delta^{34}\text{S}$ and therefore the high $\delta^{15}\text{N}$ represent protein from fresh water rather than from terrestrial fauna.

As Tomar was ruled by the Order of the Temple and later the Order of Christ (Vicente 2013) it is possible that religious dietary restrictions would be reflected in Tomar's population. Also, in Tomar, merchants, crafters and farmers participated actively at local army levels alongside with knights, raising their status (Conde 1996) and probably giving them access to similar food resources. Müldner et al (2009) found isotopically distinct diets between bishops and the general population in Scotland, the latter having higher fish intake, related to religious fasting. These dietary restrictions may have led to a higher intake of aquatic protein when meat consumption was not allowed with towns controlled by military orders likely being under increased pressure to follow religious dietary restrictions. More isotopic data from different places with similar chronologies is necessary to understand if the high intake of aquatic protein is due to the presence of the military orders at Tomar or if it was a generalised religious phenomenon.

Human diet at medieval Tomar was complex and likely included food sources from outside Tomar. The general diet was poor in terrestrial protein and rich in fresh water protein with possible terrestrial protein from other geographical locations.

5.2. Dietary differences within Tomar

Even though some historical (e.g. Adamson 2004) and anthropological (e.g. Kjellström et al. 2009; Linderholm et al. 2008; Polet and Katzenberg 2003; Schutkowski et al. 1999; Reitsema et al. 2010; Reitsema and Vercellotti 2012) sources suggest different food access based on age, sex and status in medieval times, that was not observed in Tomar. When the skeletons sampled were grouped by sex, age or inferred social status only $\delta^{15}\text{N}$ in the age groups was statistically different ($p < 0.05$; Table 2).

The bone collagen of young adults display higher $\delta^{15}\text{N}$ than the old adults (Figure 4) suggesting a higher animal protein intake for the young individuals. Since only skeletons without signs of physiological stress were sampled the higher $\delta^{15}\text{N}$ for the young adults is not related to chronic stress (Steele and Daniel 1978; Hobson et al. 1993; Gaye-Siessegger et al. 2004; Fuller et al. 2005; Deschner et al. 2012; D'Ortenzio et al. 2015) that might have resulted in premature death, due to ill health (Wood et al. 1992). These isotopic differences between young and old adults may be related to severe tooth loss that was observed in old individuals who therefore may have had increased difficulty ingesting some foods along with changes associated with metabolism in the aging, such as reductions in taste, smell and hunger, and delayed rate of absorption (Roberts and Rosenberg 2006). The amount of fresh water fish is variable and not related to sex, social status or age (Figure 4). Overall, the skeletons analysed had similar diets with smaller $\delta^{34}\text{S}$ differences compared to other European samples (e.g. Nehlich et al. 2011), despite the wide chronology estimated for Tomar's necropolis (11th – 17th century). There are no $\delta^{34}\text{S}$ differences between age, sex or social status but there could be dietary differences between chronologies. Unfortunately it was not possible to date the faunal or human remains.

The absence of statistically significant isotopic differences between sexes (Table 2) suggests that males and females had similar protein intakes at Tomar, however, sample sizes

may be too small and dispersed (sex, age, social status and chronology) to detect significant differences. The uniform spatial distribution for males and females within the graveyard and the use of structured graves for both sexes also suggest that, at least at death, social status was not dependent on sex though medieval society was male-dominant.

The only outlier analysed, a young adult male, has higher values of both $\delta^{15}\text{N}$ (12.3‰) and $\delta^{13}\text{C}$ (-15.4‰) and low $\delta^{34}\text{S}$ (9.3‰). The low $\delta^{34}\text{S}$ suggest that this individual might be an outsider, coming from a place with lower $\delta^{34}\text{S}$ in its ecosystem, but the possibility of these isotopes values being the result of a high fresh water protein intake from low $\delta^{34}\text{S}$ cannot be excluded. The high $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ can also represent a terrestrial diet rich in C_4 plants, directly or fed to the livestock, particularly if this individual was from a different geographical location.

Food can reflect social status and define social, cultural and religious boundaries (e.g. Thomas 2007; Curet and Pestle 2010), however this was not observed within the samples analysed. The $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\delta^{34}\text{S}$ distribution is uniform when comparing individuals with higher and lower social status. Individuals from lower social status may be more Susceptible to physiological stress, particularly due to malnutrition (e.g. Weston 2012). Since only individuals without skeletal signs of physiological stress (low stature, criba orbitalia, porotic hyperostosis) were sampled, the ones with lower social status could have been avoided. Adult stature is determined by genetics but also has an environmental determinant (e.g. Haviland 1967; Larsen 1997; Bogin 1999; Cardoso and Gomes 2009). The areas further away from the church (1st phase of the excavation; Figure 1) better represent the individuals from the lower social status and the ones buried inside the church would be a better example of the people from higher social status. Therefore the individuals analysed probably represent the average population and neither of the social extremes. More isotopic data from different social and

health status would help understand if the diet at Tomar was uniform or if our results were biased by selecting only apparently healthy individuals.

5.3. Other European studies

Comparing the data with other late medieval European samples (Figure 6), those with similar $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ averages are Koksijde (from a coastal Belgian monastery; Polet and Katzenberg 2003) and Rome (from an Italian mass grave; Salamon et al. 2008). Contrary to what would be expected the stable isotope values from Tomar are closer to the Belgian sample (Polet and Katzenberg 2003) than to the other Iberian samples (Lubritto et al. 2013; Alexander et al. 2015), which may be related to religious dietary requirements, particularly low meat consumption, as the Belgian sample represents a monastic community (Polet and Katzenberg 2003). The similar faunal values for Tomar and Koksijde (Annex, FigA2) allow a comparison between the two locations, despite their geographical and social differences. The impact of religious directives of the Catholic Church on the diet has been registered before (Salamon et al. 2008). This was facilitated by industrial-scale fishing in the Atlantic (Barret et al. 2004) and improvement of food preservation methods (Heinrich 1986; Robinson 2000). Müldner and Richards (2007) also associated the increased intake of aquatic protein (mostly marine fish with some freshwater fish or molluscs) at St. Andrew (Figure 6) with religious dietary habits in Later Middle Ages. Agricultural and husbandry practices used during the Middle Ages may also explain the different isotopic values between the medieval skeletons buried in Tomar and the prehistoric ones, alongside with a higher aquatic protein intake (Figure 6).

Out of the Iberian samples compared, Tomar has the highest $\delta^{15}\text{N}$ mean, particularly when compared to Zaballa (Lubritto et al. 2013), Treviño (Quirós Castillo 2013) and Zornoztegi (Quirós Castillo 2013). The high $\delta^{15}\text{N}$ mean can represent high animal protein intake, however, $\delta^{34}\text{S}$ suggests a high aquatic protein intake which can also be related with

higher $\delta^{15}\text{N}$. The faunal remains recovered from Zaballa (Lubrito et al. 2013), Treviño (Quirós Castillo 2013) and Zornoztegi (Quirós Castell, 2013) also have lower $\delta^{15}\text{N}$ when compared with the ones from Tomar. Colegiata St. Maria (Alexander et al. 2015) and Benipeixcar (Alexander et al. 2015) have similar $\delta^{15}\text{N}$ mean to Tomar's but higher $\delta^{13}\text{C}$, which the authors relate to C_4 plants consumption (directly or fed to domestic animals) or marine fish intake. It is also important to note the different locations of the Spanish collections. While Zaballa, Treviño and Zornoztegi are located at the Basque Country, Northeast of Spain, at approximately 90km to the North Atlantic Ocean; the collections from Colegiata St. Maria and Benipeixcar are from Catalonia, South East of Spain, and at approximately 5km to the Mediterranean Sea. The different locations of these collections may explain why Colegiata St. Maria and Benipeixcar may have higher aquatic protein intake than Zaballa, Treviño and Zornoztegi. Tomar's population has closer mean $\delta^{15}\text{N}$ to the ones closer to the Mediterranean Sea, suggesting also a higher intake of aquatic protein, while the different $\delta^{13}\text{C}$ may be related to C_4 plants consumption at Colegiata St. Maria and Benipeixcar (Salamon et al. 2008).

6. Conclusion

This study is part of a larger project comparing stable isotopic data from individuals without skeletal lesions compatible with diseases and/or physiological stress (presented here) and those with signs of infectious diseases. Since skeletons with lesions were not analysed, this study might better represent the diet at Tomar, instead of metabolic changes during physiological stress. The bone collagen stable isotope values ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and $\delta^{34}\text{S}$) suggest that individuals in Tomar had a complex diet, low in terrestrial animal protein and high in aquatic protein intake, despite its inland location, which could be related to the presence of the military orders in the town and more strict religious dietary restrictions. Dietary differences between sex or social status were not observed for the population of Tomar, but

the quantity of aquatic protein intake is variable, with $\delta^{34}\text{S}$ ranging from 11.4‰ to 15.6‰ (excluding the outlier). Diet appears to be very diverse in Medieval Iberia. Isotopic data from more archaeological sites are necessary to better understand how diet represents social, religious and economic factors, as well as increase our knowledge of trade, agricultural and husbandry practices in medieval times. Data from archaeological sites near Tomar would also help understanding the impact of the presence of religious orders on a town's general population.

Acknowledgments

The authors would like to thank all the anthropologists and archaeologists involved at the excavation of Tomar's, in particular to Sérgio Simões Pereira, Ricardo Ávila Ribeiro, Sónia Ferro, Cláudia Santos and Helena Santos for the maps and information about the excavation. We would like to thank Dr Angela Lamb for her support at NERC facilities. We are also thankful to Rodrigo Maia for the sulphur isotope analysis, Prof José Mirão for his help understanding Tomar's geochemistry and Dr Ana Cruz for allowing us to compare our data with unpublished isotopic data from pre-historic Tomar (Abrigo do Morgado Superior). A Curto acknowledge the financial support of the University of Kent through the University of Kent 50th Anniversary Scholarship. A-F Maurrer and C. Barrocas-Dias acknowledge the financial support of FCT throughout project "HERCULES Laboratory-Cultural Heritage Studies and Safeguard with reference UID/Multi/04449/2013.

Bibliography

Alexander MM, Gerrard CM, Gutiérrez A, Millard AR (2015) Diet, society, and economy in late medieval Spain: Stable isotope evidence from Muslims and Christians from Gandía, Valencia. *American Journal of Physical Anthropology* 156(2):263-273.

Allen LH, Uauy R (1994) Guidelines for the study of mechanisms involved in the prevention or reversal of linear growth retardation in developing countries. *European Journal of Clinical Nutrition* 48 Suppl 1:S212-6.

Ambrose SH., Norr L (1993) Experimental evidence for the relationship of the carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate. *Prehistoric human bone*, pp. 1-37 Springer.

Barber M (2012) *The new knighthood: A history of the order of the temple* Cambridge University Press.

Barber M, Bate K (2002) *The Templars: Selected sources* Manchester University Press.

Binski P (1996) In O'Connor M. (Ed.), *Medieval death: Ritual and representation*. London: British Museum Press.

Bocherens H, Drucker D (2003) Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: Case studies from recent and ancient terrestrial ecosystems. *International Journal of Osteoarchaeology* 13:(1-2), 46-53.

Bogaard A, Heaton TH, Poulton P, Merbach I (2007) The impact of manuring on nitrogen isotope ratios in cereals: Archaeological implications for reconstruction of diet and crop management practices. *Journal of Archaeological Science* 34(3):335-343.

Bogin B (1999) *Patterns of human growth* Cambridge University Press.

Böttcher ME, Oelschläger B, Höpner T, Brumsack HJ, Rullkötter J (1998). Sulfate reduction related to the early diagenetic degradation of organic matter and "black spot"

formation in tidal sandflats of the German Wadden Sea (southern North Sea): stable isotope (^{13}C , ^{34}S , ^{18}O) and other geochemical results. *Organic Geochemistry* 29(5):1517-1530.

Brooks S, Suchey, JM. (1990) Skeletal age determination based on the os pubis: A comparison of the Acsádi-Nemeskéri and Suchey-Brooks methods. *Human Evolution* 5(3):227-238.

Brown TA, Nelson DE, Vogel JS, Southon, JR (1988) Improved collagen extraction by modified Longin method. *Radiocarbon* 30(2): 171-177.

Buikstra JE, Ubelaker DH (1994) Standards for data collection from human skeletal remains.

Cardoso H, Gomes J (2009) Trends in adult stature of peoples who inhabited the modern Portuguese territory from the mesolithic to the late 20th century. *International Journal of Osteoarchaeology* 19(6): 711-725.

Chapman VJ, Chapman DJ (1980) *Seaweeds and their uses*. Chapman and Hall, London.

Chisholm BS (1989) Variation in diet reconstructions based on stable carbon isotopic evidence. *The Chemistry of Prehistoric Human Bone* 10-37.

Chisholm BS, Nelson DE, Hobson KA, Schwarcz HP, Knyf M (1983) Carbon isotope measurement techniques for bone collagen: Notes for the archaeologist. *Journal of Archaeological Science* 10(4): 355-360.

Chisholm BS, Nelson DE, Schwarcz HP (1982) Stable-carbon isotope ratios as a measure of marine versus terrestrial protein in ancient diets. *Science (New York, N.Y.)*, 216(4550): 1131-1132.

Conde MAS (1996) *Tomar medieval. O espaço e os homens*. Patrimonia, Cascais.

Curet LA, Pestle WJ (2010) Identifying high-status foods in the archeological record. *Journal of Anthropological Archaeology* 29(4):413-431.

Daniell C (1998) In Dawsonera (Ed.), Death and burial in medieval England, 1066-1550. London: Routledge.

DeNiro MJ (1985) Postmortem preservation and alteration of in vivo bone collagen isotope ratios in relation to palaeodietary reconstruction. *Nature* 317(6040):806-809.

DeNiro MJ, Epstein S (1976) You are what you eat (plus a few‰): The carbon isotope cycle in food chains. *Geological Society of America* 6: 834.

DeNiro MJ, Epstein S (1978) Influence of diet on the distribution of carbon isotopes in animals. *Geochimica Et Cosmochimica Acta* 42(5):495-506.

DeNiro MJ, Epstein S (1981) Influence of diet on the distribution of nitrogen isotopes in animals. *Geochimica Et Cosmochimica Acta* 45(3): 341-351.

Deschner T, Fuller BT, Oelze VM, Boesch C, Hublin J, Mundry R, Hohmann G (2012) Identification of energy consumption and nutritional stress by isotopic and elemental analysis of urine in bonobos (*pan paniscus*). *Rapid Communications in Mass Spectrometry* 26(1):69-77.

D'Ortenzio L, Brickley M, Schwarcz H, Prowse T (2015) You are not what you eat during physiological stress: Isotopic evaluation of human hair. *American Journal of Physical Anthropology* 157(3): 374-388.

Dufour E, Bocherens H, Mariotti A (1999) Palaeodietary implications of isotopic variability in Eurasian lacustrine fish. *Journal of Archaeological Science* 26(6):617-627.

Evans FD, Critchley AT (2014) Seaweeds for animal production use. *Journal of applied phycology* 26(2):891-899.

França JA (1994) *Cidades e Vilas de Portugal: Tomar, Vol. 18*. Lisboa, Editorial Presença.

Fry B. (2002) Conservative mixing of stable isotopes across estuarine salinity gradients: a conceptual framework for monitoring watershed influences on downstream fisheries production. *Estuaries and Coasts* 25(2):264-271.

Fuller BT, Fuller JL, Sage NE, Harris DA, O'Connell TC, Hedges RE (2005) Nitrogen balance and $\delta^{15}\text{N}$: Why you're not what you eat during nutritional stress. *Rapid Communications in Mass Spectrometry* 19(18):2497-2506.

Gaye-Siessegger J, Focken U, Abel H, Becker K (2004) Individual protein balance strongly influences $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values in Nile Tilapia, *oreochromisniloticus*. *Naturwissenschaften* 91(2):90-93.

Gonçalves I (2004) Entre a abundância e a miséria: as práticas alimentares da Idade Média portuguesa. *Estudos medievais. Quotidiano Medieval: imaginário, representação e práticas*. Lisboa: Livros Horizonte.

Grau-Sologestoa I (2017) Socio-economic status and religious identity in medieval Iberia: The zooarchaeological evidence. *Environmental Archaeology* 22(2):189-199.

Graves, C. (1989). Social space in the English medieval parish-church. *Economy and Society* 18(3):297-322.

Grumett D, Muers R (2010) *Theology on the menu: asceticism, meat and Christian diet*. Routledge.

Haviland WA (1967) Stature at Tikal, Guatemala: Implications for ancient Maya demography and social organization. *American Antiquity* 316-325.

Hobson KA (1999) Tracing origins and migration of wildlife using stable isotopes: a review. *Oecologia* 120(3):314-326.

Hobson KA, Alisauskas RT, Clark RG (1993) Stable-nitrogen isotope enrichment in avian tissues due to fasting and nutritional stress: Implications for isotopic analyses of diet. *Condor* 388-394.

Hobson KA, Clark RG (1992) Assessing avian diets using stable isotopes I: Turnover of ^{13}C in tissues. *Condor* 181-188.

Katzenberg MA, Weber A (1999) Stable isotope ecology and palaeodiet in the lake Baikal region of Siberia. *Journal of Archaeological Science* 26(6):651-659.

Kjellström A, Storå J, Possnert G, Linderholm A (2009) Dietary patterns and social structures in medieval Sigtuna, Sweden, as reflected in stable isotope values in human skeletal remains. *Journal of Archaeological Science* 36(12):2689-2699.

Larsen C (1997) *Bioarchaeology: Interpreting behavior from the human skeleton*. New York: Cambridge University Press.

Leach BF, Quinn CJ, Lyon GL (1996) A stochastic approach to the reconstruction of prehistoric human diet in the Pacific region from bone isotope signatures. *Tuhinga* 1-54.

Linderholm A, Jonson CH, Svensk O, Liden K (2008). Diet and status in Birka: stable isotopes and grave goods compared. *Antiquity* 82:446-461.

Longin R (1971) New method of collagen extraction for radiocarbon dating. *Nature*, 230(5291):241-242.

Lovejoy CO, Meindl RS, Pryzbeck TR, Mensforth RP (1985) Chronological metamorphosis of the auricular surface of the ilium: A new method for the determination of adult skeletal age at death. *American Journal of Physical Anthropology*, 68(1):15-28.

Lubritto C, Sirignano C, Ricci P, Passariello I, Castillo JQ (2013) Radiocarbon chronology and paleodiet studies on the medieval rural site of Zaballa (Spain): Preliminary insights into the social archaeology of the site. *Radiocarbon* 55(3):1222-1232.

McHugh DJ (2003) *A guide to the seaweed industry*. Rome: Food and Agriculture Organization of the United Nations.

Minagawa M, Wada E (1984) Stepwise enrichment of ^{15}N along food chains: Further evidence and the relation between $\delta^{15}\text{N}$ and animal age. *GeochimicaEtCosmochimicaActa* 48(5):1135-1140.

Moore M., Ross A (2013) Stature estimation. *Research Methods in Human Skeletal Biology*. Waltham: Elsevier 151-180.

Morris D, McAlpin M (1979) *Measuring the condition of the world's poor* Pergamons Press.

Müldner G, Richards MP (2007) Diet and diversity at later medieval Fishergate: the isotopic evidence. *American Journal of Physical Anthropology* 134(2):162-174.

Müldner G, Montgomery J, Cook G, Ellam R, Gledhill A, Lowe C (2009) Isotopes and individuals: diet and mobility among the medieval Bishops of Whithorn. *Antiquity* 83(322):1119-1133.

Nehlich O (2015) The application of sulphur isotope analyses in archaeological research: A review. *Earth-Science Reviews* 142:1-17.

Nehlich O, Borić D, Stefanović S, Richards MP (2010) Sulphur isotope evidence for freshwater fish consumption: A case study from the Danube Gorges, SE Europe. *Journal of Archaeological Science* 37(5):1131-1139.

Nehlich O, Fuller BT, Jay M, Mora A, Nicholson RA, Smith CI, Richards MP (2011) Application of sulphur isotope ratios to examine weaning patterns and freshwater fish consumption in roman Oxfordshire, UK. *GeochimicaEtCosmochimicaActa* 75(17):4963-4977.

Nehlich O, Richards MP (2009) Establishing collagen quality criteria for sulphur isotope analysis of archaeological bone collagen. *Archaeological and Anthropological Sciences* 1(1):59-75.

Nielsen H., Pilot J, Grinenko LN, Grinenko VA, Lein AY, Smith JW, Pankina RG (1991) Lithospheric sources of sulphur. In *Stable isotopes: Natural and Anthropogenic sulphur in the environment*.

Ottaway P (1992) *Archaeology in British towns: From the emperor Claudius to the Black Death*. London: Routledge.

Peterson BJ, Fry B (1987) Stable isotopes in ecosystem studies. *Annual review of ecology and systematics* 18(1):293-320.

Phenice TW (1969) A newly developed visual method of sexing the os pubis. *American Journal of Physical Anthropology* 30(2):297-301.

Platt C (1981) *The parish churches of medieval England*. London: Secker & Warburg.

Polet C, Katzenberg MA (2003) Reconstruction of the diet in a mediaeval monastic community from the coast of Belgium. *Journal of Archaeological Science* 30(5):525-533.

Quirós Castillo JA(2013) Los comportamientos alimentarios del campesinado medieval en el País Vasco y su entorno (siglos VIII–XIV). *Historia Agraria* 59:13–41.

Reitsema LJ, Crews DE, Polcyn M (2010) Preliminary evidence for medieval Polish diet from carbon and nitrogen stable isotopes. *Journal of Archaeological Science* 37(7):1413-1423.

Reitsema LJ, Vercellotti G (2012) Stable isotope evidence for sex- and status-based variations in diet and life history at medieval Trino Vercellese, Italy. *American Journal of Physical Anthropology* 148(4): 589-600.

Richards MP, Fuller BT, Hedges RE (2001) Sulphur isotopic variation in ancient bone collagen from Europe: implications for human palaeodiet, residence mobility, and modern pollutant studies. *Earth and Planetary Science Letters* 191(3):185-190.

Richards MP, Hedges RE (1999) Stable isotope evidence for similarities in the types of marine foods used by late mesolithic humans at sites along the Atlantic coast of Europe. *Journal of Archaeological Science* 26(6):717-722.

Roberts C, Manchester K (2007) *The archaeology of disease* Cornell University Press.

Roberts SB, Rosenberg I (2006) Nutrition and aging: changes in the regulation of energy metabolism with aging. *Physiological reviews* 86(2):651-667.

Salamon M, Coppa A, McCormick M, Rubini M, Vargiu R, Tuross N(2008) The consistency of historical and isotopic approaches in reconstructing the medieval Mediterranean diet. *Journal of Archaeological Science* 35(6):1667-1672.

Scheuer L, Black S (2000) Development and ageing of the juvenile skeleton. *Human Osteology in Archaeology and Forensic Science* 9-21.

Schoeninger MJ, DeNiro MJ (1984) Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochimica Et Cosmochimica Acta* 48(4):625-639.

Schoeninger MJ, DeNiro MJ, Tauber H (1983) Stable nitrogen isotope ratios of bone collagen reflect marine and terrestrial components of prehistoric human diet. *Science (New York, N.Y.)* 220(4604):1381-1383.

Schutkowski H, Herrmann B, Wiedemann F, Bocherens H, Grupe G (1999) Diet, status and decomposition at Weingarten: trace element and isotope analyses on early mediaeval skeletal material. *Journal of Archaeological Science* 26(6):675-685.

Sponheimer M, Robinson T, Ayliffe L, Roeder B, Hammer J, Passey B, West A, Cerling T, Dearing D, Ehleringer J (2003) Nitrogen isotopes in mammalian herbivores: Hair $\delta^{15}\text{N}$ values from a controlled feeding study. *International Journal of Osteoarchaeology* 13(1-2):80-87.

Steele K, Daniel RM (1978) Fractionation of nitrogen isotopes by animals: A further complication to the use of variations in the natural abundance of $\delta^{15}\text{N}$ for tracer studies. *The Journal of Agricultural Science* 90(01):7-9.

Swanson RN (1989) *Church and society in late medieval England*. Oxford: Blackwell.

Tauber H (1981) ^{13}C evidence for dietary habits of prehistoric man in Denmark. *Nature* 292(5821):332-333.

Teeri J, Schoeller D (1979) $\delta^{13}\text{C}$ values of an herbivore and the ratio of C_3 to C_4 plant carbon in its diet. *Oecologia*,39(2):197-200.

Thomas RM (2007) Food and the maintenance of social boundaries in medieval England. *The Archaeology of Food, and Identity, Occasional Papers* 34:130-151.

Tieszen LL, Boutton TW, Tesdahl K, Slade NA (1983) Fractionation and turnover of stable carbon isotopes in animal tissues: Implications for $\delta^{13}\text{C}$ analysis of diet. *Oecologia* 57(1-2):32-37.

Valente J (1998) The new frontier: The role of the knightstemplar in the establishment of Portugal as an independent kingdom. *Mediterranean Studies* 7:49-65.

Van der Merwe, Nikolaas J, Vogel JC (1978) ^{13}C content of human collagen as a measure of prehistoric diet in woodland North America. *Nature* 276(5690):815-816.

Van Klinken GJ, Richards MP, Hedges BE (2002) An overview of causes for stable isotopic variations in past European human populations: Environmental, ecophysiological, and cultural effects. *Biogeochemical approaches to paleodietary analysis* pp. 39-63 Springer.

Veiga de Oliveira E, Galhano F, Pereira B (1975) *Actividades agro-marítimas em Portugal*. Instituto de Alta Cultura, Lisboa.

Vieira VV, Santos M (1995) *Directório de aquacultura e biotecnologia marinha*. Escola Superior de Biotecnologia da Universidade Católica Portuguesa, Porto.

Vicente M(2013) Entre Zêzere e Tejo, Propriedade e Povoamento. DoutoramentoemHistoria Medieval. Universidade de Lisboa [Unpublished].

Vogel JC (1978)Isotopic assessment of the dietary habits of ungulates. South African Journal of Science 74(8):298-301.

Wakshal E, Nielsen H (1982) Variations of $\delta^{34}\text{S}$ (SO_4), $\delta^{18}\text{O}$ (H_2O) and Cl/SO_4 ratio in rainwater over northern Israel, from the Mediterranean Coast to Jordan Rift Valley and Golan Heights. Earth and Planetary Science Letters 61(2):272-282.

Walker PL, DeNiro MJ (1986) Stable nitrogen and carbon isotope ratios in bone collagen as indices of prehistoric dietary dependence on marine and terrestrial resources in southern California. American Journal of Physical Anthropology 71(1):51-61.

Weston DA (2012) Nonspecific infection in paleopathology: interpreting periosteal reactions. A companion to paleopathology 492-512.

Wood J, Milner G, Harpending H, Weiss K (1992) The osteological paradox - problems of inferring prehistoric health from skeletal samples. Current Anthropology 33(4):343-370.

Zohary T, Erez J, Gophen M, Berman-Frank I, Stiller M (1994) Seasonality of stable carbon isotopes within the pelagic food web of Lake Kinneret. Limnology and Oceanography 39(5):1030-1043.

Figures list

Figure 1. Stable isotope values of fauna and human (from different social status) bone collagen. Lines indicate the mean without the outlier ($\delta^{13}\text{C} = -18.6\text{‰}$; $\delta^{15}\text{N} = 10.8\text{‰}$) and two standard deviations ($\mu \pm 2\sigma$). Grey area indicates the expected values for the trophic level increase from the analysed fauna.

Figure 2. Stable isotope values of fauna and human bone collagen. Lines indicate the mean without the outlier ($\delta^{13}\text{C} = -18.6\text{‰}$; $\delta^{34}\text{S} = 13.1\text{‰}$) and two standard deviations ($\mu \pm 2\sigma$).

Figure 3. Stable isotope values of individuals with estimated sex and age. Line at x-axis marks the. Lines indicate the mean and two standard deviations ($\mu \pm 2\sigma$) for all the samples except the outlier ($\delta^{13}\text{C} = -18.6 \pm 1.0\text{‰}$; $\delta^{15}\text{N} = 10.8 \pm 1.7\text{‰}$), the young ($\delta^{13}\text{C} = -18.4 \pm 0.9\text{‰}$; $\delta^{15}\text{N} = 11.4 \pm 1.4\text{‰}$) and the old ($\delta^{13}\text{C} = -18.7 \pm 0.5\text{‰}$; $\delta^{15}\text{N} = 10.6 \pm 1.8\text{‰}$) adults.

Figure 4. Stable isotope values ($\delta^{34}\text{S}$, $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$) of individuals from different social status and with estimated age (young, mature and old adults).

Figure 5. Geological map of Tomar's region. Tomar's and Santarém surroundings represent the evaporites, gypsum and marl. 1 – Nabão River, 2 – Zêzere River, 3 – Tagus River.

Figure 6. Carbon and nitrogen stable isotope comparison between pre-historic and late medieval Tomar and other late medieval European samples. Portugal: Tomar (this study), Tomar prehistoric (n=2, Abrigo do Morgado Superior, unpublished data). Spain: Zaballa (n=14, 10th – 15th century, Lubritto et al., 2013); Treviño (n=15, 12th – 14th century, Quirós Castillo, 2013); Zornoztegi (n=7, 12th – 14th century, Quirós Castillo, 2013); Colegiata St. Maria (n=24, 13th – 16th century, Alexander et al., 2015); Benipeixcar (n=20, 15th – 16th century, Alexander et al., 2015). Italy: Rome (n=29 15th century Salamon et al., 2008); Trino Vercellese (n=30, 8th – 13th century, Reitsema et al., 2012). Poland: Giecz (n= 24, 11th – 12th century, Reitsema et al., 2010). Belgium: Koksijde (n=19, 12th – 15th century, Polet & Katzenberg, 2003). England: St. Andrew (n=155, 13th – 16th century, Müldner & Richards, 2007).