

Kent Academic Repository

Full text document (pdf)

Citation for published version

Loureiro, Ana F. and Xu, Kuan (2019) Volterra-type convolution of classical polynomials. *Mathematics of Computation*. ISSN 0025-5718. (In press)

DOI

doi.org/10.1090/mcom/3427

Link to record in KAR

<https://kar.kent.ac.uk/66954/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

VOLTERRA-TYPE CONVOLUTION OF CLASSICAL POLYNOMIALS

ANA F. LOUREIRO AND KUAN XU

ABSTRACT. We present a general framework for calculating the Volterra-type convolution of polynomials from an arbitrary polynomial sequence $\{P_k(x)\}_{k \geq 0}$ with $\deg P_k(x) = k$. Based on this framework, series representations for the convolutions of classical orthogonal polynomials, including Jacobi and Laguerre families, are derived, along with some relevant results pertaining to these new formulas.

1. INTRODUCTION

Volterra-type convolution, as a fundamental operation, is commonly seen in many fields of sciences and engineering, including statistics and probability theory [14], computer vision [10], image and signal processing [8], and system control [27]. Particularly, in applied mathematics, convolution operators figure in many topics: Green's function [9], Duhamel's principle [28], non-reflecting boundary condition [12], large eddy simulation [32], approximation theory [31], fractional calculus [13], among others. Convolution operators are the key building blocks of the convolution integral equations [5, 6, 20]. Let $f : [\alpha, \beta] \mapsto \mathbb{C}$ and $g : [\gamma, \delta] \mapsto \mathbb{C}$ be two continuous integrable functions defined on intervals with a same length, that is $\beta - \alpha = \delta - \gamma$, where α and γ are finite numbers while β and δ can be finite or infinite. Their convolution $h(x)$ is a third function, given by

$$(1.1) \quad h(x) = (f * g)(x) = \int_{\gamma}^{x-\alpha} f(x-t)g(t)dt, \quad x \in [\alpha + \gamma, \alpha + \delta],$$

where the domain of $h(x)$, i.e. $[\alpha + \gamma, \alpha + \delta]$, has the same length as those of $f(x)$ and $g(x)$. This operation is often denoted by an asterisk, as in (1.1).

When β and δ are finite, $f(x)$ and $g(x)$ are compactly supported and can be mapped to the interval $[-1, 1]$ via changes of variables and the convolution of the mapped versions of $f(x)$ and $g(x)$ differs from that of the original $f(x)$ and $g(x)$ by an affine transform only. Therefore, with slight abuse of our notation, we can consider exclusively the convolution of two functions $f(x)$ and $g(x)$ that are defined on $[-1, 1]$, that is,

$$(1.2) \quad h(x) = (f * g)(x) = \int_{-1}^{x+1} f(x-t)g(t)dt, \quad x \in [-2, 0],$$

2010 *Mathematics Subject Classification.* 42A85, 44A35, 33C05, 33C20, 33C45, 42A10, 41A10, 41A58 .

Key words and phrases. convolution, Volterra convolution integral, orthogonal polynomials, Jacobi polynomials, Gegenbauer polynomials, Legendre polynomials, Chebyshev polynomials, Laguerre polynomials.

where the convolution $h(x)$ is, in this case, a function on $[-2, 0]$. Analogously, when β and δ are infinities (1.1) becomes

$$(1.3) \quad h(x) = (f * g)(x) = \int_0^x f(x-t)g(t)dt, \quad x \in [0, \infty],$$

up to a real Möbius transform. Note that the domains of the transformed $f(x)$, $g(x)$, and the convolution $h(x)$ all become $[0, \infty]$ in (1.3).

A powerful working paradigm that motivates this investigation and are commonly adopted in problems where functions considered are smooth is to replace $f(x)$ and $g(x)$ by their unique series representation in terms of classical orthogonal polynomials, e.g. Chebyshev or (weighted) Laguerre series for $f(x)$ and $g(x)$ in (1.2) and (1.3) respectively. In numerical computation, such series are usually truncated at certain degrees so that the finite series accurate to machine precision can serve as good approximants. In either case, the calculation of convolution integrals boils down to the convolution of polynomial series of finite or infinite degrees, or, further, to the convolution of classical orthogonal polynomials. To see this, suppose that $f(x)$ and $g(x)$ are approximated by two series

$$(1.4) \quad f_M(x) = \sum_{m=0}^M a_m P_m(x) \quad \text{and} \quad g_N(x) = \sum_{n=0}^N b_n P_n(x),$$

of the set of polynomials $\{P_n\}_{n \geq 0}$ such that $\deg P_n = n$. The convolution of $f_M(x)$ and $g_N(x)$ results in a degree $M + N + 1$ polynomial

$$h_{M+N+1}(x) = \sum_{k=0}^{M+N+1} c_k P_k(x).$$

As discussed in [34], the convolution operator

$$V[f_M](g_N) = h_{M+N+1}(x) = \int_{-a}^{x+a} f_M(x-t)g_N(t)dt,$$

which is defined by $f_M(x)$ and applied to $g_N(x)$, can be represented as a $(M + N + 2) \times (N + 1)$ matrix R so that the coefficients vector $\underline{c} = (c_1, c_2, \dots, c_{M+N+1})^T$ equals the product of R and the column vector $\underline{b} = (b_1, b_2, \dots, b_N)^T$. That is,

$$(1.5) \quad \underline{c} = R\underline{b}.$$

In [34] this convolution matrix R is constructed numerically via a stable method. For the case of Jacobi polynomials, orthogonal on $[-1, 1]$ (therefore $a = -1$), that method is based on a four- or five-point recurrence relation satisfied by its entries alongside weighted symmetry properties of R . For the Laguerre polynomials $L_n^{(0)}$, orthogonal on $[0, +\infty]$ (therefore $a = 0$), it is shown in [34, Th. 5.2] that the convolution matrix R is constructed as a difference of two lower-triangular Toeplitz matrices. However, for neither the Jacobi polynomials nor the generalized Laguerre polynomials the entries of the convolution matrix R are known explicitly. On the one hand, the (j, n) -entry of R in (1.5) is the

coefficient $R_{j,n}$ in

$$(1.6) \quad \int_{-a}^{x+a} f_M(x-t)P_n(t)dt = \sum_{j=0}^{M+n+1} R_{j,n}P_j(x+a).$$

On the other hand, if

$$(1.7) \quad \int_{-a}^{x+a} P_m(x-t)P_n(t)dt = \sum_{j=0}^{m+n+1} \rho_{j,n}^m P_j(x+a),$$

then

$$\int_{-a}^{x+a} f_M(x-t)P_n(t)dt = \sum_{j=0}^n \sum_{m=0}^M a_m \rho_{j,n}^m P_j(x+a) + \sum_{j=n+1}^{M+n+1} \sum_{m=j-(n+1)}^M a_m \rho_{j,n}^m P_j(x+a),$$

which follows from (1.7) and (1.4), by swapping the order of the sums. Comparing (1.6) and the latter equation leads to

$$(1.8) \quad R_{j,n} = \sum_{m=\max(0,n+1-j)}^M a_m \rho_{j,n}^m.$$

It turns out that explicit expressions for the ρ -coefficients in (1.7) are not known, even when $P_n(x)$ are classical polynomials of Jacobi and (general) Laguerre type, except for Laguerre polynomials $L_n^{(0)}(x)$ [24, Eq. (18.17.2)]. The knowledge of the ρ -expressions may hint us on the rich structure of convolution matrices R and, in turn, shed light upon their fast construction as well as the design of fast algorithms for convolving polynomial series.

Our main goal in this work is to obtain explicit expressions for $\rho_{j,n}^m$ in (1.7) when $\{P_n\}_{n \geq 0}$ belongs to the family of Jacobi or Laguerre polynomials. To this end, we develop a new framework, which is universally applicable to the convolution of general polynomial sets. We approach the problem by exploiting the nature of the Volterra-type convolution operator rather than any intrinsic properties possessed by the polynomials considered, e.g. orthogonality. The results presented in this paper for the convolution coefficients of the Jacobi and Laguerre families are not seen in literature and can largely expand the collection of the existing convolution formulas comprised of those of Laguerre polynomials $L_n^{(0)}(x)$ [24, Eq. (18.17.2)] and Bessel functions of the first kind $J_n(x)$ [24, Eq. (10.22.31)].

It should be noticed that the notion of convolution of orthogonal polynomials has been explored in several works, mostly regarding a discrete convolution procedure. For instance, discrete convolution appears in the theory of irreducible Lie algebra $\mathfrak{su}(1,1)$ in [17] and, more recently, transformations of such type are discussed in [16]. In addition, there are continuous transformations involving orthogonal polynomials, which can be regarded as projection operators that somewhat resemble but do not coincide with the Volterra-type convolution transform here considered; see, for instance, [4] for Jacobi polynomials and [7] for other orthogonal polynomials.

In the next section, Theorem 2.3 gives explicit expressions for the ρ -coefficients for a general polynomial set. These expressions are given as sums depending only on the connection coefficients

between derivatives of P_n as well as those between the sequence of monomials with P_n (also known as inversion formula coefficients). For several sets of polynomials of hypergeometric type, including the classical polynomials, these connection coefficients are well known (see e.g. [1, 3, 15, 18, 19, 2, 35, 23, 26, 30, 33]). Theorem 2.4 further shows that some of the ρ -coefficients are exactly zero when $\{P_n\}_{n \geq 0}$ is a sequence of classical polynomials. In Section 3, we derive explicit and new expressions for the ρ -coefficients in (1.7), with $a = 1$, for the Jacobi family (see Theorem 3.4), including the special cases of Gegenbauer (see Corollary 3.10), Legendre (see Corollary 3.11), and Chebyshev of the second kind (see Corollary 3.13) for which the ρ -expressions become simpler. Section 4 is devoted to the Laguerre family, where we prove that the ρ -coefficients in (1.7), with $a = 0$, significantly simplify to a ratio of *Pochhammer symbols* (see Theorem 4.1). We close the paper with a few remarks in Section 5.

2. CONVOLUTION OF TWO ELEMENTS IN A POLYNOMIAL SEQUENCE

Let $\{P_n(x)\}_{n \geq 0}$ be a polynomial sequence with $\deg P_n(x) = n$, which forms a basis of the vector space of polynomials with complex coefficients. We consider the problem of finding explicit expressions for the $P_n(x)$ -series coefficients $\rho_{j,n}^m$ so that (1.7) holds for a given constant a . A change of variable $t \rightarrow x - \tau$ shows the commutativity of the convolution in (1.7), which gives a remarkable symmetry property

$$\rho_{j,n}^m = \rho_{j,m}^n,$$

for any $j, n, m \geq 0$. Therefore, there is no loss of generality if one assumes $n \geq m$ or $m \geq n$.

A sequence $\{\frac{d^s}{dx^s} P_{n+s}(x)\}_{s \geq 0}$ for $n \geq 0$, i.e. the s -th derivative of the original sequence, also spans the vector space of polynomials. Therefore, the r -th derivatives of $P_n(x)$ can be represented by a linear combination of the elements of $\{\frac{d^s}{dx^s} P_{n+s}(x)\}_{s \geq 0}$ as

$$(2.1) \quad \frac{d^r P_n(x)}{dx^r} = \sum_{k=0}^{n-r} \gamma_{n-r,k}^{(r,s)} \frac{d^s P_{k+s}(x)}{dx^s}, \quad n \geq r,$$

where the coefficients $\gamma_{n-r,k}^{(r,s)}$ are referred to as the *connection coefficients* between $\{\frac{d^r}{dx^r} P_n(x)\}_{n \geq r}$ and $\{\frac{d^s}{dx^s} P_{n+s}(x)\}_{n \geq 0}$. The unique representation of $P_n(x)$ in terms of the monomial sequence $\{(x+a)^n\}_{n \geq 0}$ can be obtained by its Taylor expansion about $x = -a$

$$P_n(x) = \sum_{k=0}^n \frac{1}{k!} \left. \frac{d^k P_n(x)}{dx^k} \right|_{x=-a} (x+a)^k, \quad n \geq 0.$$

and, reversely, a unique set of coefficients $b_{n,k}$ exist such that

$$(2.2) \quad (x+a)^n = \sum_{k=0}^n b_{n,k} P_k(x), \quad n \geq 0.$$

When $\{P_n(x)\}_{n \geq 0}$ is an orthogonal sequence, these b -coefficients can be obtained via the orthogonality measures and their moments.

Lemma 2.1. *The γ -connection coefficients in (2.1) can be expressed in terms of the connection b -coefficients in (2.2) by*

$$(2.3) \quad \gamma_{n-r,k}^{(r,s)} = \sum_{\sigma=0}^{n-(r+k)} \frac{b_{\sigma+k+s,k+s}}{(\sigma+k+s)!} \left. \frac{d^{r+k+\sigma} P_n(x)}{dx^{r+k+\sigma}} \right|_{x=-a}.$$

Proof. The Taylor expansion of $\frac{d^r}{dx^r} P_n(x)$ gives

$$\frac{d^r P_n(x)}{dx^r} = \sum_{\sigma=0}^{n-r} \frac{1}{\sigma!} \left. \frac{d^{r+\sigma} P_n(x)}{dx^{r+\sigma}} \right|_{x=-a} (x+a)^\sigma = \sum_{\sigma=0}^{n-r} \frac{1}{(\sigma+s)!} \left. \frac{d^{r+\sigma} P_n(x)}{dx^{r+\sigma}} \right|_{x=-a} \left(\frac{d^s}{dx^s} (x+a)^{\sigma+s} \right)$$

where s is an arbitrary positive integer. Substituting (2.2) into the last equation gives

$$\frac{d^r P_n(x)}{dx^r} = \sum_{\sigma=0}^{n-r} \frac{1}{(\sigma+s)!} \left. \frac{d^{r+\sigma} P_n(x)}{dx^{r+\sigma}} \right|_{x=-a} \left(\sum_{k=0}^{\sigma} b_{\sigma+k+s,k+s} \frac{d^s}{dx^s} P_{k+s}(x) \right)$$

which, after exchanging the summations, becomes

$$\frac{d^r P_n(x)}{dx^r} = \sum_{k=0}^{n-r} \left(\sum_{\sigma=0}^{n-(r+k)} \frac{b_{\sigma+k+s,k+s}}{(\sigma+k+s)!} \left. \frac{d^{r+k+\sigma} P_n(x)}{dx^{r+k+\sigma}} \right|_{x=-a} \right) \frac{d^s}{dx^s} P_{k+s}(x).$$

Matching the like terms in (2.1) leads to (2.3). \square

We omit the proof of the following lemma, which is concerned with the p -th derivative of the convolution in (1.7) and can be easily shown by repeatedly applying the Leibniz rule for differentiation under the integral sign.

Lemma 2.2. *For a positive integer p ,*

$$(2.4) \quad \frac{d^p}{dx^p} \int_{-a}^{x+a} P_m(x-t) P_n(t) dt = \int_{-a}^{x+a} \frac{d^p}{dx^p} P_m(x-t) P_n(t) dt + \sum_{k=1}^p \left. \frac{d^{p-k} P_m(x)}{dx^{p-k}} \right|_{x=-a} \frac{d^{k-1}}{dx^{k-1}} P_n(x+a).$$

With Lemmas 2.1 and 2.2, we show in the following theorem that $\rho_{j,n}^m$ in (1.7) can be represented in terms of the γ -coefficients in (2.1) and the b -coefficients in (2.2).

Theorem 2.3. *For $0 \leq j \leq m+n+1$, the coefficients $\rho_{j,n}^m$ in (1.7) can be expressed as*

$$(2.5) \quad \rho_{j,n}^m = \sum_{p=j}^{m+n+1} \frac{b_{p,j}}{p!} \sum_{v=1}^p \left(\left. \frac{d^{p-v} P_m(x)}{dx^{p-v}} \right|_{x=-a} \left. \frac{d^{v-1} P_n(x)}{dx^{v-1}} \right|_{x=-a} \right),$$

or, equivalently,

$$(2.6a) \quad \rho_{j,n}^m = \sum_{v=\max(1,j-n)}^{m+1} \left(\gamma_{n-j+v,0}^{(j-v,j)} \left. \frac{d^{v-1} P_m(x)}{dx^{v-1}} \right|_{x=-a} \right) \text{ for } j \geq m+1$$

and

$$(2.6b) \quad \rho_{j,n}^m = \sum_{v=1}^j \left(\gamma_{m-j+v,0}^{(j-v,j)} \frac{d^{v-1}P_n(x)}{dx^{v-1}} \Big|_{x=-a} \right) + \sum_{v=j+1}^{n+1} \left(\frac{d^{v-1}P_n(x)}{dx^{v-1}} \Big|_{x=-a} \sum_{p=0}^m \frac{b_{p+v,j}}{(p+v)!} \frac{d^p P_m(x)}{dx^p} \Big|_{x=-a} \right) \text{ for } 0 \leq j \leq m,$$

where the γ - and the b -coefficients are the connection coefficients given in (2.1) and (2.2), respectively.

Proof. To show (2.5), we Taylor expand the convolution integral in (1.7) about $x = -2a$

$$(2.7) \quad \int_{-a}^{x+a} P_m(x-t)P_n(t)dt = \sum_{p=0}^{m+n+1} \frac{1}{p!} \left[\frac{d^p}{d(x+a)^p} \int_{-a}^{x+a} P_m(x-t)P_n(t)dt \right]_{x=-2a} (x+2a)^p.$$

Note that the Taylor coefficients in (2.7) can be obtained using (2.4):

$$\left[\frac{d^p}{d(x+a)^p} \int_{-a}^{x+a} P_m(x-t)P_n(t)dt \right]_{x=-2a} = \sum_{v=1}^p \frac{d^{p-v}P_m(x)}{dx^{p-v}} \Big|_{x=-a} \frac{d^{v-1}P_n(x)}{dx^{v-1}} \Big|_{x=-a},$$

where the sum is assumed zero when $p = 0$, and $(x+2a)^p$ can be replaced by its expansion in $\{P_n(x+a)\}_{n \geq 0}$ as given in (2.2):

$$(x+2a)^p = \sum_{j=0}^p b_{p,j} P_j(x+a).$$

We substitute the last two equations into (2.7) and exchange the order of the summations to have

$$(2.8) \quad \int_{-a}^{x+a} P_m(x-t)P_n(t)dt = \sum_{j=0}^{m+n+1} \left(\sum_{p=j}^{m+n+1} \frac{b_{p,j}}{p!} \sum_{v=1}^p \frac{d^{p-v}P_m(x)}{dx^{p-v}} \Big|_{x=-a} \frac{d^{v-1}P_n(x)}{dx^{v-1}} \Big|_{x=-a} \right) P_j(x+a).$$

Matching terms in (2.8) and (1.7) gives (2.5).

To see (2.6a), we take the j -th derivative on both sides of (1.7) for $j \geq m+1$ to have

$$\frac{d^j}{dx^j} \int_{-a}^{x+a} P_m(x-t)P_n(t)dt = \sum_{k=0}^{m+n+1-j} \rho_{k+j,n}^m \frac{d^j}{dx^j} P_{k+j}(x+a).$$

Meanwhile, Lemma 2.2 gives

$$\frac{d^j}{dx^j} \int_{-a}^{x+a} P_m(x-t)P_n(t)dt = \sum_{\ell=1}^j \frac{d^{j-\ell}P_m(x)}{dx^{j-\ell}} \Big|_{x=-a} \frac{d^{\ell-1}}{dx^{\ell-1}} P_n(x+a),$$

where the convolution integral on the right-hand side of (2.4) vanishes here, since the integrand becomes zero for $j \geq m+1$. Combining the last two equations, we have

$$(2.9) \quad \sum_{k=0}^{m+n+1-j} \rho_{k+j,n}^m \frac{d^j}{dx^j} P_{k+j}(x+a) = \sum_{\ell=1}^j \frac{d^{j-\ell}P_m(x)}{dx^{j-\ell}} \Big|_{x=-a} \frac{d^{\ell-1}}{dx^{\ell-1}} P_n(x+a).$$

If we denote by S the sum on the right-hand side of (2.9),

$$S = \sum_{\ell=0}^{j-1} \frac{d^{j-\ell-1}P_m(x)}{dx^{j-\ell-1}} \Big|_{x=-a} \frac{d^\ell}{dx^\ell} P_n(x+a) = \sum_{\ell=j-(m+1)}^n \frac{d^{j-\ell-1}P_m(x)}{dx^{j-\ell-1}} \Big|_{x=-a} \frac{d^\ell}{dx^\ell} P_n(x+a),$$

where the last equality is obtained by noting that the summand disappear when $j - \ell - 1 > m$ and $\ell > n$. Now we use the connection formula (2.1) once again to have

$$(2.10) \quad \begin{aligned} S &= \sum_{\ell=j-(m+1)}^n \frac{d^{j-\ell-1} P_m(x)}{dx^{j-\ell-1}} \Big|_{x=-a} \sum_{k=0}^{n-\ell} \gamma_{n-\ell,k}^{(\ell,j)} \frac{d^j}{dx^j} P_{k+j}(x+a) \\ &= \sum_{k=0}^{m+n+1-j} \left(\gamma_{n-\ell,k}^{(\ell,j)} \sum_{\ell=j-(m+1)}^{n-k} \frac{d^{j-\ell-1} P_m(x)}{dx^{j-\ell-1}} \Big|_{x=-a} \right) \frac{d^j}{dx^j} P_{k+j}(x+a), \end{aligned}$$

where we have swapped the order of the sums. Combining (2.9) and (2.10) and matching terms yield

$$(2.11) \quad \rho_{k+j,n}^m = \sum_{\ell=j-(m+1)}^{n-k} \gamma_{n-\ell,k}^{(\ell,j)} \frac{d^{j-\ell-1} P_m(x)}{dx^{j-\ell-1}} \Big|_{x=-a}$$

for $0 \leq k \leq m+n+1-j$. Particularly, for $k=0$, (2.11) becomes

$$\rho_{j,n}^m = \sum_{\ell=j-(m+1)}^n \gamma_{n-\ell,0}^{(\ell,j)} \frac{d^{j-\ell-1} P_m(x)}{dx^{j-\ell-1}} \Big|_{x=-a} = \sum_{v=j-n}^{m+1} \gamma_{n-j+v,0}^{(j-v,j)} \frac{d^{v-1} P_m(x)}{dx^{v-1}} \Big|_{x=-a},$$

where we use the change of variable $\ell = j - v$ in the last step. Ensuring $v - 1 \geq 0$, we obtain (2.6a).

To see (2.6b), we swap the order of sums in (2.5) to have

$$\rho_{j,n}^m = \left(\sum_{v=1}^j \sum_{p=j}^{m+v} + \sum_{v=j+1}^{n+1} \sum_{p=v}^{m+v} \right) \left(\frac{b_{p,j}}{p!} \frac{d^{p-v} P_m(x)}{dx^{p-v}} \Big|_{x=-a} \frac{d^{v-1} P_n(x)}{dx^{v-1}} \Big|_{x=-a} \right),$$

since $\frac{d^{p-v} P_m(x)}{dx^{p-v}} \Big|_{x=-a}$ and $\frac{d^{v-1} P_n(x)}{dx^{v-1}} \Big|_{x=-a}$ vanish for $p > m+v$ and $v > n+1$, respectively. This is equivalent to

$$(2.12) \quad \begin{aligned} \rho_{j,n}^m &= \sum_{v=1}^j \frac{d^{v-1} P_n(x)}{dx^{v-1}} \Big|_{x=-a} \sum_{p=0}^{m+v-j} \left(\frac{b_{p+j,j}}{(p+j)!} \frac{d^{p-v+j} P_m(x)}{dx^{p-v+j}} \Big|_{x=-a} \right) \\ &\quad + \sum_{v=j+1}^{n+1} \frac{d^{v-1} P_n(x)}{dx^{v-1}} \Big|_{x=-a} \sum_{p=0}^m \left(\frac{b_{p+v,j}}{(p+v)!} \frac{d^p P_m(x)}{dx^p} \Big|_{x=-a} \right). \end{aligned}$$

In (2.3), we set $k=0$ and $s=j$ and replacing r and n by $j-v$ and m respectively to have

$$\gamma_{m+v-j,0}^{(j-v,j)} = \sum_{\sigma=0}^{m+v-j} \frac{b_{\sigma+j,j}}{(\sigma+j)!} \frac{d^{j-v+\sigma} P_m(x)}{dx^{j-v+\sigma}} \Big|_{x=-a},$$

by applying which the first double sum in (2.12) can be simplified as

$$\sum_{v=1}^j \frac{d^{v-1} P_n(x)}{dx^{v-1}} \Big|_{x=-a} \sum_{p=0}^{m+v-j} \frac{b_{p+j,j}}{(p+j)!} \frac{d^{p-v+j} P_m(x)}{dx^{p-v+j}} \Big|_{x=-a} = \sum_{v=1}^j \gamma_{m-j+v,0}^{(j-v,j)} \frac{d^{v-1} P_n(x)}{dx^{v-1}} \Big|_{x=-a}.$$

Hence, (2.6b) is obtained. \square

To obtain the preceding results, we have nowhere assumed the sequence of polynomials $\{P_n(x)\}_{n \geq 0}$ to be orthogonal and the expressions for the γ -connection coefficients and the b -coefficients are, in general, not easy to calculate. However, when $\{P_n(x)\}_{n \geq 0}$ is an orthogonal polynomial sequence,

these coefficients are usually explicitly known or more likely to be obtainable. In fact, for a non-decreasing, non-negative function $w(x)$ in $[a, b]$ which is measurable in the Lebesgue sense, that is, all the moments $\int_a^b x^n w(x) dx$ exist and are finite¹, there is an orthogonal sequence of polynomials $\{P_n(x)\}_{n \geq 0}$ for which

$$\langle P_m(x), P_n(x) \rangle_w = \int_a^b P_m(x) P_n(x) w(x) dx = h_m \delta_{m,n}, \quad m, n = 0, 1, 2, \dots$$

where $h_m = \langle P_m(x), P_m(x) \rangle_w \neq 0$ for all positive integers m and $\delta_{m,n}$ denotes the *Kronecker delta symbol*. Since $P_n(x) \in L_w^2(a, b)$ and $\{P_n(x)\}_{n \geq 0}$ spans the vector space of polynomials, any polynomial $p(x)$ of degree m can be written as

$$p(x) = \sum_{k=0}^m c_k P_k(x) \quad \text{with} \quad c_k = \frac{\langle p(x), P_k(x) \rangle_w}{h_k}.$$

Particularly, for classical orthogonal polynomial sequences, i.e. Jacobi, Laguerre, Hermite, and Bessel polynomials, these b - and γ - connection coefficients are well studied [2, 15, 23, 25, 26, 35]. Based on these known results, we shall explicitly calculate the ρ -coefficients in (1.7) for the Jacobi and the Laguerre polynomials in the next two sections.

We close this section with the following theorem which shows that a consecutive part of the ρ -coefficients in (1.7) could be exactly zero when $\{P_n(x)\}_{n \geq 0}$ is a classical orthogonal polynomial sequence. However, these zeros are not immediately obvious from Theorem 2.3.

Theorem 2.4. *Let m, n be two nonnegative integers such that $n \geq m$. Suppose $\{P_n(x)\}_{n \geq 0}$ is a classical polynomial sequence and the interval $(-a, x+a)$ lies within the support of the orthogonality measure of $\{P_n(x)\}_{n \geq 0}$. When $n \geq 2m + q + 2$, the series coefficients $\rho_{j,n}^m = 0$ for $m + 1 \leq j \leq n - m - q - 1$, where $q = 1, 1, 0$ and -1 for Jacobi, Bessel, Laguerre and Hermite polynomials, respectively.*

Proof. By orthogonality, we have

$$(t+a)^v P_n(t) = \sum_{k=\max\{n-v, 0\}}^{n+v} \lambda_{v,n}(k) P_k(t),$$

where $\lambda_{v,n}(k) = \frac{\langle (t+a)^v P_n(t), P_k(t) \rangle_w}{\langle P_k(t), P_k(t) \rangle_w}$, which, together with the Taylor expansion of $P_m(x-t)$ about $t = -a$

$$P_m(x-t) = \sum_{v=0}^m \frac{(-1)^v}{v!} \frac{d^v P_m(x+a)}{d(x+a)^v} (t+a)^v,$$

gives

$$(2.13) \quad P_m(x-t) P_n(t) = \sum_{v=0}^m \frac{(-1)^v}{v!} \frac{d^v P_m(x+a)}{d(x+a)^v} \sum_{k=n-v}^{n+v} \lambda_{v,n}(k) P_k(t).$$

¹In the case of $a = -\infty$ or $b = +\infty$, we require that $\lim_{x \rightarrow -\infty} x^n w(x)$ and $\lim_{x \rightarrow +\infty} x^n w(x)$ to be finite, respectively, for any positive integer n .

If, in addition, $\{P_n(x)\}_{n \geq 0}$ is a classical sequence, there exists a polynomial $\Phi(t)$ of degree at most 2 and $\xi_{n,v}$ such that

$$\Phi(t) \frac{dP_{k+1}(t)}{dt} = \sum_{r=k}^{k+\deg \Phi} \xi_{k,r} P_r(t),$$

where $\xi_{k,k+\deg \Phi} \xi_{k,k} \neq 0$, for all $n \geq 0$. Reversely, there are coefficients $\tilde{\xi}_{k,r}$ such that

$$P_k(t) = \sum_{r=k-q}^{k+1} \tilde{\xi}_{k,r} \frac{dP_r(t)}{dt},$$

where $q = \deg \Phi(x) - 1$ and $\tilde{\xi}_{k,k-q} \tilde{\xi}_{k,k+1} \neq 0$ [22, Prop. 2.4]. This can be deemed as a special case of (2.1). In particular, if $\{P_n(x)\}_{n \geq 0}$ is the classical orthogonal sequence of Jacobi, Bessel, Laguerre and Hermite polynomials, $\deg \Phi(x) = 2, 2, 1$, and 0, respectively.

We integrate (2.13) over the interval $(-a, x+a)$ to have

$$\int_{-a}^{x+a} P_m(x-t) P_n(t) dt = \sum_{v=0}^m \frac{(-1)^v}{v!} \frac{d^v P_m(x+a)}{d(x+a)^v} \sum_{k=n-v}^{n+v} \lambda_{v,n}(k) \sum_{r=k-q}^{k+1} \tilde{\xi}_{k,r} (P_r(x+a) - P_r(-a)).$$

Swapping the order of the last two sums and absorbing the innermost summation into new coefficients $\tilde{\lambda}_{v,n}(k)$, we have

$$\begin{aligned} \int_{-a}^{x+a} P_m(x-t) P_n(t) dt &= \sum_{v=0}^m \frac{(-1)^v}{v!} \frac{d^v P_m(x+a)}{d(x+a)^v} \sum_{k=n-v-q}^{n+v+1} \tilde{\lambda}_{v,n}(k) (P_k(x+a) - P_k(-a)) \\ &= \sum_{v=0}^m \frac{(-1)^v}{v!} \frac{d^v P_m(x+a)}{d(x+a)^v} \sum_{k=n-v-q}^{n+v+1} \tilde{\lambda}_{v,n}(k) P_k(x+a) - \sum_{v=0}^m \frac{(-1)^v}{v!} S_{v,n} \frac{d^v P_m(x+a)}{d(x+a)^v}, \end{aligned}$$

where $\tilde{\lambda}_{v,n}(n-v-q) \tilde{\lambda}_{v,n}(n+v+1) \neq 0$ and $S_{v,n} = \sum_{k=n-v-q}^{n+v+1} \tilde{\lambda}_{v,n}(k) P_k(-a)$. Since $\frac{d^v P_m(x+a)}{d(x+a)^v}$ is a polynomial of degree $m-v$, there are coefficients $\chi_{m,v,k}$ such that

$$\begin{aligned} \int_{-a}^{x+a} P_m(x-t) P_n(t) dt &= \sum_{v=0}^m \frac{(-1)^v}{v!} \sum_{k=n-v-q}^{n+v+1} \tilde{\lambda}_{v,n}(k) \sum_{j=k-(m-v)}^{k+(m-v)} \chi_{m,v,k}(j) P_j(x+a) \\ &\quad - \sum_{v=0}^m \frac{(-1)^v}{v!} S_{v,n} \sum_{j=0}^{m-v} \gamma_{m-v,j}^{(v,0)} P_j(x+a), \end{aligned}$$

where we have applied (2.1) to the last sum. After swapping the order of the summations, we see there are coefficients $\tilde{\chi}_{m,n}(j)$ such that

$$\int_{-a}^{x+a} P_m(x-t) P_n(t) dt = \sum_{j=n-m-q}^{m+n+1} \tilde{\chi}_{m,n}(j) P_j(x+a) - \sum_{j=0}^m \tilde{\chi}_{m,n}(j) P_j(x+a).$$

When $n \geq 2m+q+2$, this means that $\rho_{j,n}^m = 0$ for $m+1 \leq j \leq n-m-q-1$. \square

3. CONVOLUTION OF JACOBI POLYNOMIALS

In this section, we derive the ρ -coefficients in (1.7) based on the results of Section 2 for the Jacobi-family, including the subcases of Gegenbauer, Legendre, and Chebyshev. To facilitate our discussion, we denote the Jacobi-based ρ -coefficients by $\rho_{j,n}^{m;(\alpha,\beta)}$ throughout this section, that is,

$$(3.1) \quad \int_{-1}^{x+1} P_m^{(\alpha,\beta)}(x-t) P_n^{(\alpha,\beta)}(t) dt = \sum_{j=0}^{m+n+1} \rho_{j,n}^{m;(\alpha,\beta)} P_j^{(\alpha,\beta)}(x+1),$$

which corresponds to (1.7) with $a = 1$. Here, $P_n^{(\alpha,\beta)}(x)$ denotes the Jacobi polynomial of degree $n \geq 0$ with $\alpha, \beta > -1$. With the most commonly-used normalization, which can be found, for example, in [29, §4.2.1], it can be represented as a terminating hypergeometric function

$$P_n^{(\alpha,\beta)}(x) = \frac{(\alpha+1)_n}{n!} {}_2F_1 \left(\begin{matrix} -n, n+\alpha+\beta+1 \\ \alpha+1 \end{matrix}; \frac{1-x}{2} \right), \quad n \geq 0,$$

where $(z)_n$ is the Pochhammer symbol, defined as

$$(z)_0 := 1 \quad \text{and} \quad (z)_n := \prod_{\sigma=0}^{n-1} (z+\sigma) \quad \text{for } n \geq 1.$$

Here and in the rest of this paper, we will use the generalized hypergeometric series

$${}_pF_q \left(\begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix}; x \right) := \sum_{n=0}^{\infty} \frac{(a_1)_n \cdots (a_p)_n x^n}{(b_1)_n \cdots (b_q)_n n!},$$

and its detail can be found, for example, in [24, Ch. 16].

The properties of $P_n^{(\alpha,\beta)}$ that we will make use of in the rest of this section include its value at -1

$$P_n^{(\alpha,\beta)}(-1) = \frac{(-1)^n (\beta+1)_n}{n!}$$

and a symmetry property

$$(3.2) \quad P_n^{(\alpha,\beta)}(-x) = (-1)^n P_n^{(\beta,\alpha)}(x).$$

The sequence of Jacobi polynomials $\{P_n^{(\alpha,\beta)}(x)\}_{n \geq 0}$ satisfy the orthogonality condition [15, Ch. 4]

$$\int_{-1}^1 P_k^{(\alpha,\beta)}(x) P_n^{(\alpha,\beta)}(x) (1-x)^\alpha (1+x)^\beta dx = \frac{2^{\alpha+\beta+1} \Gamma(\alpha+n+1) \Gamma(\beta+n+1)}{n! \Gamma(\alpha+\beta+n+1) (\alpha+\beta+2n+1)} \delta_{k,n},$$

for any integers $n, k \geq 0$. Being a member of a classical sequence, the p -th derivative of a Jacobi polynomial is another Jacobi polynomial with shifted parameters

$$(3.3) \quad \frac{d^p}{dx^p} P_n^{(\alpha,\beta)}(x) = \frac{(\alpha+\beta+n+1)_p}{2^p} P_{n-p}^{(\alpha+p,\beta+p)}(x),$$

and, in particular,

$$(3.4) \quad \left. \frac{d^p}{dx^p} P_n^{(\alpha,\beta)}(x) \right|_{x=-1} = \frac{2^{-p} (-1)^{n+p} (p+\beta+1)_{n-p} (n+\alpha+\beta+1)_p}{(n-p)!}.$$

The properties above allow us to derive a connection formula between the derivatives of Jacobi polynomials.

Lemma 3.1. *The p -th and q -th derivatives of Jacobi polynomials are linearly connected via*

$$(3.5) \quad \frac{d^p}{dx^p} P_{n+p}^{(\alpha, \beta)}(x) = \sum_{k=0}^n \gamma_{n,k}^{(p,q)}(\alpha, \beta) \frac{d^q}{dx^q} P_{k+q}^{(\alpha, \beta)}(x)$$

with

$$(3.6) \quad \gamma_{n,k}^{(p,q)}(\alpha, \beta) = \frac{(k+p+\alpha+1)_{n-k} (n+p+\alpha+\beta+1)_p (n+2p+\alpha+\beta+1)_k}{2^{p-q} (n-k)! (k+q+\alpha+\beta+1)_q (k+2q+\alpha+\beta+1)_k} \\ \times {}_3F_2 \left(\begin{matrix} k-n, k+q+\alpha+1, k+n+2p+\alpha+\beta+1 \\ k+p+\alpha+1, 2k+2q+\alpha+\beta+2 \end{matrix}; 1 \right).$$

Proof. The following connection formula, which can be found in [1, p.357], [15, Theorem 9.1.1] or [2], relates Jacobi polynomials with distinct parameters:

$$(3.7) \quad P_n^{(\alpha, \beta)}(x) = \sum_{k=0}^n a_{n,k}^{(\alpha, \beta; \gamma, \delta)} P_k^{(\gamma, \delta)}(x),$$

where

$$(3.8) \quad a_{n,k}^{(\alpha, \beta; \gamma, \delta)} = \frac{(k+\alpha+1)_{n-k} (n+\alpha+\beta+1)_k}{(n-k)! (k+\gamma+\delta+1)_k} {}_3F_2 \left(\begin{matrix} k-n, n+k+\alpha+\beta+1, k+\gamma+1 \\ k+\alpha+1, 2k+\gamma+\delta+2 \end{matrix}; 1 \right).$$

Combining (3.3) and (3.7), we have (3.5) with

$$(3.9) \quad \gamma_{n,k}^{(p,q)}(\alpha, \beta) = \frac{(n+p+\alpha+\beta+1)_p}{2^{p-q} (k+q+\alpha+\beta+1)_q} a_{n,k}^{(\alpha+p, \beta+p; \alpha+q, \beta+q)},$$

which, with (3.8) substituted in, gives (3.6). \square

Remark 3.2. Via (3.5), the symmetry property (3.2) implies:

$$\gamma_{n,k}^{(p,q)}(\alpha, \beta) = (-1)^{n+k} \gamma_{n,k}^{(p,q)}(\beta, \alpha).$$

One last ingredient we need for deriving $\rho_{k,n}^{m;(\alpha, \beta)}$ is the b -coefficients in (2.2) for representing the monomial basis in terms of Jacobi polynomials.

Lemma 3.3. *The connection coefficients $b_{n,k}(\alpha, \beta)$ in*

$$(3.10) \quad (x+1)^n = \sum_{k=0}^n b_{n,k}(\alpha, \beta) P_k^{(\alpha, \beta)}(x),$$

are given by

$$(3.11) \quad b_{n,k}(\alpha, \beta) = 2^n n! (\beta+1)_n \frac{(\alpha+\beta+2k+1) \Gamma(\alpha+\beta+k+1)}{(\beta+1)_k \Gamma(\alpha+\beta+n+k+2) (n-k)!}.$$

Proof. See, for example, [15, (4.2.15)], [21, p.277, Eq. (30)] or [35] for the proof. \square

3.1. The Jacobi-based convolution coefficients. Though we could derive the Jacobi-based convolution coefficients directly from the orthogonality of Jacobi polynomials, we opt to find the explicit expressions for the ρ -coefficients in the expansion of the integral (3.1) from Theorem 2.3 by substituting in (2.5) and (2.6) the expressions (3.4), (3.6) and (3.11).

Theorem 3.4. *Let m and n be two positive integers with $n \geq m$. The ρ coefficients in the expansion (3.1) can be expressed as*

$$(3.12a) \quad \rho_{j,n}^{m;(\alpha,\beta)} = \sum_{v=\max(1,|j-n|)}^{m+1} \varpi_{j,v}^{m,n}(\alpha,\beta) \quad \text{for } j \geq \max(m+1, n-m-1),$$

$$(3.12b) \quad \rho_{j,n}^{m;(\alpha,\beta)} = 0 \quad \text{for } m+1 \leq j \leq n-m-2, \text{ if } n \geq 2m+3,$$

$$(3.12c) \quad \rho_{j,n}^{m;(\alpha,\beta)} = \sum_{v=1}^j \varpi_{j,v}^{n,m}(\alpha,\beta) + \sum_{v=j+1}^{n+1} d_{v,j,n}^m(\alpha,\beta) \quad \text{for } 0 \leq j \leq m,$$

where

$$(3.12d) \quad \varpi_{j,v}^{m,n}(\alpha,\beta) = \frac{2(-1)^{m+v-1}(j-v+\alpha+1)_{n-j+v}(n+\alpha+\beta+1)_{j-v}(\beta+v)_{m-v+1}(m+\alpha+\beta+1)_{v-1}}{(j+\alpha+\beta+1)_j(m+1-v)!(n-j+v)!} \\ \times {}_3F_2\left(\begin{matrix} j-n-v, j+\alpha+1, n+j-v+\alpha+\beta+1 \\ j-v+\alpha+1, 2j+\alpha+\beta+2 \end{matrix}; 1\right),$$

and

$$(3.12e) \quad d_{v,j,n}^m(\alpha,\beta) = \frac{2(-1)^{m+n+1+v}(\alpha+\beta+2j+1)(\beta+v)(j+\beta+1)_{m-j}(\beta+1)_n(n+1+\alpha+\beta)_{v-1}}{m!(n+1-v)!(v-j)!(j+\alpha+\beta+1)_{v+1}} \\ \times {}_4F_3\left(\begin{matrix} 1, -m, \beta+v+1, m+\alpha+\beta+1 \\ v-j+1, \beta+1, j+\alpha+\beta+v+2 \end{matrix}; 1\right).$$

Proof. The zero coefficients given by (3.12b) are readily known from Theorem 2.4.

Following (2.6) with $a = 1$, we have

$$(3.13a) \quad \rho_{j,n}^{m;(\alpha,\beta)} = \sum_{v=\max(1,|j-n|)}^{m+1} \gamma_{n-j+v,0}^{(j-v,j)}(\alpha,\beta) \left. \frac{d^{v-1}P_m^{(\alpha,\beta)}(x)}{dx^{v-1}} \right|_{x=-1} \quad \text{for } j \geq m+1,$$

and

$$(3.13b) \quad \rho_{j,n}^{m;(\alpha,\beta)} = \sum_{v=1}^j \gamma_{m-j+v,0}^{(j-v,j)}(\alpha,\beta) \left. \frac{d^{v-1}P_n^{(\alpha,\beta)}(x)}{dx^{v-1}} \right|_{x=-1} \\ + \sum_{v=j+1}^{n+1} \left(\left. \frac{d^{v-1}P_n^{(\alpha,\beta)}(x)}{dx^{v-1}} \right|_{x=-1} \sum_{k=0}^m \frac{b_{k+v,j}(\alpha,\beta)}{(k+v)!} \left. \frac{d^k P_m^{(\alpha,\beta)}(x)}{dx^k} \right|_{x=-1} \right) \quad \text{for } 0 \leq j \leq m.$$

Combined with (3.4) and (3.6), (3.13a) gives (3.12a) and (3.12d). Similarly, the first sum on the right hand side of (3.13b) yields the first sum in (3.12c).

To find $d_{v,j,n}^m(\alpha, \beta)$ of the second sum in (3.12c), we first calculate the inner sum of (3.13b) to have

$$(3.14) \quad \sum_{k=0}^m \frac{b_{k+v,j}(\alpha, \beta)}{(k+v)!} \frac{d^k P_m^{(\alpha, \beta)}(x)}{dx^k} \Big|_{x=-1} = \frac{2^v (-1)^m (\alpha + \beta + 2j + 1) \Gamma(m + \beta + 1)}{m! \Gamma(v - j + 1) \Gamma(j + \beta + 1)} \\ \times \frac{\Gamma(\beta + v + 1) \Gamma(j + \alpha + \beta + 1)}{\Gamma(\beta + 1) \Gamma(j + \alpha + \beta + v + 2)} \sum_{k=0}^m \frac{(-m)_k (\beta + v + 1)_k (m + \alpha + \beta + 1)_k}{(\beta + 1)_k (-j + v + 1)_k (j + \alpha + \beta + v + 2)_k}.$$

Multiplying by $\frac{d^{v-1} P_n^{(\alpha, \beta)}(x)}{dx^{v-1}} \Big|_{x=-1}$ the factors in (3.14) that are independent of index k and simplifying yields

$$\frac{2^v (-1)^m (\alpha + \beta + 2j + 1) \Gamma(m + \beta + 1) \Gamma(\beta + v + 1) \Gamma(j + \alpha + \beta + 1)}{m! \Gamma(v - j + 1) \Gamma(j + \beta + 1) \Gamma(\beta + 1) \Gamma(j + \alpha + \beta + v + 2)} \frac{d^{v-1} P_n^{(\alpha, \beta)}(x)}{dx^{v-1}} \Big|_{x=-1} \\ = (-1)^{m+n+1+v} \frac{2(\alpha + \beta + 2j + 1)(\beta + v)(j + \beta + 1)_{m-j} (\beta + 1)_n (n + 1 + \alpha + \beta)_{v-1}}{m!(n + 1 - v)! (v - j)! (j + \alpha + \beta + 1)_{v+1}}.$$

The expression of $d_{v,j,n}^m(\alpha, \beta)$ then follows from the last two equations and the fact that the k -sum in (3.14) can be concisely written as a generalized hypergeometric series:

$$\sum_{k=0}^m \frac{(-m)_k (\beta + v + 1)_k (m + \alpha + \beta + 1)_k}{(\beta + 1)_k (-j + v + 1)_k (j + \alpha + \beta + v + 2)_k} = {}_4F_3 \left(\begin{matrix} 1, -m, \beta + v + 1, m + \alpha + \beta + 1 \\ v - j + 1, \beta + 1, j + \alpha + \beta + v + 2 \end{matrix}; 1 \right).$$

□




FIGURE 3.1. The magnitude plot of the Jacobi coefficients $\rho_{j,n}^{15;(2.5,1.5)}$.

We calculate the coefficients $\rho_{j,n}^{m;(\alpha, \beta)}$ with $\alpha = 2.5$ and $\beta = 1.5$ for $m = 15$ using Theorem 3.4² and show their magnitudes for $0 \leq j, n \leq 66$ in Figure 3.1. That is, the n -th column in this matrix plot corresponds to $\rho_{j,n}^{15;(2.5,1.5)}$. There are three regions of exact zeros which are indicated by solid lines. The coefficients $\rho_{j,n}^{15;(2.5,1.5)}$ are exact zeros in Region A simply because the convolution of $P_{15}^{(2.5,1.5)}$

²In this example and the examples in the remainder of this paper, the calculation is carried out in MATHEMATICA.

and $P_n^{(2.5,1.5)}$ is a polynomial of degree $n + 16$. The zeros in Region B correspond to (3.12b). Finally, we note that $m \geq 2n + 3$ in Region C and by swapping the roles of m and n we see the exact zeros in Region C, again, from (3.12b).

Remark 3.5. A convolution formula of Volterra type makes appearance in [21, p.281, Eq. (51)]:

$$k!Q_{n,k+1}^{(\alpha,\beta,0)}(y) = \int_0^y (y-t)^k P_n^{(\alpha,\beta)}(2t-1) dt$$

which implies (see [21, p.281, Eq. (53)])

$$Q_{n,k+1}^{(\alpha,\beta,0)}(x) = \frac{(-1)^n (\beta+1)_n}{n!(k+1)!} x^{k+1} {}_3F_2 \left(\begin{matrix} -n, n+\alpha+\beta+1, 1 \\ \beta+1, k+2 \end{matrix}; x \right),$$

for a non-negative integer α and $k - \alpha = 0, 1, \dots, n-1$. Straightforwardly, one can deduce

$$\int_{-1}^{x+1} P_m^{(\alpha,\beta)}(x-t) P_n^{(\alpha,\beta)}(t) dt = \frac{2(-1)^m (\beta+1)_m}{m!} \sum_{k=0}^m \frac{(-m)_k (m+\alpha+\beta+1)_k}{(\beta+1)_k} Q_{n,k+1}^{(\alpha,\beta,0)} \left(\frac{x}{2} + 1 \right),$$

which is indeed valid for any $\alpha, \beta > -1$. However, in order to obtain the expressions for the ρ -coefficients in (3.1), it would be required to consider connection formulas between the polynomials $Q_{n,k+1}^{(\alpha,\beta,0)}$ and the Jacobi polynomials, which does not seem to be an easy task.

In [34, Th. 4.8], it is shown that the coefficients $\rho_{j,n}^{m;(\alpha,\beta)}$ are symmetric up to a scaling for $j, n \geq m+1$:

$$(3.15) \quad \rho_{n,j}^{m;(\alpha,\beta)} = (-1)^{j+n} \frac{(\alpha+\beta+2n+1)(\alpha+1)_j (\beta+1)_j \left((\alpha+\beta+1)_n \right)^2}{(\alpha+\beta+2j+1)(\alpha+1)_n (\beta+1)_n \left((\alpha+\beta+1)_j \right)^2} \rho_{j,n}^{m;(\alpha,\beta)},$$

which can be shown from (3.12d) with some tedious and lengthy work. However, this symmetry property is readily seen for the symmetric Jacobi case where $\alpha = \beta$ and we show this in the next subsection. In Figure 3.1, the entries that satisfy this symmetry property are those in the lower right part that is bordered by the dashed lines.

Bateman's formula for the expansion of Jacobi polynomials with two variables is well known (see, for example, [15, Theorem 4.3.3]). In passing, we obtain the following proposition where we show the binomial-type tensor product expansion of $P_m^{(\alpha,\beta)}(x-t)$ in $P_j^{(\alpha,\beta)}(x+1)$ and $P_k^{(\alpha,\beta)}(t)$. This way, the variables x and t in a Jacobi-polynomial-based *difference kernel* [20, p. 37] become detached.

Proposition 3.6. *For the m -th degree Jacobi polynomial $P_m^{(\alpha,\beta)}(x)$,*

$$(3.16) \quad P_m^{(\alpha,\beta)}(x-t) = \sum_{k=0}^m \sum_{j=0}^{m-k} c_{m-k,j}^{m;(\alpha,\beta)} P_j^{(\alpha,\beta)}(x+1) P_k^{(\alpha,\beta)}(t),$$

where

$$(3.17) \quad c_{m-k,j}^{m,(\alpha,\beta)} = \sum_{v=k}^{m-j} \frac{(-1)^v (\alpha + \beta + 2k + 1)}{(\beta + v + 1)_{k-v} (\alpha + \beta + k + 1)_{v+1} (v-k)!} \\ \times \frac{(j + v + \alpha + 1)_{m-v-j} (m + \alpha + \beta + 1)_v (m + v + \alpha + \beta + 1)_j}{(m - v - j)! (j + \alpha + \beta + 1)_j} \\ \times {}_3F_2 \left(\begin{matrix} j - m + v, j + \alpha + 1, j + m + v + \alpha + \beta + 1 \\ j + v + \alpha + 1, 2j + \alpha + \beta + 2 \end{matrix}; 1 \right).$$

Proof. By Taylor expansion of $P_m^{(\alpha,\beta)}(x-t)$ about $t = -1$, we have

$$P_m^{(\alpha,\beta)}(x-t) = \sum_{v=0}^m \frac{(-1)^v}{v!} \frac{d^v P_m^{(\alpha,\beta)}(x+1)}{d(x+1)^v} (t+1)^v.$$

Using (3.5) and (3.10), the latter equation becomes

$$P_m^{(\alpha,\beta)}(x-t) = \sum_{v=0}^m \frac{(-1)^v}{v!} \left(\sum_{j=0}^{m-v} \gamma_{m-v,j}^{(v,0)}(\alpha, \beta) P_j^{(\alpha,\beta)}(x+1) \right) \left(\sum_{k=0}^v b_{v,k}(\alpha, \beta) P_k^{(\alpha,\beta)}(t) \right).$$

We swap the order of the k - and the v -summations and then that of the v - and the j -summations to obtain (3.16) with

$$c_{m-k,j}^{m,(\alpha,\beta)} = \sum_{v=k}^{m-j} \frac{(-1)^v}{v!} b_{v,k}(\alpha, \beta) \gamma_{m-v,j}^{(v,0)}(\alpha, \beta).$$

Substituting in the expressions of $b_{v,k}(\alpha, \beta)$ and $\gamma_{m-v,j}^{(v,0)}(\alpha, \beta)$, given by (3.11) and (3.6) respectively, leads to (3.17). \square

3.2. Symmetric Jacobi polynomials. In this subsection, we give in Corollary 3.8 the convolution coefficients $\rho_{k,n}^{m,(\alpha,\beta)}$ for the Jacobi polynomials with $\alpha = \beta$. These coefficients could be obtained from Theorem 3.4 by simply setting $\beta = \alpha$. However, a lengthy simplification is necessary in order to obtain exactly what is given in Corollary 3.8. The route we take is to obtain the explicit expressions for $\gamma_{n,k}^{(p,q)}(\alpha, \alpha)$, $b_{n,k}(\alpha, \alpha)$, and $\left. \frac{d^p}{dx^p} P_n^{(\alpha,\alpha)}(x) \right|_{x=-1}$, from which we derive $\bar{\omega}_{j,v}^{m,n}(\alpha, \alpha)$ and $d_{v,j,n}^m(\alpha, \alpha)$ using (3.13).

Lemma 3.7. *When $n - k$ is even,*

(3.18a)

$$\gamma_{n,k}^{(p,q)}(\alpha, \alpha) = \frac{(p-q)_{\frac{n-k}{2}} (\alpha + k + q + \frac{1}{2}) \Gamma(k + q + 2\alpha + 1) \Gamma(n + p + \alpha + 1) \Gamma(\frac{k+n+1}{2} + p + \alpha)}{2^{q-p} (\frac{n-k}{2})! \Gamma(k + q + \alpha + 1) \Gamma(n + p + 2\alpha + 1) \Gamma(\frac{k+n+3}{2} + q + \alpha)},$$

and

$$(3.18b) \quad \gamma_{n,k}^{(p,q)}(\alpha, \alpha) = 0$$

otherwise.

Proof. The connection coefficients in (3.7) with $\beta = \alpha$ and $\delta = \gamma$ can be found in [1, Theorem 7.1.4]:

$$(3.19) \quad a_{n,k}^{(\alpha+p,\alpha+p;\alpha+q,\alpha+q)} = \frac{(p-q)_{\frac{n-k}{2}} (q+\alpha+\frac{3}{2})_k (2(q+\alpha)+1)_k (p+\alpha+1)_n (p+\alpha+\frac{1}{2})_{\frac{k+n}{2}}}{(\frac{n-k}{2})! (q+\alpha+\frac{1}{2})_k (q+\alpha+1)_k (2(p+\alpha)+1)_n (q+\alpha+\frac{3}{2})_{\frac{k+n}{2}}},$$

when $n-k$ is even, while

$$a_{n,k}^{(\alpha+p,\alpha+p;\alpha+q,\alpha+q)} = 0,$$

when $n-k$ is odd, implied by $P_n^{(\alpha,\alpha)}(-x) = (-1)^n P_n^{(\alpha,\alpha)}(x)$ [1, Theorem 7.1.4]. Now the relation (3.9) between the a - and the γ -coefficients readily show that $\gamma_{n,k}^{(p,q)}(\alpha,\alpha) = 0$ when $n-k$ is odd. For the case where $n-k$ is even, we substitute (3.19) in (3.9) and simplify using the Legendre duplication formula [24, Eq. (5.5.5)] to obtain (3.18). \square

Corollary 3.8. *Let m and n be two positive integers with $n \geq m$ and suppose $\alpha > -1$ to be nonzero. For $\alpha = \beta$, the ρ -coefficients in the expansion (3.1) become*

$$(3.20a) \quad \rho_{j,n}^{m;(\alpha,\alpha)} = \sum_{v=\max(1,|j-n|)}^{m+1} \varpi_{j,v}^{m,n}(\alpha,\alpha) \quad \text{for } j \geq \max(m+1, n-m-1),$$

$$(3.20b) \quad \rho_{j,n}^{m;(\alpha,\alpha)} = 0 \quad \text{for } m+1 \leq j \leq n-m-2, \text{ if } n \geq 2m+3,$$

$$(3.20c) \quad \rho_{j,n}^{m;(\alpha,\alpha)} = \sum_{v=1}^j \varpi_{j,v}^{n,m}(\alpha,\alpha) + \sum_{v=j+1}^{n+1} d_{v,j,n}^m(\alpha,\alpha) \quad \text{for } 0 \leq j \leq m,$$

where

$$(3.20d) \quad \varpi_{j,v}^{m,n}(\alpha,\alpha) = \frac{2(-1)^{m+v+1} (\alpha+v)_{m+1-v} (m+2\alpha+1)_{v-1} (n+2\alpha+1)_{j-v}}{(m-v+1)! \left(\frac{n+v-j}{2}\right)!} \\ \times \frac{(-v)_{\frac{n+v-j}{2}} (j+\alpha-v+\frac{1}{2})_{\frac{n+v-j}{2}} (j+\alpha-v+1)_{n+v-j}}{(j+2\alpha+1)_j (j+\alpha+\frac{3}{2})_{\frac{n+v-j}{2}} (2j+2\alpha-2v+1)_{n+v-j}}$$

when $n+v-j$ is even and

$$(3.20e) \quad \varpi_{j,v}^{m,n}(\alpha,\alpha) = 0$$

otherwise. Here,

$$(3.20f) \quad d_{v,j,n}^m(\alpha,\alpha) = \frac{2(-1)^{m+n+1+v} (2\alpha+2j+1)(\alpha+v)(j+\alpha+1)_{m-j} (\alpha+1)_n (n+1+2\alpha)_{v-1}}{m!(n+1-v)!(v-j)!(j+2\alpha+1)_{v+1}} \\ \times {}_4F_3 \left(\begin{matrix} 1, -m, \alpha+v+1, m+2\alpha+1 \\ v-j+1, \alpha+1, j+2\alpha+v+2 \end{matrix}; 1 \right).$$

Proof. The ρ -coefficients in (3.20) inherit from (3.12).

From (3.4), we have

$$(3.21) \quad \frac{d^p}{dx^p} P_n^{(\alpha,\alpha)}(x) \Big|_{x=-1} = \frac{2^{-p} (-1)^{n+p} (p+\alpha+1)_{n-p} (n+2\alpha+1)_p}{(n-p)!}.$$

Substituting (3.18) and (3.21) into (3.13a), we have (3.20d) and (3.20e) after some algebraic simplifications.

Replacing β by α in (3.12e) gives (3.20f). \square

The following proposition instantiates the symmetry property for the symmetric Jacobi-based coefficients, which can be easily derived from Corollary 3.8.

Proposition 3.9. *For $j, n \geq m + 1$, the ρ -coefficients in (3.1) with $\beta = \alpha$ satisfy*

$$(3.22) \quad \rho_{j,n}^{m;(\alpha,\alpha)} = (-1)^{j+n} \frac{(2\alpha + 2j + 1)((2\alpha + 1)_j)^2 ((\alpha + 1)_n)^2}{(2\alpha + 2n + 1)((\alpha + 1)_j)^2 ((2\alpha + 1)_n)^2} \rho_{n,j}^{m;(\alpha,\alpha)}.$$

Proof. When $(n + v - j)$ is even, so is $(j + v - n)$ and (3.20d) gives

$$(3.23) \quad \frac{\varpi_{j,v}^{m,n}(\alpha, \alpha)}{\varpi_{n,v}^{m,j}(\alpha, \alpha)} = (-1)^{j+n} \frac{(2\alpha + 2j + 1)((2\alpha + 1)_j)^2 ((\alpha + 1)_n)^2}{(2\alpha + 2n + 1)((\alpha + 1)_j)^2 ((2\alpha + 1)_n)^2},$$

which is independent from v . If $|n - j| \leq m + 1$, (3.23) and (3.20a) imply (3.22).

If $n \geq 2m + 3$, for $m + 1 \leq j \leq n - m - 2$ we have $\rho_{j,n}^{m;(\alpha,\alpha)} = 0$, as indicated by (3.20b). Also, $j \leq n - m - 2$ suggests $n \geq j + m + 2$, in which case $\rho_{n,j}^{m;(\alpha,\alpha)} = 0$. Therefore, (3.22) holds too for $m + 1 \leq j \leq n - m - 2$.

Hence, (3.22) is true for $j, n \geq m + 1$. \square




FIGURE 3.2. The magnitude plot of the symmetric Jacobi coefficients $\rho_{j,n}^{15;(2.5,2.5)}$.

Figure 3.2 shows the magnitudes of the coefficients $\rho_{j,n}^{m;(\alpha,\alpha)}$ with $\alpha = 2.5$ and $m = 15$. The regions A, B, and C with exact zeros are inherited from those of the Jacobi-based convolution coefficients. The symmetric coefficients are bordered by the dashed lines.

3.2.1. *Gegenbauer case.* Gegenbauer polynomials $C_n^{(\lambda)}$ are symmetric Jacobi polynomials with a different normalization:

$$(3.24) \quad C_n^{(\lambda)}(x) = \frac{(2\lambda)_n}{(\lambda + \frac{1}{2})_n} P_n^{(\lambda - \frac{1}{2}, \lambda - \frac{1}{2})}(x), \quad n \geq 0,$$

with $\lambda > -\frac{1}{2}$ and $\lambda \neq 0$. If we denote by $\tilde{\rho}_{k,n}^{m;(\lambda)}$ the series coefficients of the convolution of two Gegenbauer polynomials, that is,

$$(3.25) \quad \int_{-1}^{x+1} C_m^{(\lambda)}(x-t) C_n^{(\lambda)}(t) dt = \sum_{k=0}^{m+n+1} \tilde{\rho}_{j,n}^{m;(\lambda)} C_j^{(\lambda)}(x+1),$$

the relation (3.24) gives

$$(3.26) \quad \tilde{\rho}_{j,n}^{m;(\lambda)} = \frac{(\lambda + \frac{1}{2})_j (2\lambda)_m (2\lambda)_n}{(2\lambda)_j (\lambda + \frac{1}{2})_m (\lambda + \frac{1}{2})_n} \rho_{j,n}^{m;(\lambda-1/2; \lambda-1/2)},$$

where $\rho_{j,n}^{m;(\lambda-1/2; \lambda-1/2)}$ are the coefficients given in Corollary 3.8. Combining Corollary 3.8 and (3.26) leads to the following corollary, the proof of which is omitted.

Corollary 3.10. *Let m and n be two positive integers with $n \geq m$ and suppose $\lambda > -\frac{1}{2}$ to be nonzero. The $\tilde{\rho}$ -coefficients in the expansion (3.25) can be expressed as*

$$\begin{aligned} \tilde{\rho}_{j,n}^{m;(\lambda)} &= \sum_{v=\max(1, |j-n|)}^{m+1} \tilde{\omega}_{j,v}^{m,n}(\lambda) && \text{for } j \geq \max(m+1, n-m-1), \\ \tilde{\rho}_{j,n}^{m;(\lambda)} &= 0 && \text{for } m+1 \leq j \leq n-m-2, \text{ if } n \geq 2m+3, \\ \tilde{\rho}_{j,n}^{m;(\lambda)} &= \sum_{v=1}^j \tilde{\omega}_{j,v}^{n,m}(\lambda) + \sum_{v=j+1}^{n+1} \tilde{d}_{v,j,n}^m(\lambda) && \text{for } 0 \leq j \leq m, \end{aligned}$$

where

$$\tilde{\omega}_{j,v}^{m,n}(\lambda) = \frac{(j+\lambda)(-1)^{m+v+1}(\lambda)_{v-1}(2\lambda+2v-2)_{m-v+1}(-v)_{\frac{-j+n+v}{2}}}{2(m-v+1)! \left(\frac{-j+v+n}{2}\right)! \left(\lambda + \frac{j+n-v}{2}\right)_{v+1}}$$

when $n+v-j$ is even and $\tilde{\omega}_{j,v}^{m,n}(\lambda) = 0$ otherwise. Here,

$$\begin{aligned} \tilde{d}_{v,j,n}^m(\lambda) &= \frac{2(-1)^{m+v+n+1}(j+\lambda)(2\lambda+2v-1)(2\lambda)_m(j+2\lambda+v+1)_{-j+n-2}}{m!(v-j)!(-v+n+1)!} \\ &\quad \times {}_4F_3 \left(\begin{matrix} 1, -m, m+2\lambda, \lambda+v+\frac{1}{2} \\ \lambda+\frac{1}{2}, -j+v+1, j+2\lambda+v+1 \end{matrix}; 1 \right). \end{aligned}$$

3.2.2. *Legendre case.* Legendre polynomials $P_n^{(0,0)}(x)$ are the symmetric Jacobi polynomials with $\alpha = \beta = 0$ or, equivalently, the special case of Gegenbauer polynomials with $\lambda = 1/2$. We show in the following corollary that the convolution coefficients of Legendre polynomials become significantly simpler than those of symmetric Jacobi or Gegenbauer.

Corollary 3.11. *Let m and n be two positive integers with $n \geq m$. The coefficients $\rho_{j,n}^{m;(0,0)}$ in (3.1) can be expressed as*

$$\begin{aligned} \rho_{j,n}^{m;(0,0)} &= \sum_{v=\max(1,|j-n|)}^{m+1} \omega_{j,v}^{m,n}(0,0) && \text{for } j \geq \max(m+1, n-m-1), \\ \rho_{j,n}^{m;(0,0)} &= 0 && \text{for } m+1 \leq j \leq n-m-2, \text{ if } n \geq 2m+3, \\ \rho_{j,n}^{m;(0,0)} &= \sum_{v=1}^j \omega_{j,v}^{n,m}(0,0) + \sum_{v=j+1}^{n+1} d_{v,j,n}^m && \text{for } 0 \leq j \leq m, \end{aligned}$$

where

$$(3.27a) \quad \omega_{j,v}^{m,n}(0,0) = \frac{(-1)^{m+v+1} (2j+1)(m+v-1)! (-v)_{\frac{n-j+v}{2}}}{4^v (v-1)! (m-v+1)! \left(\frac{n-j+v}{2}\right)! \left(\frac{n+j-v+1}{2}\right)_{v+1}}$$

for even $n+v-j$ and $\omega_{j,v}^{m,n}(0,0) = 0$ otherwise. Here,

$$(3.27b) \quad d_{v,j,n}^m(0,0) = \frac{\sqrt{\pi} (2j+1) v 2^{j+m-v} (-1)^{m+v+n+1} (n+v-1)! \left(\frac{-j-m+v+1}{2}\right)_{j+m} \left(\frac{j-m+v+2}{2}\right)_m}{(n-v+1)! (j+m+v)! \Gamma\left(\frac{-j+m+v+2}{2}\right) \Gamma\left(\frac{j+m+v+3}{2}\right)}.$$

Proof. Since $P_n(x) = P_n^{(0,0)}(x)$, (3.27a) can be obtained by setting $\alpha = 0$ in (3.20d) and simplifying using the Legendre duplication formula.

Setting $\alpha = 0$ in (3.20f), we have

$$d_{v,j,n}^m(0,0) = \frac{2(-1)^{v+m+n+1} v (2j+1) (n+v-1)!}{(n-v+1)! (v-j)! (j+v+1)!} {}_3F_2 \left(\begin{matrix} -m, m+1, v+1 \\ -j+v+1, j+v+2 \end{matrix}; 1 \right),$$

where the generalized hypergeometric series ${}_3F_2$ can be represented in terms of Gamma functions using the Whipple's sum [24, Eq. (16.4.7)]. Further algebraic simplification using the Legendre duplication formula gives (3.27b). \square

The following theorem shows that the symmetry property (3.15) holds for all $j, n \geq 0$ in the Legendre case, which is difficult to see directly from Theorem 3.11. Our proof employs the fact that the Legendre polynomials are L^2 orthogonal on $[-1, 1]$.

Theorem 3.12. *For any $m, n, j \geq 0$, the coefficients $\rho_{j,n}^{m;(0,0)}$ in (3.1) satisfy*

$$(3.28) \quad \rho_{j,n}^{m;(0,0)} = (-1)^{n+j} \frac{2j+1}{2n+1} \rho_{n,j}^{m;(0,0)}.$$

Proof. As indicated in (1.2), $x \in [-2, 0]$ in (3.1). By letting $y = x + 1$, we have

$$\int_{-1}^y P_m^{(0,0)}(y-t-1) P_n^{(0,0)}(t) dt = \sum_{k=0}^{m+n+1} \rho_{k,n}^{m;(0,0)} P_k^{(0,0)}(y),$$

where $y \in [-1, 1]$. Since the Legendre polynomials are L^2 orthogonal, i.e. for $j, n \geq 0$,

$$\int_{-1}^1 P_j^{(0,0)}(y) P_k^{(0,0)}(y) dy = \frac{2}{2j+1} \delta_{j,k},$$

we have

$$(3.29) \quad \int_{-1}^1 P_j^{(0,0)}(y) \int_{-1}^y P_m^{(0,0)}(y-t-1) P_n^{(0,0)}(t) dt dy = \frac{2}{2j+1} \rho_{j,n}^{m;(0,0)}$$

for $0 \leq j \leq m+n+1$.

Now, we swap the order of integration to have

$$\int_{-1}^1 P_j^{(0,0)}(y) \int_{-1}^y P_m^{(0,0)}(y-t-1) P_n^{(0,0)}(t) dt dy = \int_{-1}^1 P_n^{(0,0)}(t) \int_t^1 P_j^{(0,0)}(y) P_m^{(0,0)}(y-t-1) dy dt,$$

where $t \in [-1, 1]$. Applying the changes of variables $t = -T$ and $y = -Y$ gives

$$(3.30) \quad \int_{-1}^1 P_j^{(0,0)}(y) \int_{-1}^y P_m^{(0,0)}(y-t-1) P_n^{(0,0)}(t) dt dy \\ = (-1)^{n+j} \int_{-1}^1 P_n^{(0,0)}(T) \int_{-1}^T P_j^{(0,0)}(Y) P_m^{(0,0)}(T-Y-1) dY dT,$$

where we have used $P_n(y) = (-1)^n P_n(-y)$.

Recognizing the inner integral in (3.30) as the convolution of $P_j(T)$ and $P_m(T)$ with $T \in [-1, 1]$, we can replace it by its series representation

$$(3.31) \quad \int_{-1}^1 P_j^{(0,0)}(y) \int_{-1}^y P_m^{(0,0)}(y-t-1) P_n^{(0,0)}(t) dt dy \\ = (-1)^{n+j} \sum_{k=0}^{j+m+1} \rho_{k,j}^{m;(0,0)} \int_{-1}^1 P_n^{(0,0)}(T) P_k^{(0,0)}(T) dT = (-1)^{n+j} \frac{2}{2n+1} \rho_{n,j}^{m;(0,0)},$$

where the second equality follows from the orthogonality. Finally, (3.29) and (3.31) leads to (3.28). \square




FIGURE 3.3. The magnitude plot of the Legendre coefficients $\rho_{j,n}^{15;(0,0)}$.

The Legendre-based symmetry property (3.28) suggests extended regions of exact zeros in the magnitude plot, as seen in Figure 3.3. Now, the non-zero coefficients are confined in a tilted rectangle-shaped band surrounded by zeros in Regions A, B, and C.

3.2.3. *Chebyshev case.* Chebyshev polynomials of second kind $T_n(x)$ are symmetric Jacobi polynomials with $\alpha = \beta = -1/2$ and a different normalization:

$$(3.32) \quad P_n^{(-1/2, -1/2)}(x) = \frac{(1/2)_n}{n!} T_n(x), \quad n \geq 0.$$

Corollary 3.13. *Let m and n be two positive integers with $n \geq m$. The coefficients $\tilde{\rho}_{k,n}^m$ in the expansion*

$$\int_{-1}^{x+1} T_m(x-t)T_n(t)dt = \sum_{k=0}^{m+n+1} \tilde{\rho}_{k,n}^m T_k(x+1),$$

can be expressed as

$$(3.33a) \quad \tilde{\rho}_{j,n}^m = \sum_{v=\max(1,|j-n|)}^{m+1} \tilde{\omega}_{j,v}^{m,n} \quad \text{for } j \geq \max(m+1, n-m-1),$$

$$(3.33b) \quad \tilde{\rho}_{j,n}^m = 0 \quad \text{for } m+1 \leq j \leq n-m-2, \text{ if } n \geq 2m+3,$$

$$(3.33c) \quad \tilde{\rho}_{j,n}^m = \sum_{v=1}^j \tilde{\omega}_{j,v}^{n,m} + \sum_{v=j+1}^{n+1} \tilde{d}_{v,j,n}^m \quad \text{for } 0 \leq j \leq m,$$

where

$$(3.33d) \quad \tilde{\omega}_{j,v}^{m,n} = \frac{(-1)^m 2^{1-2v} n(-m)_{v-1} (m)_{v-1} (-v)_{\frac{n+v-j}{2}} \left(\frac{n+v-j+2}{2}\right)_{j-v-1}}{(1/2)_{v-1} \left(\frac{n+v+j}{2}\right)!}$$

for even $(n+v-j)$ and $\tilde{\omega}_{j,v}^{m,n} = 0$ otherwise. Here,

$$(3.33e) \quad \tilde{d}_{v,j,n}^m = \frac{2^{2-\delta_{0,j}} (-1)^{m+n} (-n)_{v-1} (n)_{v-1} (v-1/2)}{(v-j)!(j+v)!} {}_4F_3 \left(\begin{matrix} 1, -m, m, v+1/2 \\ 1/2, -j+v+1, j+v+1 \end{matrix}; 1 \right).$$

Proof. Equation (3.32) suggests

$$(3.34) \quad \tilde{\rho}_{j,n}^m = \frac{m!n!(1/2)_j}{(1/2)_m(1/2)_n j!} \rho_{j,n}^{m;(-1/2, -1/2)},$$

which leads to (3.33a)-(3.33c) with

$$\tilde{\omega}_{j,v}^{m,n} = \frac{m!n!(1/2)_j}{(1/2)_m(1/2)_n j!} \omega_{j,v}^{m,n}(-1/2, -1/2).$$

Setting $\alpha = -1/2$ in (3.20d) and simplifying give (3.33d).

Similarly, (3.34) implies

$$(3.35) \quad \tilde{d}_{v,j,n}^m = \frac{m!n!(1/2)_j}{(1/2)_m(1/2)_n j!} d_{v,j,n}^m(-1/2, -1/2),$$

where

$$(3.36) \quad d_{\nu,j,n}^m(-1/2, -1/2) = \frac{4(-1)^{m+n+1+\nu}(-1/2+\nu)(j+1/2)_{m-j}(1/2)_n(n)_{\nu-1}}{m!(n+1-\nu)!(\nu-j)!(j+1)_{\nu}} \\ \times {}_4F_3 \left(\begin{matrix} 1, -m, 1/2+\nu, m \\ \nu-j+1, 1/2, j+\nu+1 \end{matrix}; 1 \right)$$

for $j \geq 1$. When $j = 0$, (3.20f) gives

$$(3.37) \quad d_{\nu,0,n}^m(-1/2, -1/2) = \lim_{\alpha \rightarrow -1/2} d_{\nu,0,n}^m(\alpha, \alpha) \\ = \frac{2(-1)^{m+n+1+\nu}(\nu-1/2)(1/2)_m(1/2)_n(n)_{\nu-1}}{m!(n+1-\nu)!(\nu!)^2} {}_4F_3 \left(\begin{matrix} 1, -m, 1/2+\nu, m \\ \nu+1, 1/2, \nu+1 \end{matrix}; 1 \right).$$

By combining (3.36) and (3.37), we have

$$d_{\nu,j,n}^m(-1/2, -1/2) = \frac{2^{2-\delta_{0,j}}(-1)^{m+n+1+\nu}(-1/2+\nu)(j+1/2)_{m-j}(1/2)_n(n)_{\nu-1}}{m!(n+1-\nu)!(\nu-j)!(j+1)_{\nu}} \\ \times {}_4F_3 \left(\begin{matrix} 1, -m, 1/2+\nu, m \\ \nu-j+1, 1/2, j+\nu+1 \end{matrix}; 1 \right),$$

which, along with (3.35), yields (3.33e). □




FIGURE 3.4. The magnitude plot of the Chebyshev coefficients $\tilde{\rho}_{j,n}^{15}$.

Figure 3.4 shows the magnitudes of the coefficients $\rho_{j,n}^m$ for $m = 15$ and the region A, B, and C where $\rho_{j,n}^m$ are exactly zero. The coefficients that satisfy the symmetry are encompassed by the dashed lines.

4. LAGUERRE CASE

Laguerre polynomials parameterized by α with $\Re(\alpha) > -1$ can be defined in terms of terminating hypergeometric functions with the following commonly-used normalization

$$L_n^{(\alpha)}(x) = \frac{(1+\alpha)_n}{n!} {}_1F_1\left(\begin{matrix} -n \\ \alpha+1 \end{matrix}; x\right), \quad n \geq 0,$$

and they satisfy the orthogonality relations

$$\int_0^{+\infty} x^\alpha e^{-x} L_k^{(\alpha)}(x) L_n^{(\alpha)}(x) dx = \frac{\Gamma(\alpha+n+1)}{n!} \delta_{n,k}.$$

Laguerre polynomials with different parameters are linearly related via [4, (3.46)]

$$(4.1) \quad L_n^{(\alpha)}(x) = \sum_{k=0}^n \frac{(\alpha-\beta)_{n-k}}{(n-k)!} L_k^{(\beta)}(x),$$

and the p -th derivatives of $L_{n+p}^{(\alpha)}$ equals to $L_n^{(\alpha+p)}$ up to a sign:

$$(4.2) \quad \frac{d^p}{dx^p} L_{n+p}^{(\alpha)}(x) = (-1)^p L_n^{(\alpha+p)}(x).$$

We combine (4.1) and (4.2) to have

$$(4.3) \quad \frac{d^p}{dx^p} L_{n+p}^{(\alpha)}(x) = \sum_{k=0}^n \gamma_{n,k}^{(p,q)}(\alpha; L) \frac{d^q}{dx^q} L_{k+q}^{(\alpha)}(x), \quad \text{where } \gamma_{n,k}^{(p,q)}(\alpha; L) = \frac{(-1)^{p+q} (p-q)_{n-k}}{(n-k)!}.$$

The Laguerre representation of monomials can be found in, for example, [25, p.207]:

$$(4.4) \quad x^n = \sum_{k=0}^n b_{n,k}(\alpha) L_k^{(\alpha)}(x), \quad \text{where } b_{n,k}(\alpha) = \frac{(-n)_k \Gamma(n+\alpha+1)}{\Gamma(k+\alpha+1)} = (-1)^k n! \left. \frac{d^k L_n^{(\alpha)}}{dx^k} \right|_{x=0}.$$

It also follows from (4.2) the value of $L_n^{(\alpha)}(x)$ or that of its derivatives at $x=0$:

$$(4.5) \quad \left. \frac{d^p L_n^{(\alpha)}}{dx^p} \right|_{x=0} = (-1)^p \frac{(1+\alpha+p)_{n-p}}{(n-p)!}.$$

The Chu-Vandermonde's identity used repeatedly in the rest of this section:

$$(4.6) \quad \frac{(\alpha+\beta+2)_n}{n!} = \sum_{k=0}^n \frac{(\alpha+1)_k}{k!} \frac{(\beta+1)_{n-k}}{(n-k)!},$$

is valid for any complex numbers α and β . Observe that for $\alpha, \beta > -1$, it corresponds to (4.1) evaluated at $x=0$.

For the case of $\alpha = 0$, convolution of Laguerre polynomials $L_n^{(0)}(x)$ and $L_m^{(0)}(x)$ is sparse in the sense that only two coefficients of its $L_n^{(0)}$ -series representation are nonzero:

$$(4.7) \quad \int_0^x L_m^{(0)}(x-t)L_n^{(0)}(t)dt = -L_{m+n+1}^{(0)}(x) + L_{m+n}^{(0)}(x).$$

This well-known result can be found in, for example, [24, Eq. (18.17.2)] or [11, Eq. (7.411.4)]. There does not seem to be any similar formula addressing the cases with other values of α and this is what we present in Theorem 4.1 below.

Theorem 4.1. *Let m and n be two positive integers with $n \geq m$ and α be a complex number with $\Re(\alpha) > -1$. When $n \geq m+1$, the $\widehat{\rho}$ -coefficients in the expansion*

$$\int_0^x L_m^{(\alpha)}(x-t)L_n^{(\alpha)}(t)dt = \sum_{j=0}^{m+n+1} \widehat{\rho}_{j,n}^{m;(\alpha)} L_j^{(\alpha)}(x)$$

are given by

$$(4.8a) \quad \widehat{\rho}_{j,n}^{m;(\alpha)} = \begin{cases} -\frac{(\alpha-1)_{m+n+1-j}}{(m+n+1-j)!} & \text{for } n+1 \leq j \leq m+n+1, \\ \frac{(\alpha)_m}{m!} & \text{for } j = n, \\ 0 & \text{for } m+1 \leq j \leq n-1 \quad (\text{when } n \geq m+2), \\ \frac{(\alpha)_{n+1}}{(n+1)!} & \text{for } j = m, \\ \frac{(\alpha-1)_{m+n+1-j}}{(m+n+1-j)!} & \text{for } 0 \leq j \leq m-1. \end{cases}$$

In case of $n = m$, the $\widehat{\rho}$ -coefficients become

$$(4.9a) \quad \widehat{\rho}_{j,n}^{m;(\alpha)} = \begin{cases} -\frac{(\alpha-1)_{2m+1-j}}{(2m+1-j)!} & \text{for } m+1 \leq j \leq 2m+1, \\ \frac{(\alpha)_{m+1}}{(m+1)!} + \frac{(\alpha)_m}{m!} & \text{for } j = m, \\ \frac{(\alpha-1)_{2m+1-j}}{(2m+1-j)!} & \text{for } 0 \leq j \leq m-1. \end{cases}$$

Proof. With $a = 0$, Theorem 2.3 gives

$$(4.10a) \quad \widehat{\rho}_{j,n}^{m;(\alpha)} = \sum_{v=\max(1, j-n)}^{m+1} \gamma_{n-j+v,0}^{(j-v,j)} \left. \frac{d^{v-1} L_m^{(\alpha)}}{dx^{v-1}} \right|_{x=0} \quad \text{for } j \geq m+1$$

and

$$(4.10b) \quad \widehat{\rho}_{j,n}^{m;(\alpha)} = \sum_{v=1}^j \gamma_{m-j+v,0}^{(j-v,j)} \left. \frac{d^{v-1} L_n^{(\alpha)}}{dx^{v-1}} \right|_{x=0} + \sum_{v=j+1}^{n+1} \left(\left. \frac{d^{v-1} L_n^{(\alpha)}}{dx^{v-1}} \right|_{x=0} \sum_{p=0}^m \frac{b_{p+v,j}}{(p+v)!} \left. \frac{d^p L_m^{(\alpha)}}{dx^p} \right|_{x=0} \right) \quad \text{for } 0 \leq j \leq m,$$

where the γ -coefficients, the b -coefficients, and the derivatives of Laguerre polynomials at $x = 0$ are given by (4.3), (4.4), and (4.5), respectively. The main task now boils down to the simplification of (4.10a) and (4.10b). Our discussion branches for different ranges of j .

For $j \geq m + 1$:

Equation (4.10a) gives

$$(4.11) \quad \widehat{\rho}_{j,n}^{m;(\alpha)} = \sum_{v=\max(1,j-n)}^{m+1} \gamma_{n-j+v,0}^{(j-v,j)} \left. \frac{d^{v-1} L_m^{(\alpha)}}{dx^{v-1}} \right|_{x=0} = - \sum_{v=\max(1,j-n)}^{m+1} \frac{(-v)_{-j+n+v} (\alpha+v)_{m-v+1}}{(n-j+v)! (m-v+1)!}.$$

For $n+1 \leq j \leq m+n+1$, the summation index v in (4.11) runs from $j-n$. Making a change of variable $v \rightarrow v+j-n$ and using the fact that $(-j+n-v)_v = (-1)^v (j-n+1)_v$ and $(\alpha+j-n+v)_{m+n+1-j-v} = (-1)^{m+n+1-j-v} (-\alpha-m)_{m+n+1-j-v}$ leads to

$$\widehat{\rho}_{j,n}^{m;(\alpha)} = (-1)^{m+n+j} \sum_{v=0}^{m+n+1-j} \frac{(j-n+1)_v (-\alpha-m)_{m+n+1-j-v}}{(m+n+1-j-v)! v!},$$

from which we obtain (4.8a) by applying (4.6) and noting $(-\alpha-n-m+j+1)_{m+n+1-j} = (-1)^{m+n-j+1} (\alpha-1)_{m+n+1-j}$. In case of $m = n$, this gives (4.9a).

For $j = n$, by noting that $(-v)_v = (-1)^v v!$ we simplify (4.11) to find

$$\widehat{\rho}_{n,n}^{m;(\alpha)} = - \sum_{v=1}^{m+1} \frac{(-1)^v (\alpha+v)_{m-v+1}}{(m-v+1)!} = (-1)^m \sum_{v=1}^{m+1} \frac{(-\alpha-m)_{m-v+1}}{(m-v+1)!} = (-1)^m \sum_{v=0}^m \frac{(-\alpha-m)_v}{v!} = \frac{(\alpha)_m}{m!},$$

where the first equality follows the sign-flip trick $(\alpha+v)_{m-v+1} = (-1)^{m-v+1} (-\alpha-m)_{m-v+1}$ and the second is due to a change of variable with $m+1-v$ in place of v . The last equality is obtained by using (4.6). This proves (4.8b).

If $n \geq m+2$, it is possible that $m+1 \leq j \leq n-1$. In this case, $(-v)_{-j+n+v} = 0$, which leads to (4.8c).

For $0 \leq j \leq m$:

We denote the two v -sums in (4.10b) by S_1 and S_2 , respectively. The first sum

$$S_1 = \sum_{v=1}^j \gamma_{m-j+v,0}^{(j-v,j)} \left. \frac{d^{v-1} L_n^{(\alpha)}}{dx^{v-1}} \right|_{x=0} = - \sum_{v=1}^j \frac{(-v)_{m-j+v} (\alpha+v)_{n-v+1}}{(m-j+v)! (n-v+1)!}$$

vanishes for any $j \leq m-1$, since $(-v)_\tau = 0$ for $\tau \geq v+1$. Therefore, we only have to consider the case of $j = m$, for which

$$(4.12) \quad \begin{aligned} S_1 &= \sum_{v=1}^m \frac{(-1)^{v+1} (\alpha + v)_{n-v+1}}{(n+1-v)!} = (-1)^n \sum_{v=n-m+1}^n \frac{(-\alpha - n)_v}{v!} \\ &= (-1)^n \left(\sum_{v=0}^n - \sum_{v=0}^{n-m} \right) \frac{(-\alpha - n)_v}{v!} = \frac{(\alpha)_n}{n!} - \frac{(-1)^m (\alpha + m)_{n-m}}{(n-m)!}, \end{aligned}$$

where the first equality follows from a change of variable and the last is obtained by applying (4.6) to each of the two sums.

The second sum in (4.10b) reads

$$\begin{aligned} S_2 &= \sum_{v=j+1}^{n+1} \left(\frac{d^{v-1} L_n^{(\alpha)}}{dx^{v-1}} \Big|_{x=0} \sum_{p=0}^m \frac{b_{p+v,j}}{(p+v)!} \frac{d^p L_m^{(\alpha)}}{dx^p} \Big|_{x=0} \right) \\ &= \sum_{v=j+1}^{n+1} \left(\frac{(-1)^{v-1} (\alpha + v)_{n-v+1}}{(n-v+1)!} \sum_{p=0}^m \frac{(-1)^p (-p-v)_j \Gamma(p + \alpha + v + 1) (p + \alpha + 1)_{m-p}}{(p+v)! \Gamma(j + \alpha + 1) (m-p)!} \right) \\ &= (-1)^{m+n+j} \sum_{v=j+1}^{n+1} \left(\frac{(-\alpha - n)_{n-v+1}}{(n-v+1)!} \sum_{p=0}^m \frac{(\alpha + j + 1)_{p+v-j} (-\alpha - m)_{m-p}}{(p+v-j)! (m-p)!} \right), \end{aligned}$$

where, to obtain the last equality, we have used the identities $(\alpha + v)_{n-v+1} = (-1)^{n-v+1} (-\alpha - n)_{n-v+1}$ and $(p + \alpha + 1)_{m-p} = (-1)^{m-p} (-\alpha - m)_{m-p}$. Making the changes of variables $v \rightarrow n - v + 1$ and $p \rightarrow m - p$, we have

$$\begin{aligned} S_2 &= (-1)^{m+n+j} \sum_{v=0}^{n-j} \left(\frac{(-\alpha - n)_v}{v!} \sum_{p=0}^m \frac{(j + \alpha + 1)_{m+n+1-j-v-p} (-\alpha - m)_p}{(m+n+1-j-v-p)! p!} \right) \\ &= (-1)^{m+n+j} \sum_{v=0}^{n-j} \left[\frac{(-\alpha - n)_v}{v!} \left(\sum_{p=0}^{m+n+1-j-v} - \sum_{p=m+1}^{m+n+1-j-v} \right) \frac{(j + \alpha + 1)_{m+n+1-j-v-p} (-\alpha - m)_p}{(m+n+1-j-v-p)! p!} \right]. \end{aligned}$$

By virtue of (4.6), the first p -summation in the last equation simplifies to $\frac{(j-m+1)_{m+n+1-j-v}}{(m+n+1-j-v)!}$, which vanishes for $0 \leq j \leq m-1$ and equals 1 for $j = m$. Now we change the variable $p \rightarrow p + m + 1$ in the second p -summation to obtain

$$\begin{aligned} S_2 &= (-1)^{m+n+j} \sum_{v=0}^{n-j} \left[\frac{(-\alpha - n)_v}{v!} \left(\delta_{j,m} - \sum_{p=0}^{n-j-v} \frac{(j + \alpha + 1)_{n-j-p-v} (-\alpha - m)_{m+1+p}}{(n-v-j-p)! (m+1+p)!} \right) \right] \\ &= (-1)^{m+n+j} \left[\sum_{v=0}^{n-j} \frac{(-\alpha - n)_v}{v!} \delta_{j,m} - \sum_{p=0}^{n-j} \left(\frac{(-\alpha - m)_{m+1+p}}{(m+1+p)!} \sum_{v=0}^{n-j-p} \frac{(j + \alpha + 1)_{n-j-p-v} (-\alpha - n)_v}{(n-v-j-p)! v!} \right) \right], \end{aligned}$$

where we have swapped the order of v - and p -sums. Now, both the v -sums can be simplified using (4.6):

$$(4.13) \quad S_2 = (-1)^{m+n+j} \left[\frac{(-\alpha - n + 1)_{n-j}}{(n-j)!} \delta_{j,m} - \sum_{p=0}^{n-j} \frac{(j-n+1)_{n-j-p} (-\alpha - m)_{m+1+p}}{(n-j-p)! (m+1+p)!} \right]$$

$$= (-1)^{m+n+j} \left[(-1)^{n-j} \frac{(\alpha + j)_{n-j}}{(n-j)!} \delta_{j,m} - \sum_{p=0}^{n-j} \frac{(j-n+1)_p (-\alpha - m)_{m+1+n-j-p}}{p! (m+1+n-j-p)!} \right],$$

where we have used the sign-flip trick for the first term in the brackets and changed variable $p \rightarrow n - j - p$ in the p -sum.

When $j = m = n$, only the zeroth summand is left for the p -sum in (4.13):

$$S_2 = (-1)^m \left[1 - \frac{(-\alpha - m)_{m+1}}{(m+1)!} \right] = (-1)^m + \frac{(\alpha)_{m+1}}{(m+1)!},$$

which, combined with (4.12) for $m = n$, gives (4.9b). For all other cases, the upper limit of the p -sum in (4.13) can be bumped to $m + n + 1 - j$ as $(j - n + 1)_p$ vanishes for $p \geq n - j + 1$. Simplifying the sum by (4.6), followed by using the sign-flip trick again, we finally obtain

$$S_2 = \frac{(-1)^m (\alpha + m)_{n-m}}{(n-m)!} \delta_{j,m} + \frac{(\alpha - 1)_{m+n+1-j}}{(m+n+1-j)!},$$

which, together with (4.12), yields (4.8d), (4.8e), and (4.9c). \square

Remark 4.2. When $\alpha = 0$, $\widehat{\rho}_{j,n}^{m;(0)}$ given by (4.8) becomes zero for any $0 \leq j \leq m + n - 1$, while $\widehat{\rho}_{m+n+1,n}^{m;(0)} = -1$ and $\widehat{\rho}_{m+n,n}^{m;(0)} = 1$, which recovers (4.7). Same for $\widehat{\rho}_{j,m}^{m;(0)}$ given by (4.9).

Remark 4.3. When $\alpha = 1$, $\widehat{\rho}$ -coefficients also enjoy sparsity, suggested by Theorem 4.1:

$$\int_0^x L_m^{(1)}(x-t) L_n^{(1)}(t) dt = -L_{m+n+1}^{(1)}(x) + L_n^{(1)}(x) + L_m^{(1)}(x).$$




FIGURE 4.5. The magnitude plot of the Laguerre coefficients $\widehat{\rho}_{j,n}^{15;(\alpha)}$ for $0 \leq n \leq 50$.

Figure 4.5 shows the magnitudes of the coefficients $\widehat{\rho}_{j,n}^{m;(\alpha)}$ with $m = 15$ for (a) $\alpha = 0$, (b) $\alpha = 1$, and (c) $\alpha = 2.5$. The n -th column in this matrix plot corresponds to $\rho_{j,n}^{15;(\alpha)}$. Panes (a) and (b) confirm Remarks 4.2 and 4.3, respectively. With $\alpha = 2.5$, the plot in pane (c) is representative for a general case where three regions of exact zeros are indicated by solid lines. The exact zeros in Region A is again due to the fact that the convolution of $L_{15}^{(2.5)}$ and $L_n^{(2.5)}$ is a polynomial of degree $n + 16$, while the zeros in Region B corresponds to (4.8c). The zeros in region C are also due to (4.8c) but with the roles of n and m exchanged.

5. CLOSING REMARKS

In this paper, we have derived the explicit formulas for the coefficients in the series representation for the convolution of the elements in a polynomial sequence. Particularly, the results are significantly simplified when the polynomial sequence is formed by classical orthogonal polynomials of Jacobi- or Laguerre families.

As seen from the magnitude plots in Section 3, many of the ρ -coefficients, though they are non-zero in an exact sense, can be deemed as zeros in floating point arithmetic due to their tiny magnitudes. An exciting extension of the results in this paper is the investigation of the asymptotic behavior of the ρ -coefficients for large j and n using the newly derived explicit formulas. The asymptotics will help us identify via (1.8) the entries of R that can be safely zeroed in numerical computation and, therefore, enable a faster construction of the convolution matrix R .

On the other hand, the explicit formulas for the convolution coefficients may also reveal the numerical rank of the convolution matrix R . The low rank property of R or its subparts, if any, could lead to potential speed-up in either construction of R or fast algorithms for convolution. We save these possibilities for a future work.

Acknowledgment We would like to thank Erik Koelink for sharing his insights on the convolution product of two Jacobi polynomials and Alfredo Deaño and Walter Van Assche for useful discussions. We are also grateful to the two anonymous referees for their careful reading and suggestions.

REFERENCES

1. George E Andrews, Richard Askey, and Ranjan Roy, *Special functions*, Cambridge University Press, Cambridge, 1999.
2. I. Area, E. Godoy, A. Ronveaux, and A. Zarzo, *Solving connection and linearization problems within the askey scheme and its q -analogue via inversion formulas*, J. Comput. Appl. Math. **133** (2001), 151–162.
3. Richard Askey, *Orthogonal polynomials and special functions*, vol. 21, SIAM, 1975.
4. Richard Askey and James Fitch, *Integral representations for jacobi polynomials and some applications*, J. Math. Anal. Appl. **26** (1969), no. 2, 411–437.
5. Kendall Atkinson, *The numerical solution of integral equations of the second kind*, Cambridge University Press, 1997.

6. H. Brunner and P. J. van der Houwen, *The numerical solution of volterra equations*, Elsevier, 1986.
7. W. Bryc, W. Matysiak, R. Szwarz, and J. Wesolowski, *Projection formulas for orthogonal polynomials*, arXiv:math/0606092v2, 2007.
8. Steven B Damelin and Willard Miller Jr, *The mathematics of signal processing*, Cambridge University Press, 2012.
9. Dean G Duffy, *Green's functions with applications*, CRC Press, 2015.
10. David Forsyth and Jean Ponce, *Computer vision: A modern approach*, Prentice Hall, 2011.
11. Izrail Solomonovich Gradshteyn and Iosif Moiseevich Ryzhik, *Table of integrals, series, and products*, Academic press, 2014.
12. Thomas Hagstrom, *Radiation boundary conditions for the numerical simulation of waves*, Acta Numerica **8** (1999), 47–106.
13. Rudolf Hilfer, *Applications of fractional calculus in physics*, World Scientific, 2000.
14. Robert V Hogg, Joseph McKean, and Allen T Craig, *Introduction to mathematical statistics*, 7th ed., Pearson Education, 2013.
15. Mourad E H Ismail, *Classical and quantum orthogonal polynomials in one variable*, Encyclopedia of Mathematics and its Applications, vol. 98, Cambridge University Press, Cambridge, 2009.
16. Mourad E.H. Ismail, Erik Koelink, and Pablo Román, *Generalized burchnell-type identities for orthogonal polynomials and expansions*, SIGMA **14** (2018), no. 072, 24 pages.
17. H.T. Koelink and J. Van der Jeugt, *Convolutions for orthogonal polynomials from lie and quantum algebra representations*, SIAM J. Math. Anal. **29** (1998), 794–822.
18. Wolfram Koepf, *Representations of orthogonal polynomials.*, J. Comput. Appl. Math. **90** (1998), no. 1, 57–94.
19. S. Lewanowicz, *The hypergeometric functions approach to the connection problem for the classical orthogonal polynomials*, Technical Report, 1998.
20. Peter Linz, *Analytical and numerical methods for volterra equations*, SIAM, 1985.
21. Yudell L. Luke, *The special functions and their approximations*, vol. I, Academic Press, New York-London, 1969.
22. Pascal Maroni, *Semi-classical character and finite-type relations between polynomial sequences*, Appl. Numer. Math. **31** (1999), no. 3, 295–330.
23. Pascal Maroni and Zélia da Rocha, *Connection coefficients between orthogonal polynomials and the canonical sequence: an approach based on symbolic computation*, Numer. Algorithms **47** (2008), no. 3, 291–314.
24. Frank W J Olver, Daniel W Lozier, Ronald F Boisvert, and Charles W Clark, *Nist handbook of mathematical functions*, Cambridge Univeristy Press, 2010.
25. Earl D Rainville, *Special functions*, Chelsea Publishing Co., Bronx, N.Y., 1971.
26. Jorge Sánchez-Ruiz, Pedro López Artés, Andrei Martínez-Finkelshtein, and Jesús Dehesa, *General linearization formulae for products of continuous hypergeometric-type polynomials*, J. Phys. A-Math. Gen. **32** (1999), no. 42, 7345.
27. Eduardo D Sontag, *Mathematical control theory: Deterministic finite dimensional systems*, Springer, 2013.
28. Ivar Stakgold and Michael J Holst, *Green's functions and boundary value problems*, John Wiley & Sons, 2011.
29. Gábor Szegő, *Orthogonal polynomials*, American Mathematical Society, 1939.
30. Daniel D. Tcheutia, *On connection, linearization and duplication coefficients of classical orthogonal polynomials*, Ph.D. thesis, Kassel: Universitätsbibliothek, 2014.
31. Lloyd N. Trefethen, *Approximation theory and approximation practice*, SIAM, Philadelphia, 2013.
32. Claus Wagner, Thomas Hüttl, and Pierre Sagaut, *Large-eddy simulation for acoustics*, Cambridge University Press, 2007.
33. Jet Wimp, *Connection coefficients, orthogonal polynomials and the wz-algorithms.*, Numer. Algorithms **21** (1999), no. 1-4, 377–386.
34. Kuan Xu and Ana F Loureiro, *Spectral approximation of convolution operators*, SIAM J. Sci. Comput. **40** (2018), no. 4, A2336–A2355.

35. A. Zarzo, I. Area, E. Godoy, and A. Ronveaux, *Results for some inversion problems for classical continuous and discrete orthogonal polynomials*, J. Phys. A: Math. Gen. **30** (1997), no. 3, L35–L40.

Ana F Loureiro: SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCES, UNIVERSITY OF KENT, SIBSON BUILDING, CT2 7FS, U.K.

E-mail address: `A.Loureiro@kent.ac.uk`

Kuan Xu: SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, 96 JINZHAI ROAD, HEFEI, 230026, ANHUI, CHINA

E-mail address: `kuanxu@ustc.edu.cn`