

Kent Academic Repository

Full text document (pdf)

Citation for published version

Lawrence, Andrew D. and Nemoto-Smith, Emi H and Deery, Evelyne and Baker, Joseph A. and Schroeder, Susanne and Brown, David G. and Tullet, Jennifer M.A. and Howard, Mark J. and Brown, Ian R. and Smith, Alison G. and Boshoff, Helena I. and Barry, Clifton E. and Warren, Martin J. (2018) Construction of Fluorescent Analogs to Follow the Uptake and Distribution

DOI

<https://doi.org/10.1016/j.chembiol.2018.04.012>

Link to record in KAR

<http://kar.kent.ac.uk/67083/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Construction of fluorescent analogues to follow the uptake and distribution of cobalamin
(vitamin B₁₂) in bacteria, worms and plants.

Andrew D. Lawrence^{1,2} Emi Nemoto-Smith^{1,2,3}, Evelyne Deery¹, Joseph A. Baker¹, Susanne Schroeder¹, David G. Brown¹, Jennifer M. A. Tullet¹, Mark J. Howard¹, Ian R. Brown¹, Alison G. Smith⁴, Helena I. Boshoff³, Clifton E. Barry III³, Martin J. Warren¹

¹School of Biosciences, University of Kent, Canterbury, Kent, CT2 7NJ, UK.

²Contributed equally to this work

³National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20850, USA

⁴Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA, UK.

*Lead Contact: Martin J Warren; email m.j.warren@kent.ac.uk

SUMMARY

Vitamin B₁₂ is made by only certain prokaryotes yet is required by a number of eukaryotes such as mammals, fish, birds, worms and Protista, including algae. There is still much to learn about how this nutrient is trafficked across the domains of life. Herein, we describe ways to make a number of different corrin analogues with fluorescent groups attached to the main tetrapyrrole-derived ring. A further range of analogues were also constructed by attaching similar fluorescent groups to the ribose ring of cobalamin, thereby generating a range of complete and incomplete corrinoids to follow uptake in bacteria, worms and plants. By using these fluorescent derivatives we were able to demonstrate that *Mycobacterium tuberculosis* is able to acquire both cobyrinic acid and cobalamin analogues, that *Caenorhabditis elegans* takes up only the complete corrinoid, and that seedlings of higher plants such as *Lepidium sativum* are also able to transport B₁₂.

INTRODUCTION

The cobamides encompass a group of closely structurally related nutrients, cofactors and coenzymes that harbour a cobalt-containing corrin ring, and are often loosely referred to as vitamin B₁₂ (Figure 1)(Renz, 1999; Warren, et al., 2002). These molecules vary in the nature of the upper ligand attached to the cobalt and the character of the lower nucleotide loop. Cobalamin, for instance, contains dimethylbenzimidazole as the base in the nucleotide loop but this is replaced with adenine in pseudocobalamin (Figure 1) (Degnan, et al., 2014; Maggio-Hall and Escalante-Semerena, 2003). In the biologically active forms of cobalamin the cobalt ion is normally either adenosylated (adenosylcobalamin) or methylated (methylcobalamin) (Banerjee and Ragsdale, 2003) whilst vitamin B₁₂ represents the cyanolated derivative of cobalamin that is produced during the commercial extraction and isolation of the nutrient (Martens, et al., 2002). However, the term B₁₂ is frequently used to refer to all forms of cobalamin.

Cobamides are produced exclusively by certain prokaryotes (Roth, et al., 1996). Apart from cobalamin and pseudocobalamin, further variants with shorter nucleotide chain lengths and/or alternative nucleotide bases are synthesized by different bacteria (Degnan, et al., 2014; Krautler, et al., 2003; Renz, 1999). Cobamides are made via either an anaerobic or aerobic biochemical pathway, both involving around thirty enzyme-mediated steps (Blanche, et al., 1995; Deery, et al., 2012; Moore, et al., 2013; Warren, et al., 2002). The key steps of the aerobic pathway are shown in Figure 1, which concisely depicts contraction of the tetrapyrrole ring, insertion of the cobalt ion and building of the nucleotide loop. Variations in the construction of the lower nucleotide loop, mediated by different substrate preferences for bases and linkers (Figure 1) during its biosynthesis, give rise to the assorted cobamides that are found in nature (Chan and Escalante-Semerena, 2011; Crofts, et al., 2013; Krautler, et al., 2003; Maggio-Hall and Escalante-Semerena, 2003; Zayas, et al., 2007). Many prokaryotic microorganisms (Allen and Stabler, 2008; Degnan, et al., 2014; Gray and Escalante-Semerena, 2009) and microbial eukaryotes (Croft, et al., 2005; Helliwell, et al., 2016) rely on other producers of cobamides, where both bacteria and archaea can be counted, and therefore have to recover the nutrient from the environment they inhabit. Such salvage systems often involve remodelling of the lower nucleotide loop to produce the corrin variant that suits the new host.

In this respect bacteria, as well as archaea, can be divided into a range of classes with regards to their B₁₂ requirements. There are bacteria that make their own cobalamin, those that are auxotrophic for the nutrient, those that require cobalamin for some non-essential processes and those that do not require the nutrient at all. In this way the nutrient plays an important role

in bacterial ecology, especially in complex microbial communities (Degnan, et al., 2014; Helliwell, et al., 2016). However, little is known about the movement of cobamides in such communities, whether it is specifically shared or exchanged in mutualistic or symbiotic relationships.

Bacteria are also the source of cobalamin for a large number of eukaryotes that require the nutrient for either specific methylation or isomerisation reactions. For instance, approximately 50% of all microalgal species require cobalamin for growth, and it has been shown that bacteria can provide this compound in course of a symbiotic exchange for photosynthate (Croft, et al., 2005). Similarly, enteric bacteria are the source of cobalamin (Roth, et al., 1996) in ruminants such as cattle and, as dairy products and meat represent a major form of human dietary cobalamin, are therefore also responsible for most of the cobalamin found in humans (Brito, et al., 2015; Girard, et al., 2009). Other eukaryotes, including nematode worms (Yilmaz and Walhout, 2014), fish (Greibe, et al., 2012) and birds (Zaman and Zak, 1989) have to acquire cobalamin as part of their diets from B₁₂-producing bacteria. Within eukaryotic cells cobalamin is further compartmentalised, with methylcobalamin found in the cytoplasm associated with methionine synthase and adenosylcobalamin located within the mitochondria with its cognate enzyme methylmalonyl CoA mutase (Gherasim, et al., 2013; Watkins and Rosenblatt, 2013). There is the added complication of translocation through the body of multicellular organisms, requiring some form of vascular transport.

Therefore, from their prokaryotic origins, cobamides such as cobalamin undergo a complex journey through many different domains and kingdoms of life, a journey that involves scavenging, recycling, modification and compartmentalisation. In terms of understanding the role played by cobamides in ecology, health and disease there is a need to devise sensitive methods to follow the nutrient on this biological voyage. Following the movement of cobamides has proved difficult as the nutrient is normally required in small quantities. One way of following cobamides along their transformation and transportation chain is to use fluorescent derivatives coupled to the cobalamin molecule to allow its detection via fluorescence microscopy (Lee and Grissom, 2009; Smeltzer, et al., 2001). The most common of these fluorescent cobalamin derivatives involve attachment of a fluorophore to either the cobalt ion or to the ribose ring of the lower nucleotide. However, attachment to the cobalt ion involves modifications that are usually light sensitive whereas conjugation to the ribose can be easily removed through nucleotide loop remodelling when such molecules are taken up by bacteria. Attachment of fluorophores to the side chains of the corrin macrocycle have also been reported. However, this attachment requires the hydrolysis of one of the amide side chain and the separation of a range of derivatives prior to the modification. This is low yielding process. Alternatively, a fluorophore could also be attached to the carboxylic acid side chain of cobyrinic acid, but this

would prevent the molecule from being converted into a cobalamin analogue in vivo. Herein, we describe ways to attach fluorophores directly to the C5 position of the corrin ring according to our understanding of how the corrin ring is constructed. We also report on which of these C5-modified corrin variants, together with some ribose-modified fluorophores, are taken up by a range of different organisms.

RESULTS

Synthesis of C5-allyl HBA. The transformation of uroporphyrinogen III into a corrin involves, amongst other events, the addition of 8 S-adenosyl-L-methionine (SAM) derived methyl groups, the last of which is added to the C5 position (Deery, et al., 2012; Warren, et al., 2002). We hypothesized that it may be possible to use SAM analogues, with extended chain lengths, to alkylate the C5 position (Figure 2). This requires the isolation of precorrin-7, the substrate for the final methyltransferase, CobL. We recently reported the construction of a recombinant *E. coli* strain from which precorrin-7 can be purified (Deery, et al., 2012). Methylation of precorrin-7 by CobL generates precorrin-8, which is isomerised by CobH to generate hydrogenobyric acid (HBA). We sought to modify the C5 position by employing an allyl analogue of SAM (Dalhoff, et al., 2006; Dalhoff, et al., 2006) with a view to making the respective C5-modified HBA derivative (C5-allyl HBA) (Figure 2).

When precorrin-7 was incubated with CobL, CobH and allyl-SAM the colour of the solution changed from yellow to orange consistent with the expected enzymatic transformation. The observed shift in the UV-visible spectrum of the starting material with a maximum absorption at 400 nm to a new species with maxima at 328, 520 and 555 nm is characteristic of the formation of a HBA-like molecule. Analysis of the reaction mixture by HPLC-MS identified a new compound with m/z 907.4 (Figure S1). The observed increase in mass and change in UV-visible spectrum confirms the addition of an allyl group to the macrocycle and indicates that migration of the C11 methyl group to C12 has occurred.

The yield of allyl-HBA obtained from this initial reaction was less than 50%, with the other major product being allyl-precorrin-8. The presence of allyl-precorrin-8 suggested that CobH is the rate limiting step in this reaction sequence. We considered that the lower activity of CobH could be due to spatial constraints resulting from the larger substrate and therefore we decided to make a number of structure-based (Shipman, et al., 2001) mutations that would introduce more space into the active site, especially around the region of the protein where the C5 position of the substrate is found. One of these mutations, T85A, did indeed significantly improve the yield, such that the C5-allyl-HBA was obtained in greater than 85% yield.

The structure of the product was further confirmed as being C5-allyl-HBA by X-ray crystallography. This was achieved by determining the structure of the CobH^{T85A}-C5-allyl-HBA product complex at 1.6 Å resolution (Figure 3, Table S1). The extended alkyl group at the C5

position is clearly seen in the structure. Although the T85A mutation of CobH does increase the size of the active site around the C5 position, the allyl group is still constrained, suggesting that further enhancement could be afforded with more rational engineering. The PDB coordinates are deposited under the code 5N0G.

We also explored the synthesis of C5-propargyl-HBA using a propargyl SAM derivative (Dalhoff, et al., 2006). To achieve this outcome it was necessary to prepare the respective SAM analogue in situ from Se-propargyl-L-homocysteine using a modified SAM synthetase (I117A mutant of hMAT 2A) as the propargyl-SAM analogue is not stable at physiological pH (Wang, et al., 2013; Wang, et al., 2014). Incubation of precorrin-7 with Se-propargyl-Hcy, Mg-ATP, hMAT 2A I117A, CobL, CobH^{T85A}, resulted in the same characteristic shift in the UV-visible spectrum associated with the formation of a HBA like chromophore. Analysis of the purified product by HPLC-MS confirmed the addition of a propargyl group with an observed m/z of 905.4 (Figure S1). This approach demonstrated that a number of C5-derivatives could be made using activated SAM variants. The propargyl-compounds have the advantage of being able to be used with click chemistry approaches for further functionalisation.

Conversion of the C5-allyl-HBA to its cobyric acid form. The conversion of HBA to HBA a,c-diamide (HBAD) is mediated by CobB, which amidates the acetic acid side chains attached to the C2 and C7 positions of the corrin framework (Figure 2) (Crouzet, et al., 1990). Incubation of CobB with C5-allyl-HBA, in the presence of glutamine and Mg-ATP, resulted in the production of C5-allyl-HBAD. The product of the reaction was found to have a shorter retention time on the HPLC column and a 2-mass unit decrease in comparison to the starting material, all of which is consistent with the amidation of the a and c sidechains.

In the aerobic biosynthesis of cobalamin, HBAD undergoes cobalt chelation and adenosylation prior to the completion of the amidation reactions by CobQ (Blanche, et al., 1995). However, the light sensitive nature of the adenosylated intermediate make this a difficult compound with which to do further modifications. We thus sought a way to by-pass the metal-insertion stage of the pathway in order to complete the amidation of HBAD to generate hydrogenobyric acid (Hby), the cobalt-free analogue of cobyric acid. This was achieved by using the CobQ from *Allochromatium vinosum*, an organism that was shown to accumulate cobalt-free corrinoids when grown in the absence of cobalt (Koppenhagen, et al., 1973). The purified recombinant *A. vinosum* CobQ, in the presence of glutamine and Mg-ATP, did indeed catalyse the amidation of the b, d, e and g side chains of HBAD, to generate Hby. Similarly, the enzyme also accepted the C5-allyl-HBAD as a substrate and converted it into C5-allyl Hby (Figure 2),

whose structure was confirmed by HPLC-MS (m/z 901) (Figure S1) and 2D heteronuclear NMR spectroscopy (Figure S2 and Table S2).

Transformation of the C5-allyl HBA analogues into cobyric acid analogues. The allyl-group was modified to a primary amine through a thiol-ene reaction by reacting the C5-allyl-Hby with cysteamine in the presence of the radical initiator VA-044 (2,2'-Azobis[2-(2-imidazolin-2-yl)propane]dihydrochloride) (Figure 2). The reaction product, C5-(2-thiopropyl-1-aminoethane (TPEA))-Hby, was confirmed by HPLC-MS (Figure S1). Cobalt was inserted chemically using CoCl_2 in dilute aqueous ammonia at elevated temperature. The reaction was quenched with KCN and the product was purified by RP18 chromatography. The structure of the C5-(TPEA)-cobyric acid was confirmed by HPLC-MS (Figure S1) and NMR (Figure S3 and Table S2). Together, all of these steps demonstrate how a combination of in vitro enzyme-mediated transformations coupled with chemical approaches allow the construction of a regioselectively derivatised form of cobyric acid (Figure 2).

C5 Fluorophore attachment. To attach the fluorophores BODIPY® TR-X (BoD) and Oregon Green® 514 (OG) to the terminal amine of the C5 extension, the amine was reacted with the N-hydroxysuccinimide ester of the fluorophore (Figure 2). All of the products were analysed by HPLC-MS and purified by semi-preparative HPLC. As a result it was possible to generate both BoD and OG C5-linked forms of cobyric acid (Figure 4). These compounds were further characterised in terms of their functionality and uptake in biological systems as described below.

Synthesis of ribose linked analogues. The ribose linked analogues were synthesised by carbonyldiimidazole coupling of ethylene diamine to the C5-OH of the ribose moiety of vitamin B₁₂, then the primary amine was linked to the N-hydroxysuccinimide ester of either OG or BoD (Figure 4). These were made following previously published methods for the construction of ribose-linked analogues of cobalamin (Horton, et al., 2003; McEwan, et al., 1999). The products were analysed by HPLC-MS, and purified by semi-preparative HPLC. Making both fluorescently labelled cobyric acid and cobalamin derivatives provided the opportunity to compare their respective activities in terms of their ability to be transported into biological systems and ability to act as functionalised nutrients.

Biological activity of the corrin analogues. The biological activity of the C5-cobyric acid analogues as well as the ribose-linked derivatives was investigated initially by comparing their effect in a B₁₂-dependent microbial bioassay (Raux, et al., 1996). In this case, an agar plate embedded with B₁₂-dependent reporter strain of *Salmonella enterica* (AR3612; *cysG*, *metE*)

was used to detect growth around an application point, where the size of any resulting growth circle can be measured against growth observed with known amounts of B₁₂. With the microbial bioassay we looked at the efficacy of the C5-(TPEA) cobyric acid, the BoD and OG C5-cobyric acid as well as the BoD and OG ribose analogues of cobalamin to support growth of the strain. The observed growth circles can be seen in Figure 5. We were initially surprised to see that the C5-(TPEA)-cobyric acid and the two fluorescently labelled versions of cobyric acid and cobalamin all promoted growth, albeit they needed to be added at much higher concentrations than cyanocobalamin. The requirement for a higher concentration of the analogue not only likely reflects poorer molecular recognition by the uptake mechanism but also recognition by methionine synthase. The cobalamin-derivatives were more active than the C5-analogues, and the OG derivatives were more active than the corresponding BoD compounds. The reduced efficacy of the cobyric acid analogues in comparison to the cobalamin analogues is probably because the cobyric acid intermediates have to be converted into cobalamin forms before growth can occur.

Uptake of analogues into bacteria. We next investigated the uptake of the fluorescent forms of cobyric acid and cobalamin into *E. coli* BL21 (DE3). The BL21 strain has a mutation in *btuB*, which encodes the outer membrane transport component of the cobalamin uptake system (Studier, et al., 2009). To overcome this limitation, the BL21 cells were transformed with *pLysS-btuB* and induced overnight with IPTG at which point 1 μ M of the OG analogues were added. The bacteria were then imaged by widefield fluorescence microscopy and the resulting images demonstrated that whilst *E. coli* is able to take up both OG fluorescent analogues, significantly greater fluorescence was observed with the cobalamin derivative suggesting that it is taken up more efficiently.

For the BoD analogues, *E. coli* OP50 cells were transformed with a plasmid harbouring both *btuB* and *btuF* (*pET-BAD-btuB-btuF*), where *btuF* encodes the periplasmic cobalamin binding protein of the B₁₂ transporter. These were cloned in order to ensure maximal uptake of the corrin analogues, which were added to a final concentration of 1 μ M at the same time as the cells were induced. Cells were incubated separately with either C5-BoD-cobyric acid, ribose-BoD-cobalamin or the unattached BoD fluorophore. As judged by the internal fluorescence of the bacteria, both the C5-BoD and ribose-BoD analogues were seen to accumulate within the cells, indicating that these compounds can be used to monitor transport into *E. coli* (Figure 5). No internal fluorescence was observed with the unattached BoD fluorophore. The significance of using *E. coli* OP50 is that this strain can also be used as a source of food for nematodes such as *C. elegans*.

The BoD analogues of cobalamin were also incubated with *E. coli* BL21 (DE3) and JM109 cells. No fluorescence was observed with the BL21(DE3) cells indicating that the defective *btuB* gene prevent uptake of the analogue even at high (1 μ M) concentrations. In contrast, uptake of the C5-BoD-cobyric acid analogue into JM109 cells was detectable at levels of 10 nM, whilst the ribose-BoD-cobalamin analogues was detectable at a concentration of 100 nM (Figure S4). These results demonstrate that the uptake of the analogues is dependent upon the presence of a functional *BtuB* and that there is some variance between the analogues in regard to molecular recognition by the uptake mechanism.

There is always the possibility that the fluorescently labelled compounds undergo breakdown within the cell. To investigate the possibility of the breakdown of the analogues, we extracted both the ribose-OG-cobalamin and the C5-BoD-cobyric acid analogue from *E. coli* that had been grown in their presence. In both cases we were able to extract the analogues intact from the cells after several days growth. No evidence for degradation of the compounds was observed (Figure S5)

We also investigated the uptake of the BoD analogues into *Mycobacterium tuberculosis*. Although *M. tuberculosis* has a B_{12} pathway it is missing *CobF* (Gopinath, et al., 2013; Rodionov, et al., 2003) and cannot make its own cobalamin. Moreover, *M. tuberculosis* has a different B_{12} uptake mechanism, employing *BacA* as a transporter (Gopinath, et al., 2013). We investigated B_{12} uptake into a B_{12} -dependent strain of *M. tuberculosis*, H37Rv, which carries a mutation in *metE* (H37Rv-*metE*) (Warner, et al., 2007). After growth in Sauton's defined medium, supplemented with either the C5-BoD-cobyric acid or the ribose-BoD-cobalamin, the cells were found to be fluorescent after 24 and 48 hours respectively using fluorescence microscopy (Figure 5). It is clear, therefore, that *M. tuberculosis* can take up analogues of both coobyric acid and cobalamin and that both of these can support growth.

B_{12} uptake into *C. elegans*. Several recent papers have demonstrated that a free-living nematode *C. elegans* represents a good eukaryotic model system to study B_{12} as a nutrient (Bito, et al., 2013; Bito, et al., 2017; Watson, et al., 2014). Like humans, *C. elegans* requires B_{12} for the activities of both B_{12} -dependent methionine synthase and methylmalonyl CoA mutase. Thereby, B_{12} deficiency causes infertility, retarded growth and reduced lifespan (Bito, et al., 2013). Bioinformatic analysis has revealed the presence in the *C. elegans* genome of genetic orthologues associated with human inherited B_{12} disorders but, interestingly, there is no evidence of the main B_{12} binding and trafficking proteins such as intrinsic factor or transcobalamin (Yilmaz and Walhout, 2014). To investigate B_{12} uptake into *C. elegans*, nematodes of the N2 Bristol strain were fed with *E. coli* OP50 that had been grown in the

presence of either C5-BoD-cobyric acid, ribose-BoD-cobalamin, or unattached BoD. L4 worms that had been grown in the presence of these analogues since their embryonic stage were imaged on a confocal microscope. Only worms fed on *E. coli* containing the ribose-BoD-cobalamin revealed any fluorescence within the body. Interestingly, confocal microscopy revealed that in these worms the fluorescence was localised to discrete spots in the head, vulva and tail regions, corresponding to the coelomocytes (Figure 6). Even within this discrete cell type, the fluorescence appears to show a degree of sub-cellular localisation or compartmentalisation (Figure 6). This result demonstrates that *C. elegans* is able to absorb and retain the BoD-labelled cobalamin (which was concentrated in some organs), whereas retention of the C5-cobyric acid analogue was clearly absent.

B₁₂ uptake into higher plants.

It is generally accepted that land plants neither make nor require B₁₂ and live in a B₁₂-less world. Early reports suggesting that some plants contained B₁₂-dependent enzyme activity were later put down to bacterial contamination. However, there does appear to be some evidence suggesting that some plants can at least take up and transport cobalamin if the nutrient is cultivated with organic fertilisers or if plants are grown in B₁₂-enriched media (Mozafar, 1994; Sato, et al., 2004).

To investigate this observation further, we instigated an outreach project with a local secondary school, where pupils grew *Lepidium sativum* (garden cress) aseptically on an agar medium containing increasing concentrations of vitamin B₁₂. After growth for 7 days the cotyledons from the germinating seedlings were removed, washed and homogenised. The soluble supernatant after centrifugation was then applied to bioassay plates, which demonstrated that the seedlings were able to absorb cobalamin from the growth medium in a concentration dependent manner (Figure S6). To confirm the ability of *L. sativum* to absorb cobalamin we repeated the uptake experiment using ribose-OG-cobalamin. *L. sativum* was therefore grown with ribose-OG-cobalamin and, after sectioning, was imaged by confocal microscopy. This approach confirmed that *L. sativum* absorbs ribose-OG-cobalamin, which was found to localise to the vacuoles of the cotyledons (Figure 7). These results provide definitive evidence that some plants can absorb and transport cobalamin, at least by passive diffusion. The linear shape of the concentration response and high amounts of Cbl used in the experiment (see supplementary Figure S6) indicate an unspecific uptake mechanism.

DISCUSSION.

In this research we have demonstrated that it is possible to functionalise the main corrin framework by substituting the methyl group of SAM with either an allyl or propargyl group.

Although this approach was used to modify the C5 position, in theory it may be possible to modify any of the positions on the corrin ring that are methylated by SAM-dependent methyl transferases. Through this development we have shown that it is possible to generate either C5-allyl or C5-propargyl derivatives of HBA, and outlined how these can be transformed into cobyrinic acid analogues. By attaching fluorophores to the activated C5 allyl group it was possible to make C5-BoD and C5-OG cobyrinic acid derivatives. This has been achieved through the application of enzymatic and chemical tools to functionalize the comparatively inert corrin framework of B₁₂. In doing so, we have been able to make analogues that can be used to investigate specific biological processes, expanding upon our initial research that allowed us to make different metal isosteres of cobalamin (Widner, et al., 2016). Our chemical and biological synthesis approach allows for a regio-specific modification to the corrin molecule. Other have developed methods to generate a reactive group on the corrin molecule through non-specific acid hydrolysis of cobalamin in order to isolate variants with a free acid side chain, which was then subsequently modified (Waibel, et al., 2008). This involves elaborate purification procedures and overall low yields.

The success of our approach relies on combining biosynthetic enzymes with chemical synthesis to produce new functional analogues. We wondered whether the additional chemical space occupied around the C5 position would present a steric hindrance especially with the downstream biosynthetic enzymes. Indeed, the presence of the C5-allyl group on precorrin-8 did reduce the rate of reaction with CobH, which catalyses the methyl migration from C11 to C12 in the formation of HBA. However, with knowledge of the 3-dimensional structure of the enzyme (Shipman, et al., 2001) we were able to rationally engineer more space into the active site of CobH to make it significantly more active with the C5-analogues.

The amide groups in cobalamin are added by CobB and CobQ. CobB accepts the C5-allyl modified substrate well and transforms it into C5-HBA a,c-diamide in high yields. Normally, the next steps in the aerobic cobalamin pathway involve metal chelation and adenosylation. The amidation of the remaining side chains is only completed after the chelation and adenosylation processes have taken place. We wanted to change sequence of pathway reactions by completing the amidations prior to metal chelation, ie to transform HBAD into Hby. We were aware that some organisms must be able to make Hby as an intermediate, as reports from 40 years ago had highlighted that bacteria such as *A. vinosum*, when grown in the absence of cobalt, are able to produce cobalt-free corrinoids such as hydrogenobalamin (Kopenhagen, et al., 1973). We thus cloned cobQ from *A. vinosum* and demonstrated that the encoded enzyme is indeed able to convert HBA-diamide into Hby and similarly convert the C5 analogues into the C5-Hby equivalents. With all these enzyme tools in hand we were able to produce the C5-allyl Hby on the tens of mg scale. After coupling of the fluorophore to the

C5-allyl position cobalt was then inserted to generate the C5-cobyric acid analogues containing either an OG or BoD fluorophore. These coobyric acid analogues were then compared to cobalamin analogues where the fluorophores were attached to the ribose position.

The functional analysis of the various corrin and cobalamin analogues was surprising. Addition of the C5-TPEA-cobyric acid to a microbial B₁₂ bioassay plate promoted growth around the application point, indicating that the C5-material must be taken up by the *S. enterica* reporter strain, which then presumably converted it into a functional cobalamin analogue. The other analogues such as the C5-OG and C5-BoD variants also promoted growth, albeit less well. The high levels of the analogues required to promote growth reflect several constraints on their molecular recognition in terms of their uptake and incorporation into methionine synthase. For instance, it is clear from our results that the ribose-OG-cobalamin is taken up more efficiently than the C5-linked coobyric acid analogue, whilst, conversely, the C5-BoD-cobyric acid analogue is taken up more easily than the ribose-linked cobalamin equivalent. What is most surprising of all is that all these analogues appear to fit into the active site of methionine synthase. The crystal structure of the *E. coli* methionine synthase suggests that there is a certain amount of room to accommodate some extra bulk on the ribose group (Drennan, et al., 1994). However, within the corrin binding site there is less room for the further chemical space required for the C5-attachment. Nonetheless, as mentioned above, the same is true with CobH, the enzyme required to convert the modified C5-allyl-precobalamin-8 intermediate into C5-allyl-HBA, where the enzyme still catalyses the reaction albeit at a reduced rate in comparison to the natural substrate. So, with methionine synthase the extra group on the corrin ring may force some conformational change within the protein so that the extra bulk is accommodated. There is always the possibility that the fluorescent analogues are broken down in the cell to release the fluorophore from the corrin. However, extraction of C5-BoD-cobyric acid from *E. coli* that had been grown with the compound resulted only in the identification of intact C5-BoD-cobyric acid. Overall, this tells us that the *S. enterica* and *E. coli* uptake mechanisms will accept corrin and B₁₂ analogues with significant extra chemical bulk and that these compounds are relatively active inside the cell.

We investigated whether these fluorescent corrin analogues were also taken up by *M. tuberculosis*. Interestingly, both the C5-BoD-cobyric acid and the ribose-BoD-cobalamin analogues were taken up by cultured *M. tuberculosis* cells with the C5-BoD analogue appearing to be taken up more quickly than the ribose analogue. Significantly, both analogues appear to support growth of this B₁₂-dependent TB strain indicating that the molecules are able to get inside the cells. The B₁₂ uptake mechanism in *M. tuberculosis*, which is different to that found in most other bacteria (Gopinath, et al., 2013), is clearly able to take up incomplete

corrinoids. As humans are not able to take up cobalamin, the use of cobalamin conjugates could prove a useful method for specifically labelling and targeting TB infections.

There is still much to learn about how B₁₂ makes its journey from bacteria to specific compartments in eukaryotic cells. Several recent reports have highlighted the fact that *C. elegans* may prove to be a good model to learn more about the movement, distribution and storage of B₁₂ (Bito, et al., 2013; Bito, et al., 2017; Watson, et al., 2014; Yilmaz and Walhout, 2014). Indeed, we were able to show that a fluorescent B₁₂ derivative is absorbed from bacteria by *C. elegans* and is observed to accumulate within coelomocytes. Interestingly, only the ribose-linked cobalamin analogue was absorbed indicating that the worm is able to differentiate between complete and incomplete corrinoids. It is not clear whether this is due to the B₁₂ analogue being recognised as foreign material or whether the cells represent a storage area for excess cobalamin. Nonetheless, the results demonstrate that this is a powerful model to learn about how B₁₂ is absorbed and, as this system is different to mammalian systems, there is the possibility of exploiting this difference to try and treat worm-based parasites such as hook worms.

It is well documented that vegetarians are more prone to B₁₂ deficiency as plants lack B₁₂ (Stabler and Allen, 2004). However, it has been suggested that some plants are able to absorb the nutrient if added exogenously to their growth media (Mozafar, 1994; Sato, et al., 2004). Through an outreach project we demonstrated that garden cress, *L. sativum*, can indeed take up cobalamin. The amount of B₁₂ absorbed by garden cress is dependent upon the amount present in the growth medium, and displayed a linear rather than a hyperbolic relationship. This would suggest an unspecific form of uptake. Nonetheless, we were able to confirm B₁₂ uptake using our OG analogue of cobalamin, showing that the nutrient ends up in the vacuole in the leaf. The uptake of B₁₂ into plants is irrelevant in terms of plant metabolism, since plants do not possess any B₁₂-dependent enzymes, but the finding that some plants are able to accumulate B₁₂ is important as such nutrient-enriched plants would be important in helping overcome dietary limitations in countries such as India, which have a high proportion of vegetarians.

SIGNIFICANCE.

Vitamin B₁₂, cobalamin, is an essential dietary component that is synthesised solely by certain prokaryotes. Although some eukaryotes, including mammals, require the nutrient, kingdoms such as Plantae and Fungi have evolved to live in a B₁₂-less environment, which has implications for people on a vegan diet. In order to follow more closely the journey undertaken by cobalamin, from its prokaryotic inception, through sharing in microbial communities and its

uptake by animals, we devised ways to construct a range of fluorescent analogues of cobalamin. By manipulation of the B₁₂ biosynthetic pathway we were able to modify the C5 position of the corrin framework so as to incorporate an allyl substituent, where the vinyl group could be used to attach fluorophores to give tagged cobalamin analogues. Fluorescent cobalamin variants were also constructed through attachment of fluorophores to the ribose group in the lower nucleotide loop. Together, these fluorescent analogues were used to show that *M. tuberculosis* is able to absorb both complete and incomplete corrinoids, demonstrating that cobalamin can be used to carry cargo into the cell. *C. elegans* was also found to absorb cobalamin, demonstrating that this approach can be used to follow intracellular trafficking in a model organism. Moreover, it also represents a target for parasites such as hook worms. Finally, and very surprisingly, certain plants were observed to be able to absorb B₁₂ from their growth medium, accumulating the molecule in the leaf vacuole. This latter finding may be important as a way to address the global challenge of providing a nutrient-complete vegetarian diet, a valuable development as the world becomes increasingly meat-free due to population expansion.

STAR*METHODS

Detailed methods are provided in the online version of this paper and include the following:

SUPPLEMENTAL INFORMATION

Supplemental information includes five figures and one table and can be with this article online.

AUTHOR CONTRIBUTION

ADL, EN-S, ED, JAB and SS all undertook aspects of the experimental work including cloning, protein purification and synthesis, and helped with the interpretation of the data. DGB collected the X-ray data and assisted ADL and EN-S with the structure determination of CobH. MJH performed all NMR data acquisition and assisted ADL with analyses. JMAT, ED and EN-S designed the *C. elegans* experiments. AGS helped in the design of the plant-based investigations. IRB undertook some of the imaging experiments and provided guidance on experiment design. HIB and CEB designed the TB work. ADL, ED, EN-S and MJW designed the experiments and wrote the paper.

DECLARATION OF INTERESTS

The authors declare no competing interests.

ACKNOWLEDGEMENTS

This work was supported, in part, by the Intramural Research Program of NIAID and in part by grants from the Biotechnology and Biological Sciences Research Council (BBSRC; BB/L010208/1, BB/K009249/1, BB/I012079 and the Wellcome Trust 097627/Z/11/Z). We thank Prof Jason Micklefield (Manchester) and Prof Udo Oppermann (Oxford) for the gift of SAHH and hMATIIA. Our research into B₁₂ uptake into plants was done as part of an outreach programme to engage pupils in real scientific research, inspired by the Authentic Biology project (www.authentic-biology.org). The detection and measurement of B₁₂ in the cotyledons of garden cress was performed with the help of the biology teachers and year 11 and 12 pupils at Sir Roger Manwood School in Sandwich, Kent. Their names are given below.

Teachers: Dr Jackie Wilson and Vicki Beale.

Pupils: Megan Battenfield, Megan Betts, Alex Gallagher, Inga Loo, Shiree Kinder, George Morris, Emma Rudge, James Russell, Rebekah Smith, Maddy Stock, Pema Tamang, Elizabeth Westbrook, Himmy Wu, Katie Fidock

REFERENCES

- Allen, R.H., and Stabler, S.P. (2008). Identification and quantitation of cobalamin and cobalamin analogues in human feces. *Am J Clin Nutr* 87, 1324-1335.
- Banerjee, R., and Ragsdale, S.W. (2003). The many faces of vitamin B₁₂: catalysis by cobalamin-dependent enzymes. *Annu Rev Biochem* 72, 209-247.
- Bito, T., Matsunaga, Y., Yabuta, Y., Kawano, T., and Watanabe, F. (2013). Vitamin B₁₂ deficiency in *Caenorhabditis elegans* results in loss of fertility, extended life cycle, and reduced lifespan. *FEBS Open Bio* 3, 112-117.
- Bito, T., Misaki, T., Yabuta, Y., Ishikawa, T., Kawano, T., and Watanabe, F. (2017). Vitamin B₁₂ deficiency results in severe oxidative stress, leading to memory retention impairment in *Caenorhabditis elegans*. *Redox Biol* 11, 21-29.
- Blanche, F., Cameron, B., Crouzet, J., Debussche, L., Thibaut, D., Vuilhorgne, M., Leeper, F.J., and Battersby, A.R. (1995). Vitamin B₁₂ - How the Problem of Its Biosynthesis Was Solved. *Angewandte Chemie-International Edition in English* 34, 383-411.

Brito, A., Chiquette, J., Stabler, S.P., Allen, R.H., and Girard, C.L. (2015). Supplementing lactating dairy cows with a vitamin B₁₂ precursor, 5,6-dimethylbenzimidazole, increases the apparent ruminal synthesis of vitamin B₁₂. *Animal* 9, 67-75.

Chan, C.H., and Escalante-Semerena, J.C. (2011). ArsAB, a novel enzyme from *Sporomusa ovata* activates phenolic bases for adenosylcobamide biosynthesis. *Mol Microbiol* 81, 952-967.

Croft, M.T., Lawrence, A.D., Raux-Deery, E., Warren, M.J., and Smith, A.G. (2005). Algae acquire vitamin B₁₂ through a symbiotic relationship with bacteria. *Nature* 438, 90-93.

Crofts, T.S., Seth, E.C., Hazra, A.B., and Taga, M.E. (2013). Cobamide structure depends on both lower ligand availability and CobT substrate specificity. *Chem Biol* 20, 1265-1274.

Crouzet, J., Cauchois, L., Blanche, F., Debussche, L., Thibaut, D., Rouyez, M.C., Rigault, S., Mayaux, J.F., and Cameron, B. (1990). Nucleotide sequence of a *Pseudomonas denitrificans* 5.4 kilobase DNA fragment containing 5 cob genes and identification of structural genes encoding S-adenosyl-L-methionine - uroporphyrinogen III methyltransferase and cobyrinic acid a,c-diamide synthase. *J bacteriol* 172, 5968-5979.

Dalhoff, C., Lukinavicius, G., Klimasauskas, S., and Weinhold, E. (2006). Direct transfer of extended groups from synthetic cofactors by DNA methyltransferases. *Nat Chem Biol* 2, 31-32.

Dalhoff, C., Lukinavicius, G., Klimasauskas, S., and Weinhold, E. (2006). Synthesis of S-adenosyl-L-methionine analogs and their use for sequence-specific transalkylation of DNA by methyltransferases. *Nat Protoc* 1, 1879-1886.

Deery, E., Schroeder, S., Lawrence, A.D., Taylor, S.L., Seyedarabi, A., Waterman, J., Wilson, K.S., Brown, D., Geeves, M.A., Howard, M.J., et al. (2012). An enzyme-trap approach allows isolation of intermediates in cobalamin biosynthesis. *Nat Chem Biol* 8, 933-940.

Degnan, P.H., Taga, M.E., and Goodman, A.L. (2014). Vitamin B₁₂ as a modulator of gut microbial ecology. *Cell Metab* 20, 769-778.

Drennan, C.L., Huang, S., Drummond, J.T., Matthews, R.G., and Ludwig, M.L. (1994). How a protein binds B₁₂ - a 3.0-Angstrom X-ray structure of B₁₂-binding domains of methionine synthase. *Science* 266, 1669-1674.

Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of Coot. *Acta Crystallogr D* 66, 486-501.

Gherasim, C., Lofgren, M., and Banerjee, R. (2013). Navigating the B₁₂ road: assimilation, delivery, and disorders of cobalamin. *J Biol Chem* 288, 13186-13193.

Girard, C.L., Berthiaume, R., Stabler, S.P., and Allen, R.H. (2009). Identification of cobalamin and cobalamin analogues along the gastrointestinal tract of dairy cows. *Arch Anim Nutr* 63, 379-388.

Gopinath, K., Moosa, A., Mizrahi, V., and Warner, D.F. (2013). Vitamin B₁₂ metabolism in *Mycobacterium tuberculosis*. *Future Microbiol* 8, 1405-1418.

Gopinath, K., Venclovas, C., Ioerger, T.R., Sacchettini, J.C., McKinney, J.D., Mizrahi, V., and Warner, D.F. (2013). A vitamin B₁₂ transporter in *Mycobacterium tuberculosis*. *Open Biol* 3, 120175.

Gray, M.J., and Escalante-Semerena, J.C. (2009). In vivo analysis of cobinamide salvaging in *Rhodobacter sphaeroides* strain 2.4.1. *J Bacteriol* 191, 3842-3851.

Greibe, E., Fedosov, S., and Nexø, E. (2012). The cobalamin-binding protein in zebrafish is an intermediate between the three cobalamin-binding proteins in human. *PLoS One* 7, e35660.

Helliwell, K.E., Lawrence, A.D., Holzer, A., Kudahl, U.J., Sasso, S., Krautler, B., Scanlan, D.J., Warren, M.J., and Smith, A.G. (2016). Cyanobacteria and eukaryotic algae use different chemical variants of vitamin B₁₂. *Curr Biol* 26, 999-1008.

Horton, R.A., Bagnato, J.D., and Grissom, C.B. (2003). Structural determination of 5'-OH alpha-ribofuranoside modified cobalamins via ¹³C and DEPT NMR. *J Org Chem* 68, 7108-7111.

Kopenhagen, V.B., Wagner, F., and Pfiffner, J.J. (1973). Alpha-(5,6-dimethylbenzimidazolyl)rhodibamide and rhodibinamide, the rhodium analogues of vitamin B₁₂ and cobinamide. *J Biol Chem* 248, 7999-8002.

Krautler, B., Fieber, W., Ostermann, S., Fasching, M., Ongania, K.H., Gruber, K., Kratky, C., and Mikl, C. (2003). The cofactor of tetrachloroethene reductive dehalogenase of *Dehalospirillum multivorans* is norpseudo B₁₂, a new type of a natural corrinoid. *Helv Chim Acta* 86, 3698-3716.

Lee, M., and Grissom, C.B. (2009). Design, synthesis, and characterization of fluorescent cobalamin analogues with high quantum efficiencies. *Org Lett* 11, 2499-2502.

Maggio-Hall, L.A., and Escalante-Semerena, J.C. (2003). Alpha-5,6-dimethylbenzimidazole adenine dinucleotide (alpha-DAD), a putative new intermediate of coenzyme B₁₂ biosynthesis in *Salmonella typhimurium*. *Microbiology* 149, 983-990.

Martens, J.H., Barg, H., Warren, M.J., and Jahn, D. (2002). Microbial production of vitamin B₁₂. *Appl Microbiol Biotechnol* 58, 275-285.

- McEwan, J.F., Veitch, H.S., and Russell-Jones, G.J. (1999). Synthesis and biological activity of ribose-5'-carbamate derivatives of vitamin B₁₂. *Bioconjug Chem* 10, 1131-1136.
- Moore, S.J., Lawrence, A.D., Biedendieck, R., Deery, E., Frank, S., Howard, M.J., Rigby, S.E., and Warren, M.J. (2013). Elucidation of the anaerobic pathway for the corrin component of cobalamin (vitamin B₁₂). *Proc Natl Acad Sci USA* 110, 14906-14911.
- Mozafar, A. (1994). Enrichment of some B-vitamins in plants with application of organic fertilizers. *Plant Soil* 167, 305-311.
- Potterton, E., Briggs, P., Turkenburg, M., and Dodson, E. (2003). A graphical user interface to the CCP4 program suite. *Acta Cryst.* D59, 1131-1137.
- Raux, E., Lanois, A., Levillayer, F., Warren, M.J., Brody, E., Rambach, A., and Thermes, C. (1996). Salmonella typhimurium cobalamin (vitamin B₁₂) biosynthetic genes: functional studies in S. typhimurium and Escherichia coli. *J Bacteriol* 178, 753-767.
- Renz, P. (1999). Biosynthesis of the 5,6-dimethylbenzimidazole moiety of cobalamin and of the other bases found in natural corrinoids. In *Chemistry and biochemistry of B₁₂*, R. Banerjee, ed. (New York: John Wiley and sons), pp. 557-575.
- Rodionov, D.A., Vitreschak, A.G., Mironov, A.A., and Gelfand, M.S. (2003). Comparative genomics of the vitamin B₁₂ metabolism and regulation in prokaryotes. *J Biol Chem* 278, 41148-41159.
- Roth, J.R., Lawrence, J.G., and Bobik, T.A. (1996). Cobalamin (coenzyme B₁₂): synthesis and biological significance. *Annu Rev Microbiol* 50, 137-181.
- Sato, K., Kudo, Y., and Muramatsu, K. (2004). Incorporation of a high level of vitamin B₁₂ into a vegetable, kaiware daikon (Japanese radish sprout), by the absorption from its seeds. *BBA-Gen Subjects* 1672, 135-137.
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). FIJI: an open-source platform for biological-image analysis. *Nat Methods* 9, 676-682.
- Shipman, L.W., Li, D., Roessner, C.A., Scott, A.I., and Sacchettini, J.C. (2001). Crystal structure of precorrin-8x methyl mutase. *Structure* 9, 587-596.
- Singh, S., Zhang, J., Huber, T.D., Sunkara, M., Hurley, K., Goff, R.D., Wang, G., Zhang, W., Liu, C., Rohr, J., et al. (2014). Facile chemoenzymatic strategies for the synthesis and utilization of S-adenosyl-(L)-methionine analogues. *Angew Chem Int Ed Engl* 53, 3965-3969.

Smeltzer, C.C., Cannon, M.J., Pinson, P.R., Munger, J.D., Jr., West, F.G., and Grissom, C.B. (2001). Synthesis and characterization of fluorescent cobalamin (CobalaFluor) derivatives for imaging. *Org Lett* 3, 799-801.

Stabler, S.P., and Allen, R.H. (2004). Vitamin B₁₂ deficiency as a worldwide problem. *Annu Rev Nutr* 24, 299-326.

Studier, F.W., Daegelen, P., Lenski, R.E., Maslov, S., and Kim, J.F. (2009). Understanding the differences between genome sequences of *Escherichia coli* B strains REL606 and BL21(DE3) and comparison of the *E. coli* B and K12 genomes. *J Mol Biol* 394, 653-680.

Vranken, W.F., Boucher, W., Stevens, T.J., Fogh, R.H., Pajon, A., Llinas, M., Ulrich, E.L., Markley, J.L., Ionides, J., and Laue, E.D. (2005). The CCPN data model for NMR spectroscopy: development of a software pipeline. *Proteins* 59: 687-96.

Waibel, R., Treichler, H., Schaefer, N.G., van Staveren, D.R., Mundwiler, S., Kunze, S., Kuenzi, M., Alberto, R., Nuesch, J., Knuth, A., et al. (2008). New derivatives of vitamin B₁₂ show preferential targeting of tumors. *Cancer Res* 68, 2904-2911.

Wang, R., Islam, K., Liu, Y., Zheng, W., Tang, H., Lailier, N., Blum, G., Deng, H., and Luo, M. (2013). Profiling genome-wide chromatin methylation with engineered posttranslation apparatus within living cells. *J Am Chem Soc* 135, 1048-1056.

Wang, R., Zheng, W., and Luo, M. (2014). A sensitive mass spectrum assay to characterize engineered methionine adenosyltransferases with S-alkyl methionine analogues as substrates. *Anal Biochem* 450, 11-19.

Wang, R., Zheng, W., Yu, H., Deng, H., and Luo, M. (2011). Labeling substrates of protein arginine methyltransferase with engineered enzymes and matched S-adenosyl-L-methionine analogues. *J Am Chem Soc* 133, 7648-7651.

Warner, D.F., Savvi, S., Mizrahi, V., and Dawes, S.S. (2007). A riboswitch regulates expression of the coenzyme B₁₂-independent methionine synthase in *Mycobacterium tuberculosis*: implications for differential methionine synthase function in strains H37Rv and CDC1551. *J Bacteriol* 189, 3655-3659.

Warren, M.J., Raux, E., Schubert, H.L., and Escalante-Semerena, J.C. (2002). The biosynthesis of adenosylcobalamin (vitamin B₁₂). *Nat Prod Rep* 19, 390-412.

Watkins, D., and Rosenblatt, D.S. (2013). Lessons in biology from patients with inborn errors of vitamin B₁₂ metabolism. *Biochimie* 95, 1019-1022.

Watson, E., MacNeil, L.T., Ritter, A.D., Yilmaz, L.S., Rosebrock, A.P., Caudy, A.A., and Walhout, A.J. (2014). Interspecies systems biology uncovers metabolites affecting *C. elegans* gene expression and life history traits. *Cell* 156, 759-770.

Widner, F.J., Lawrence, A.D., Deery, E., Heldt, D., Frank, S., Gruber, K., Wurst, K., Warren, M.J., and Krautler, B. (2016). Total Synthesis, Structure, and Biological Activity of Adenosylrhodibalamin, the Non-Natural Rhodium Homologue of Coenzyme B₁₂. *Angew Chem Int Ed Engl* 55, 11281-11286.

Winter, G. (2010). XIA2: an expert system for macromolecular crystallography data reduction. *J Appl Crystallogr* 43, 186-190.

Wishart, D.S., and Case, D.A. (2001). Use of chemical shifts in macromolecular structure determination. *Methods Enzymol* 338, 3-34.

Yilmaz, L.S., and Walhout, A.J. (2014). Worms, bacteria, and micronutrients: an elegant model of our diet. *Trends Genet* 30, 496-503.

Zaman, K., and Zak, Z. (1989). Purification and partial characterization of a cobalamin-binding protein from chicken egg yolk. *Biochim Biophys Acta* 998, 102-104.

Zayas, C.L., Claas, K., and Escalante-Semerena, J.C. (2007). The CbiB protein of *Salmonella enterica* is an integral membrane protein involved in the last step of the de novo corrin ring biosynthetic pathway. *J Bacteriol* 189, 7697-7708.

FIGURE LEGENDS

Figure 1. Biosynthesis of adenosylcobalamin from uroporphyrinogen III. Initially, uroporphyrinogen III is acted upon by the enzymes in the sequence of CobA, I, G, J, M, F, K, L and H to generate a cobalt-free corrinoid called HBA. Amidation of the a and c side chains followed by metal insertion, adenosylation and the remaining amidations by CobB, NST, R, O and Q gives rise to adenosylcobyrinic acid. The final stages of the biosynthesis see the attachment of an aminopropanol side chain and the alpha ribazole nucleotide, which contains the unusual base dimethylbenzimidazole. These reactions are mediated by the enzymes CobD, P, V and U. The numbering of the corrin macrocycle (1-20) is shown as is the lettering associated with the side chains (a-g). Different variants of cobalamin are found through changes in the highlighted areas of the molecule. The upper ligand (orange) can be a methyl, cyano or water group, the linking aminopropanol (purple) is an ethanolamine molecule in nor-cobalamin, and a range of different bases can replace the dimethylbenzimidazole (blue).

Figure 2. The ChemBio synthesis of a C5 analogue of cobyrinic acid. Precorrin-7 is modified with an allyl-SAM derivative by the action of the C5 methyltransferase, CobL. This results in the production of C5-allyl-precorrin-8, which is acted upon by the enzyme CobH, B and Q to generate C5-allyl-Hby. Cysteamine is next chemically attached to the allyl group to generate C5-TPEA-Hby prior to the chemical insertion of cobalt to give C5-TPEA-cobyric acid. The coupling of a fluorophore via the free amine group on C5 generates the C5-fluorophore conjugate of cobyrinic acid. The fluorophore is represented by the filled red circle.

Figure 3. Product complex of CobH^{T85A} with C5-allyl-HBA. The figure shows the position of the reaction product bound at the active site of the enzyme. The proximity of the C5-group to Ala85 demonstrates how the removal of the threonine side chain has provided extra space to accommodate the allyl group.

Figure 4. Chemical structures of the four fluorescent analogues of cobalamin. The top two structures show the attachment of Orgeon Green and BoDIPY to the C5 position of the corrin ring, whereas the lower two structures show how these fluorophores are attached to the ribose moiety of the lower nucleotide loop.

Figure 5. Effect of cobyrinic acid and cobalamin analogues on bacteria. Panel A shows a bioassay plate containing a *S. enterica* strain that requires cobyrinic acid or later pathway intermediates for growth. Addition of a known amount of vitamin B₁₂ (spot 3) allows growth. Spots 1, 2 and 4 represent C5-Allyl-hydrogenobyric acid a,c-diamide (C5-Allyl-HBAD), C5-Allyl-Hby and C5-TPEA-Hby, none of which promote growth. Spot 5 contains the C5-TPEA-cobyric acid analogue and promotes growth. Panel B is the same bioassay plate assay as in panel A, with spot 3 containing a standard vitamin B₁₂ solution. Spots 1, 2, 4 and 5 are solutions of C5-BoD-cobyric acid, ribose-BoD-cobalamin, C5-OG-cobyric acid and ribose-OG-cobalamin, respectively. All the analogues support growth although the C5 analogues are weaker than the respective ribose-linked analogues. Panels C and D are overlay pictures of *E. coli* cells that have been grown in the presence of ribose-BoD-cobalamin and C5-BoD-cobyric acid, respectively. The co-localisation of the red fluorescence with the cells indicates that *E. coli* has taken up both compounds. Panels E and F show cultures of *M. tuberculosis*, grown in the presence of ribose-BoD-cobalamin and C5-BoD-cobyric acid, respectively. Again, the presence of the red fluorescence with the cells indicates that *M. tuberculosis* is able to take up both analogues.

Figure 6. Localisation of ribose-BoD-cobalamin to coelomocytes in *C. elegans*. *C. elegans* were fed with *E. coli* OP50 that had been grown in the presence of ribose-BoD-cobalamin and were imaged after several days. The red fluorescence associated with the

fluorophore was found localised in the 6 coelomocytes, two of which are found at the head (Panel A), two near the vulva (Panel B) and two at the anus (Panel C).

Figure 7. Localisation of ribose-OG-cobalamin to vacuoles in *L. sativum* (garden cress).

Seeds were germinated in sterile media containing ribose-OG-cobalamin and were imaged after 5 days. Thin sections of the cotyledons of cress grown in the presence of the ribose-OG-cobalamin had highly fluorescent vacuoles (Panel A), whereas control plants grown in the absence of the cobalamin derivative lacked this fluorescence (Panel B).

STAR*METHODS

CONTACT FOR REAGENT AND RESOURCES SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Professor Martin J. Warren (m.j.warren@kent.ac.uk)

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All bacterial strains used in this study are listed in the Key Resources Table together with their source and genotypes. For *E. coli*, unless stated otherwise, the strains were grown in LB media supplemented with the appropriate antibiotics. All *Mycobacterium tuberculosis* cultures were grown in cobalamin deficient Sauton's defined medium (0.5 g KH_2PO_4 , 0.5 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g citric acid, 0/05 g ferric ammonium citrate, 60 mLs glycerol, 4 g asparagine and 500 μL TWEEN 80 in 1 L pH 7.0. Worm studies were carried out with *Caenorhabditis elegans* var. Bristol (strain N2) and maintained on nematode growth medium (NGM) agar.

METHODS DETAILS

Chemicals and reagents. Oregon Green® 514 (OG) and BODIPY® TR-X (BoD) succinimidyl ester were both purchased from Thermo Fisher Scientific, Inc., as was the P-Per for plant extraction. Allyl-Bromide, Cysteamine hydrochloride, Seleno-L-methionine, Dowex 50WX4 hydrogen form, Diethylaminoethyl–Sephacel, and N, N-Diisopropylethylamine Reagent Plus® were bought from Sigma Aldrich, Ltd. Other chemicals and equipment include LICHROPREP RP-18 from VWR, VA-044 2,2'-Azobis[2-(2-imidazolin-2-yl)propane] dihydrochloride from alpha laboratories Ltd. and crystal screens from Molecular Dimensions Ltd.

Protein production and purification. All cobalamin biosynthetic genes were amplified by PCR using genomic DNA from either *Rhodobacter capsulatus* SB1003 (cobL, H and B), or *Allochrochromatium vinosum* ATCC 17899 (cobQ). The cob genes were amplified individually with primers then cloned into modified pET14b plasmids. These were then transformed into BL21 star (DE3) pLysS. hMATIIA I117A was transformed into Rosetta 2 (DE3) pLysS. The recombinant strains were grown in LB at 37°C and protein overexpression was induced with 0.4 mM of isopropyl-1-thio- β -D-galactopyranoside. The cells were resuspended in buffer A (20 mM Tris pH 8, 500 mM NaCl, 10 mM Imidazole) and sonicated. The protein was purified using Chelating Sepharose™ charged with NiSO_4 . Unbound proteins were washed off with buffer A, buffer A containing 30 mM imidazole and buffer A containing 80 mM Imidazole. Proteins were eluted with buffer A containing 400 mM imidazole and passed through a PD10 column equilibrated with buffer B (20 mM Tris, pH 8.0, containing 100 mM NaCl) for CobL and H, or buffer C (20 mM Tris, pH 8.0, containing 500 mM NaCl) for CobB and Q.

Production and purification of precorrin-7. *E. coli* BL21 star (DE3) pLysS was transformed with pET3a/cobA-I-G-J-M-F-K-L^C-E and grown in 2TY at 28 °C for 24 hours (Deery, et al., 2012). The cells were resuspended in 20 mM Tris, pH 8.0, and sonicated. Proteins were precipitated with acetonitrile and following clarification by centrifugation precorrin-7 was purified over DEAE-Sepharose. The column was washed with 20 mM Tris, pH 8.0, containing 100 mM NaCl, before elution with 500 mM NaCl. The pH of the eluent was lowered to pH 4.0 with TFA and the precorrin-7 was further purified over RP18. The column was washed with a stepwise gradient of methanol in 0.1% (v/v) TFA and elution was achieved with 50% methanol.

Synthesis of SAM analogues. Allyl-SAM was prepared from homocysteine and allylbromide as described previously (Wang, et al., 2011). Allyl bromide (44 µL, 0.214 mmol) and AgClO₄ (6 mg, 0.025 mmol) were added to S-adenosyl-L-homocysteine (5 mg, 0.013 mmol) dissolved in a mixture of formic acid and acetic acid (1:1, 0.6 mL) at 0 °C. The mixture was allowed to warm up to room temperature and stirred continuously for 2.5 h. A further portion of allyl bromide (44 µL, 0.214 mmol) and AgClO₄ (6 mg, 0.025 mmol) were added and the reaction mixture was stirred for another 2.5 h. following centrifugation to remove the precipitate, H₂O (3 mL) was added and the residual allyl bromide was extracted with diethyl ether (5 mL × 3). The aqueous phase was then concentrated under vacuum.

Synthesis of Se-propargyl-L-selenohomocysteine. Se-propargyl-L-selenohomocysteine was synthesized from L-selenomethionine as described previously (Singh, et al., 2014). L-selenomethionine (100 mg) was dissolved in 20 mL liquid ammonia at -78 °C and stirred. Small fragments of sodium metal (34 mg) were added until a deep blue colour persisted for at least 1 min. Propargyl bromide (46.5 µL) was then added followed by ammonium chloride (20 mg). The reaction removed from the cooling bath and the flask was left open to the atmosphere for the ammonia to evaporate. The resulting residue was dissolved in H₂O (5 mL) and the pH was adjusted to pH 5 - 7 by the addition of HCl (1 M). The solvent was removed under reduced pressure.

Synthesis C5-allyl-HBA. Precorrin-7 (60 µM) was incubated with CobL (6 µM), CobH (30 µM) and allyl-SAM (170 µM) in buffer B at 28°C for 16 hours in the dark. The reaction mixture was then heated to 65°C and the precipitate was removed by centrifugation. The resulting supernatant was applied to a DEAE-Sepharose column equilibrated in 20 mM Tris, pH 8.0. The column was washed with 20 mM Tris, pH 8.0, and the product was eluted with 20 mM Tris, pH 8.0, containing 500 mM NaCl.

Synthesis of C5-propargyl-HBA. C5-propargyl-HBA was prepared from precorrin-7 and Se-propargyl-L-selenohomocysteine. Precorrin-7 (60 μ M), Se-propargyl-L-selenohomocysteine (1.5 mM), ATP (2 mM), $MgCl_2$ (5 mM), hMAT2A I117A (60 μ M), CobL (6 μ M), CobH (30 μ M), Se-propargyl-Hcy (1.5 mM) and SAHH (50 μ M) were incubated in 20 mM Tris, pH 8.0, containing 100 mM NaCl at 28°C in the dark for 16 h. The reaction mixture was heated to 80°C for 15 minutes and the precipitate was removed by centrifugation. The supernatant was applied to DEAE column pre-equilibrated in 20 mM Tris, pH 8.0. The column was washed with 20 mM Tris, pH 8.0, containing 100 mM NaCl, and eluted with 20 mM Tris, pH 8.0, containing 500 mM NaCl.

Synthesis of C5-allyl-Hby. C5-allyl-HBA (30 μ M), CobB (2 μ M), CobQ (5 μ M), ATP (8 mM), L-Glutamine (8 mM), and $MgCl_2$ (20 mM) were incubated in buffer C at 28°C for 16 hours in the dark. The reaction mixture was heated to 80°C, acidified to pH 4 with trifluoroacetic acid (TFA) and the precipitate removed by centrifugation. The supernatant was applied to an RP18 column equilibrated in 0.1% TFA, washed with 0.1% TFA followed by 20% methanol before elution in 50% methanol. The product was then dried in a vacuum centrifuge.

C5 allyl group extension. The C5-allyl group was modified through a thiol-ene coupling reaction with cysteamine. A solution of C5-allyl-Hby (30 μ M) was incubated at 55°C in acetate buffer (pH 4.0, 0.25 M) containing cysteamine (100 mM) and VA-044 (20 mM) in the dark for 15 mins. The product was purified over RP18 and analyzed by HPLC-MS.

Cobalt insertion. Cobalt insertion was achieved through incubation of the desired metal-free corrinoid (20 μ M) with $CoCl_2$ (10 mM) in ammonium hydroxide (0.2 M) at 80 °C under an N_2 atmosphere. The reaction was followed UV-visible spectroscopy until deemed complete (~ 60 mins) after which it was quenched with KCN and the product purified over RP18 and eluted in methanol.

Synthesis of fluorescent C5-cobyric acid derivatives. CN_2 -C5-TPEA-cobyric acid was dissolved in DMSO and either the succinimidyl ester of OG or BoD along with 2 equivalents of DIPEA were added. The reaction was stirred at room temperature for 16 hours in the dark. The solvent was removed in vacuo and the compound was purified by semi-preparative HPLC.

Ribose linked fluorescent vitamin B₁₂ derivatives. The ribose-5'-hydroxyl group of vitamin B₁₂ was activated with carbonyldiimidazole and coupled with 1,2-diaminoethane as described previously (McEwan, et al., 1999). The derivative was added to either the succinimidyl ester of OG or BoD with 2 equivalents of DIPEA in DMSO. The reaction was stirred for 16 hours in

the dark at room temperature. The product was precipitated with acetone and collected by filtration. Fluorescent derivatives were dissolved in 50% acetonitrile and purified by semi-preparative HPLC.

Crystal structure determination of CobH^{T85A} with bound C5-allyl-HBA. Recombinantly produced CobH^{T85A} was purified on a nickel affinity column. After affinity purification CobH^{T85A} was subject to gel filtration on a G200 Superdex column (GE Healthcare). Protein fractions were collected and concentrated to 7 mg mL⁻¹. Allyl-HBA was added to the protein in a 1:1 ratio prior to crystallisation at 19°C using the hanging drop vapour diffusion method. Initial crystals were obtained in 0.2 M calcium acetate hydrate, 0.1 M sodium cacodylate pH 6.5, 18 % (w/v) PEG 8000 (Molecular Dimensions, MD01 condition 20). Crystal optimisation required the addition of 20% methanol to the well condition and increasing the PEG 8000 to 20%. This resulted in multiple square shaped crystals which formed within three days. A single crystal was picked, transferred to a drop of in 0.2 M calcium acetate hydrate, 0.1 M sodium cacodylate pH 6.5, 20 % (w/v) PEG 8000 containing 20% glycerol as a cryoprotectant and then flash frozen in liquid nitrogen. The crystal diffracted to 1.6 Å and was found to belong to space group C 1 2 1 which is the same as the published structure of *R. capsulatus* CobH co-crystallised with HBA (PDB 4AU1) (Deery, et al., 2012). Data for the crystal was collected using beamline IO4-1 at the Diamond Light Source (UK). The data was auto-integrated and scaled using the Xia2 package using XDS (X-ray data software) and XSCALE (3dii) at the beamline. An initial model was generated by refinement of the scaled and merged data against PDB:4AU1 in Refmac 5. The model was then refined through cycles of manual model building in Coot and refinement with Refmac 5. The final model was validated in Coot and with wwPDB. The PDB coordinates were deposited with the code 5N0G.

Bioassay plates. A quantitative bioassay using *S. typhimurium* AR3612 was performed as described previously (Raux, et al., 1996). The bioassay plates containing the indicator strain AR3612 were prepared from lawns of the corresponding bacteria. The bacteria were scraped from an overnight minimal medium (M9) plate containing methionine (50 mg/L) and cysteine (50 mg/L) and washed with 0.9 % (w/v) NaCl to remove traces of methionine. The cells were mixed with 300 mL of minimal medium (M9) containing cysteine (50 mg/L) agar at 47 °C, and poured into plates.

Imaging in *Escherichia coli*. The OP50 *E. coli* was transformed with pET-BAD-btuBF and grown in a 4 mL LB culture after inoculation with 16 µL of starter culture. This was grown with 1 µM of C5-BoD-cobyric acid or ribose-BoD-cobalamin overnight with induction with 0.02% arabinose. Cultures (1 mL) were spun down and washed three times with fresh LB to remove

external fluorophore. The final pellet was resuspended in 1 mL of fresh LB, 5 μ L of this culture was dried on to a 1 % (w/v) LB-agarose pad and imaged on an Olympus IX81 widefield microscope with PlanApo 150 x OTIRFM-SP 1.49 numerical aperture lens mounted on ASI stage (Applied Scientific), and illuminated using LED light sources (Cairn Research Ltd) with appropriate filters (YFP and mCherry, Chroma). The samples were visualised using a Princeton ProEM 1024 back-thinned EMCCD camera (Princeton Instruments) on Metamorph software (Molecular Devices). Each 3D-maximum projection of volume data was calculated from 13 z-plane images, each 0.2 μ m apart.

Imaging in *Mycobacterium tuberculosis*. Relevant strains of *M. tuberculosis* were grown in 150 μ L cultures in a 96 well plate containing various concentrations of the BODIPY-labelled analogues of cobalamin and cobyric acid. The cells were grown for up to 48 h. After growth, the cells were spun down and washed three times to remove the fluorophore. Cells were pipetted directly onto a slide and imaged live on a Leica SP5 microscope using 594 nm excitation and imaging between 600-700 nm. The images were processed using Leica Lite confocal software and FIJI (Schindelin, et al., 2012).

Extraction of cobalamin from *Lepidium sativum* using P-Per. *Lepidium sativum* seeds were sterilised by washing with 70 % ethanol three times then rinsed five times with sterile water. The seeds were placed on Murashige-Skoog agar containing different concentrations of cobalamin, and grown for one week in sunlight at room temperature. The cotyledons of the cress were collected and washed 5 times with water. The residual water was removed by pipette, after centrifugation. The P-Per was prepared as instructed in the enclosed manual (Thermo Fisher): 283 μ L reagent A, 2.9 μ L of reagent B, and 214 μ L of reagent C. A little sand was added to the cotyledons along with 100 μ L of the P-Per. The cotyledons were ground using a hand-held pellet pestle for 2 minutes. The suspension was vortexed and then centrifuged for 3 minutes at 15,000 rpm rotation. The lower aqueous phase was collected and applied to the bioassay plate (see Section 2.6.1).

Imaging in *L. sativum*. *L. sativum* seeds were sterilised by washing with 70 % ethanol three times, then rinsed five times with sterile water. The seeds were placed on the Murashige-Skoog agar with 0.5 μ M ribose-OG cobalamin, and grown at room temperature in the dark for five days. Whole cotyledons or sectioned cress were placed directly on the glass slide with water. The samples were imaged with a Leica TCS SP2 laser scanning confocal microscope (Leica Microsystems, Germany) with AOBS (Acoust-Optical Beam Splitter) detected using PMTs (photomultiplier tubes). Both the 40 x and 63 x HCX PL APO oil lenses, numerical aperture 1.25 and 1.4 respectively, were used. Samples were excited at 514 nm from an Argon-Krypton-mixed gas laser and images were acquired in the green/yellow region of the

light spectrum (525–590 nm). The software used to image was LCS (Leica Confocal Software) and the images were processed using Leica Lite confocal software and FIJI (Schindelin, et al., 2012).

Imaging in *Caenorhabditis elegans*. *C. elegans* were maintained on normal *E. coli* OP50 at 20 °C unless otherwise stated. A food source carrying the fluorescent analogues was made by transforming the OP50 with pET-BAD-btuBF and were grown in a 4 mL LB culture after inoculation with 16 µL of starter culture. This was incubated with 1 µM of C5-BoD-cobyrinic acid or ribose-BoD-cobalamin overnight with induction with 0.02% arabinose. 1 mL of culture was spun down and washed three times with fresh LB to remove external fluorophore. The final pellet was resuspended in 1 mL of fresh LB and 200 µL of the OP50 culture was pipetted on to the centre of NGM agar plates. These plates were left to dry in a sterile culture hood for 4 hours and then stored at 4 °C until used. Three fourth larval stage (L4) *C. elegans* were transferred to the seeded plates and left to grow at 20 °C for four days before imaging. Three third larval stage (L3) *C. elegans* were transferred to the seeded plates and allowed to lay eggs. These progeny were then imaged at the L4 stage.

For each slide, 3-5 L4 stage *C. elegans* were mounted in M9 + 0.2 % levamisole on a 2 % agarose pad and imaged within 30 min at the University of Bristol on a Leica SP8X AOBS confocal laser scanning microscope attached to a Leica DMI8 inverted epifluorescence microscope with 'Adaptive Focus Control'. The sample was excited with a white light laser at 594 nm and detected between 599-712 nm using a hybrid gated detector for the fluorophore, and excited with a 405 nm laser, with images acquired between 410-505 nm for the gut granule autofluorescence. All images were taken using a 20 x numerical aperture 0.75 dry lens. Again the images were processed using Leica Lite confocal software and FIJI (Schindelin, et al., 2012).

HPLC MS analysis. Samples were separated on an Agilent 1000 series HPLC coupled to a micrOTOF-Q II (Bruker) mass spectrometer using an Ace 5 AQ column (2.1 x 150 mm; Advanced Chromatography Technologies) maintained at 30°C and with a flow rate of 0.2 mL/min. The mobile phase consisted of 0.1% TFA (v/v) in water (solvent A) and 100% acetonitrile (solvent B). Three different HPLC gradients were employed. Gradient 1 (methionine and SAM and analogues): Linear gradient from 0-30% solvent B over 30 mins. Gradient 2 (pathway intermediates): 5-20% B over 6 mins, 20-30% B over 19 mins followed by 30-90% B over 5 mins. Gradient 3 (fluorescent cobalamin analogues): Linear gradient from 0-100% solvent B over 45 mins. Semi-preparative HPLC was performed on an Agilent 1000 series HPLC using a CS SIL C18 column (250 mm x 10 mm; 5 µm; Charlton Scientific) at a flow rate of 5 mL/min running the gradients described above.

NMR analysis. NMR experiments were performed using a 600 MHz (^1H) Bruker Avance III spectrometer with a 5 mm QCI-F cryoprobe. Assignments were completed using DQF-COSY, TOCSY (20 ms and 80 ms mixing times), NOESY (500 ms mixing time), ^{13}C -HSQC-TOCSY, ^{13}C , ^1H -HSQC and ^{13}C , ^1H -HMBC experiments with WaterGATE or excitation sculpting water suppression on 0.1-1 mM samples dissolved in deuterium oxide. All ^{13}C data was obtained using natural isotopic abundance. Data were processed and analysed using Bruker Topspin version 3.2 and CCPN analysis. ^1H Chemical shift referencing was based on the position of the water resonance, ^{13}C referencing used $^1\text{H}/^{13}\text{C}$ gyromagnetic ratios to define indirect carrier position (Wishart and Case, 2001) and all data were obtained at 25°C.

QUANTIFICATION AND STATISTICAL ANALYSIS

Bioassays (Figure 5 and Supplementary Figure 6) were performed three times ($n=3$) and the values for the amount of B_{12} present in the samples was generated from a standard curve. For B_{12} uptake into plants the averages of the separate measurements were then calculated and plotted using Microsoft Excel. The structure of CobH was refined with internal statistical analysis as reported in the deposition in the PDB

No additional statistical tests were performed.

DATA AND SOFTWARE AVAILABILITY

All software used in this study is reported in Method Details and indicated in the Key Resources Table. The coordinates for the CobH T85A crystal structure were deposited in the Protein Data Bank PDB:5N0G (<https://www.rcsb.org/structure/5n0g>).