

Kent Academic Repository

Full text document (pdf)

Citation for published version

Lampe, P. (2013) Quantum cluster algebras of type A and the dual canonical basis. Proceedings of the London Mathematical Society, 108 (1). pp. 1-43. ISSN 00246115.

DOI

<https://doi.org/10.1112/plms/pds098>

Link to record in KAR

<http://kar.kent.ac.uk/67634/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Quantum cluster algebras of type A and the dual canonical basis

Philipp Lampe

May 24, 2013

Abstract

The article concerns the subalgebra $U_v^+(w)$ of the quantized universal enveloping algebra of the complex Lie algebra \mathfrak{sl}_{n+1} associated with a particular Weyl group element of length $2n$. We verify that $U_v^+(w)$ can be endowed with the structure of a quantum cluster algebra of type A_n . The quantum cluster algebra is a deformation of the ordinary cluster algebra Geiß-Leclerc-Schröer attached to w using the representation theory of the preprojective algebra. Furthermore, we prove that the quantum cluster variables are, up to a power of v , elements in the dual of Lusztig's canonical basis under Kashiwara's bilinear form.

1 Introduction

Cluster algebras are commutative algebras created in 2000 by Fomin-Zelevinsky [18] in the hope to obtain a combinatorial description of the dual of Lusztig's *canonical basis* of a quantum group.

A cluster algebra comes with a distinguished set of generators called *cluster variables*. Each cluster variable belongs to several overlapping *clusters*. Every cluster, and hence every cluster variable, is obtained from an initial cluster by a sequence of *mutations*. Every mutation replaces an element in a cluster by an explicitly defined rational function in the variables of that cluster. We refer to Fomin-Zelevinsky [18] for definitions and to Fomin-Zelevinsky [22] for a good survey about cluster algebras.

It quickly turned out that Fomin-Zelevinsky's theory of cluster algebras has many interesting applications and coheres with various mathematical objects. Let us mention the representation theory of quivers and finite-dimensional algebras, the representation theory of preprojective algebras, root systems of Kac-Moody algebras, Calabi-Yau categories, quantum groups, and Lusztig's canonical basis of universal enveloping algebras.

A momentous step in the development was the *additive categorification* of *acyclic* cluster algebras by *cluster categories*. Cluster categories were defined by Buan-Marsh-Reineke-Reiten-Todorov [5] and independently by Caldero-Chapoton-Schiffler [11] for type A . The cluster category \mathcal{C}_Q associated with a quiver Q is an orbit category of the bounded derived category of the category of representations of Q . Keller [35] proved that cluster categories are triangulated categories. Key ingredients for the verification of the additive categorification of acyclic cluster algebras by cluster categories are due to Buan-Marsh-Reiten-Todorov [8], Buan-Marsh-Reiten [6, 7], Geiß-Leclerc-Schröer [25, 27], and Caldero-Keller [12, 13]. The process of mutation in the cluster algebra

resembles the process of *tilting* in the cluster category. Hence, we obtain a link between quiver representations and triangulated categories on one side and a large class of cluster algebras on the other side.

Furthermore, Geiß-Leclerc-Schröer [28] provided an additive categorification by categories arising from the study of *Kac-Moody groups* and *unipotent cells*. In this construction the categorified cluster algebras are not necessarily acyclic. Let \mathfrak{g} be the Kac-Moody Lie algebra attached to Q and let $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}$ be its triangular decomposition. Geiß-Leclerc-Schröer's construction is related to the *preprojective algebra* Λ associated with Q . Buan-Iyama-Reiten-Scott [4] attached to every element w in the Weyl group of \mathfrak{g} a subcategory $\mathcal{C}_w \subset \text{mod}(\Lambda)$. Geiß-Leclerc-Schröer [28] endow the coordinate ring $\mathbb{C}[N(w)]$ of the unipotent group $N(w)$ with the structure of a cluster algebra $\mathcal{A}(w)$. Here, N denotes the pro-unipotent pro-group associated with the completion $\hat{\mathfrak{n}}$ and $N(w) = N \cap (w^{-1}N_-w)$. The coordinate ring $\mathbb{C}[N(w)]$ is naturally isomorphic to a subalgebra of the graded dual $U(\mathfrak{n})_{gr}^*$ of the universal enveloping algebra of \mathfrak{n} . The cluster variables are δ -functions of rigid modules over the preprojective algebra. All cluster monomials lie in the dual semicanonical basis.

Let us mention that cluster algebras also gained popularity in other branches of mathematics; for example, for Poisson geometry, see Gekhtman-Shapiro-Vainshtein [30], for Teichmüller theory, see Fock-Goncharov [16], for combinatorics, see Musiker-Propp [43], for integrable systems, see Fomin-Zelevinsky [20], for Donaldson-Thomas invariants and mathematical physics see Kontsevich-Soibelman [37], etc.

In this article we transfer to the quantized setup. We consider the case $\mathfrak{g} = \mathfrak{sl}_{n+1}$ (for some natural number n) and choose a particular Weyl group element w of length $2n$ which is equal to the square of a Coxeter element. The reasons for this particular choice are the following three: First, in this case the stable category $\underline{\mathcal{C}}_w$ is triangle equivalent to the corresponding cluster category \mathcal{C}_Q by a result of Geiß-Leclerc-Schröer [28, Theorem 11.1]; second, there exist recursions for quantum cluster variables (simpler than the mutation relations) which allow to connect quantum cluster variables with canonical basis elements; third, cluster algebras of almost all types can be realized as $\mathcal{A}(w)$ for some square w of a Coxeter element in the Weyl group of a Kac-Moody Lie algebra of the same type, see Geiß-Leclerc-Schröer [29, Section 2.6].

We establish a quantum cluster algebra structure on a subalgebra $U_v^+(w)$ of the *quantized* universal enveloping algebra $U_v(\mathfrak{n})$ of \mathfrak{n} . Quantum cluster algebras were introduced by Berenstein-Zelevinsky [10]. The quivers corresponding to $\mathfrak{g} = \mathfrak{sl}_{n+1}$ are *Dynkin* quivers of type A_n . We choose a particular orientation: let $Q = (Q_0, Q_1)$ be the Dynkin quiver of type A_n with an alternating orientation beginning with a source. We denote the set of vertices by $Q_0 = \{1, 2, \dots, n\}$. Figure 1 illustrates the example $n = 13$. The choice of the orientation matches the choice of the Weyl group element w . The reduced expression of w (that is used to construct $U_v^+(w)$) and its initial subsequences (that are used to construct the generators $U_v^+(w)$) are related to the indecomposable injective modules over the *path algebra* of Q and their *Auslander-Reiten translates*, respectively.




Figure 1: The quiver Q of type A_{13}

The construction of $U_v^+(w)$ is due to Lusztig [42]. The algebra is generated by $2n$ elements that satisfy straightening relations; it degenerates to a commutative algebra in the classical limit $v = 1$. The generators are constructed via Lusztig's T -automorphisms. The quantized universal enveloping algebra $U_v(\mathfrak{n})$ is a self-dual Hopf algebra. Also, the subalgebra $U_v^+(w)$ is isomorphic to its dual, i.e., it is isomorphic to the quantized coordinate ring $\mathbb{C}_v[N(w)]$. The algebra $U_v^+(w)$ possesses several distinguished bases, including a *Poincaré-Birkhoff-Witt basis* for every reduced expression for w , a *canonical basis*, and their duals. The article concerns the dual of Lusztig's canonical basis under Kashiwara's bilinear form [33].

It is conjectured (see for example Kimura [36]) that the quantized coordinate rings $\mathbb{C}_v[N(w)]$ are *quantum cluster algebras* $\mathcal{A}_v(w)$ in general and that the set \mathcal{M}_v of all quantum cluster monomials, taken up to powers of v , is a subset of the dual canonical basis \mathcal{B}^* , i.e., the following diagram commutes:

$$\begin{array}{ccc} \mathcal{A}_v(w) & \longrightarrow & \mathbb{C}_v[N(w)] \subset U_v(\mathfrak{n})_{gr}^* \\ \uparrow & & \uparrow \\ \mathcal{M}_v & \hookrightarrow & \mathcal{B}^* \end{array}$$

The conjecture has only been verified in very few cases, see Berenstein-Zelevinsky [9] for type A_2 and A_3 , and the author [38] for an example of Kronecker type.

We verify that in our case the integral form $U_v^+(w)_{\mathbb{Z}}$ of $U_v^+(w)$ is (after extending coefficients) a quantum cluster algebra. The proof relies on the exact form of the straightening relations. The description of the straightening relations features (besides Lusztig's T -automorphisms) Leclerc's embedding [39] of $U_v(\mathfrak{n})$ in the *quantum shuffle algebra*. The exact form of the straightening relations enables us to verify that recursively defined variables satisfy a lattice property and an invariance property so that they are elements in the dual canonical basis.

The cluster algebra $\mathcal{A}(w)$, just as the quantum cluster algebra $\mathcal{A}_v(w)$, is of type A_n . Every cluster contains n *frozen* and n *mutable* cluster variables. Altogether there are $n + \frac{n(n+1)}{2}$ mutable and n frozen cluster variables. Most of the cluster variables can be realized as minors of certain matrices, see Section 2.5. The structure of these minors implies that there is (besides the usual cluster exchange relation) a recursive way to compute these cluster variables avoiding denominators. Theorem 4.11, the main theorem, asserts that the recursion can be quantized to a recursion for the corresponding quantum cluster variables. The quantized recursions imply quantum exchange relations so that the integral form $U_v^+(w)_{\mathbb{Z}}$ of $U_v^+(w)$ becomes (after extending coefficients) a quantum cluster algebra.

Furthermore, it follows from our construction that the quantum cluster variables are (up to a power of v) elements in the dual of Lusztig's canonical basis under Kashiwara's bilinear form [33].

2 Representation theory of the quiver of type A and cluster algebras

2.1 The indecomposable modules over the path algebra

Let k be a field. In what follows we study the category $\text{rep}_k(Q)$ of finite-dimensional k -representations of Q over the field k . (For more detailed information on represen-




Figure 2: The quiver Q' of type A_{12}

tations of quivers see for example Crawley-Boevey [14].) The category $\text{rep}_k(Q)$ is equivalent to the category $\text{mod}(kQ)$ of finite-dimensional modules over the *path algebra* kQ . Gabriel's theorem [23] asserts that the quiver Q admits (up to isomorphism) only finitely many indecomposable representations. In fact there are $\frac{(n+1)n}{2}$ indecomposable representations (up to isomorphism) and they are in bijection with the set of intervals $[i, j] = \{i, i+1, i+2, \dots, j\} \subset \mathbb{Z}$ with $1 \leq i \leq j \leq n$. The indecomposable representation associated with the interval $[i, j]$ is $V_{[i, j]} = ((V_s)_{s \in Q_0}, (V_a)_{a \in Q_1})$ defined by k -vector spaces

$$V_s = \begin{cases} k, & \text{if } i \leq s \leq j; \\ 0, & \text{otherwise;} \end{cases}$$

associated with vertices s , and k -linear maps

$$V_a = \begin{cases} 1, & \text{if } V_s = V_t = k; \\ 0, & \text{otherwise;} \end{cases}$$

associated with arrows $a: s \rightarrow t$.

All further considerations will basically depend on the parity of n . For a compact and effective handling of all cases we make the assumption that n is odd. Denote by $Q' = (Q'_0, Q'_1)$ to be the quiver obtained from Q by removing the vertex n . The quiver Q' is of type A_{n-1} , and the examination of both Q' and Q covers all cases. Note that every representation of Q' can be viewed as a representation of Q supported on the first $n-1$ vertices. An example of the quiver Q' is shown in Figure 2.

If $n = 1$, i.e., the quiver has one vertex and no arrows, then the category $\text{rep}_k(Q)$ can easily be described. In this case modules over the path algebra kQ are k -vector spaces, and the k -vector space k of dimension 1 is the only indecomposable module. In the other cases, the most suggestive way to illustrate the category $\text{rep}_k(Q)$ is given by its *Auslander-Reiten quiver*. For an introduction to Auslander-Reiten theory we refer to Assem-Simson-Skowronski [1, Chapter IV].

The simplest non-trivial example is the Auslander-Reiten quiver of type A_2 which can be seen in Figure 3. In this case there are (up to isomorphism) three indecomposable representations, two of which are injective. The representations are displayed by numbers that represent basis vectors and composition series; cf. Geiß-Leclerc-Schröer [28, Section 7.5]. The solid arrows represent *irreducible maps*; the dashed arrow represents the *Auslander-Reiten translation*. Note that the Auslander-Reiten translate of the injective representation associated with vertex 2 is the zero representation.

In what follows we are interested in the indecomposable injective kQ -modules I_i associated with vertices $i \in Q_0$ and their Auslander-Reiten translates $\tau_{kQ}(I_i)$. Similarly, we are interested in the indecomposable injective kQ' -modules I'_i associated with vertices $i \in Q'_0$ and their Auslander-Reiten translates $\tau_{kQ'}(I'_i)$. (For simplicity, we drop from now on the index attached to τ whenever it is clear which algebra we are




Figure 3: The Auslander-Reiten quiver for $n = 2$

referring to.) The choice of the alternating orientations of the quivers Q and Q' ensure that from type A_3 onwards we have $\tau(I) \neq 0$ for every indecomposable injective kQ -module. (This would not be true for the linear orientation of the Dynkin diagram A_n . The Auslander-Reiten translate of the indecomposable injective representation corresponding to the sink would be zero in this case.) The direct sum $M = \bigoplus_{i=1}^n I_i \oplus \tau(I_i)$ is a *terminal* kQ -module in the sense of Geiß-Leclerc-Schröer [28, Section 2.2], and so is the kQ' module $M' = \bigoplus_{i=1}^{n-1} I'_i \oplus \tau(I'_i)$.




Figure 4: The Auslander-Reiten quiver for $n = 3$

The small cases A_3 and A_4 will have to be treated separately. Figure 4 and Figure 5 display the indecomposable injective modules (red), their Auslander-Reiten translates (blue), and irreducible maps between them for the case $n = 3$ and $n = 4$, respectively.

If $n = 3$, then M is the direct sum of all indecomposable kQ -modules, i.e., $\text{mod}(kQ) = \text{add}(M)$.




Figure 5: A part of the Auslander-Reiten quiver for $n = 4$

From A_5 onwards a uniform description is possible. For type A_n (remember that n is assumed to be odd) the indecomposable components of M can be written down

explicitly:

$$\begin{aligned}
I_i &= V_{[i,i]}, & \text{if } i \text{ is odd and } 1 \leq i \leq n, \\
I_i &= V_{[i-1,i+1]}, & \text{if } i \text{ is even and } 2 \leq i \leq n, \\
\tau(I_1) &= V_{[2,3]}, \\
\tau(I_i) &= V_{[i-2,i+2]}, & \text{if } i \text{ is odd and } 3 \leq i \leq n-2, \\
\tau(I_n) &= V_{[n-2,n-1]}, \\
\tau(I_2) &= V_{[2,5]}, \\
\tau(I_i) &= V_{[i-3,i+3]}, & \text{if } i \text{ is even and } 4 \leq i \leq n-3, \\
\tau(I_{n-1}) &= V_{[n-4,n-1]}.
\end{aligned}$$

We display the relevant part of the Auslander-Reiten quiver of A_n in Figure 6 for the case $n = 13$. As above, the indecomposable injective modules are colored red, their Auslander-Reiten translates blue.

There are only a few changes if we restrict Q to Q' . Observe that $I'_i = I_i$ for $i \in \{1, 2, \dots, n-3\}$, and that $\tau(I'_i) = \tau(I_i)$ for $i \in \{1, 2, \dots, n-3\}$. Note that the latter modules are kQ -modules supported on the first $n-1$ vertices and may therefore be viewed as kQ' -modules. Furthermore, we have

$$\begin{aligned}
I'_{n-1} &= V_{[n-2,n-1]}, \\
\tau(I'_{n-3}) &= V_{[n-6,n-1]}, \\
\tau(I'_{n-2}) &= V_{[n-4,n-1]}, \\
\tau(I'_{n-1}) &= V_{[n-4,n-3]}.
\end{aligned}$$

An example of type A_{12} is illustrated in Figure 7.

2.2 The preprojective algebra and rigid modules

The representation theory of the path algebra kQ is closely related to the representation theory of the corresponding *preprojective algebra* Λ defined as follows. For every arrow $a: s \rightarrow t$ in Q_1 introduce an additional arrow $a^*: t \rightarrow s$ in reverse direction and denote by $Q_1^* = \{a^*: a \in Q_1\}$ the set of all reversed arrows. The *double quiver* of Q is defined to be the quiver $\bar{Q} = (\bar{Q}_0, \bar{Q}_1)$ given by a vertex set $\bar{Q}_0 = Q_0$ and an arrow set $\bar{Q}_1 = Q_1 \cup Q_1^*$. The preprojective algebra is defined to be

$$\Lambda = k\bar{Q}/(c)$$

where the ideal (c) is the two-sided ideal generated by the element

$$c = \sum_{a \in Q_1} (a^*a - aa^*) \in k\bar{Q}.$$

The algebra Λ is finite-dimensional, since Q is an orientation of a Dynkin diagram, see Reiten [45, Theorem 2.2a]. The category $\text{mod}(\Lambda)$ of finite-dimensional Λ -modules is equivalent to the category $\text{rep}_k(\bar{Q}, (c))$ of finite-dimensional representations $M = ((M_s)_{s \in Q_0}, (M_a)_{a \in \bar{Q}_1})$ of \bar{Q} such that for any two vertices $s, t \in Q_0$ and any linear combination $\sum_{i=1}^m \lambda_i p_i \in (c)$ of paths $p_i: s \rightarrow t$ with scalars $\lambda_i \in k$ the associated linear map $\sum_{i=1}^m \lambda_i M_{p_i}$ is zero.




Figure 6: A part of the Auslander-Reiten quiver of $\text{mod}(kQ)$




Figure 7: A part of the Auslander-Reiten quiver of $\text{mod}(kQ')$

$$T_{i,[0,0]} = \begin{cases} \begin{array}{c} i \end{array} & \text{if } i \text{ is odd and } 1 \leq i \leq n \\ \begin{array}{cc} i-1 & i+1 \\ & i \end{array} & \text{if } i \text{ is even and } 2 \leq i \leq n-1 \end{cases}$$

Figure 8: The modules $T_{i,[0,0]}$

There is a *restriction functor* $\pi_Q: \text{mod}(\Lambda) \rightarrow \text{mod}(kQ)$ given by forgetting the linear maps associated with a^* for all $a \in Q_1$ in the corresponding representation of the quiver \bar{Q} . Ringel [46, Theorem B] proved that the category $\text{mod}(\Lambda)$ is equivalent to the category $C(1, \tau)$ whose objects are pairs (X, f) consisting of a kQ -module X and a kQ -module homomorphism $f: X \rightarrow \tau(X)$ from X to its translate $\tau(X)$ and where a morphism $h: (X, f) \rightarrow (Y, g)$ is given by a kQ -module homomorphism $h: X \rightarrow Y$ for which the diagram

$$\begin{array}{ccc} X & \xrightarrow{h} & Y \\ \downarrow f & & \downarrow g \\ \tau(X) & \xrightarrow{\tau(h)} & \tau(Y) \end{array}$$

commutes.

Using the correspondence from above Geiß-Leclerc-Schröer [28, Section 7.1] constructed for every $i \in Q_0$ and any natural numbers a, b satisfying $0 \leq a \leq b \leq 1$ a Λ -module $T_{i,[a,b]} = (I_{i,[a,b]}, e_{i,[a,b]})$ where $I_{i,[a,b]} = \bigoplus_{j=a}^b \tau^j(I_i)$ and the map

$$e_{i,[a,b]}: I_{i,[a,b]} \rightarrow \tau(I_{i,[a,b]}) = \bigoplus_{j=a+1}^{b+1} \tau^j(I_i)$$

is identity on every $\tau^j(I_i)$ for $a+1 \leq j \leq b$ and zero otherwise. We study Λ -modules $T_{i,[a,b]}$ for $i \in Q_0$ and $0 \leq a, b \leq 1$. We display the modules in Figures 8, 9, 10.

The modules $T_{i,[a,b]}$ for $i \in Q_0$ and $0 \leq a, b \leq 1$ are *rigid* and *nilpotent*. Recall that a Λ -module T is said to be rigid if $\text{Ext}_{\Lambda}^1(T, T) = 0$ and it is said to be nilpotent if there exists an integer $N > 0$ such that for each path $a_1 a_2 \cdots a_N$ of length N in \bar{Q} the associated linear map $T_{a_1} T_{a_2} \cdots T_{a_N}$ is zero. Rigidity follows from Geiß-Leclerc-Schröer [28, Lemma 7.1]; nilpotency follows from Lusztig [41, Proposition 14.2].

Similarly, the representation theory of the path algebra kQ' is closely related to the representation theory of the corresponding preprojective algebra Λ' .

2.3 Notations from Lie theory

The representation theory of the quiver Q is related with Lie theory. Let $k = \mathbb{C}$. The Lie algebra associated with the Dynkin diagram A_n is $\mathfrak{g} = \mathfrak{sl}_{n+1}$, i.e., the Lie algebra of $(n+1) \times (n+1)$ matrices with complex entries and vanishing trace. It admits a triangular decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}$. Here, \mathfrak{n} and \mathfrak{n}_- denote the Lie algebras of strictly upper and strictly lower triangular $(n+1) \times (n+1)$ matrices, respectively, and \mathfrak{h} denotes the Lie algebra of $(n+1) \times (n+1)$ diagonal matrices. The Lie algebra \mathfrak{n} is called the *positive part* of \mathfrak{g} .




Figure 9: The modules $T_{i,[0,1]}$




Figure 10: The modules $T_{i,[1,1]}$

Let $C = (a_{ij})_{1 \leq i, j \leq n}$ be the *Cartan matrix* associated with the quiver Q ; its entries are:

$$a_{ij} = \begin{cases} 2, & \text{if } i = j; \\ -1, & \text{if } |i - j| = 1; \\ 0, & \text{otherwise.} \end{cases}$$

The Lie algebra $\mathfrak{g} = \mathfrak{sl}_{n+1}$ is studied by its *roots*. The *root lattice* R is defined to be the free abelian group generated by $\alpha_1, \alpha_2, \dots, \alpha_n$. Here, the elements $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathfrak{h}^*$ are defined by $\alpha_i(\text{diag}(d_1, d_2, \dots, d_n)) = d_i - d_{i+1}$ and they are called *simple roots*. By $R^+ \subset R$ we denote the set of all linear combinations $\sum_{i=1}^n d_i \alpha_i$ with $d_i \in \mathbb{N}$. There is a symmetric bilinear form $(\cdot, \cdot): R \times R \rightarrow \mathbb{R}$ which satisfies $(\alpha_i, \alpha_j) = a_{ij}$ for $1 \leq i, j \leq n$. By $\Delta^+ \subseteq R$ we denote the set of *positive roots* of the corresponding root system. Then $\Delta^+ = \{\alpha_i + \alpha_{i+1} + \dots + \alpha_j : 1 \leq i \leq j \leq n\}$. Under the bijection of Gabriel's theorem, a positive root $\alpha_i + \alpha_{i+1} + \dots + \alpha_j$ with $1 \leq i \leq j \leq n$ is mapped to the indecomposable representation $V_{[i,j]}$ from Section 2.1.

The simple reflections $s_1, s_2, \dots, s_n: \mathfrak{h}^* \rightarrow \mathfrak{h}^*$ act on the simple roots by

$$s_i(\alpha_j) = \begin{cases} -\alpha_i, & \text{if } i = j; \\ \alpha_i + \alpha_j, & \text{if } |i - j| = 1; \\ \alpha_j, & \text{otherwise.} \end{cases}$$

The group W generated by the simple reflections is called the *Weyl group* of type \mathfrak{g} . The simple reflection satisfy the following relations

$$s_i s_j = s_j s_i, \quad \text{if } |i - j| \geq 2; \quad (1)$$

$$s_i s_j s_i = s_j s_i s_j, \quad \text{if } |i - j| = 1; \quad (2)$$

$$s_i^2 = 1, \quad (3)$$

for all $1 \leq i, j \leq n$. Therefore, the Weyl group W is isomorphic to the symmetric group S_n .

To every terminal kQ -module Geiß-Leclerc-Schröer [28, Section 3.7] attach a Q^{op} -adapted Weyl group element. The Q^{op} -adapted Weyl group element associated with the terminal module M from Section 2.1 is $w = s_1 s_3 s_5 \dots s_n s_2 s_4 s_6 \dots s_{n-1} s_1 s_3 s_5 \dots s_n s_2 s_4 s_6 \dots s_{n-1}$. The given expression for w is reduced. Let $j_1, j_2, \dots, j_{2n} \in [1, n]$ such that for the reduced expression for w from above we have $w = s_{j_1} s_{j_2} \dots s_{j_{2n}}$. We put $\beta_1 = \alpha_{j_1}$ and $\beta_k = s_{j_1} s_{j_2} \dots s_{j_{k-1}}(\alpha_{j_k})$ for $2 \leq k \leq 2n$. Denote by $\Delta_w^+ = \{\beta_1, \beta_2, \dots, \beta_{2n}\} \subseteq \Delta^+$ the set of all β_k with $1 \leq k \leq 2n$. Note that the notation is well-defined. If we choose another reduced expression $w = s_{j'_1} s_{j'_2} \dots s_{j'_{2n}}$ for w , then

$$\left\{ s_{j'_1} s_{j'_2} \dots s_{j'_{k-1}}(\alpha_{j'_k}) : 1 \leq k \leq 2n \right\} = \{\beta_1, \beta_2, \dots, \beta_{2n}\}.$$

Furthermore, notice that under the bijection of Gabriel's theorem, the $2n$ positive roots β_k with $1 \leq k \leq 2n$, correspond to the dimension vectors of the indecomposable

direct summands of M (compare Figure 1). More precisely, for $n \geq 5$,

$$\begin{aligned} \Delta_w^+ &= \{\alpha_i : \text{is odd and } 1 \leq i \leq n\} \\ &\cup \{\alpha_{i-1} + \alpha_i + \alpha_{i+1} : \text{is even and } 2 \leq i \leq n-1\} \\ &\cup \{\alpha_2 + \alpha_3\} \cup \{\alpha_{n-2} + \alpha_{n-1}\} \\ &\cup \{\alpha_{i-2} + \cdots + \alpha_{i+2} : \text{is odd and } 3 \leq i \leq n-3\} \\ &\cup \{\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5\} \cup \{\alpha_{n-4} + \alpha_{n-3} + \alpha_{n-2} + \alpha_{n-1}\} \\ &\cup \{\alpha_{i-3} + \cdots + \alpha_{i+3} : \text{is even and } 4 \leq i \leq n-4\}. \end{aligned}$$

The universal enveloping algebra $U(\mathfrak{n})$ of \mathfrak{n} is the associative \mathbb{C} -algebra generated by E_i ($1 \leq i \leq n$) subject to the relations

$$E_i E_j = E_j E_i, \quad \text{for } |i - j| \geq 2, \quad (4)$$

$$E_i^2 E_j - 2E_i E_j E_i + E_j E_i^2 = 0, \quad \text{for } |i - j| = 1. \quad (5)$$

The last relation is called *Serre relation*.

Similarly, the representation theory of the quiver Q' of type A_{n-1} is linked with the Lie algebra $\mathfrak{g}' = \mathfrak{sl}_n$ with Weyl group W' . The Lie algebra $\mathfrak{g}' = \mathfrak{sl}_n$ similarly admits a triangular decomposition $\mathfrak{g}' = \mathfrak{n}'_- \oplus \mathfrak{h}' \oplus \mathfrak{n}'_+$. The Weyl group element associated with M' is $w' = s_1 s_3 s_5 \cdots s_{n-2} s_2 s_4 s_6 \cdots s_{n-1} s_1 s_3 s_5 \cdots s_{n-2} s_2 s_4 s_6 \cdots s_{n-1} \in W'$. The universal enveloping algebra $U(\mathfrak{n}')$ may be viewed as the subalgebra of $U(\mathfrak{n})$ generated by E_i ($1 \leq i \leq n-1$).

2.4 The cluster algebra attached to the terminal module

To the terminal $\mathbb{C}Q$ -module M from Section 2 Geiß-Leclerc-Schröer ([28, Section 4]) attached the subcategory $\mathcal{C}_M = \pi_Q^{-1}(\text{add}(M)) \subseteq \text{nil}(\Lambda)$ of the category of nilpotent Λ -modules. Here, $\text{add}(M) \subseteq \text{mod}(\mathbb{C}Q)$ is the subcategory consisting of all modules isomorphic to direct summands of direct sums of finitely many copies of M .

The projective and injective objects in \mathcal{C}_M coincide, so \mathcal{C}_M is a *Frobenius category* and the stable category $\underline{\mathcal{C}}_M$ is triangulated according to Happel [31, Section 2.6]. Furthermore, Geiß-Leclerc-Schröer [28, Theorem 11.1] showed that there is an equivalence of triangulated categories $\underline{\mathcal{C}}_M \simeq \mathcal{C}_Q$ between $\underline{\mathcal{C}}_M$ and the *cluster category* \mathcal{C}_Q as defined by Buan-Marsh-Reineke-Reiten-Todorov [5] to be the orbit category $\mathcal{D}^b(\text{mod}(\mathbb{C}Q)) / \tau_{\mathcal{D}}^{-1} \circ [1]$.

With every \mathcal{C}_M Geiß-Leclerc-Schröer [28, Section 4] associated a cluster algebra $\mathcal{A}(\mathcal{C}_M)$; it is constructed as a subalgebra of the graded dual of the universal enveloping algebra of the positive part of the corresponding Lie algebra, i.e., $\mathcal{A}(\mathcal{C}_M) \subseteq U(\mathfrak{n})_{gr}^*$. For a definition of and a general introduction to cluster algebras see Fomin-Zelevinsky [21]. The cluster algebra $\mathcal{A}(\mathcal{C}_M)$ is also called $\mathcal{A}(w)$.

There is an isomorphism between $U(\mathfrak{n})$ and an algebra \mathcal{M} of \mathbb{C} -valued functions on $\text{mod}(\Lambda)$. We refer to Geiß-Leclerc-Schröer [28] for a precise definition of \mathcal{M} . It is generated by functions d_i that map a Λ -module X to the Euler characteristic of the flag variety of X of type \mathbf{i} . Prominent elements in $\mathcal{A}(\mathcal{C}_M)$ are (under the described isomorphism) the δ -functions of the rigid Λ -modules $T_{i,[a,b]}$ with $i \in Q_0$ and $0 \leq a \leq b \leq 1$. For $1 \leq i \leq n$ put

$$\begin{aligned} P_i &= \delta_{T_{i,[0,1]}}; \\ Y_i &= \begin{cases} \delta_{T_{i,[0,0]}} & \text{if } i \text{ is odd;} \\ \delta_{T_{i,[1,1]}} & \text{if } i \text{ is even;} \end{cases} \end{aligned}$$




Figure 11: The initial seed for the case $n = 9$

$$Z_i = \begin{cases} \delta_{T_{i,[0,0]}}, & \text{if } i \text{ is even;} \\ \delta_{T_{i,[1,1]}}, & \text{if } i \text{ is odd.} \end{cases}$$

Note that the module $T_{i,[0,1]}$ corresponding to the variable P_i (for $1 \leq i \leq n$) is a projective object in the category \mathcal{C}_M , but it is in general not projective in $\text{mod}(\Lambda)$.

The initial seed of the cluster algebra $\mathcal{A}(\mathcal{C}_M)$ for the case $n = 9$ is shown in Figure 11. The vertices represent the cluster variables in the initial cluster, the arrows describe the initial exchange matrix. Just as in Keller's mutation applet [34], the blue vertices are frozen, the red vertices are mutable. The frozen variables P_1, P_2, \dots, P_n may be viewed as coefficients in the sense of Fomin-Zelevinsky [22]. The cluster algebra $\mathcal{A}(\mathcal{C}_M)$ is of type A_n , and therefore of finite type. Besides the n frozen variables there are $n + \frac{n(n+1)}{2}$ mutable cluster variables grouped into $C_{n+1} = \frac{1}{n+2} \binom{2n+2}{n+1}$ clusters, where C_{n+1} denotes the $(n+1)$ th Catalan number (see Fomin-Zelevinsky [19, Section 12]). The Catalan number C_{n+1} is the number of triangulations of a convex polygon with $n+3$ sides using only diagonals.

The δ -functions of P_i, Y_i , and Z_i for $i \in Q_0$ are not algebraically independent. For example, the equation

$$P_i = Y_i Z_i - Z_{i-1} Z_{i+1} \tag{6}$$

holds for every $i \in Q_0$. The equations are due to Geiß-Leclerc-Schröer [28, Theorem 18.1] and called *determinantal identities*. Here and in what follows we use the convention $Z_0 = Z_{n+1} = 1$.

Similarly, we can construct a cluster algebra $\mathcal{A}(\mathcal{C}_{M'})$ associated with the terminal $\mathbb{C}Q'$ -module M' from Section 2. The initial seed of $\mathcal{A}(\mathcal{C}_{M'})$ is obtained from the initial seed of $\mathcal{A}(\mathcal{C}_M)$ by ignoring the vertices Y_n and P_n and all incident arrows. We denote the corresponding cluster variables of $\mathcal{A}(\mathcal{C}_{M'})$ by P'_i, Y'_i , and Z'_i (for $1 \leq i \leq n-1$).

2.5 The description of cluster variables

In this subsection we describe the cluster variables explicitly. Note that our description of cluster variables differs from the explicit description of Geiß-Leclerc-Schröer [28, Section 18.2] due to a different choice of orientation of the quiver. Put $c_i = \frac{Z_{i-1} Z_{i+1}}{Z_i}$ for $1 \leq i \leq n$.

Definition 2.1. For two natural numbers i, j with $1 \leq i \leq j \leq n$ let $M_{ij} = ((M_{ij})_{rs})_{i \leq r, s \leq j}$ be the $(j-i+1) \times (j-i+1)$ matrix defined by

$$(M_{ij})_{rs} = \begin{cases} \frac{Y_r}{c_r}, & \text{if } r = s; \\ 1, & \text{if } s > r \text{ or } r = s + 1; \\ 0, & \text{otherwise.} \end{cases}$$

Put $\Delta_{i,j} = c_i c_{i+1} \cdots c_j \cdot \det(M_{ij})$, i.e., $\Delta_{i,j}$ is given by the following determinant:

$$\Delta_{i,j} = c_i c_{i+1} \cdots c_j \begin{vmatrix} \frac{Y_i}{c_i} & 1 & 1 & 1 & \cdots & 1 & 1 & 1 \\ 1 & \frac{Y_{i+1}}{c_{i+1}} & 1 & 1 & \cdots & 1 & 1 & 1 \\ 0 & 1 & \frac{Y_{i+2}}{c_{i+2}} & 1 & \cdots & 1 & 1 & 1 \\ 0 & 0 & 1 & \frac{Y_{i+3}}{c_{i+3}} & \cdots & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \frac{Y_{j-2}}{c_{j-2}} & 1 & 1 \\ 0 & 0 & 0 & 0 & \cdots & 1 & \frac{Y_{j-1}}{c_{j-1}} & 1 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 & \frac{Y_j}{c_j} \end{vmatrix}.$$

Remark 2.2. Note that

$$\Delta_{i,j} = \begin{vmatrix} Y_i & c_i & c_i & c_i & \cdots & c_i & c_i & c_i \\ c_{i+1} & Y_{i+1} & c_{i+1} & c_{i+1} & \cdots & c_{i+1} & c_{i+1} & c_{i+1} \\ 0 & c_{i+2} & Y_{i+2} & c_{i+2} & \cdots & c_{i+2} & c_{i+2} & c_{i+2} \\ 0 & 0 & c_{i+3} & Y_{i+3} & \cdots & c_{i+3} & c_{i+3} & c_{i+3} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & Y_{j-2} & c_{j-2} & c_{j-2} \\ 0 & 0 & 0 & 0 & \cdots & c_{j-1} & Y_{j-1} & c_{j-1} \\ 0 & 0 & 0 & 0 & \cdots & 0 & c_j & Y_j \end{vmatrix}$$

for $1 \leq i \leq j \leq n$. It follows that each $\Delta_{i,j}$ ($1 \leq i, j \leq n$) is actually a polynomial in Y_i ($1 \leq i \leq n$) and Z_i ($1 \leq i \leq n$), i.e., $\Delta_{i,j} \in \mathbb{Z}[Y_k, Z_k : 1 \leq k \leq n]$ for all $i \leq j$. Polynomiality follows from Geiß-Leclerc-Schröer [28, Theorem 3.4], but it also follows directly from the formula above once we notice that $c_i c_{i+1} \cdots c_j \in \mathbb{Z}[Z_k : 1 \leq k \leq n]$ for all i, j with $1 \leq i < j \leq n$.

Proposition 2.3. For all i, j with $1 \leq i \leq j \leq n$ and $j - i \geq 3$ the equation

$$\Delta_{i,j} = Y_j \Delta_{i,j-1} - Z_{j+1} P_{j-2} \Delta_{i,j-3}$$

holds.

Proof. Perform a Laplace expansion of the determinant on the last row. The last row has only two non-zero entries and it is easy to see that the two occurring summands in the Laplace expansion are the two summands in the recursion formula. \square

For $1 \leq i \leq n$ let $\Delta_{i,i-1}$, $\Delta_{i,i-2}$, and $\Delta_{i,i-3}$ be the unique elements from $\mathbb{Q}(Y_k, Z_k : 1 \leq k \leq n)$ such that the recursion formula from Proposition 2.3 also holds for $j = i+2$, $j = i+1$, and $j = i$. Explicitly, we put $\Delta_{i,i-1} = 1$, $\Delta_{i,i-2} = \frac{1}{Y_{i-1} - c_{i-1}}$, and $\Delta_{i,i-3} = 0$. The next lemma follows easily from Proposition 2.3.




Figure 12: The base seed for the case $n = 9$

Lemma 2.4. For all i, j with $1 \leq i \leq j \leq n$ the equation

$$\Delta_{i,j}Z_j = P_j\Delta_{i,j-1} + Z_{j+1}P_{j-1}\Delta_{i,j-2}$$

holds.

Proof. Fix i . We prove Lemma 2.4 by induction on j . The statement is true for $j = i$ since $Y_iZ_i = P_i + Z_{i+1}Z_{i-1}$. If the statement is true for $j - 1$, then by Proposition 2.3

$$\begin{aligned} \Delta_{i,j}Z_j &= Y_jZ_j\Delta_{i,j-1} - Z_{j+1}Z_jP_{j-2}\Delta_{i,j-3} \\ &= P_j\Delta_{i,j-1} + Z_{j+1}Z_{j-1}\Delta_{i,j-1} - Z_{j+1}Z_jP_{j-2}\Delta_{i,j-3} \\ &= P_j\Delta_{i,j-1} + Z_{j+1}P_{j-1}\Delta_{i,j-2}, \end{aligned}$$

and the statement is true for j . \square

Lemma 2.5. The mutable cluster variables are $Z_1, Z_2, Z_3, \dots, Z_n$ and $\Delta_{i,j}$ for $1 \leq i \leq j \leq n$.

Proof. Starting with the initial seed (which is shown in Figure 11 for the case $n = 9$) perform mutations at the odd vertices $1, 3, 5, \dots, n$, consecutively. In each step, because of the equation $Y_iZ_i = P_i + Z_{i-1}Z_{i+1}$, the cluster variable Y_i for odd i with $1 \leq i \leq n$ is replaced by the cluster variable Z_i . Therefore, the mutations generate a seed whose mutable cluster variables are $Z_1, Z_2, Z_3, \dots, Z_n$. We refer to that seed as the *base seed*. The exchange matrix of the base seed is described by the associated quiver. By the rules of quiver mutation the mutable vertices of the base seed form an alternating quiver of type A_n isomorphic to Q . The only other arrows are the following. For every i with $1 \leq i \leq n$ there is an arrow between Z_i and P_i starting in P_i if i is odd and starting in Z_i if i is even. The quiver of the base seed for the example $n = 9$ is shown in Figure 12.

We now claim that starting from the base seed the cluster variable obtained by consecutive mutations at $i, i+1, i+2, \dots, j$ is $\Delta_{i,j}$ for all $1 \leq i \leq j \leq n$. The equation $\Delta_{i,j}Z_j = P_j\Delta_{i,j-1} + Z_{j+1}P_{j-1}\Delta_{i,j-2}$ from Lemma 2.4 is the exchange relation. For a proof consider the mutation of the quiver of the base seed. Fix i . We assume that i is odd. (If i is even, then reverse all arrows in the following argumentation.) We

We have an arrow $Z_k \rightarrow P_k$ in the base seed. The adjacency relations for P_k remain unaffected by mutations at $i, i+1, \dots, k-1$. After mutation at k the arrows reverse (and Z_k is replaced by $\Delta_{i,k}$) and we get an additional arrow $Z_{k+1} \rightarrow P_k$. Mutation at $k+1$ cancels the arrow $P_k \rightarrow \Delta_{i,k}$ whereas the arrow $Z_{k+1} \rightarrow P_k$ is replaced by an arrow $P_k \rightarrow \Delta_{i,k+1}$. Afterwards all adjacency relations for P_k with vertices Z_l for $k \leq l$ remain unaffected.

The adjacency relations for the vertices together with the induction hypothesis and the mutation rule for cluster variables imply that $(P_j \Delta_{i,j-1} + Z_{j+1} P_{j-1} \Delta_{i,j-2})/Z_j$ is the cluster variable obtained from consecutive mutation at $i, i+1, \dots, j$. By Lemma 2.4 it is equal to $\Delta_{i,j}$.

The number of mutable cluster variables of a cluster algebra of finite type is the sum of the rank of the cluster algebra and the number of positive roots of the associated root system. Since the $n + \frac{n(n+1)}{2}$ cluster variables $Z_1, Z_2, Z_3, \dots, Z_n$ and $\Delta_{i,j}$ for $1 \leq i \leq j \leq n$ are all distinct these must be all mutable cluster variables. \square

By Lemma 2.5 the recursion provided by Proposition 2.3 allows to compute iteratively every cluster variable in terms of the Y_i and Z_i ($1 \leq i \leq n$).

Example 2.6. Let us look at an example. We put $n = 3$. The initial cluster contains three mutable and three frozen variables. It is $(P_1, P_2, P_3, Y_1, Z_2, Y_3)$. One can check, by hand or by using Keller's mutation applet [34], that the following figure describes the exchange graph of the cluster algebra $\mathcal{A}(\mathcal{C}_M)$ in the case $n = 3$. This particular exchange graph is known as *associahedron* or *Stasheff polytope* K_5 , see Fomin-Reading [17, Section 3.1]. The mutable cluster variables are colored red, the frozen cluster variables blue. Beside the $3+3$ initial cluster variables there are 6 further cluster variables, namely Z_1, Y_2, Z_3 ,

$$\begin{aligned} \Delta_{1,2} &= \begin{vmatrix} Y_1 & c_1 \\ c_2 & Y_2 \end{vmatrix} = Y_1 Y_2 - Z_3, & \Delta_{2,3} &= \begin{vmatrix} Y_2 & c_2 \\ c_3 & Y_3 \end{vmatrix} = Y_2 Y_3 - Z_1, \\ \Delta_{1,3} &= \begin{vmatrix} Y_1 & c_1 & c_1 \\ c_2 & Y_2 & c_2 \\ 0 & c_3 & Y_3 \end{vmatrix} = Y_1 Y_2 Y_3 - Y_1 Z_1 - Y_3 Z_3 + Z_2. \end{aligned}$$

The cluster variables are grouped into $C_4 = 14$ clusters.

Remark 2.7. Formulae for cluster variables in $\mathcal{A}(\mathcal{C}_{M'})$ can be obtained from these formulae by setting $Y_n = Z_n = P_n = 1$.

Remark 2.8. The exchange relation (6) implies that mutation of the base seed at Z_i for some $i \in \{1, 2, \dots, n\}$ yields the cluster variable Y_i . Note that the base seed is *acyclic*, i.e., the corresponding quiver does not contain oriented cycles. Hence, by Berenstein-Fomin-Zelevinsky [3, Theorem 1.16] the cluster algebra $\mathcal{A}(\mathcal{C}_M)$ is equal to its *lower bound*, i.e., the set $\{Y_i, Z_i : 1 \leq i \leq n\}$ generates $\mathcal{A}(\mathcal{C}_M)$.

Remark 2.9. The cluster variables correspond to δ -functions of indecomposable rigid objects in \mathcal{C}_M . These objects have been classified by Rohleder [47, Theorem 7.3]. Besides the $3n$ objects of the form $T_{i,[a,b]}$ for $1 \leq i \leq n$ and $0 \leq a \leq b \leq 1$, these are (when viewed as elements in $C(1, \tau)$) the objects $M_{i,j}$

$$\bigoplus_{\substack{i < r < j \\ r \text{ odd}}} I_i \oplus \bigoplus_{\substack{i < r < j \\ r \text{ even}}} \tau(I_i) \xrightarrow{f} \bigoplus_{\substack{i < r < j \\ r \text{ odd}}} \tau(I_i) \oplus \bigoplus_{\substack{i < r < j \\ r \text{ even}}} \tau^2(I_i)$$




Figure 14: The exchange graph for $n = 3$: a Stasheff polyhedron

for $1 < j \leq n$ where $f|_{\tau(I_i)}^{\tau(I_{i\pm 1})} = 1$ for all even i and $f|_x^y = 0$ for all other direct summands X, Y .

3 The quantized universal enveloping algebra

3.1 Definition of the quantized enveloping algebra

Let v be an indeterminate. The quantized universal enveloping algebra $U_v(\mathfrak{g})$ is a deformation of the ordinary universal enveloping algebra $U(\mathfrak{g})$. To describe this construction we introduce quantized integers and quantized binomial coefficients.

Definition 3.1. For a natural number k , denote by $[k] = \frac{v^k - v^{-k}}{v - v^{-1}} \in \mathbb{Q}(v)$ the quantum integer and by $[k]! = [k][k-1] \cdots [1]$ the quantized factorial. For two natural numbers k and l , define the quantum binomial coefficient by

$$\begin{bmatrix} k \\ l \end{bmatrix} = \frac{[k]!}{[l]![k-l]!} \in \mathbb{Q}(v).$$

Remark 3.2. Both $[k]$ and $\begin{bmatrix} n \\ k \end{bmatrix}$ are actually Laurent polynomials in v . If we specialize $v = 1$, then $[k] = k$, $\begin{bmatrix} n \\ k \end{bmatrix} = \binom{n}{k}$, and $[k]! = k!$. Some authors such as Kac-Cheung [32] use a different convention for quantum integers.

Definition 3.3. The *quantized enveloping algebra* $U_v(\mathfrak{g})$ is the $\mathbb{Q}(v)$ -algebra generated by E_i, F_i, K_i, K_i^{-1} for $i = 1, 2, \dots, n$, subject to the following relations

$$K_i K_j = K_j K_i, \quad (i \neq j) \quad (7)$$

$$K_i K_i^{-1} = K_i^{-1} K_i = 1, \quad (i = 1, 2, \dots, n) \quad (8)$$

$$K_i E_j K_i^{-1} = v^{a_{ij}} E_j, \quad (1 \leq i, j \leq n) \quad (9)$$

$$K_i F_j K_i^{-1} = v^{-a_{ij}} F_j, \quad (1 \leq i, j \leq n) \quad (10)$$

$$E_i F_j - F_j E_i = \delta_{ij} \frac{K_i - K_i^{-1}}{v - v^{-1}}, \quad (1 \leq i, j \leq n) \quad (11)$$

$$E_i^2 E_j - [2] E_i E_j E_i + E_j E_i^2 = 0, \quad |i - j| = 1, \quad (12)$$

$$F_i^2 F_j - [2] F_i F_j F_i + F_j F_i^2 = 0, \quad |i - j| = 1, \quad (13)$$

$$E_i E_j = E_j E_i, \quad |i - j| \geq 2, \quad (14)$$

$$F_i F_j = F_j F_i, \quad |i - j| \geq 2, \quad (15)$$

where δ_{ij} is the Kronecker delta function. Note that $[2] = v + v^{-1}$, so we may write equation (12) as $E_i^2 E_j - (v + v^{-1}) E_i E_j E_i + E_j E_i^2 = 0$.

Definition 3.4. The subalgebra generated by E_i for $i = 1, 2, \dots, n$ is called the *quantized enveloping algebra* $U_v(\mathfrak{n})$.

The only relations in $U_v(\mathfrak{n})$ remain (12) and (14). These are called *quantized Serre relations*. The algebra $U_v(\mathfrak{n})$ specializes to $U(\mathfrak{n})$ in the limit $v = 1$.

Remark 3.5. The algebra $U_v(\mathfrak{g})$ is a *graded algebra*. It is graded by the root lattice R if we set $\deg(E_i) = \alpha_i$, $\deg(F_i) = -\alpha_i$, and $\deg(K_i) = 0$ for all $1 \leq i \leq n$. Note that $\deg(A) \in R^+$ for all $A \in U_v(\mathfrak{n})$. We also use the abbreviation $\deg(A) = |A|$ for $A \in U_v(\mathfrak{n})$.

Remark 3.6. Put $\sigma(v) = v^{-1}$ and $\sigma(E_i) = E_i$ for all i with $1 \leq i \leq n$. By the symmetry of the relations (12) and (14) in $U_v(\mathfrak{n})$ the map σ extends to an algebra anti-homomorphism $\sigma: U_v(\mathfrak{n}) \rightarrow U_v(\mathfrak{n})$, i.e., a \mathbb{Q} -linear map $\sigma: U_v(\mathfrak{n}) \rightarrow U_v(\mathfrak{n})$ such that $\sigma(AB) = \sigma(B)\sigma(A)$ for all $A, B \in U_v(\mathfrak{n})$. By construction σ is an antiautomorphism and an involution, i.e., $\sigma^2(A) = A$ for all $A \in U_v(\mathfrak{n})$.

Remark 3.7. In literature the deformation parameter v is sometimes called q . There are also different sign conventions for the exponent of the deformation parameter. We adopt Lusztig's convention [42]. It matches Leclerc's usage [39] if we set $q = v^{-1}$.

Remark 3.8. The quantized enveloping algebra $U_v(\mathfrak{g}') = U_v(\mathfrak{sl}_n)$ associated with Q' is defined similarly and may be regarded as the subalgebra of $U_v(\mathfrak{g}) = U_v(\mathfrak{sl}_{n+1})$ generated by the elements E_i, F_i, K_i, K_i^{-1} for $1 \leq i \leq n-1$.

3.2 The subalgebra $U_v^+(w)$ and the Poincaré-Birkhoff-Witt basis

We introduce Lusztig's T-automorphisms. For $1 \leq i \leq j$ put

$$T_i(E_j) = \begin{cases} -K_i^{-1}F_i, & \text{if } i = j; \\ E_jE_i - v^{-1}E_iE_j, & \text{if } |i - j| = 1; \\ E_j, & \text{if } |i - j| \geq 2; \end{cases}$$

$$T_i(F_j) = \begin{cases} -E_iK_i, & \text{if } i = j; \\ F_iF_j - vF_jF_i, & \text{if } |i - j| = 1; \\ F_j, & \text{if } |i - j| \geq 2; \end{cases}$$

$$T_i(K_j) = K_jK_i^{-a_{ij}}.$$

Lusztig [42, Chapter 37] shows that every T_i can be extended to an $\mathbb{Q}(v)$ -algebra homomorphism $T_i: U_v(\mathfrak{g}) \rightarrow U_v(\mathfrak{g})$. (In Lusztig's book [42] it is called $T'_{i,-1}$.) In fact, every T_i is an $\mathbb{Q}(v)$ -algebra automorphism. The images of the generators of $U_v(\mathfrak{g})$ under the inverse T_i^{-1} are given by

$$T_i^{-1}(E_j) = \begin{cases} -F_iK_i, & \text{if } i = j; \\ E_iE_j - v^{-1}E_jE_i, & \text{if } |i - j| = 1; \\ E_j, & \text{if } |i - j| \geq 2; \end{cases}$$

$$T_i^{-1}(F_j) = \begin{cases} -K_i^{-1}E_i, & \text{if } i = j; \\ F_jF_i - vF_iF_j, & \text{if } |i - j| = 1; \\ F_j, & \text{if } |i - j| \geq 2; \end{cases}$$

$$T_i^{-1}(K_j) = K_jK_i^{-a_{ij}}.$$

Remark 3.9. If $g \in U_v(\mathfrak{g})$ is homogeneous of degree β , then $T_i(g)$ is homogeneous of degree $s_i(\beta)$.

Remark 3.10. Furthermore, the T_i satisfy *braid relations*. For brevity we write T_iT_j for $T_i \circ T_j$ for all $i, j \in Q_0$. The braid relations are

$$T_iT_j = T_jT_i, \quad \text{if } |i - j| \geq 2;$$

$$T_iT_jT_i = T_jT_iT_j, \quad \text{if } |i - j| = 1.$$

Definition 3.11. To $w = s_1 s_3 s_5 \cdots s_n s_2 s_4 s_6 \cdots s_{n-1} s_1 s_3 s_5 \cdots s_n s_2 s_4 s_6 \cdots s_{n-1}$ we attach elements in $U_v(\mathfrak{g})$. If $j_1, j_2, \dots, j_{2n} \in [1, n]$ are indices such that for the reduced expression from above we have $w = s_{j_1} s_{j_2} \cdots s_{j_{2n}}$, then we consider the elements $T_{j_1} T_{j_2} \cdots T_{j_{k-1}}(E_{j_k})$ for $1 \leq k \leq 2n$. Since $\deg(T_{j_1} T_{j_2} \cdots T_{j_{k-1}}(E_{j_k})) = s_{j_1} s_{j_2} \cdots s_{j_{k-1}}(\alpha_{j_k}) = \beta_k$ for all k , we introduce the shorthand notation $E(\beta_k) = T_{j_1} T_{j_2} \cdots T_{j_{k-1}}(E_{j_k})$ for all $1 \leq k \leq 2n$. Here, $E(\beta_1)$ is assumed to be $E(\beta_1) = E_{j_1}$.

Definition 3.12. For $i \in Q_0$ and $a \in \mathbb{N}$ put $E_i^{(a)} = \frac{1}{[a]!} E_i^a \in U_v(\mathfrak{n})$. For a natural number k with $1 \leq k \leq 2n$ and $a \in \mathbb{N}$ put

$$E^{(a)}(\beta_k) = T_{j_1} T_{j_2} \cdots T_{j_{k-1}}(E_{j_k}^{(a)}) = \frac{1}{[a]!} E(\beta)^a.$$

The following theorem is due to Lusztig [42, Theorem 40.2.1]. It also contains the definition of the subalgebra $U_v^+(w)$ which is crucial for our further studies; moreover, it enables us to define the Poincaré-Birkhoff-Witt basis of $U_v^+(w)$. For an idea of a proof different from Lusztig's [42] see Bergman's diamond lemma [2].

Theorem 3.13. The set

$$\mathcal{P} = \left\{ E^{(a_1)}(\beta_1) E^{(a_2)}(\beta_2) \cdots E^{(a_{2n})}(\beta_{2n}) : (a_1, a_2, \dots, a_{2n}) \in \mathbb{N}^{2n} \right\}$$

is linearly independent over the field $\mathbb{Q}(v)$. It forms a $\mathbb{Q}(v)$ -basis of a $\mathbb{Q}(v)$ -subalgebra $U_v^+(w) \subset U_v(\mathfrak{n})$. Moreover, $U_v^+(w)$ is well-defined in the sense that it is independent of the choice of the reduced expression for w . If we choose another reduced expression $w = s_{j'_1} s_{j'_2} \cdots s_{j'_{2n}}$ for w , then the set of all

$$E_{j'_1}^{(a_1)} \cdot T_{j'_1}(E_{j'_2}^{(a_2)}) \cdots T_{j'_1} T_{j'_2} \cdots T_{j'_{2n-1}}(E_{j'_{2n}}^{(a_{2n})})$$

for all sequences $(a_1, a_2, \dots, a_{2n}) \in \mathbb{N}^{2n}$ is also a basis of the same subalgebra $U_v^+(w) \subset U_v(\mathfrak{n})$.

Remark 3.14. The basis \mathcal{P} is called the *Poincaré-Birkhoff-Witt basis* of $U_v^+(w)$ associated with the reduced expression. Unlike the canonical basis which we will define later the Poincaré-Birkhoff-Witt basis \mathcal{P} depends on the choice of the reduced expression for w . Every choice of a reduced expression for w induces a bijection between \mathbb{N}^{2n} and a basis for $U_v^+(w)$. In this sense \mathcal{P} is not canonical. The various bijections are called *Lusztig parametrizations*.

Remark 3.15. Theorem 3.13 particularly implies that $E(\beta_k) \in U_v(\mathfrak{n})$ for every $1 \leq k \leq 2n$ which is not obvious from the definition of the T -automorphisms.

For any vector $\mathbf{a} = (a_1, a_2, \dots, a_{2n}) \in \mathbb{N}^{2n}$ we introduce the shorthand notation $E[\mathbf{a}]$ for $E^{(a_1)}(\beta_1) E^{(a_2)}(\beta_2) \cdots E^{(a_{2n})}(\beta_{2n})$. We also use a different notation for $E(\beta_k)$ with $1 \leq k \leq 2n$; namely we put

$$\begin{aligned} u_i &= T_1 T_3 T_5 \cdots T_{i-2}(E_i), & \text{for odd } i \text{ with } 1 \leq i \leq n, \\ v_i &= T_1 T_3 T_5 \cdots T_n T_2 T_4 \cdots T_{i-2}(E_i), & \text{for even } i \text{ with } 2 \leq i \leq n-1, \\ w_i &= T T_1 T_3 T_5 \cdots T_{i-2}(E_i), & \text{for odd } i \text{ with } 1 \leq i \leq n, \\ x_i &= T T_1 T_3 T_5 \cdots T_n T_2 T_4 \cdots T_{i-2}(E_i), & \text{for even } i \text{ with } 2 \leq i \leq n-1, \end{aligned}$$

where $T = T_1 T_3 T_5 \cdots T_n T_2 T_4 T_6 \cdots T_{n-1}$. In what follows we use the convention $T_i = \text{id}_{U_v(\mathfrak{g})}$ for $i \notin Q_0$. Because of the braid relation $T_i T_j = T_j T_i$ for $|i-j| \geq 2$ and $T_i(E_j) = E_j$, $T_i(F_j) = F_j$, and $T_i(K_j) = K_j$ for $|i-j| \geq 2$ the formulae simplify to

$$\begin{aligned} u_i &= E_i, & \text{for odd } i \text{ s.t. } 1 \leq i \leq n, \\ v_i &= T_{i-1} T_{i+1}(E_i), & \text{for even } i \text{ s.t. } 2 \leq i \leq n-1, \\ w_i &= T_{i-2} T_i T_{i+2} T_{i-1} T_{i+1}(E_i), & \text{for odd } i \text{ s.t. } 1 \leq i \leq n, \\ x_i &= T_{i-3} T_{i-1} T_{i+1} T_{i+3} T_{i-2} T_i T_{i+2} T_{i-1} T_{i+1}(E_i), & \text{for even } i \text{ s.t. } 2 \leq i \leq n-1. \end{aligned}$$

Note that $w_i = T u_i$ for all odd i with $1 \leq i \leq n$ and $x_i = T v_i$ for all even i with $2 \leq i \leq n-1$.

Remark 3.16. The degrees of these variables are in bijection with the dimension vectors of the indecomposable direct summands of the terminal module T , compare Figure 6. To be more precise, let $1 \leq i \leq n$. If i is odd, then we have $E(\beta_i) = u_i$ and $E(\beta_{n+i}) = w_i$; if i is even, then we have $E(\beta_i) = v_i$ and $E(\beta_{n+i}) = x_i$.

Remark 3.17. Similarly, we can associate elements $E(\beta_k) \in U_v(\mathfrak{n}') \subset U_v(\mathfrak{n})$ for $1 \leq k \leq 2(n-1)$ to the reduced expression $s_1 s_3 s_5 \cdots s_{n-2} s_2 s_4 s_6 \cdots s_{n-1} s_1 s_3 s_5 \cdots s_{n-2} \cdots s_2 s_4 s_6 \cdots s_{n-1}$ of the Weyl group element $w' \in W'$. The elements generate an algebra $U_v^+(w') \subset U_v(\mathfrak{n}')$, and the set of all ordered products of the $E(\beta_k)$ is a Poincaré-Birkhoff-Witt basis of $U_v^+(w')$ just as above. Elements u'_i, w'_i (for odd i with $1 \leq i \leq n-2$) and v'_i, x'_i (for even i with $2 \leq i \leq n-1$) in $U_v(\mathfrak{n}')$ are defined analogously. Under the inclusion $U_v(\mathfrak{n}') \subset U_v(\mathfrak{n})$ they are literally the same as the corresponding elements except for

$$\begin{aligned} v'_{n-1} &= T_{n-2}(E_{n-1}), \\ w'_{n-2} &= T_{n-4} T_{n-2} T_{n-3} T_{n-1}(E_{n-2}), \\ x'_{n-3} &= T_{n-6} T_{n-4} T_{n-2} T_{n-5} T_{n-3} T_{n-1} T_{n-4} T_{n-2}(E_{n-3}), \\ x'_{n-1} &= T_{n-4} T_{n-2} T_{n-3} T_{n-1} T_{n-2}(E_{n-1}). \end{aligned}$$

3.3 The quantum shuffle algebra and Euler numbers

Rosso [48] noted that we may embed the quantized enveloping algebra $U_v(\mathfrak{n}) \hookrightarrow \mathcal{F}$ into the *quantum shuffle algebra* \mathcal{F} . Hence we may view $U_v(\mathfrak{n})$ as a subalgebra of \mathcal{F} . As we will see below \mathcal{F} is defined in purely combinatorial terms. As Leclerc [39, Section 2.5, 2.6] observes, the embedding $U_v(\mathfrak{n}) \hookrightarrow \mathcal{F}$ is very useful for explicit calculations.

Definition 3.18. Let r, s be natural numbers. A permutation $\pi \in S_{r+s}$ is called a *shuffle* of type (r, s) if $\pi(1) < \pi(2) < \cdots < \pi(r)$ and $\pi(r+1) < \pi(r+2) < \cdots < \pi(r+s)$.

The following definition is due to Leclerc [39, Section 2.5].

Definition 3.19. For every sequence $(i_1, i_2, \dots, i_r) \in Q_0^r$ of elements in Q_0 of length $r \geq 0$ define a symbol $w[i_1, i_2, \dots, i_r]$. (Especially, we have a symbol $w[]$ for the empty sequence.) Let \mathcal{F} be the $\mathbb{Q}(v)$ -vector space generated by all $w[i_1, i_2, \dots, i_r]$ for all $r \geq 0$. Define the *quantum shuffle product* on two basis elements by

$$w[i_1, i_2, \dots, i_r] * w[i_{r+1}, i_{r+2}, \dots, i_s] = \sum_{\substack{\pi \text{ shuffle} \\ \text{of type } (r,s)}} v^{e(\pi)} w[i_{\pi(1)}, i_{\pi(2)}, \dots, i_{\pi(r+s)}],$$

where the function $e: S_{r+s} \rightarrow \mathbb{Z}$ is defined as

$$e(\pi) = \sum_{k \leq r < l, \pi(k) < \pi(l)} (\alpha_{i_{\pi k}}, \alpha_{i_{\pi(l)}}).$$

It is easy to see that the product is associative. We extend the product bilinearly to a map $*$: $\mathcal{F} \times \mathcal{F} \rightarrow \mathcal{F}$. The algebra $(\mathcal{F}, *)$ is called the *quantum shuffle algebra*.

Remark 3.20. The shuffle product $*$ on the quantum shuffle algebra \mathcal{F} is not commutative. Hence, \mathcal{F} is a non-commutative $\mathbb{Q}(v)$ -algebra, but it degenerates to the classical commutative shuffle algebra when we specialize $v = 1$. Furthermore, the quantum shuffle algebra \mathcal{F} is graded by the root lattice if we set $\deg(w[i]) = \alpha_i$ for all $i \in \{1, 2, \dots, n\}$.

Lemma 3.21. The map $E_i \rightarrow w[i]$ extends to an embedding of graded algebras $U_v(\mathfrak{n}) \hookrightarrow \mathcal{F}$. In other words, $U_v(\mathfrak{n})$ is isomorphic to the subalgebra of \mathcal{F} generated by all $w[i]$ for $i \in Q_0$.

Proof. See Leclerc [39, Theorem 4]. □

From now on we view $U_v(\mathfrak{n})$ as a subalgebra of \mathcal{F} . In the rest of the section we expand the generators $u_i, v_i, w_i, x_i \in U_v^+(w)$ in the shuffle basis. The following elements will be important for the description.

Definition 3.22. For integers i, j such that $1 \leq i \leq j \leq n$ put

$$X_{i,j} = T_i^{(-1)^{i-1}} T_{i+1}^{(-1)^i} \dots T_{j-1}^{(-1)^{j-2}} (E_j).$$

By definition, $X_{i,j} \in U_v(\mathfrak{n}) \subset \mathcal{F}$.

Example 3.23. Let us give some examples of $X_{i,j}$ expanded in the shuffle basis: First of all, we have $X_{1,1} = E_1 = u_1 = w[1]$. Moreover,

$$\begin{aligned} X_{1,2} &= T_1(E_2) = E_2 E_1 - v^{-1} E_1 E_2 = w[2] * w[1] - v^{-1} w[1] * w[2] \\ &= w[1, 2] + v^{-1} w[2, 1] - v^{-1} (w[2, 1] + v^{-1} w[1, 2]) = (1 - v^{-2}) w[1, 2] \end{aligned}$$

is a second example.

The next lemma shows that we can compute the expansion of $X_{i,j}$ for all pairs (i, j) in the shuffle basis explicitly.

Lemma 3.24. Let i, j be integers such that $1 \leq i \leq j \leq n$. Then

$$X_{i,j} = (1 - v^{-2})^{j-i} \sum_{\pi} w[\pi(i), \pi(i+1), \dots, \pi(j)]$$

where the sum runs over all permutations π of $\{i, i+1, \dots, j\}$ such that for every even number k with $i \leq k \leq j-1$ we have $\pi^{-1}(k) > \pi^{-1}(k+1)$ and for every even number k with $i+1 \leq k \leq j$ we have $\pi^{-1}(k) > \pi^{-1}(k-1)$.

Proof. By backwards induction on i we see that the $X_{i,j}$ (for $1 \leq i < j \leq n$) satisfy the following recursion:

$$X_{i,j} = \begin{cases} E_j X_{i,j-1} - v^{-1} X_{i,j-1} E_j, & \text{if } j \text{ is even;} \\ X_{i,j-1} E_j - v^{-1} E_j X_{i,j-1}, & \text{if } j \text{ is odd.} \end{cases}$$

Now fix i . We proceed by induction on $j - i$. The statement is trivial for $j = i$. Suppose that $j > i$ and that $X_{i,j-1} = (1 - v^{-2})^{j-1-i} \sum_{\pi} w[\pi(i), \pi(i+1), \dots, \pi(j-1)]$, where the sum is taken over all permutations of $\{i, i+1, \dots, j-1\}$ such that for every even number k with $i \leq k \leq j-2$ we have $\pi^{-1}(k) > \pi^{-1}(k+1)$ and for every even number k with $i+1 \leq k \leq j-1$ we have $\pi^{-1}(k) > \pi^{-1}(k-1)$.

We consider two cases. First of all, assume that j is even. Let π be a permutation of $\{i, i+1, \dots, j-1\}$ as above. When shuffling the sequence (j) of length 1 with the sequence $(\pi(i), \pi(i+1), \dots, \pi(j-1))$ of length $j-i$, we get $j-i+1$ permutations of $\{i, i+1, \dots, j\}$. Among these we distinguish two kinds of permutations. The permutations π_1 where j comes after $j-1$ satisfy $\pi_1^{-1}(k) > \pi_1^{-1}(k+1)$ and $\pi_1^{-1}(k) > \pi_1^{-1}(k-1)$ for all even numbers k such that $i \leq k \leq j-1$ or $i+1 \leq k \leq j$, respectively. Conversely, every permutation π_1 of $\{i, i+1, \dots, j\}$ satisfying these conditions is uniquely obtained from shuffling (j) with a such a sequence $(\pi(i), \pi(i+1), \dots, \pi(j-1))$ such that j comes after $j-1$.

We also get permutations π_2 where j occurs before $j-1$. Now we see that

$$\begin{aligned} & w[j] * w[\pi(i), \pi(i+1), \dots, \pi(j-1)] \\ &= \sum_{\pi_1} w[\pi_1(i), \dots, \pi_1(j)] + v^{-1} \sum_{\pi_2} w[\pi_2(i), \dots, \pi_2(j)], \\ & w[\pi(i), \pi(i+1), \dots, \pi(j-1)] * w[j] \\ &= v^{-1} \sum_{\pi_1} w[\pi_1(i), \dots, \pi_1(j)] + \sum_{\pi_2} w[\pi_2(i), \dots, \pi_2(j)]. \end{aligned}$$

It follows by induction hypothesis that

$$X_{i,j} = w[j] * X_{i,j-1} - v^{-1} X_{i,j-1} * w[j] = (1 - v^{-2})^{j-i} \sum_{\pi_1} w[\pi_1].$$

The other case where j is odd is proved similarly. \square

Remark 3.25. The number $a(i, j)$ of permutations of $\{i, i+1, \dots, j-1\}$ such that for every even number k with $i \leq k \leq j-2$ we have $\pi^{-1}(k) > \pi^{-1}(k+1)$ and for every even number k with $i+1 \leq k \leq j-1$ we have $\pi^{-1}(k) > \pi^{-1}(k-1)$ only depends on $j-i$. The table displays some values of $a(i, j)$.

$j-i$	0	1	2	3	4	5	6
$a(i, j)$	1	1	2	5	16	61	272

The numbers are called *Euler numbers*. The sequence of Euler numbers is listed as A000111 in Sloane's Encyclopedia of Integer Sequences [50]. Its exponential generating function is $\sec(x) + \tan(x)$, see Stanley [51, Proposition 1.61].

Lemma 3.26. The following formulae for the generators of $U_v^+(w)$ are valid:

$$\begin{aligned}
u_i &= E_i, & \text{for odd } i \text{ with } 1 \leq i \leq n; \\
v_i &= T_{i-1}T_i^{-1}(E_{i+1}), & \text{for even } i \text{ with } 2 \leq i \leq n-2; \\
w_1 &= T_2^{-1}(E_3); \\
w_i &= T_{i-2}T_{i-1}^{-1}T_iT_{i+1}^{-1}(E_{i+2}), & \text{for odd } i \text{ with } 3 \leq i \leq n-3; \\
w_n &= T_{n-2}(E_{n-1}); \\
x_2 &= T_2^{-1}T_3T_4^{-1}(E_5); \\
x_i &= T_{i-3}T_{i-2}^{-1}T_{i-1}T_i^{-1}T_{i+1}T_{i+2}^{-1}(E_{i+3}), & \text{for even } i \text{ with } 4 \leq i \leq n-4; \\
x_{n-1} &= T_{n-4}T_{n-3}^{-1}T_{n-2}(E_{n-1});
\end{aligned}$$

Proof. The equation $u_i = E_i$ for odd i follows from definition. Note that by definition of Lusztig's T -automorphisms we have $T_{i+1}(E_i) = T_i^{-1}(E_{i+1})$ for $1 \leq i \leq n-1$ and that $T_{i-1}(E_i) = T_i^{-1}(E_{i-1})$ for $2 \leq i \leq n$. The first equation is equivalent to $T_iT_{i+1}(E_i) = E_{i+1}$, the second one is equivalent to $T_iT_{i-1}(E_i) = E_{i-1}$. Therefore, for even i we have $v_i = T_{i-1}T_{i+1}(E_i) = T_{i-1}T_i^{-1}(E_{i+1})$.

For all further calculations the formula

$$T_iT_{i-1}T_{i+1}(E_i) = T_{i-1}^{-1}T_i(E_{i+1}) = T_{i+1}^{-1}T_i(E_{i-1})$$

which holds for $2 \leq i \leq n-1$ will be crucial. To verify the formula note that by the braid relation we have $T_{i-1}T_iT_{i-1}T_{i+1}(E_i) = T_iT_{i-1}T_iT_{i+1}(E_i) = T_iT_{i-1}(E_{i+1}) = T_i(E_{i+1})$. Application of T_{i-1}^{-1} gives the first part of the equation, the second part is proved analogously.

Now we compute $w_1 = T_1T_3T_2(E_1) = T_3T_1T_2(E_1) = T_3(E_2) = T_2^{-1}(E_3)$. Similarly, $w_n = T_{n-2}T_nT_{n-1}(E_n) = T_{n-2}(E_{n-1})$. Furthermore, for odd i with $3 \leq i \leq n-2$ we compute

$$\begin{aligned}
w_i &= T_{i-2}T_iT_{i+2}T_{i-1}T_{i+1}(E_i) = T_{i-2}T_{i+2}T_iT_{i-1}T_{i+1}(E_i) \\
&= T_{i-2}T_{i+2}T_{i-1}^{-1}T_i(E_{i+1}) = T_{i-2}T_{i-1}^{-1}T_iT_{i+2}(E_{i+1}) = T_{i-2}T_{i-1}^{-1}T_iT_{i+1}^{-1}(E_{i+2}).
\end{aligned}$$

Moreover, we see that $x_2 = T_1T_3T_5T_2T_4T_1T_3(E_2) = T_1T_3T_5T_4T_1^{-1}T_2(E_3) = T_3T_5T_4T_2(E_3) = T_5T_2^{-1}T_3(E_4) = T_2^{-1}T_3T_5(E_4) = T_2^{-1}T_3T_4^{-1}(E_5)$, and

$$\begin{aligned}
x_{n-1} &= T_{n-4}T_{n-2}T_nT_{n-3}T_{n-1}T_{n-2}T_n(E_{n-1}) \\
&= T_{n-4}T_{n-2}T_nT_{n-3}T_{n-1}^{-1}T_{n-1}(E_{n-2}) \\
&= T_{n-4}T_{n-2}T_{n-3}T_{n-1}(E_{n-2}) = T_{n-4}T_{n-3}^{-1}T_{n-2}(E_{n-1}).
\end{aligned}$$

Finally, for even i with $4 \leq i \leq n-3$, the equation

$$\begin{aligned}
x_i &= T_{i-3}T_{i-1}T_{i+1}T_{i+3}T_{i-2}T_iT_{i+2}T_{i-1}T_{i+1}(E_i) \\
&= T_{i-3}T_{i-2}^{-1}T_{i-2}T_{i-1}T_{i-2}T_{i+1}T_{i+3}T_{i+2}T_iT_{i-1}T_{i+1}(E_i) \\
&= T_{i-3}T_{i-2}^{-1}T_{i-1}T_{i-2}T_{i-1}T_{i+1}T_{i+3}T_{i+2}T_{i-1}^{-1}T_i(E_{i+1}) \\
&= T_{i-3}T_{i-2}^{-1}T_{i-1}T_{i-2}T_{i+1}T_{i+3}T_{i+2}T_i(E_{i+1}) \\
&= T_{i-3}T_{i-2}^{-1}T_{i-1}T_{i-2}T_{i+3}T_i^{-1}T_{i+1}(E_{i+2}) \\
&= T_{i-3}T_{i-2}^{-1}T_{i-1}T_i^{-1}T_{i-2}T_{i+3}T_{i+1}(E_{i+2}) \\
&= T_{i-3}T_{i-2}^{-1}T_{i-1}T_i^{-1}T_{i+1}T_{i+2}^{-1}(E_{i+3}).
\end{aligned}$$

holds which is the last equation to be checked. \square

Remark 3.27. Lemma 3.26 shows all generators u_i, v_i, w_i, x_i (for appropriate i) of $U_v^+(w)$ are of the form $X_{i',j'}$ (for appropriate i', j'). Hence, the formula of Lemma 3.24 applies. In each case, $V_{[i',j']}$ is the associated kQ -module from Figure 1. In other words, in each case $\deg(X_{i',j'}) = \alpha_{i'} + \alpha_{i'+1} + \dots + \alpha_{j'}$.

Remark 3.28. With the same argument we can conclude that for Q' the equations

$$\begin{aligned} v'_{n-1} &= T_{n-2}(E_{n-1}) \\ w'_{n-2} &= T_{n-4}T_{n-3}^{-1}T_{n-2}(E_{n-1}) \\ x'_{n-1} &= T_{n-4}(E_{n-3}) \\ x'_{n-3} &= T_{n-6}T_{n-5}^{-1}T_{n-4}T_{n-3}^{-1}T_{n-2}(E_{n-1}) \end{aligned}$$

hold. Hence, Lemma 3.26 and Remark 3.27 also hold true for the generators u'_i, v'_i, w'_i, x'_i (for appropriate i) of $U_v^+(w')$.

3.4 The straightening relations for the generators of $U_v^+(w)$

The following lemma expands every $E(\beta_j)E(\beta_i)$ with $1 \leq i < j \leq 2n$ in the Poincaré-Birkhoff-Witt basis \mathcal{P} . The relations of Lemma 3.29 are called *straightening relations*. Iterative use of the straightening relations from Lemma 3.29 allows us to write an arbitrarily ordered monomial in the generators $E(\beta_k)$ with $1 \leq k \leq 2n$ (and hence every element in $U_v^+(w)$) as a linear combination of Poincaré-Birkhoff-Witt basis elements $E[\mathbf{a}]$ with $\mathbf{a} \in \mathbb{N}^{2n}$.

Lemma 3.29. The generators of $U_v^+(w)$ satisfy the following relations

$$\begin{aligned} v_{i+1}u_i &= vu_iv_{i+1}, & \text{for } i = 1, 3, \dots, n-2, \\ v_{i-1}u_i &= vu_iv_{i-1}, & \text{for } i = 3, 5, \dots, n, \\ \\ w_{i+2}u_i &= vu_iw_{i+2}, & \text{for } i = 1, 3, \dots, n-2, \\ w_{i-2}u_i &= vu_iw_{i-2}, & \text{for } i = 3, 5, \dots, n, \\ w_1u_1 &= v^{-1}u_1w_1 + v_2, \\ w_iu_i &= u_iw_i + (v - v^{-1})v_{i-1}v_{i+1}, & \text{for } i = 3, 5, \dots, n-2, \\ w_nu_n &= v^{-1}u_nw_n + v_{n-1}, \\ \\ x_{i+3}u_i &= vu_ix_{i+3}, & \text{for } i = 1, 3, \dots, n-4, \\ x_{i-3}u_i &= vu_ix_{i-3}, & \text{for } i = 5, 7, \dots, n, \\ x_{i-1}u_i &= u_ix_{i-1} + (v - v^{-1})v_{i+1}w_{i-2}, & \text{for } i = 3, 5, \dots, n-2, \\ x_{n-1}u_n &= v^{-1}u_nx_{n-1} + w_{n-2}, \\ x_2u_1 &= v^{-1}u_1x_2 + w_3, \\ x_{i+1}u_i &= u_ix_{i+1} + (v - v^{-1})v_{i-1}w_{i+2}, & \text{for } i = 3, 5, \dots, n-2, \\ \\ w_{i+1}v_i &= vv_iw_{i+1}, & \text{for } i = 2, 4, \dots, n-1, \\ w_{i-1}v_i &= vv_iw_{i-1}, & \text{for } i = 2, 4, \dots, n-1, \end{aligned}$$

$$\begin{aligned}
x_{i+2}v_i &= vv_ix_{i+2}, & \text{for } i = 2, 4, \dots, n-3, \\
x_{i-2}v_i &= vv_ix_{i-2}, & \text{for } i = 4, 6, \dots, n-1, \\
x_iv_i &= v_ix_i + (v - v^{-1})w_{i-1}w_{i+1}, & \text{for } i = 2, 4, \dots, n-1, \\
x_{i+1}w_i &= vw_ix_{i+1}, & \text{for } i = 1, 3, \dots, n-2, \\
x_{i-1}w_i &= vw_ix_{i-1}, & \text{for } i = 3, 5, \dots, n.
\end{aligned}$$

For every i, j with $1 \leq i < j \leq 2n$ such that $E(\beta_j)E(\beta_i)$ is not listed on the left-hand side above the commutativity relation $E(\beta_j)E(\beta_i) = E(\beta_i)E(\beta_j)$ holds.

Proof. Let i be an integer with $1 \leq i \leq n-1$. We have

$$\begin{aligned}
T_i T_{i+2}(E_{i+1}) &= T_i(E_{i+1}E_{i+2} - v^{-1}E_{i+2}E_{i+1}) \\
&= (E_{i+1}E_i - v^{-1}E_iE_{i+1})E_{i+2} - v^{-1}E_{i+2}(E_{i+1}E_i - v^{-1}E_iE_{i+1}) \\
&= E_{i+1}E_iE_{i+2} - v^{-1}E_iE_{i+1}E_{i+2} \\
&\quad - v^{-1}E_{i+2}E_{i+1}E_i + v^{-2}E_iE_{i+2}E_{i+1}. \tag{16}
\end{aligned}$$

Now let i be an odd integer with $1 \leq i \leq n-2$. Then $u_i = E_i$ and $v_{i+1} = T_i T_{i+2}(E_{i+1})$. By equation (16) the following relations hold:

$$\begin{aligned}
v_{i+1}u_i &= (E_{i+1}E_i^2 - v^{-1}E_iE_{i+1}E_i)E_{i+2} + E_{i+2}(v^{-2}E_iE_{i+1}E_i - v^{-1}E_{i+1}E_i^2), \\
u_iv_{i+1} &= (E_iE_{i+1}E_i - v^{-1}E_i^2E_{i+1})E_{i+2} + E_{i+2}(v^{-2}E_i^2E_{i+1} - v^{-1}E_iE_{i+1}E_i).
\end{aligned}$$

By equation (12) we get $v_{i+1}u_i - vu_iv_{i+1} = 0$. The equation $v_{i-1}u_i = vu_iv_{i-1}$, for odd i with $3 \leq i \leq n$, is proved analogously. Application of T to the last two equations yields $x_{i+1}w_i = vw_ix_{i+1}$, for odd integers i with $1 \leq i \leq n-2$, and $x_{i-1}w_i = vw_ix_{i-1}$, for odd integers i with $3 \leq i \leq n$.

Now let i be an even integer with $2 \leq i \leq n-1$. Then $v_i = T_{i-1}T_{i+1}(E_i)$ and $w_{i+1} = T_{i-3}T_{i-1}T_{i+1}T_{i+3}T_iT_{i+2}(E_{i+1})$. With the same argument as above we have $T_iT_{i+2}(E_{i+1})E_i = vE_iT_iT_{i+2}(E_{i+1})$. Apply the automorphism $T_{i-3}T_{i-1}T_{i+1}T_{i+3}$ to the last equation. Using the equation $T_{i-3}T_{i-1}T_{i+1}T_{i+3}(E_i) = T_{i-1}T_{i+1}(E_i) = v_i$ we get $w_{i+1}v_i = vv_iw_{i+1}$. Similarly, the equation $w_{i-1}v_i = vv_iw_{i-1}$ holds.

Let i be an integer with $1 \leq i \leq n-1$. We have

$$\begin{aligned}
-T_{i+1}(E_{i+2})F_iK_i &= -v^{-1}(E_{i+2}E_{i+1} - v^{-1}E_{i+1}E_{i+2})F_iK_i \\
&= -F_iK_i(E_{i+2}E_{i+1} - v^{-1}E_{i+1}E_{i+2}) = -F_iK_iT_{i+1}(E_{i+2}).
\end{aligned}$$

Application of the composition of automorphisms $T_iT_{i+2}T_{i+4}T_{i+3}$ yields

$$T_iT_{i+2}T_{i+4}T_{i+1}T_{i+3}(E_{i+2})E_i = vE_iT_iT_{i+2}T_{i+4}T_{i+1}T_{i+3}(E_{i+2}). \tag{17}$$

Now let i be more specifically an odd integer with $1 \leq i \leq n-2$. The previous equation (17) asserts that $w_{i+2}u_i = vu_iw_{i+2}$. The equation $w_{i-2}u_i = vu_iw_{i-2}$, for odd i with $3 \leq i \leq n$, is proved analogously. Now let i be an even integer with $2 \leq i \leq n-3$. From (17) we see after application of $T_{i-1}T_{i+1}T_{i+3}T_{i+5}$ that $x_{i+2}v_i = vv_ix_{i+2}$. The equation $x_{i-2}v_i = vv_ix_{i-2}$, for even i with $4 \leq i \leq n-1$, is proved analogously.

Furthermore, there holds:

$$\begin{aligned}
& -F_1K_1(E_1E_2 - v^{-1}E_2E_1) \\
& = -vF_1E_1E_2K_1 + F_1E_2E_1K_1 \\
& = v\left(\frac{K_1 - K_1^{-1}}{v - v^{-1}} - E_1F_1\right)E_2K_1 + E_2\left(E_1F_1 - \frac{K_1 - K_1^{-1}}{v - v^{-1}}\right)K_1 \\
& = -v(E_1E_2 - v^{-1}E_2E_1)F_1K_1 + \frac{1}{v - v^{-1}}(E_2K_1^2 - v^2E_2 - E_2K_1^2 + E_2) \\
& = -v(E_1E_2 - v^{-1}E_2E_1)F_1K_1 - vE_2.
\end{aligned}$$

Application of the map T_1T_3 yields to $u_1w_1 = vw_1u_1 - vv_2$ which is equivalent to $w_1u_1 = v^{-1}u_1w_1 + v_2$. The next equation $w_nu_n = v^{-1}u_nw_n + v_{n-1}$ is proved analogously.

Let i be an integer with $2 \leq i \leq n - 1$. Introduce the abbreviation

$$\begin{aligned}
S & = T_{i-1}T_{i+1}(E_i) \\
& = E_iE_{i-1}E_{i+1} - v^{-1}E_{i-1}E_iE_{i+1} - v^{-1}E_{i+1}E_iE_{i-1} + v^{-2}E_{i-1}E_{i+1}E_i.
\end{aligned}$$

We have $K_iS = SK_i$, so $-F_iK_iS$ is equal to

$$\begin{aligned}
& (-F_iE_iE_{i-1}E_{i+1} + v^{-1}E_{i-1}F_iE_iE_{i+1} \\
& \quad + v^{-1}E_{i+1}F_iE_iE_{i-1} - v^{-2}E_{i-1}E_{i+1}F_iE_i)K_i \\
& = -SF_iK_i + \frac{1}{v - v^{-1}}\left[(K_i - K_i^{-1})E_{i-1}E_{i+1} - v^{-1}E_{i-1}(K_i - K_i^{-1})E_{i+1} \right. \\
& \quad \left. - v^{-1}E_{i+1}(K_i - K_i^{-1})E_{i-1} + v^{-2}E_{i-1}E_{i+1}(K_i - K_i^{-1})\right] \\
& = -SF_iK_i + \frac{1}{v - v^{-1}}\left[(v^{-2} - v^{-2} - v^{-2} + v^{-2})E_{i-1}E_{i+1}K_i^2 \right. \\
& \quad \left. + (-v^2 + 2 - v^{-2})E_{i-1}E_{i+1}\right] \\
& = -SF_iK_i + (v^{-1} - v)E_{i+1}E_{i-1}. \tag{18}
\end{aligned}$$

Now let i be more specifically an odd integer with $3 \leq i \leq n - 2$. After application of $T_{i-2}T_iT_{i+2}$ the previous equation (18) asserts that $u_iw_i = w_iu_i + (v^{-1} - v)v_{i+1}v_{i-1}$. If i is an even integer with $2 \leq i \leq n - 1$, then application of the composition $T_{i-3}T_{i-1}T_{i+1}T_{i+3}T_{i-2}T_iT_{i+2}$ to equation (18) yields $v_ix_i = x_iv_i + (v^{-1} - v)w_{i+1}w_{i-1}$.

Let i be an odd integer with $1 \leq i \leq n - 4$. Since $T_{i+1}T_{i+2}T_{i+4}(E_{i+3})$ is a linear combination of monomials in E_{i+1} , E_{i+2} , E_{i+3} , and E_{i+4} with each factor appearing once, we see that

$$-T_{i+1}T_{i+2}T_{i+4}(E_{i+3})F_iK_i = -vF_iK_iT_{i+1}T_{i+2}T_{i+4}(E_{i+3}). \tag{19}$$

Applying $T_iT_{i+2}T_{i+4}T_{i+6}T_{i+3}T_{i+5}$ to (19) yields $x_{i+3}u_i = vu_ix_{i+3}$. The equation $x_{i-3}u_i = vu_ix_{i-3}$, for odd i with $5 \leq i \leq n$, is proved analogously.

Consider the three elements $T_2^{-1}(-F_1K_1)$, $T_1T_3(E_2)$, and E_3 . We introduce the

abbreviation $X = T_2^{-1}(-F_1K_1) = (vF_2F_1 - F_1F_2)K_1K_2$. We have

$$\begin{aligned} (vF_2F_1 - F_1F_2)E_1 &= vF_2 \left(E_1F_1 - \frac{K_1 - K_1^{-1}}{v - v^{-1}} \right) - \left(E_1F_1 - \frac{K_1 - K_1^{-1}}{v - v^{-1}} \right) F_2 \\ &= E_1(vF_2F_1 - F_1F_2) \\ &\quad + \frac{1}{v - v^{-1}} [-vF_2(K_1 - K_1^{-1}) + (K_1 - K_1^{-1})F_2] \\ &= E_1(vF_2F_1 - F_1F_2) + F_2K_1^{-1}. \end{aligned}$$

Therefore, $XE_1 = vE_1X + vF_2K_2$. Similarly, $XE_2 = vE_2X + vF_1K_1K_2^2$. Furthermore, $XE_3 = v^{-1}E_3X$. Hence

$$\begin{aligned} X(E_2E_1 - v^{-1}E_1E_2) &= (vE_2X + vF_1K_1K_2^2)E_1 - (E_1X + F_2K_2)E_2 \\ &= v(vE_1X + vF_2K_2) + v^2F_1K_1E_1 - E_1(vE_2X + vF_1K_1K_2^2) - F_2K_2E_2 \\ &= v^2(E_2E_1 - v^{-1}E_1E_2)X + v^2(E_2F_2 - F_2E_2)K_2 + v(F_1E_1 - E_1F_1)K_1K_2^2. \end{aligned}$$

Introduce the abbreviations $Y = E_2E_1 - v^{-1}E_1E_2$ and $R = v^2(E_2F_2 - F_2E_2)K_2 + v(F_1E_1 - E_1F_1)K_1K_2^2 = \frac{1}{v-v^{-1}} [v^2K_2^2 - v^2 - vK_1^2K_2^2 + vK_2^2]$. It follows that $RE_3 - v^{-2}E_3R = -vE_3$. Note that $T_1T_3(E_2) = YE_3 - v^{-1}E_3Y$ is equal to a v -commutator. Hence

$$\begin{aligned} XT_1T_3(E_2) &= X(YE_3 - v^{-1}E_3Y) \\ &= (v^2YX + R)E_3 - v^{-2}E_3(v^2YX + R) \\ &= v(YE_3 - v^{-1}E_3Y) - RE_3 - v^{-2}E_3R \\ &= vT_1T_3(E_2) - vE_3. \end{aligned} \tag{20}$$

Application of $T_1T_3T_5T_2T_4$ to equation (20) yields to $u_1x_2 = vx_2u_1 - vw_3$ which is equivalent to $x_2u_1 = v^{-1}u_1x_2 + w_3$. The equation $x_{n-1}u_n = v^{-1}u_nx_{n-1} + w_{n-2}$ is proved analogously.

Now let i be an odd integer such that $3 \leq i \leq n-2$. Let us consider the four elements $T_{i+1}^{-1}(-F_iK_i)$, $T_{i-1}T_iT_{i+2}(E_{i+1})$, E_{i+2} , and E_{i-1} . Now we denote by X the element $T_{i+1}^{-1}(-F_iK_i)$. Similarly as above, the equations $XE_i = vE_iX + vF_{i+1}K_{i+1}$, $XE_{i+1} = vE_{i+1}X + vF_iK_iK_{i+1}^2$, and $XE_{i+2} = v^{-1}E_{i+2}X$ hold. Furthermore, we see that $XE_{i-1} = (vF_{i+1}F_i - F_iF_{i+1})K_iK_{i+1}E_{i-1} = v^{-1}E_{i-1}X$. Note that

$$\begin{aligned} T_{i-1}T_iT_{i+2}(E_{i+1}) &= T_{i-1}(E_{i+1}E_iE_{i+2} - v^{-1}E_iE_{i+1}E_{i+2} - v^{-1}E_{i+2}E_{i+1}E_i + v^{-2}E_iE_{i+2}E_{i+1}) \\ &= T_iT_{i+2}(E_{i+1})E_{i-1} - v^{-1}E_{i-1}T_iT_{i+2}(E_{i+1}). \end{aligned}$$

With the same argument as above one can prove that

$$XT_iT_{i+2}(E_{i+1}) = vT_iT_{i+2}(E_{i+1})X - vE_{i+2}.$$

From this equation it follows that

$$\begin{aligned} XT_{i-1}T_iT_{i+2}(E_{i+1}) &= (vT_iT_{i+2}(E_{i+1})X - vE_{i+2})E_{i-1} - v^{-2}E_{i-1}(vT_iT_{i+2}(E_{i+1})X - vE_{i+2}) \\ &= (T_iT_{i+2}(E_{i+1})E_{i-1} - v^{-1}E_{i-1}T_iT_{i+2}(E_{i+1}))X + (v^{-1} - v)E_{i+2}E_{i-1}. \end{aligned} \tag{21}$$

Application of the automorphism $T_{i-2}T_iT_{i+2}T_{i+4}T_{i+1}T_{i+3}$ to equation (21) gives $u_i x_{i+1} = x_{i+1} u_i + (v^{-1} - v)w_{i+2}v_{i-1}$. The equation $u_i x_{i-1} = x_{i-1} u_i + (v^{-1} - v)w_{i-2}v_{i+1}$ is proved analogously.

After multiplying with appropriate T -automorphisms, all others pairs $E(\beta_i)$ and $E(\beta_j)$ of generators become $\mathbb{Q}(v)$ -linear combinations of monomials $E_{i_1}E_{i_2}\cdots E_{i_k}$ and $E_{i'_1}E_{i'_2}\cdots E_{i'_k}$, respectively, where the two occurring sequences (i_1, i_2, \dots, i_k) and $(i'_1, i'_2, \dots, i'_k)$ of indices come from two intervals of distance at least two. Hence they commute. \square

Remark 3.30. In the straightening relations of Lemma 3.29, for all i, j with $1 \leq i < j \leq 2n$, the coefficient in front of $E(\beta_i)E(\beta_j)$ in the expansion of $E(\beta_j)E(\beta_i)$ in the Poincaré-Birkhoff-Witt basis is $v^{(\beta_i, \beta_j)}$.

Remark 3.31. Similarly, there are straightening relations for the generators u'_i, v'_i, w'_i, x'_i (for appropriate i) of $U_v^+(w')$ that enable us to expand every element of $U_v^+(w')$ in the Poincaré-Birkhoff-Witt basis. Let us describe them. First of all, note that $v'_{n-1} = T_n^{-1}(v_{n-1})$, $w'_{n-1} = T_n^{-1}w_{n-1}$, $x'_{n-3} = T_n^{-1}x_{n-3}$, and that T_n^{-1} leaves all generators invariant except for $v'_{n-1}, w'_{n-1}, x'_{n-3}$, and x'_{n-1} . Therefore, the straightening relations of $U_v^+(w')$ are the same as the ones for $U_v^+(w)$ except for the straightening relations involving x'_{n-1} .

Calculations similar to those in Lemma 3.29 show that the straightening relations involving x'_{n-1} are commutativity relations except for:

$$\begin{aligned} x'_{n-1}w'_{n-2} &= vw'_{n-2}x'_{n-1}, \\ x'_{n-1}v'_{n-3} &= vv'_{n-3}x'_{n-1}, \\ x'_{n-1}v'_{n-1} &= v^{-1}v'_{n-1}x'_{n-1} + w'_{n-2}, \\ x'_{n-1}u'_{n-2} &= v^{-1}u'_{n-2}x'_{n-1} + v'_{n-3}, \\ x'_{n-1}u'_{n-4} &= vu'_{n-4}x'_{n-1}. \end{aligned}$$

Note that Remark 3.30 is also true in this case.

Remark 3.32. The commutation exponent of Remark 3.30 and a weaker (non-explicit) form the straightening relations of Lemma 3.29 is given by the Lemma of Levendorkii-Soibelman [40, Proposition 5.5.2]. See also Kimura [36, Theorem 4.24].

3.5 The dual Poincaré-Birkhoff-Witt basis

Kashiwara [33] introduced operators $E'_i \in \text{End}(U_v(\mathfrak{n}))$ for $1 \leq i \leq n$ such that the following two properties hold: First of all, $E'_i(E_j) = \delta_{i,j}$ for all i, j . Secondly, the Leibniz rule $E'_i(xy) = E'_i(x)y + v^{(\alpha_i, |x|)}xE'_i(y)$ holds for all i and all homogeneous elements $x, y \in U_v(\mathfrak{n})$. Furthermore, Kashiwara [33] introduced a non-degenerate symmetric bilinear form

$$(\cdot, \cdot) : U_v(\mathfrak{n}) \times U_v(\mathfrak{n}) \rightarrow \mathbb{Q}(v).$$

It is characterized by the assumption that the endomorphism E'_i of $U_v(\mathfrak{n})$ is adjoint to the left multiplication with E_i , i.e., $(E'_i(x), y) = (x, E_i y)$ for all $x, y \in U_v(\mathfrak{n})$ and $i \in \{1, 2, \dots, n\}$.

The algebra $U_v(\mathfrak{n})$ is a Hopf algebra. The E'_i may be viewed as elements in the graded dual Hopf algebra $U_v(\mathfrak{n})_{gr}^*$. We refer to Berenstein-Zelevinsky [9, Appendix] for details.

Lusztig [42, Section 1.2] defined a different non-degenerate symmetric bilinear form $(\cdot, \cdot)_L : U_v(\mathfrak{n}) \times U_v(\mathfrak{n}) \rightarrow \mathbb{Q}(v)$. In this paper we use Kashiwara's form. Both forms can be compared, see Leclerc [39, Section 2.2].

According to Lusztig [42, Proposition 38.2.3] the Poincaré-Birkhoff-Witt basis is orthogonal with respect to Lusztig's bilinear form. The comparison between both forms shows that it is also orthogonal with respect to Kashiwara's bilinear form. More precisely, for all $\beta, \gamma \in \Delta^+$, we have (see Leclerc [39, Equation 21] where the author uses the variable q^{-1} instead of v) $(E(\beta), E(\gamma)) = 0$ for $\beta \neq \gamma$, and

$$(E(\beta), E(\beta)) = \frac{\prod_{i=1}^n (1 - v^{-(\alpha_i, \alpha_i)})^{d_i}}{1 - v^{-(\beta, \beta)}}$$

for $\beta = \sum_{i=1}^n d_i \alpha_i$ with $d_i \in \mathbb{N}$. Compare also with Kimura [36, Proposition 4.18].

Remark 3.33. Every $\beta \in \Delta^+$ fulfills $(\beta, \beta) = 2$; every α_i with $i \in Q_0$ fulfills $(\alpha_i, \alpha_i) = 2$. Hence, if $\beta = \alpha_i + \alpha_{i+1} + \dots + \alpha_j \in \Delta^+$, then $(E(\beta), E(\beta)) = (1 - v^{-2})^{j-i}$.

Definition 3.34. The dual Poincaré-Birkhoff-Witt basis \mathcal{P}^* of $U_v^+(w)$ is defined to be the basis adjoint to the Poincaré-Birkhoff-Witt basis with respect to Kashiwara's form. For every natural number k with $1 \leq k \leq 2n$ we denote by $E^*(\beta_k) \in \mathcal{P}^*$ the dual of $E(\beta_k) \in \mathcal{P}$, i.e., the unique scalar multiple of $E(\beta_k)$ such that $(E(\beta_k), E^*(\beta_k)) = 1$.

Remark 3.35. Assume that k is an integer with $1 \leq k \leq 2n$. Let i, j be the integers with $1 \leq i \leq j \leq n$ such that we can write $\beta_k \in \Delta^+$ as $\beta_k = \alpha_i + \alpha_{i+1} + \dots + \alpha_j$. By Lemma 3.24 we have

$$E^*(\beta_k) = (1 - v^{-2})^{i-j} X_{i,j} = \sum_{\pi} w[\pi(i), \pi(i+1), \dots, \pi(j)]$$

where the sum runs over all permutations π of $\{i, i+1, \dots, j\}$ such that for every even number k with $i \leq k \leq j-1$ we have $\pi^{-1}(k) > \pi^{-1}(k+1)$ and for every even number k with $i+1 \leq k \leq j$ we have $\pi^{-1}(k) > \pi^{-1}(k-1)$.

Definition 3.36. We also introduce a shorthand notation for $E^*(\beta_k)$ with $1 \leq k \leq 2n$. To reflect similarities with the cluster algebra from Section 2.4 we put

$$\begin{aligned} y_i &= u_i^*, & \text{for odd } i \text{ with } 1 \leq i \leq n, \\ z_i &= v_i^*, & \text{for even } i \text{ with } 2 \leq i \leq n-1, \\ z_i &= w_i^*, & \text{for odd } i \text{ with } 1 \leq i \leq n, \\ y_i &= x_i^*, & \text{for even } i \text{ with } 2 \leq i \leq n-1. \end{aligned}$$

Remark 3.37. Let $1 \leq i \leq n$. If i is odd, then we have $E^*(\beta_i) = y_i$ and $E^*(\beta_{n+i}) = z_i$; if i is even, then we have $E^*(\beta_i) = z_i$ and $E^*(\beta_{n+i}) = y_i$.

Remark 3.38. The straightening relations of Lemma 3.29 now become

$$\begin{aligned} z_{i+1}y_i &= vy_i z_{i+1}, & \text{for } i \text{ odd with } 1 \leq i \leq n-2, \\ z_{i-1}y_i &= vy_i z_{i-1}, & \text{for } i \text{ odd with } 3 \leq i \leq n, \\ z_{i+2}y_i &= vy_i z_{i+2}, & \text{for } i \text{ odd with } 1 \leq i \leq n-2, \\ z_{i-2}y_i &= vy_i z_{i-2}, & \text{for } i \text{ odd with } 3 \leq i \leq n, \\ z_1y_1 &= v^{-1}y_1z_1 + (1 - v^{-2})z_2, \end{aligned}$$

$$\begin{aligned} z_i y_i &= y_i z_i + (v - v^{-1}) z_{i-1} z_{i+1}, & \text{for } i \text{ odd with } 3 \leq i \leq n-2, \\ z_n y_n &= v^{-1} y_n z_n + (1 - v^{-2}) z_{n-1}, \end{aligned}$$

$$\begin{aligned} y_{i+3} y_i &= v y_i y_{i+3}, & \text{for } i \text{ odd with } 1 \leq i \leq n-4, \\ y_{i-3} y_i &= v y_i y_{i-3}, & \text{for } i \text{ odd with } 5 \leq i \leq n, \\ y_{i-1} y_i &= y_i y_{i-1} + (v - v^{-1}) z_{i+1} z_{i-2}, & \text{for } i \text{ odd with } 3 \leq i \leq n-2, \\ y_{n-1} y_n &= v^{-1} y_n y_{n-1} + (1 - v^{-2}) z_{n-2}, \\ y_2 y_1 &= v^{-1} y_1 y_2 + (1 - v^{-2}) z_3, \\ y_{i+1} y_i &= y_i y_{i+1} + (v - v^{-1}) z_{i-1} z_{i+2}, & \text{for } i \text{ odd with } 3 \leq i \leq n-2, \end{aligned}$$

$$\begin{aligned} z_{i+1} z_i &= v z_i z_{i+1}, & \text{for } i \text{ even with } 2 \leq i \leq n-1, \\ z_{i-1} z_i &= v z_i z_{i-1}, & \text{for } i \text{ even with } 2 \leq i \leq n-1, \end{aligned}$$

$$\begin{aligned} y_{i+2} z_i &= v z_i y_{i+2}, & \text{for } i \text{ even with } 2 \leq i \leq n-3, \\ y_{i-2} z_i &= v z_i y_{i-2}, & \text{for } i \text{ even with } 4 \leq i \leq n-1, \\ y_i z_i &= z_i y_i + (v - v^{-1}) z_{i-1} z_{i+1}, & \text{for } i \text{ even with } 2 \leq i \leq n-1, \end{aligned}$$

$$\begin{aligned} y_{i+1} z_i &= v z_i y_{i+1}, & \text{for } i \text{ odd with } 1 \leq i \leq n-2, \\ y_{i-1} z_i &= v z_i y_{i-1}, & \text{for } i \text{ odd with } 3 \leq i \leq n. \end{aligned}$$

Definition 3.39. For every element $\mathbf{a} \in \mathbb{N}^{2n}$ denote by $E[\mathbf{a}]^* \in U_v^+(w)$ the dual of $E[\mathbf{a}] \in \mathcal{P}$, i.e., the unique scalar multiple of $E[\mathbf{a}]$ such that $(E[\mathbf{a}], E[\mathbf{a}]^*) = 1$.

Consider $U_v^+(w)_{\mathbb{Z}} = \bigoplus_{\mathbf{a} \in \mathbb{N}^{2n}} \mathbb{Z}[v, v^{-1}] E[\mathbf{a}]^*$, the integral form of $U_v^+(w)$. Furthermore, put $\mathcal{A}(w)_1 = \mathbb{Q} \otimes_{\mathbb{Z}[v, v^{-1}]} U_v^+(w)_{\mathbb{Z}}$. Here, the action of v in the tensor product is given by 1. We call the algebra $\mathcal{A}(w)_1$ the classical limit of $U_v^+(w)$ or the specialization of $U_v^+(w)$ at $v = 1$. Furthermore, the $\mathbb{Z}[v^{\pm \frac{1}{2}}]$ -algebra $\mathcal{A}_v(w) = \bigoplus_{\mathbf{a} \in \mathbb{N}^{2n}} \mathbb{Z}[v^{\pm \frac{1}{2}}] E[\mathbf{a}]^*$ is the integral form of the algebra $\mathbb{Q}[v^{\pm \frac{1}{2}}] \otimes_{\mathbb{Z}[v, v^{-1}]} \mathcal{A}(w)$ and will be useful in further considerations.

Remark 3.40. Note that, by the form of the straightening relations for the dual variables from above, $\mathcal{A}(w)_1$ is a commutative algebra.

Definition 3.41. Define a function $b: \mathbb{N}^{2n} \rightarrow \mathbb{Z}$ by $b(\mathbf{a}) = \sum_{k=1}^{2n} \binom{a_k}{2}$ for a sequence $\mathbf{a} = (a_1, a_2, \dots, a_{2n}) \in \mathbb{N}^{2n}$.

Proposition 3.42. For every sequence $\mathbf{a} = (a_1, a_2, \dots, a_{2n}) \in \mathbb{N}^{2n}$ the equation $E[\mathbf{a}]^* = v^{-b(\mathbf{a})} E^*(\beta_1)^{a_1} E^*(\beta_2)^{a_2} \dots E^*(\beta_{2n})^{a_{2n}}$ is true.

Proof. The proposition follows from Lusztig's evaluation [42, Proposition 38.2.3] for of the bilinear form at Poincaré-Birkhoff-Witt basis elements together with Leclerc's conversion formula [39, Section 2.2], see Leclerc [39, Section 5.5.3]. \square

3.6 The dual canonical basis

In this section we present the dual canonical basis of $U_v^+(w)$. It is the dual of Lusztig's canonical basis from Lusztig [42, Theorem 14.2.3]. We need some auxiliary notations.

The following definitions are special cases of Leclerc's definitions [39, Section 2.7] where \mathfrak{g} is a general complex semisimple Lie algebra.

Definition 3.43. For $\mathbf{a} = (a_1, a_2, \dots, a_{2n}) \in \mathbb{N}^{2n}$ write $\deg(E[\mathbf{a}]^*) = \sum_{k=1}^{2n} a_k \beta_k \in Q^+$ as a \mathbb{N} -linear combination in the simple roots, i.e., $\deg(E[\mathbf{a}]^*) = \sum_{k=1}^{2n} a_k \beta_k = \sum_{i=1}^n d_i \alpha_i$ with $d_i \in \mathbb{Z}$. Put

$$N(\mathbf{a}) = \frac{1}{2} \left(\deg(E[\mathbf{a}]^*), \deg(E[\mathbf{a}]^*) \right) - \sum_{i=1}^n d_i.$$

We call N the *norm* of the sequence $\mathbf{a} \in \mathbb{N}^{2n}$. We also use the convention

$$N\left(\sum_{i=1}^n d_i \alpha_i\right) = \frac{1}{2} \left(\sum_{i=1}^n d_i \alpha_i, \sum_{i=1}^n d_i \alpha_i \right) - \sum_{i=1}^n d_i$$

for elements in the root lattice.

Proposition 3.44. For every natural number k with $1 \leq k \leq 2n$ we have $\sigma(E^*(\beta_k)) = v^{N(\beta_k)} E^*(\beta_k)$.

Proof. Leclerc [39, Lemma 7] proves that a homogeneous element $f \in \mathcal{F}$ satisfies $\sigma(f) = v^{N(|f|)} f$ if and only if all coefficients in the expansion of f in the basis of shuffles are invariant under σ . By Remark 3.35 all coefficients are 0 or 1 in the case of $E^*(\beta_k)$. \square

Definition 3.45. We define a partial order \triangleleft on the parametrizing set \mathbb{N}^{2n} of dual Poincaré-Birkhoff-Witt basis elements. For every k with $1 \leq k \leq 2n$ let $\mathbf{e}_k \in \mathbb{N}^{2n}$ be the vector satisfying $(\mathbf{e}_k)_l = \delta_{k,l}$ for all l . For every integer i with $1 \leq i \leq n$ let $k_i, l_i, m_i, n_i \in \{1, 2, \dots, 2n\}$ be the indices for which $|y_i| = \beta_{k_i}$, $|z_{i-1}| = \beta_{l_i}$, $|z_{i+1}| = \beta_{m_i}$, $|z_i| = \beta_{n_i}$. (Note that k_1 and n_n are not defined. We put $\mathbf{e}_{k_1} = \mathbf{e}_{n_n} = 0$.) Put $\mathbf{v}_i = \mathbf{e}_{k_i} - \mathbf{e}_{l_i} - \mathbf{e}_{m_i} + \mathbf{e}_{n_i}$. Now we say that $\mathbf{a}, \mathbf{b} \in \mathbb{N}^{2n}$ satisfy $\mathbf{a} \triangleleft \mathbf{b}$ if and only if $\mathbf{b} - \mathbf{a} \in \bigoplus_{i=1}^n \mathbb{N} \mathbf{v}_i$.

Remark 3.46. The straightening relations of Lemma 3.29 imply the following fact: If $\mathbf{a} \in \mathbb{N}^{2n}$ and we expand $E^*(\beta_{2n})^{a_{2n}} E^*(\beta_{2n-1})^{a_{2n-1}} \dots E^*(\beta_1)^{a_1}$ in the dual Poincaré-Birkhoff-Witt basis, then we get a $\mathbb{Q}(v)$ -linear combination of $E[\mathbf{b}]^*$ with $\mathbf{b} \in S(\mathbf{a}) \cup \{\mathbf{a}\}$.

Definition 3.47. For $\mathbf{a} \in \mathbb{N}^{2n}$ put $S(\mathbf{a}) = \{\mathbf{b} \in \mathbb{N}^{2n} : \mathbf{a} \triangleleft \mathbf{b}, \mathbf{a} \neq \mathbf{b}\}$.

Theorem 3.48. There exist elements $B[\mathbf{a}]^* \in U_v^+(w)$ parametrized by sequences $\mathbf{a} \in \mathbb{N}^{2n}$ such that the set $\mathcal{B}^* = \{B[\mathbf{a}]^* : \mathbf{a} \in \mathbb{N}^{2n}\}$ is a basis of $U_v^+(w)$ and the following two properties hold.

- (1) For every $\mathbf{a} \in \mathbb{N}^{2n}$ we have $B[\mathbf{a}]^* - E[\mathbf{a}]^* \in \bigoplus_{\mathbf{b} \in S(\mathbf{a})} v^{-1} \mathbb{Z}[v^{-1}] E[\mathbf{b}]^*$.
- (2) For every $\mathbf{a} \in \mathbb{N}^{2n}$ we have $\sigma(B[\mathbf{a}]^*) = v^{N(\mathbf{a})} B[\mathbf{a}]^*$.

The elements $B[\mathbf{a}]^* \in U_v^+(w)$ are uniquely determined by these two properties.

Proof. Note that if $\mathbf{a} \triangleleft \mathbf{b}$, then $|E[\mathbf{a}]^*| = \sum_{k=1}^{2n} a_k \beta_k = \sum_{k=1}^{2n} b_k \beta_k = |E[\mathbf{b}]^*|$ since the straightening relations are relations in a R -graded algebra. For $\gamma \in R$ in the root lattice, consider the (finite) set $S_\gamma \subset \mathbb{N}^{2n}$ of all $\mathbf{a} = (a_i)_{1 \leq i \leq 2n}$ with $\sum_{k=1}^{2n} a_k \beta_k = \gamma$

γ . We extend the partial order \triangleleft on S_γ to a total order $<$. Let $\mathbf{a}_1 < \mathbf{a}_2 < \dots < \mathbf{a}_m$ be the elements of S_γ written in increasing order. Now we prove by backward induction that for every $k = m, m-1, \dots, 2, 1$ there exist linearly independent $B[\mathbf{a}_k], B[\mathbf{a}_{k+1}], \dots, B[\mathbf{a}_m]$ satisfying (1) and (2).

Put $B[\mathbf{a}_m] = E[\mathbf{a}_m]$. It clearly satisfies property (1). Let $\mathbf{a}_m = (a_1, a_2, \dots, a_{2n})$. Since there are no $\mathbf{b} \in S_\gamma$ such that $\mathbf{a}_m < \mathbf{b}$, the dual Poincaré-Birkhoff-Witt element $E[\mathbf{a}_m]^*$ cannot be straightened, i.e., all $E^*(\beta_k)$ for which $a_k \neq 0$ are v -commutative. Therefore, by Remark 3.30 we have

$$\begin{aligned} \sigma(E[\mathbf{a}_m]^*) &= \sigma(v^{-b(\mathbf{a})} E^*(\beta_1)^{a_1} E^*(\beta_2)^{a_2} \dots E^*(\beta_{2n})^{a_{2n}}) \\ &= v^{b(\mathbf{a})} v^{\sum_{k=1}^{2n} a_k N(\beta_k)} E^*(\beta_{2n})^{a_{2n}} \dots E^*(\beta_2)^{a_2} E^*(\beta_1)^{a_1} \\ &= v^{b(\mathbf{a})} v^{\sum_{k=1}^{2n} \frac{1}{2} a_k (\beta_k, \beta_k) - \sum_{k=1}^{2n} a_k \|\beta_k\|} \cdot v^{\sum_{k < l} (\beta_k, \beta_l)} E^*(\beta_1)^{a_1} E^*(\beta_2)^{a_2} \dots E^*(\beta_{2n})^{a_{2n}} \\ &= v^{\frac{1}{2} (\sum_{k=1}^{2n} \beta_k, \sum_{k=1}^{2n} \beta_k) - \sum_{k=1}^{2n} a_k \|\beta_k\|} E[\mathbf{a}_m]^* \\ &= v^{N(\mathbf{a}_m)} E[\mathbf{a}_m]^*. \end{aligned}$$

Here $\|\beta_k\|$ denotes the sum of the coefficients of β_k when expanded as a \mathbb{Z} -linear combination of simple roots as in Definition 3.43. Hence, the variable $E[\mathbf{a}_m]$ also satisfies property (2).

Now let $1 \leq k < m$ and assume that properties (1) and (2) hold for $\mathbf{a}_{k+1}, \mathbf{a}_{k+2}, \dots, \mathbf{a}_m$. We expand $\sigma(E[\mathbf{a}_k]^*)$ in the dual Poincaré-Birkhoff-Witt basis. Note that by the same argument as above (and ignoring terms of lower order) we see that the coefficient of the leading term $E[\mathbf{a}_k]^*$ is $v^{N(\mathbf{a}_k)}$. Thus, we have

$$\sigma(E[\mathbf{a}_k]^*) = v^{N(\mathbf{a}_k)} E[\mathbf{a}_k]^* + \sum_{k < l \leq m} f_l E[\mathbf{a}_l]^*$$

for some $f_l \in \mathbb{Z}[v, v^{-1}]$. By induction hypothesis every $B[\mathbf{a}_l]^*$ with $l > k$ is a $\mathbb{Z}[v, v^{-1}]$ -linear combination of $E[\mathbf{a}_l']^*$ with $l' > l$. By solving an upper triangular linear system of equations we see that every $E[\mathbf{a}_l]^*$ with $l > k$ is a $\mathbb{Z}[v, v^{-1}]$ -linear combination of $B[\mathbf{a}_l']^*$ with $l' > l$. Hence, we may write

$$\sigma(E[\mathbf{a}_k]^*) = v^{N(\mathbf{a}_k)} E[\mathbf{a}_k]^* + \sum_{k < l \leq m} g_l B[\mathbf{a}_l]^*$$

for some $g_l \in \mathbb{Z}[v, v^{-1}]$. We apply the antiinvolution σ to the last equation:

$$E[\mathbf{a}_k]^* = v^{-N(\mathbf{a}_k)} \sigma(E[\mathbf{a}_k]^*) + \sum_{k < l \leq m} v^{N(\mathbf{a}_l)} \sigma(g_l) B[\mathbf{a}_l]^*.$$

Note that $N(\mathbf{a}_k) = N(\mathbf{a}_l)$ for all l . Comparing coefficients yields $v^{2N(\mathbf{a}_l)} \sigma(g_l) = -g_l$. It follows that $\sigma(v^{-N(\mathbf{a}_l)} g_l) = -v^{-N(\mathbf{a}_l)} g_l$. Thus, we may write $v^{-N(\mathbf{a}_l)} g_l = h_l - \sigma(h_l)$ for some $h_l \in v^{-1} \mathbb{Z}[v^{-1}]$. Now put

$$B[\mathbf{a}_k]^* = E[\mathbf{a}_k]^* + \sum_{k < l \leq m} h_l B[\mathbf{a}_l]^*.$$

It is easy to see that properties (1) and (2) are true for $B[\mathbf{a}_k]^*$ and that $B[\mathbf{a}_k]^*, B[\mathbf{a}_{k+1}]^*, \dots, B[\mathbf{a}_m]^*$ are linearly independent.

For the uniqueness, suppose that k is some index such that there are variables $B[\mathbf{a}_k]_1^*$ and $B[\mathbf{a}_k]_2^*$ fulfilling the two properties of the theorem. Then their difference $B[\mathbf{a}_k]_1^* - B[\mathbf{a}_k]_2^* \in \bigoplus_{l > k} v^{-1} \mathbb{Z}[v^{-1}] B[\mathbf{a}_l]^*$. Application of σ and multiplication with $v^{-N(\mathbf{a}_k)}$ afterwards yields $B[\mathbf{a}_k]_1^* - B[\mathbf{a}_k]_2^* \in \bigoplus_{l > k} v \mathbb{Z}[v] B[\mathbf{a}_l]^*$, so $B[\mathbf{a}_k]_1^* = B[\mathbf{a}_k]_2^*$. \square

Remark 3.49. It is known that the dual of Lusztig's canonical basis under Kashiwara's bilinear form obeys the two properties of Theorem 3.48, compare Leclerc [39, Proposition 39]. By uniqueness, the set $\mathcal{B}^* = \{B[\mathbf{a}]^* : \mathbf{a} \in \mathbb{N}^{2n}\}$ is the dual of Lusztig's canonical basis, or the *dual canonical basis* for short.

We call $E[\mathbf{a}]^*$ (for $\mathbf{a} \in \mathbb{N}^{2n}$) the *leading term* in the expansion of $B[\mathbf{a}]^*$ in the Poincaré-Birkhoff-Witt basis. In what follows we use the convention $z_0 = z_{n+1} = 1$. Prominent elements in \mathcal{B}^* are

$$\begin{aligned} p_i &= y_i z_i - v^{-1} z_{i-1} z_{i+1}, & \text{for } i \text{ odd with } 1 \leq i \leq n, \\ p_i &= z_i y_i - v^{-1} z_{i-1} z_{i+1}, & \text{for } i \text{ even with } 2 \leq i \leq n-1. \end{aligned}$$

The first property of Theorem 3.48 is obvious and the second follows easily from a calculation using the straightening relations. The variables are v -deformations of the δ -functions of the \mathcal{C}_M -projective rigid Λ -modules from Section 2.2. The non-deformed δ -function associated with these modules are frozen cluster variables in Geiß-Leclerc-Schröer's cluster algebra $\mathcal{A}(\mathcal{C}_M)$, compare Section 2.4.

Lemma 3.50. For every pair (i, k) of natural numbers such that $1 \leq i \leq n$ and $1 \leq k \leq 2n$ the elements p_i and $E^*(\beta_k) \in U_v^+(w)$ are v -commutative, i.e., there is an integer a such that $p_i E^*(\beta_k) = v^a E^*(\beta_k) p_i$.

Proof. First of all, we claim that $(|z_i|, |y_i|)$ is equal to -1 if $i \in \{1, n\}$ and equal to 0 if $2 \leq i \leq n-1$. For a proof, note that $(|z_1|, |y_1|) = (\alpha_2 + \alpha_3, \alpha_1) = -1$, and similarly $(|z_n|, |y_n|) = -1$. In the same way we see that $(|z_2|, |y_2|) = (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5, \alpha_1 + \alpha_2 + \alpha_3) = 0$, and similarly $(|z_{n-1}|, |y_{n-1}|) = 0$. Furthermore, we have $(\alpha_i + \alpha_{i+1} + \dots + \alpha_j, \alpha_k) = 0$ for all $i+1 \leq k \leq j-1$ from which it easily follows that $(|z_i|, |y_i|) = 0$ for $i \in \{3, 4, \dots, n-2\}$.

The claim implies that for odd i in the straightening relation for $z_i y_i$ the coefficient in front of $y_i z_i$ is $v^{-(|z_i|, |y_i|)}$. The same is true for the coefficient in front of $z_i y_i$ in the straightening relation for $y_i z_i$. Let i be an odd integer such that $1 \leq i \leq n$. Note that

$$p_i = y_i z_i - v^{-1} z_{i-1} z_{i+1} = v^{-(|z_i|, |y_i|)} z_i y_i - v z_{i-1} z_{i+1}$$

by property (2) of Theorem 3.48. From the straightening relations of Lemma 3.29 it is clear that p_i commutes with every y_j with $|j-i| \geq 4$ and with every z_j with $|j-i| \geq 3$. If $i \geq 5$, then $y_{i-3} p_i = y_{i-3} y_i z_i - v^{-1} y_{i-3} z_{i-1} z_{i+1} = v y_i z_i y_{i-3} - z_{i-1} z_{i+1} y_{i-3} = v p_i y_{i-3}$. Now assume that $i \geq 3$. We have $y_{i-2} p_i = y_{i-2} y_i z_i - v^{-1} y_{i-2} z_{i-1} z_{i+1} = v^{-1} y_i z_i y_{i-2} - v^{-2} z_{i-1} z_{i+1} y_{i-2} = v^{-1} p_i y_{i-2}$. Furthermore, we see that $z_{i-2} p_i = z_{i-2} y_i z_i - v^{-1} z_{i-2} z_{i-1} z_{i+1} = v y_i z_i z_{i-2} - z_{i-1} z_{i+1} z_{i-2} = v p_i z_{i-2}$. The calculation

$$\begin{aligned} y_{i-1} p_i &= y_{i-1} y_i z_i - v^{-1} y_{i-1} z_{i-1} z_{i+1} \\ &= v^{(|y_{i-1}|, |y_i|)} \left(y_i y_{i-1} + (v - v^{-1}) z_{i+1} z_{i-2} \right) z_i \\ &\quad - v^{-1} \left(z_{i-1} y_{i-1} + (v - v^{-1}) z_i z_{i-2} \right) z_{i+1} \\ &= v^{(|y_{i-1}|, |y_i|)} (v y_i z_i y_{i-1} - z_{i-1} z_{i+1} y_{i-1}) = v^{1+(|y_{i-1}|, |y_i|)} p_i y_{i-1} \end{aligned}$$

shows that p_i also v -commutes with y_{i-1} . It also v -commutes with z_{i-1} as the calculation shows: $z_{i-1} p_i = z_{i-1} y_i z_i - v^{-1} z_{i-1}^2 z_{i+1} = y_i z_i z_{i-1} - v^{-1} z_{i-1} z_{i+1} z_{i-1} =$

$p_i z_{i-1}$. Now assume that $i \geq 1$. The following equation is true:

$$\begin{aligned}
y_i p_i &= y_i^2 z_i - v^{-1} y_i z_{i-1} z_{i+1} \\
&= y_i \left(v^{-(|y_i|, |z_i|)} z_i y_i + (v^{-1} - v) z_{i-1} z_{i+1} \right) - v^{-1} y_i z_{i-1} z_{i+1} \\
&= v^{-(|y_i|, |z_i|)} y_i z_i z_i - v y_i z_{i-1} z_{i+1} \\
&= v^{-(|y_i|, |z_i|)} (y_i z_i - v^{-1} z_{i-1} z_{i+1}) y_i = v^{-(|y_i|, |z_i|)} p_i y_i.
\end{aligned}$$

Finally we see that $z_i p_i = z_i y_i z_i - v^{-1} z_i z_{i-1} z_{i+1} = v^{(|y_i|, |z_i|)} (v^{-(|y_i|, |z_i|)} z_i y_i - v z_{i-1} z_{i+1}) z_i = v^{(|y_i|, |z_i|)} p_i z_i$. The v -commutativity relations of p_i with elements with index $j > i$ are proved in the same way.

The case of even i can be handled with similar arguments. \square

Remark 3.51. 1. The v -commutativity relations of Lemma 3.50 will be crucial for the construction of the initial quantum seed of $\mathcal{A}(w)_v$. Another verification of v -commutativity relations for the initial seed is due to Kimura [36, Section 6].

2. Multiplicative properties of (dual) canonical basis elements have also been studied by Reineke [44].

3. As observed by Leclerc, the v -deformations of the δ -functions of \mathcal{C}_M -projective rigid Λ -modules also v -commute with the generators of $U_v^+(w)$ in the Kronecker cases for w of length 4, see the author [38, Section 4.1].

4. A consideration of the leading terms of the two occurring variables in Lemma 3.50 is sufficient to determine the integer a .

Remark 3.52. The techniques in this section work with the same proofs for the case $U_v^+(w')$ as well. We use a similar notation $y'_i, z'_i \in U_v^+(w')$ for the elements dual to $u'_i, v'_i, w'_i, x'_i \in U_v^+(w)$ (for appropriate indices i). The straightening relations for the dual variables can be computed using the same methods.

4 The quantum cluster algebra structure induced by the dual canonical basis

In this section we are going to prove that the integral form $\bigoplus_{\mathbf{a} \in \mathbb{N}^{2n}} \mathbb{Z}[v^{\pm \frac{1}{2}}] E[\mathbf{a}]^*$ is a quantum cluster algebra in the sense of Bereinstein-Zelevinsky [10]. The corresponding non-quantized cluster algebra is Geiß-Leclerc-Schröer's [28] cluster algebra $\mathcal{A}(w)$.

Natural Quantum cluster algebra structures have only been observed in very few cases, see for example Grabowski-Launois [24], Rupel [49], and the author [38]. For a study of bases of quantum cluster algebras of type $\tilde{A}_1^{(1)}$ see Ding-Xu [15].

Definition 4.1. For $1 \leq i \leq j \leq n$ define $\Delta_{i,j}^v \in \mathcal{B}^*$ to be the dual canonical basis element with leading term $\prod_{i \leq r \leq j, r \text{ odd}} y_r \prod_{i \leq r \leq j, r \text{ even}} y_r$.

Remark 4.2. We provide some examples of elements in the dual canonical basis \mathcal{B}^* of the form $\Delta_{i,j}^v$ with $1 \leq i \leq j \leq n$. We focus on examples where the interval $[i, j]$ is small, i.e., $j \leq i + 2$. First of all, we clearly have $\Delta_{i,i}^v = y_i$ for all i . Furthermore, an elementary calculation using the straightening relations shows that: $\Delta_{1,2}^v = y_1 y_2 -$

$v^{-1}z_3 = vy_2y_1 - vz_3$, $\Delta_{n-1,n}^v = y_ny_{n-1} - v^{-1}z_{n-2} = vy_{n-1}y_n - vz_{n-2}$, and that for $2 \leq i \leq n-2$

$$\Delta_{i,i+1}^v = \begin{cases} y_iy_{i+1} - v^{-1}z_{i-1}z_{i+2} = y_{i+1}y_i - vz_{i+2}z_{i-1}, & \text{if } i \text{ is odd;} \\ y_{i+1}y_i - v^{-1}z_{i+2}z_{i-1} = y_iy_{i+1} - vz_{i-1}z_{i+2}, & \text{if } i \text{ is even.} \end{cases}$$

Recall the convention $z_0 = z_{n+1} = 1$. The formulae simplify to $\Delta_{i,i+1}^v = y_iy_{i+1} - v^{-1}z_{i-1}z_{i+2}$ for odd i and $\Delta_{i,i+1}^v = y_{i+1}y_i - v^{-1}z_{i+2}z_{i-1}$ for even i . With the same convention we can compute:

$$\begin{aligned} \Delta_{1,3} &= y_1y_3y_2 - v^{-1}y_1z_4z_1 - v^{-1}y_3z_3 + v^{-2}z_2z_4 \\ &= vy_2y_1y_3 - vz_3y_3 - v^2z_1z_4y_1 + v^2z_2z_4, \\ \Delta_{n-2,n} &= y_ny_{n-2}y_{n-1} - v^{-1}y_nz_{n-3}z_n - v^{-1}y_{n-2}z_{n-2} + v^{-2}z_{n-1}z_{n-3} \\ &= vy_{n-1}y_ny_{n-2} - vz_{n-2}y_{n-2} - v^2z_nz_{n-3}y_n + v^2z_{n-1}z_{n-3}. \end{aligned}$$

For odd i such that $3 \leq i \leq n-4$ we have:

$$\begin{aligned} \Delta_{i,i+2} &= y_iy_{i+2}y_{i+1} - v^{-1}y_iz_{i+3}z_i - v^{-1}y_{i+2}z_{i-1}z_{i+2} + v^{-2}z_{i-1}z_{i+1}z_{i+3} \\ &= y_{i+1}y_{i+2}y_i - vz_iz_{i+3}y_i - vz_{i+2}z_{i-1}y_{i+2} + v^2z_{i-1}z_{i+1}z_{i+3}. \end{aligned}$$

For odd i such that $3 \leq i \leq n-4$ we have:

$$\begin{aligned} \Delta_{i,i+2} &= y_{i+1}y_iy_{i+2} - v^{-1}z_iz_{i+3}y_i - v^{-1}z_{i+2}z_{i-1}y_{i+2} + v^2z_{i-1}z_{i+1}z_{i+3} \\ &= y_iy_{i+2}y_{i+1} - vy_iz_{i+3}z_i - vy_{i+2}z_{i+2}z_{i-1} + v^2z_{i-1}z_{i+1}z_{i+3}. \end{aligned}$$

Remark 4.3. In what follows we prove recursions for the $\Delta_{i,j}^v$ with $1 \leq i \leq j \leq n$. The formulae turn out to be quantized versions of the formulae for the $\Delta_{i,j}$ from Section 2.5. The quantized recursions will be crucial for the verification of the quantum cluster algebra structure on $U_v^+(w)$ in the next section.

To formulate the recursion effectively the following definitions are helpful. First of all we introduce the convention $\Delta_{i,i-1}^v = 1$ for all i . Furthermore, we give the following two definitions.

Definition 4.4. For $1 \leq i, j \leq n$ put

$$s_{i,j} = \sum_{i \leq s \leq j} |y_s|; \quad o_{i,j} = \sum_{\substack{i \leq s \leq j \\ s \text{ odd}}} |y_s|; \quad e_{i,j} = \sum_{\substack{i \leq s \leq j \\ s \text{ even}}} |y_s|.$$

Definition 4.5. For all i, j with $1 \leq i, j \leq n$ and $j - i \geq 2$ define

$$A_{i,j} = \begin{cases} -1, & \text{if } j - i \leq 3; \\ -2, & \text{if } j - i \geq 4. \end{cases}$$

Lemma 4.6. For all pairs (i, j) of integers such that $1 \leq i \leq j \leq n$ the following equation is true:

$$N(s_{i,j}) - N(s_{i,j-1}) - N(|y_j|) = \begin{cases} (|y_j|, o_{i,j-1}), & \text{if } j \text{ is even;} \\ (|y_j|, e_{i,j-1}), & \text{if } j \text{ is odd.} \end{cases}$$

Proof. The proposition follows easily from the observation $\frac{1}{2}(s_{i,j}, s_{i,j}) = \frac{1}{2}(e_{i,j} + o_{i,j}, e_{i,j} + o_{i,j}) = (j - i + 1) + (e_{i,j}, o_{i,j})$. \square

Lemma 4.7. Let i, j be integers such that $1 \leq i \leq j \leq n$ and $j \geq i + 2$. The following equation holds:

$$N(s_{i,j}) - N(s_{i,j-3}) - N(|p_{j-2}|) - N(|z_{j+1}|) = \begin{cases} (|y_j| + |y_{j-2}|, o_{i,j-3}) + (|y_{j-1}|, e_{i,j-4}), & \text{if } j \text{ is even;} \\ (|y_j| + |y_{j-2}|, e_{i,j-3}) + (|y_{j-1}|, o_{i,j-4}), & \text{if } j \text{ is odd.} \end{cases}$$

Proof. If j is odd, then the elements $y_j, y_{j-1}, z_{j+1}, z_{j-2}$ have degrees $\beta_j, \beta_{n+j-1}, \beta_{j+1}$ and β_{n+j-2} , respectively. If j is even, then the elements $y_j, y_{j-1}, z_{j+1}, z_{j-2}$ have degrees $\beta_{n+j}, \beta_{j-1}, \beta_{n+j+1}$ and β_{j-2} , respectively. It is easy to see in Figure 6 that in all cases the equation $|y_j| + |y_{j-1}| = |z_{j+1}| + |z_{j-2}|$ holds. With this fact the proposition follows just as above from the observation $\frac{1}{2}(s_{i,j}, s_{i,j}) = \frac{1}{2}(e_{i,j} + o_{i,j}, e_{i,j} + o_{i,j}) = (j - i + 1) + (e_{i,j}, o_{i,j})$. \square

Remark 4.8. By definition, the leading term of the variable $\Delta_{i,j}^v$ (where $1 \leq i \leq j \leq n$) is $\prod_{i \leq r \leq j, r \text{ odd}} y_r \prod_{i \leq r \leq j, r \text{ even}} y_r$. Therefore, $\Delta_{i,j}^v$ is a $\mathbb{Z}[v, v^{-1}]$ -linear combination of terms of the form

$$\prod_{\substack{i \leq r \leq j \\ r \text{ odd}}} y_r^{1-a_r} \prod_{\substack{i-1 \leq r \leq j+1 \\ r \text{ even}}} z_r^{a_{r+1}+a_{r-1}-a_r} \prod_{\substack{i-1 \leq r \leq j+1 \\ r \text{ odd}}} z_r^{a_{r+1}+a_{r-1}-a_r} \prod_{\substack{i \leq r \leq j \\ r \text{ even}}} y_r^{1-a_r}$$

for some $\mathbf{a} = (a_1, a_2, \dots, a_n) \in \{0, 1\}^n$ such that $a_1 = a_2 = \dots = a_{i-1} = 0$, $a_{j+1} = a_{j+2} = \dots = a_n = 0$, and $a_{r+1} + a_{r-1} - a_r \geq 0$ for all r . We denote this term by $\Delta_{i,j}^v[\mathbf{a}]$.

We know that p_{j+1} commutes, up to a power of v , with every $\Delta_{i,j}^v[\mathbf{a}]$. The following proposition shows that much more is true: The commutation exponent only depends on the pair (i, j) , but not on $\mathbf{a} \in \{0, 1\}^n$.

Proposition 4.9. Let i, j be integers such that $1 \leq i \leq j \leq n - 1$. The variables $\Delta_{i,j}^v[\mathbf{a}]$ and p_{j+1} are v -commutative. More precisely: If j is even, then

$$\Delta_{i,j}^v[\mathbf{a}]p_{j+1} = v^{(|y_{j+1}|, e_{i,j}) + (|z_{j+1}|, e_{i,j}) - (|z_{j+1}|, o_{i,j-1})} p_{j+1} \Delta_{i,j}^v[\mathbf{a}],$$

and if j is odd, then

$$\Delta_{i,j}^v[\mathbf{a}]p_{j+1} = v^{-(|y_{j+1}|, o_{i,j}) + (|z_{j+1}|, e_{i,j}) - (|z_{j+1}|, o_{i,j-1})} p_{j+1} \Delta_{i,j}^v[\mathbf{a}].$$

Proof. Assume that j be even. By Lemma 3.50 we know that p_{j+1} commutes, up to a power of v , with every $\Delta_{i,j}^v[\mathbf{a}]$. To determine the appropriate power of v , we compare the leading terms. The leading term of p_{j+1} is $y_{j+1}z_{j+1}$. By Remark 3.30 we see that

$$\begin{aligned} \Delta_{i,j}^v[\mathbf{a}]p_{j+1} &= v^{a_j(|y_{j+1}|, |z_{j+1}| + |z_{j-1}| - |y_j| - |-z_j|)} \cdot v^{a_{j-1}(|y_{j+1}|, |z_j| + |z_{j-2}| - |z_{j-1}|)} \\ &\quad \cdot v^{a_{j-2}(|y_{j+1}|, |z_{j-1}| - |z_{j-2}|)} \cdot v^{a_j(|z_{j+1}|, |y_j| - |z_j|)} \cdot v^{a_{j-1}(|z_{j+1}|, |y_{j-1}| - |z_j| + |z_{j-2}|)} \\ &\quad \cdot v^{(|y_{j+1}|, e_{i,j}) + (|z_{j+1}|, e_{i,j}) - (|z_{j+1}|, o_{i,j-1})} p_{j+1} \Delta_{i,j}^v[\mathbf{a}]. \end{aligned}$$

Notice that first of all $(|y_k|, |y_{k-2}|) = (|y_k|, |y_{k-4}|) = 0$ for all k , that secondly $(|y_k|, |y_l|) = 0$ for $|k - l| \geq 4$, that thirdly $(|y_k|, |z_l|) = 0$ for $|k - l| \geq 3$, and that

finally $|y_j| + |z_j| = |z_{j+1}| + |z_{j-1}|$ for all k . Furthermore, note that $(|z_{k+1}|, |y_k|) = (|z_{k+1}|, |z_k|)$ for all k . Hence, we have

$$\Delta_{i,j}^v[\mathbf{a}]p_{j+1} = v^{(|y_{j+1}|, e_{i,j}) + (|z_{j+1}|, e_{i,j}) - (|z_{j+1}|, o_{i,j-1})} p_{j+1} \Delta_{i,j}^v[\mathbf{a}].$$

By a similar argument we can show that for odd numbers j the equation

$$\Delta_{i,j}^v[\mathbf{a}]p_{j+1} = v^{-(|y_{j+1}|, o_{i,j}) + (|z_{j+1}|, e_{i,j}) - (|z_{j+1}|, o_{i,j-1})} p_{j+1} \Delta_{i,j}^v[\mathbf{a}].$$

□

The independence of the commutation exponent on $\mathbf{a} \in \{0, 1\}^n$ implies the corollary.

Corollary 4.10. Let i, j be integers as above. Then the variables $\Delta_{i,j}^v$ and p_{j+1} are v -commutative. More precisely: If j is even, then

$$\Delta_{i,j}^v p_{j+1} = v^{(|y_{j+1}|, e_{i,j}) + (|z_{j+1}|, e_{i,j}) - (|z_{j+1}|, o_{i,j-1})} p_{j+1} \Delta_{i,j}^v,$$

and if j is odd, then

$$\Delta_{i,j}^v p_{j+1} = v^{-(|y_{j+1}|, o_{i,j}) + (|z_{j+1}|, e_{i,j}) - (|z_{j+1}|, o_{i,j-1})} p_{j+1} \Delta_{i,j}^v.$$

Having established these conventions, definitions and propositions, we are now able to formulate a theorem that provides a recursion for the $\Delta_{i,j}^v$ and a quantized version of the exchange relation. These are the parts (a) and (b) of Theorem 4.11. For a proof of the theorem, we proceed by induction. For a functioning induction step we include also the v -commutator relations in part (c) and (d).

Theorem 4.11. Let i, j be integers such that $1 \leq i, j \leq n$ and $j - i \geq 2$.

(a) The dual canonical basis element $\Delta_{i,j}^v$ can be computed recursively from elements $\Delta_{i,j'}^v$, with $j' < j$. More precisely, if j is even, then we have:

$$\begin{aligned} \Delta_{i,j}^v &= \Delta_{i,j-1}^v y_j - v^{A_{i,j}} \Delta_{i,j-3}^v p_{j-2} z_{j+1} \\ &= v^{-(|y_j|, o_{i,j-1})} y_j \Delta_{i,j-1}^v \\ &\quad - v^{-A_{i,j} - (|y_j| + |y_{j-2}|, o_{i,j-3}) - (|y_{j-1}|, e_{i,j-4})} z_{j+1} p_{j-2} \Delta_{i,j-3}^v. \end{aligned}$$

If j is odd, then we have:

$$\begin{aligned} \Delta_{i,j}^v &= y_j \Delta_{i,j-1}^v - v^{A_{i,j}} z_{j+1} p_{j-2} \Delta_{i,j-3}^v \\ &= v^{-(|y_j|, e_{i,j-1})} \Delta_{i,j-1}^v y_j \\ &\quad - v^{-A_{i,j} - (|y_j| + |y_{j-2}|, e_{i,j-3}) - (|y_{j-1}|, o_{i,j-4})} \Delta_{i,j-3}^v p_{j-2} z_{j+1}. \end{aligned}$$

(b) Furthermore, the following quantum cluster exchange relation holds:

$$\Delta_{i,j}^v z_j = \begin{cases} \Delta_{i,j-1}^v p_j + v^{1 - (|y_{j-1}|, e_{i,j-2})} \Delta_{i,j-2}^v p_{j-1} z_{j+1}, & \text{if } j \text{ is even;} \\ v^{-(|y_j|, e_{i,j-1})} \Delta_{i,j-1}^v p_j + v^{-1 - (|y_j|, e_{i,j-1})} \Delta_{i,j-2}^v p_{j-1} z_{j+1}, & \text{if } j \text{ is odd.} \end{cases}$$

(c) If $j + 1 \leq n$, then the following v -commutator relation holds. If j is even, then

$$y_{j+1} \Delta_{i,j}^v = v^{-(|y_{j+1}|, e_{i,j})} \Delta_{i,j}^v y_{j+1} + v^{-1 - (|y_{j-1}|, e_{i,j-2})} (v^{-1} - v) \Delta_{i,j-2}^v p_{j-1} z_{j+2}.$$

If j is odd, then

$$y_{j+1} \Delta_{i,j}^v = v \Delta_{i,j}^v y_{j+1} + v^{1 - (|y_j|, e_{i,j-3})} (v - v^{-1}) \Delta_{i,j-2}^v p_{j-1} z_{j+2}.$$

- (d) If $j+2 \leq n$, then $\Delta_{i,j}^v$ and y_{j+2} are v -commutative. More precisely, if j is even, then $\Delta_{i,j}^v y_{j+2} = v^{-1} y_{j+2} \Delta_{i,j}^v$, and if j is odd, then $\Delta_{i,j}^v y_{j+2} = v y_{j+2} \Delta_{i,j}^v$.

Proof. We proceed by induction on $j-i$. Using the explicit formulae provided by Remark 4.2 it is easy to see that Theorem 4.11 is true for $j-i=2$. Now let $j-i \geq 3$ and assume that Theorem 4.11 is true for all smaller values of $j-i$. We distinguish two cases.

First of all assume that j is even. It follows that $4 \leq j \leq n-1$. Put $D = \Delta_{i,j-1}^v y_j - v^{A_{i,j}} \Delta_{i,j-3}^v p_{j-2} z_{j+1}$. We have to prove that $D = \Delta_{i,j}^v$, i.e., we have to show that D satisfies properties (1) and (2) of Theorem 3.48. First of all, we verify property (1). We expand the dual canonical basis elements $\Delta_{i,j-1}^v, \Delta_{i,j-3}^v$ according to Remark 4.8. We see that

$$\begin{aligned} \Delta_{i,j-1}^v = \sum_{\mathbf{a}} f_{\mathbf{a}} v^{\sum -\binom{a_{r+1}+a_{r-1}-a_r}{2}} & \prod_{\substack{i \leq r \leq j-3 \\ r \text{ odd}}} y_r^{1-a_r} \prod_{\substack{i-1 \leq r \leq j \\ r \text{ even}}} z_r^{a_{r+1}+a_{r-1}-a_r} \\ & \cdot \prod_{\substack{i-1 \leq r \leq j \\ r \text{ odd}}} z_r^{a_{r+1}+a_{r-1}-a_r} \prod_{\substack{i \leq r \leq j-1 \\ r \text{ even}}} y_r^{1-a_r} \end{aligned}$$

where the sum is taken over all the admissible sequences $\mathbf{a} = (a_1, a_2, \dots, a_n) \in \{0, 1\}^n$ as in Remark 4.8 and $f_{\mathbf{a}} \in v^{-1}\mathbb{Z}[v^{-1}]$ except for $f_0 = 1$. Here, we have used that $-\binom{1-a_r}{2} = 0$ for all terms a_r in such a sequence. It is clear that $\Delta_{i,j-1}^v y_j - \Delta_{i,j-1}^v [0] y_j \in \bigoplus_{\mathbf{b} \in S(\mathbf{a})} v^{-1}\mathbb{Z}[v^{-1}] E[\mathbf{b}]^*$ and that $\Delta_{i,j-1}^v [0] y_j$ is the dual Poincaré-Birkhoff-Witt basis element from Definition 4.1 that serves as leading term.

Furthermore, we have

$$\begin{aligned} \Delta_{i,j-3}^v = \sum_{\mathbf{a}} g_{\mathbf{a}} v^{\sum -\binom{a_{r+1}+a_{r-1}-a_r}{2}} & \prod_{\substack{i \leq r \leq j-3 \\ r \text{ odd}}} y_r^{1-a_r} \prod_{\substack{i-1 \leq r \leq j-2 \\ r \text{ even}}} z_r^{a_{r+1}+a_{r-1}-a_r} \\ & \cdot \prod_{\substack{i-1 \leq r \leq j-3 \\ r \text{ odd}}} z_r^{a_{r+1}+a_{r-1}-a_r} \prod_{\substack{i \leq r \leq j-4 \\ r \text{ even}}} y_r^{1-a_r} \end{aligned}$$

where the sum is taken over all the admissible sequences $\mathbf{a} = (a_1, a_2, \dots, a_n) \in \{0, 1\}^n$ as in Remark 4.8 and $g_{\mathbf{a}} \in v^{-1}\mathbb{Z}[v^{-1}]$ except for $g_0 = 1$. Now we consider $v^{A_{i,j}} \Delta_{i,j-3}^v p_{j-2} z_{j+1}$. Note that the generator z_{j+1} commutes with all occurring terms in this expansion. For p_{j-2} , by Lemma 3.50 the v -commutativity relation

$$\left(\prod_{\substack{i \leq r \leq j-4 \\ r \text{ even}}} y_r^{1-a_r} \right) p_{j-2} = v^{(1-a_{j-4})(|z_{j-2}|, |y_{j-4}|)} p_{j-2} \left(\prod_{\substack{i \leq r \leq j-4 \\ r \text{ even}}} y_r^{1-a_r} \right)$$

holds. Hence, in each summand we can transfer p_{j-2} to the left of the product. We get a factor $v^{(1-a_{j-4})(|z_{j-2}|, |y_{j-4}|)}$ since $(|z_{j-2}|, |y_{j-4}|) = 1$ for all j . We concentrate on a single summand. Write $p_{j-2} = z_{j-2} y_{j-2} - v^{-1} z_{j-3} z_{j-1}$. This decomposition splits the sum into two parts. Consider the summand coming from $z_{j-2} y_{j-2}$. To write this term in the dual PBW basis we have to transfer z_{j-2} to the left of the product of the odd z_r . We get a factor $v^{a_{j-4}-a_{j-3}}$. Now all the monomials are in the right order. The generators z_{j+1} and y_{j-2} do not occur in the expansion of $\Delta_{i,j-3}^v$ in the dual PBW basis, but z_{j-2} may. In the summands where z_{j-2} occurs, we have increased the exponent from 1 to 2. So all coefficients in the dual PBW expansion of these summands have the form

$$g_{\mathbf{a}} v^{A_{i,j}} v^{1-a_{j-4}} v^{a_{j-4}-a_{j-3}} v^{\binom{1+a_{j-3}}{2}}.$$

Note that $A_{i,j} \leq -1$ and that $\binom{1+a_{j-3}}{2} - a_{j-3} = 0$ for $a_{j-3} \in \{0, 1\}$. Hence, all coefficients are in $v^{-1}\mathbb{Z}[v^{-1}]$. Now consider the summand coming from $z_{j-3}z_{j-1}$. The monomials are already in the right order. We may have increased the exponent of z_{j-3} from 1 to 2. It is easy to see that all coefficients are in $v^{-1}\mathbb{Z}[v^{-1}]$. This shows that D satisfies property (1).

To conclude $D = \Delta_{i,j}^v$, we have to verify property (2) of Theorem 4.11. We use parts (c) and (d) of the induction hypothesis for the pair $(i, j-1)$ to obtain the following equation. Note that if $j-i \leq 3$, then $(|y_{j-1}|, |e_{i,j-4}|) = 0$, and if $j-i \geq 4$, then $(|y_{j-1}|, |e_{i,j-4}|) = 1$, therefore

$$\begin{aligned} D &= \Delta_{i,j-1}^v y_j - v^{A_{i,j}} \Delta_{i,j-3}^v p_{j-2} z_{j+1} \\ &= v^{-1} y_j \Delta_{i,j-1}^v + v^{-(|y_{j-1}|, |e_{i,j-4}|)} (v^{-1} - v) \Delta_{i,j-3}^v p_{j-2} z_{j+1} \\ &\quad - v^{A_{i,j}} \Delta_{i,j-3}^v p_{j-2} z_{j+1} \\ &= v^{-1} y_j \Delta_{i,j-1}^v + v^{1-(|y_{j-1}|, |e_{i,j-4}|)} \Delta_{i,j-3}^v p_{j-2} z_{j+1}. \end{aligned}$$

It is easy to see that z_{j+1} commutes with p_{j-2} . Furthermore, it commutes with $\Delta_{i,j-3}^v$ since it commutes with every $E^*(\beta_k)$ in every summand in the dual PBW expansion of $\Delta_{i,j-3}^v$ according to Remark 4.8. Lemma 3.50 implies

$$\Delta_{i,j-3}^v p_{j-2} = v^{-(|y_{j-2}|, o_{i,j-3}) - (|z_{j-2}|, o_{i,j-3}) + (|z_{j-2}|, e_{i,j-4})} p_{j-2} \Delta_{i,j-3}^v.$$

The assumptions $j-i \geq 3$ and $j < n$ imply the equations

$$\begin{aligned} (|y_j|, o_{i,j-1}) &= (|y_j|, o_{i,j-3}) = 1, \\ (|z_{j-2}|, o_{i,j-3}) &= 1. \end{aligned}$$

The observation $(|z_{j-2}|, e_{i,j-4}) = 0$ for $j-i \leq 3$, and $(|z_{j-2}|, e_{i,j-4}) = 1$ for $j-i \geq 4$ yields $(|z_{j-2}|, e_{i,j-4}) = -A_{i,j} - 1$. It follows that

$$D = v^{-(|y_j|, o_{i,j-1})} y_j \Delta_{i,j-1}^v - v^{-A_{i,j} - (|y_j| + |y_{j-2}|, o_{i,j-3}) - (|y_{j-1}|, e_{i,j-4})} z_{j+1} p_{j-2} \Delta_{i,j-3}^v.$$

By Lemmas 4.6 and 4.7 the last equation is equivalent to property (2). Thus $A = \Delta_{i,j}^v$. Incidentally, we have verified part (a) for the pair (i, j) .

Now we prove that the new defined $\Delta_{i,j}^v$ satisfies the quantized cluster recursion (b) of Theorem 4.11 using the induction hypothesis for part (b):

$$\begin{aligned} \Delta_{i,j}^v z_j &= \Delta_{i,j-1}^v (p_j + v z_{j-1} z_{j+1}) - v^{A_{i,j}} \Delta_{i,j-3}^v p_{j-2} z_{j+1} z_j \\ &= \Delta_{i,j-1}^v p_j + \left(v \Delta_{i,j-1}^v z_{j-1} - v^{1+A_{i,j}} \Delta_{i,j-3}^v p_{j-2} z_j \right) z_{j+1} \\ &= \Delta_{i,j-1}^v p_j + \left(v \Delta_{i,j-1}^v z_{j-1} - v^{-(|y_{j-1}|, e_{i,j-2})} \Delta_{i,j-3}^v p_{j-2} z_j \right) z_{j+1} \\ &= \Delta_{i,j-1}^v p_j + v^{1-(|y_{j-1}|, e_{i,j-2})} \Delta_{i,j-3}^v p_{j-2} z_{j+1}. \end{aligned}$$

Now we prove that the new defined $\Delta_{i,j}^v$ satisfies property (c) of Theorem 4.11. By induction hypothesis we know that part (c) is true for the pair $(i, j-1)$. Note that $(|y_{j+1}|, |z_{j+1}|) = (|y_j|, |y_{j+1}|) = -\delta_{j,n-1}$. The following calculation verifies part (c)

of the theorem:

$$\begin{aligned}
y_{j+1}\Delta_{i,j}^v &= y_{j+1}\Delta_{i,j-1}^v y_j - v^{A_{i,j}} y_{j+1}\Delta_{i,j-3}^v p_{j-2} z_{j+1} \\
&= v^{-1}\Delta_{i,j-1}^v \left(v^{-(|y_j|, |y_{j+1}|)} y_j y_{j+1} + (v^{-1} - v) z_{j+2} z_{j-1} \right) \\
&\quad - v^{A_{i,j-1}} \Delta_{i,j-3}^v p_{j-2} \left(v^{-(|y_{j+1}|, |z_{j+1}|)} z_{j+1} y_{j+1} + (v^{-1} - v) z_j z_{j+2} \right) \\
&= v^{-(|y_{j+1}|, e_{i,j})} \Delta_{i,j}^v y_{j+1} \\
&\quad + v^{-1}(v^{-1} - v) \left(\Delta_{i,j-1}^v z_{j-1} - v^{A_{i,j}} \Delta_{i,j-3}^v p_{j-2} z_j \right) z_{j+2} \\
&= v^{-(|y_{j+1}|, e_{i,j})} \Delta_{i,j}^v y_{j+1} + (v^{-1} - v) v^{-1-(|y_{j-1}|, e_{i,j-2})} \Delta_{i,j-2}^v p_{j-1} z_{j+1}.
\end{aligned}$$

It remains to verify part (d). Note that in each monomial in the dual PBW expansion of $\Delta_{i,j-1}^v$ either the variable y_{j-1} or the variable z_j occurs (depending on whether a_{j-1} in Remark 4.8 is zero or one). The variable y_{j+2} commutes with all other terms. Thus, we see that $\Delta_{i,j-1}^v y_{j+2} = v^{-1} y_{j+2} \Delta_{i,j-1}^v$. It follows that

$$\begin{aligned}
\Delta_{i,j}^v y_{j+2} &= y_j \Delta_{i,j-1}^v y_{j+2} - v^{A_{i,j}} z_{j+1} p_{j-2} \Delta_{i,j-3}^v y_{j+2} \\
&= v^{-1} y_j y_{j+2} \Delta_{i,j-1}^v - v^{-1+A_{i,j}} y_{j+2} z_{j+1} p_{j-2} \Delta_{i,j-3}^v = v^{-1} y_{j+2} \Delta_{i,j}^v.
\end{aligned}$$

Now assume that j is odd. Put $D = y_j \Delta_{i,j-1}^v - v^{A_{i,j}} z_{j+1} p_{j-2} \Delta_{i,j-3}^v$. We prove that $D = \Delta_{i,j}^v$. By the same arguments as above D fulfills property (1) of the theorem. To conclude to $D = \Delta_{i,j}^v$, we have to verify property (2) of Theorem 4.11. We use parts (c) and (d) of the induction hypothesis for the pair $(i, j-1)$ to obtain the following equation:

$$\begin{aligned}
D &= y_j \Delta_{i,j-1}^v - v^{A_{i,j}} z_{j+1} p_{j-2} \Delta_{i,j-3}^v \\
&= v^{-(|y_j|, e_{i,j-1})} \Delta_{i,j-1}^v y_j + v^{-1-(|y_{j-2}|, e_{i,j-3})} (v^{-1} - v) \Delta_{i,j-3}^v p_{j-2} z_{j+1} \\
&\quad - v^{A_{i,j}} v^{-(|y_{j-2}|, e_{i,j-3}) - (|z_{j-2}|, e_{i,j-3}) + (|z_{j-2}|, o_{i,j-4})} \Delta_{i,j-3}^v p_{j-2} z_{j+1}.
\end{aligned}$$

Note that $(|z_{j-2}|, e_{i,j-3}) = 1$, and that $(|z_{j-2}|, o_{i,j-4}) = 0$ for $j - i \leq 3$ and $(|z_{j-2}|, o_{i,j-4}) = 1$ for $j - i \geq 4$. Hence, $A_{i,j} - (|z_{j-2}|, o_{i,j-4})(|z_{j-2}|, e_{i,j-3}) = -2$ yielding

$$D = v^{-(|y_j|, e_{i,j-1})} \Delta_{i,j-1}^v y_j - v^{-(|y_{j-2}|, e_{i,j-3})} \Delta_{i,j-3}^v p_{j-2} z_{j+1}.$$

The equation $-A_{i,j} - (|y_j|, e_{i,j-3}) - (|y_{j-1}|, o_{i,j-4})$ finishes the proof of the second equation of part (a). By Lemmas 4.6 and 4.7 the last equation is equivalent to property (2). Hence, $D = \Delta_{i,j}^v$.

Now we prove that the new defined $\Delta_{i,j}^v$ satisfies property (b) of Theorem 4.11. First of all assume that $j < n$. We obtain:

$$\begin{aligned}
\Delta_{i,j}^v z_j &= v^{-(|y_j|, e_{i,j-1})} \Delta_{i,j-1}^v (p_j + v^{-1} z_{j-1} z_{j+1}) \\
&\quad - v^{-(|y_{j-2}|, e_{i,j-3})} \Delta_{i,j-3}^v p_{j-2} z_{j+1} z_{j-1} \\
&= v^{-(|y_j|, e_{i,j-1})} \Delta_{i,j-1}^v p_j \\
&\quad + \left(v^{-1-(|y_j|, e_{i,j-1})} \Delta_{i,j-1}^v z_{j-1} - v^{-1-(|y_{j-2}|, e_{i,j-3})} \Delta_{i,j-3}^v p_{j-2} z_{j-1} \right) z_{j+1} \\
&= v^{-(|y_j|, e_{i,j-1})} \Delta_{i,j-1}^v p_j + v^{-1-(|y_j|, e_{i,j-1})} \Delta_{i,j-2}^v p_{j-1} z_{j+1}.
\end{aligned}$$

Here we used the fact that $(|y_j|, e_{i,j-1}) = 1$ to adopt the induction hypothesis. For $j = n$ we have $(|y_j|, e_{i,j-1}) = 0$, but this defect is compensated by the relation $z_{j-1}z_{j+1} = z_{j+1}z_{j-1}$ (due to $z_{j+1} = 1$) instead of $z_{j-1}z_{j+1} = v^{-1}z_{j+1}z_{j-1}$.

Now we prove that the new defined $\Delta_{i,j}^v$ satisfies property (c) of Theorem 4.11. By induction hypothesis we know that part (b) is true for the pair $(i, j-1)$. Note that $(|y_j|, e_{i,j-3}) = (|y_j|, e_{i,j-1})$ and that $1 = -A_{i,j} - (|y_{j-1}|, o_{i,j-4})$. We also use the fact that z_{j+1} commutes with $\Delta_{i,j-3}^v$ since it commutes with every factor of every monomial in the PBW expansion of $\Delta_{i,j-3}^v$. Thus, we see that $y_{j+1}\Delta_{i,j}^v$ is equal to:

$$\begin{aligned}
& v^{-(|y_j|, e_{i,j-1})} y_{j+1} \Delta_{i,j-1}^v y_j \\
& \quad - v^{-A_{i,j} - (|y_j| + |y_{j-2}|, e_{i,j-3}) - (|y_{j-1}|, o_{i,j-4})} y_{j+1} \Delta_{i,j-3}^v p_{j-2} z_{j+1} \\
& = v^{-(|y_j|, e_{i,j-3})+1} \left[\Delta_{i,j-1}^v \left(y_j y_{j+1} + (v - v^{-1}) z_{j-1} z_{j+2} \right) \right. \\
& \quad \left. - v^{1 - (|y_{j-2}|, e_{i,j-3})} \Delta_{i,j-3}^v p_{j-2} \left(z_{j+1} y_{j+1} + (v - v^{-1}) z_j z_{j+2} \right) \right] \\
& = v \Delta_{i,j}^v y_{j+1} + v^{-(|y_j|, e_{i,j-3})+1} (v - v^{-1}) \left[\Delta_{i,j-1}^v z_{j-1} \right. \\
& \quad \left. - v^{1 - (|y_{j-2}|, e_{i,j-3})} \Delta_{i,j-3}^v p_{j-2} z_j \right] z_{j+2} \\
& = v \Delta_{i,j}^v y_{j+1} + v^{-(|y_j|, e_{i,j-3})+1} (v - v^{-1}) \Delta_{i,j-2}^v p_{j-1} z_{j+2}.
\end{aligned}$$

Part (d) is proved similarly as in the case where j is even. \square

Remark 4.12. By symmetry there is also a recursion for every $\Delta_{i,j}^v$ (with $j - i \geq 2$) in terms of various $\Delta_{i',j}^v$ with $i' > i$.

Quantum cluster algebras were introduced by Berenstein-Zelevinsky [10]. We ready to conclude with our main theorem which establishes a quantum cluster algebra structure on the $\mathbb{Z}[v^{\pm \frac{1}{2}}]$ -algebra $\mathcal{A}_v(w) = \bigoplus_{\mathbf{a} \in \mathbb{N}^{2n}} \mathbb{Z}[v^{\pm \frac{1}{2}}] E[\mathbf{a}]^*$.

Theorem 4.13. The $\mathbb{Z}[v^{\pm \frac{1}{2}}]$ -algebra $\mathcal{A}_v(w) = \bigoplus_{\mathbf{a} \in \mathbb{N}^{2n}} \mathbb{Z}[v^{\pm \frac{1}{2}}] E[\mathbf{a}]^*$ is a quantum cluster algebra of type A_n . Every mutable quantum cluster variables is up to a power of v an element in the dual canonical basis \mathcal{B}^* . The following properties hold:

- (a) The quantum cluster variables $v^{\frac{1}{2}} z_i$ for $1 \leq i \leq n$ together with the frozen quantum cluster variables are $v^{\frac{1}{4}(|y_i| + |z_i|, |y_i| + |z_i|)} p_i$ for $1 \leq i \leq n$ form an initial cluster of $\mathcal{A}_v(w)$ whose B -matrix is encoded in the quiver from Figure 12.
- (b) The remaining quantum cluster variables are $v^{\frac{1}{4}(s_{i,j}, s_{i,j})} \Delta_{i,j}^v$ for $1 \leq i \leq j \leq n$.

Note that $\frac{1}{2}(s_{i,j}, s_{i,j}) = j - i + 1 + (e_{i,j-1}, o_{i,j-1}) \in \mathbb{Z}$.

Proof. We quantize the proof of Lemma 2.5. We construct a seed of a quantum cluster algebra similar to the base seed in Figure (12). The cluster variables Z_i, P_i (for $1 \leq i \leq n$) are replaced by the quantum cluster variables $v^{\frac{1}{2}} z_i, v^{\frac{1}{4}(|y_i| + |z_i|, |y_i| + |z_i|)} p_i$ (for $1 \leq i \leq n$). Notably, every pair of quantum cluster variables in the base seed forms a quantum torus, i.e., it satisfies a v -commutativity relation. (The v -commutativity relations among the z_i follow from the straightening relations; the v -commutativity relations between the P_i and the z_i follow from Lemma 3.50; the v -commutativity relations among the P_i can be checked using the straightening relations.) These relations

are strict commutativity relations except for the following v -commutativity relations:

$$\begin{aligned}
z_i z_j &= v^{-(|z_i|, |z_j|)} z_j z_i, & i \text{ even}, j \text{ odd}, \\
z_i p_j &= v^{(|z_i|, |y_j|)} p_j z_i, & i, j \text{ even}, \\
z_i p_j &= v^{-(|z_i|, |y_j|)} p_j z_i, & i, j \text{ odd}, \\
p_i p_j &= v^{-(|z_i|, |z_j|) + (|z_i|, |y_j|) + (|y_i|, |z_j|) + (|y_i|, |y_j|)} p_j z_i, & i \text{ even}, j \text{ odd}. \\
p_i p_j &= v^{-(|z_i|, |y_j|) + (|y_i|, |z_j|)} p_j z_i, & i, j \text{ even}, \\
p_i p_j &= v^{(|z_i|, |y_j|) - (|y_i|, |z_j|)} p_j z_i, & i, j \text{ odd}.
\end{aligned}$$

Now it is easy to see that the B -matrix induced from the quiver in Figure 12 and the Λ -matrix induced from the v -commutativity relations form a *compatible pair* in the sense of Berenstein-Zelevinsky [10, Definition 3.1]. Hence, the base seed is a valid initial quantum cluster.

Now fix an integer i such that $1 \leq i \leq n$. Beginning with the base we perform mutations at vertices $i, i+1, \dots, j$, consecutively, as in the proof of Lemma 2.5. We prove by induction on j that the new quantum cluster variable X that occurs after the sequence of mutations from above is equal to $\Delta_{i,j}^v$. The case $i = j$ is trivial. Note that part (c) of Theorem 4.11 makes also sense for $j = i+1$. We distinguish two cases. First of all, assume that j is even. By Berenstein-Zelevinsky [10, Proposition 4.9] we have

$$X = M' + M''$$

where definition of M' and M'' involves v -commutativity relations among the quantum cluster variables in the previous seed. To describe the variable M' we compute $\Delta_{i,j-1}^v p_j z_j^{-1} = v^{-(|y_j|, o_{i,j-1})} z_j^{-1} p_j \Delta_{i,j-1}^v$. Therefore, the summand M' is given by the following equation:

$$\begin{aligned}
M' &= v^{\frac{1}{2}(|y_j|, o_{i,j-1})} \cdot v^{\frac{1}{2}(j-i) + \frac{1}{2}(e_{i,j-2}, o_{i,j-1})} \Delta_{i,j-1}^v \cdot v p_j \cdot v^{-\frac{1}{2}} z_j^{-1} \\
&= v^{\frac{1}{2}(j-i+1) + \frac{1}{2}(e_{i,j}, o_{i,j-1})} \Delta_{i,j-1}^v p_j z_j^{-1}.
\end{aligned}$$

Furthermore, to describe M'' we obtain:

$$\begin{aligned}
\Delta_{i,j-2}^v p_{j-1} z_{j+1} z_j^{-1} &= v^{-(|z_j|, |z_{j+1}|) - (|z_j|, |z_{j-1}|) + (|z_j|, |y_{j-1}|) - (|z_{j-1}|, |y_{j-1}|)} \\
&\quad \cdot v^{(e_{i,j-2}, |y_{j-1}| + |z_{j-1}| - |z_j|)} \cdot v^{(o_{i,j-3}, -|y_{j-1}| - |z_{j-1}| + |z_j|)} \\
&\quad \cdot z_j^{-1} z_{j+1} p_{j-1} \Delta_{i,j-2}^v \\
&= v^{-2 + (e_{i,j-2}, |y_{j-1}|) - (o_{i,j-3}, |y_j|)} z_j^{-1} z_{j+1} p_{j-1} \Delta_{i,j-2}^v
\end{aligned}$$

Therefore, the summand M'' is given by the following equation:

$$\begin{aligned}
M'' &= v^{1 - \frac{1}{2}(e_{i,j-2}, |y_{j-1}|) + \frac{1}{2}(o_{i,j-3}, |y_j|)} \cdot v^{\frac{1}{2}(j-i-1) + \frac{1}{2}(e_{i,j-2}, o_{i,j-3})} \Delta_{i,j-2}^v \\
&\quad \cdot v^{1 - \frac{1}{2}(|y_j|, |y_{j-1}|)} \cdot v^{\frac{1}{2}} z_{j+1} \cdot v^{-\frac{1}{2}} z_j^{-1} \\
&= v^{1 - (|y_{j-1}|, e_{i,j-2})} v^{\frac{1}{2}(j-i+1) + \frac{1}{2}(e_{i,j}, o_{i,j-1})} \Delta_{i,j-2}^v p_{j-1} z_{j+1} z_j^{-1}.
\end{aligned}$$

A comparison with the formula in part (c) of Theorem 4.11 shows that $X = \Delta_{i,j}^v$ which completes the induction step.

The case with odd j is treated similarly. \square

Remark 4.14. Similarly, by adjusting the bilinear form we can equip the algebra $U_v^+(w')$ attached to Q' with the structure of a quantum cluster algebra.

Remark 4.15. Note that $\deg(\Delta_{i,j}) = \underline{\dim}(M_{i,j}) = s_{i,j}$ for all $1 < i < j < n$. Thus, the object $M_{i,j}$ from Remark 2.9 is the indecomposable rigid object in $\mathcal{A}(\mathcal{C}_M)$ corresponding to the cluster variable $\Delta_{i,j}$. The corresponding quantum cluster variable is the dual canonical basis element $\Delta_{i,j}^v$, scaled by a factor $v^{\frac{1}{4}(s_{i,j}, s_{i,j})}$. By [26, Lemma 3.12] the exponent can be interpreted as $\frac{1}{4}(s_{i,j}, s_{i,j}) = \frac{1}{2}\dim(\text{End}(M_{i,j}))$. The same relation is true for the other (mutable and frozen) quantum cluster variables.

Acknowledgements

This paper is a part of my Ph.D. studies at the University of Bonn supervised by Jan Schröer whom I would like to thank for many productive discussions and steady encouragement. I am also grateful to Bernard Leclerc for valuable explanations. I would like to thank an anonymous referee for various comments and remarks.

References

- [1] I. Assem, D. Simson and A. Skowronski, *Elements of the representation theory of associative algebras* (Cambridge University Press, Cambridge, 2006).
- [2] G. Bergman, ‘The diamond lemma for ring theory’, *Advances in Mathematics* 29, no. 2 (1978) 178-218.
- [3] A. Berenstein, S. Fomin and A. Zelevinsky, ‘Cluster Algebras III: Upper bounds and double Bruhat cells.’ *Duke Mathematical Journal* 126, no. 1 (2005) 1-52.
- [4] A. Buan, O. Iyama, I. Reiten and J. Scott, ‘Cluster structures for 2-Calabi-Yau categories and unipotent groups’, *Compositio Mathematica* 145, (2009) 1035-1079.
- [5] A. Buan, R. Marsh, M. Reineke, I. Reiten, and G. Todorov, ‘Tilting theory and cluster combinatorics’, *Advances in Mathematics* 204, no. 2 (2006) 572-618.
- [6] A. Buan, R. Marsh and I. Reiten, ‘Cluster-tilted algebras’, *Transactions of the AMS* 359, (2007) 323-332.
- [7] A. Buan, R. Marsh and I. Reiten, ‘Cluster mutation via quiver representations’, *Commentarii Mathematici Helvetici* 83, no. 1 (2008) 143-177.
- [8] A. Buan, R. Marsh, I. Reiten and G. Todorov, ‘Clusters and seeds in acyclic cluster algebras’, *Proceeding of the AMS* 135, no. 10 (2007) 3049-3060.
- [9] A. Berenstein and A. Zelevinsky, ‘String bases for quantum groups of type A_r ’, *I. M. Gel'fand Seminar, Advances in Soviet Mathematics* 16, Part 1 (Providence, 1993) 51-89.
- [10] A. Berenstein and A. Zelevinsky, ‘Quantum cluster algebras’, *Advances in Mathematics* 195, no. 2 (2005) 405-455.
- [11] P. Caldero, F. Chapoton and R. Schiffler, ‘Quivers with relations arising from clusters (A_n case)’, *Transactions of the AMS* 358, (2006) 1347-1364.

- [12] P. Caldero and B. Keller, ‘From triangulated categories to cluster algebras II’, *Annales Scientifiques de l’École Normale Supérieure* (4) 39, no. 6 (2006) 983-1009.
- [13] P. Caldero and B. Keller, ‘From triangulated categories to cluster algebras’, *Inventiones Mathematicae* 172, no. 1 (2008) 169-211.
- [14] W. Crawley-Boevey, ‘Lectures on Representations of Quivers’, *Lecture notes available at the author’s webpage* <http://www.amsta.leeds.ac.uk/~pmtwc/quivlecs.pdf>.
- [15] M. Ding and F. Xu, ‘Bases of Quantum cluster algebra of type $\tilde{A}_1^{(1)}$ ’, *Preprint arXiv:1004.2349* (2010).
- [16] V. Fock and A. Goncharov, ‘Moduli spaces of local systems and higher Teichmüller theory’, *Publications Mathématiques. Institut de Hautes Études Scientifiques* 103 (2006) 1-211.
- [17] S. Fomin and N. Reading, ‘Root systems and generalized associahedra’, *Lecture notes for IAS Park-City* (2008).
- [18] S. Fomin and A. Zelevinsky, ‘Cluster algebras I: Foundations’, *Journal of the American Mathematical Society* 15, no. 2 (2002) 497-529.
- [19] S. Fomin and A. Zelevinsky, ‘Cluster algebras II: Finite type classification’, *Inventiones Mathematicae* 154, no. 1 (2003) 63-121.
- [20] S. Fomin and A. Zelevinsky, ‘Y -systems and generalized associahedra’, *Annals of Mathematics* 158, no. 3 (2003) 977-1018.
- [21] S. Fomin and A. Zelevinsky, ‘Cluster algebras: notes for the CDM-03 conference’, *Current developments in mathematics* (International Press, Somerville, 2003) 1-34.
- [22] S. Fomin and A. Zelevinsky, ‘Cluster algebras IV: Coefficients’, *Compositio Mathematica* 143, no. 1 (2007) 112-164.
- [23] P. Gabriel, ‘Unzerlegbare Darstellungen. I’, *Manuscripta Mathematica* 6 (1972) 71-103.
- [24] J. Grabowski and S. Launois, ‘Quantum cluster algebra structures on quantum Grassmannians and their quantum Schubert cells: the finite type cases’, *Preprint arXiv:0912.4397* (2009).
- [25] C. Geiß, B. Leclerc and J. Schröer, ‘Semicanonical bases and preprojective algebras’, *Annales Scientifiques de l’École Normale Supérieure* (4) 38, no. 2 (2005) 193-253.
- [26] C. Geiß, B. Leclerc and J. Schröer, ‘Rigid modules over preprojective algebras’, *Inventiones Mathematicae* 165 (2006) 589-632.
- [27] C. Geiß, B. Leclerc and J. Schröer, ‘Semicanonical bases and preprojective algebras II: A multiplication formula’, *Compositio Mathematica* 143, (2007) 1313-1334.

- [28] C. Geiß, B. Leclerc and J. Schröer, ‘Cluster algebra structures and semicanonical bases for unipotent groups’, *Preprint arXiv:math/0703039* (2007).
- [29] C. Geiß, B. Leclerc and J. Schröer, ‘Kac-Moody groups and cluster algebras’, *Advances in Mathematics* 228 (2011), 329-433.
- [30] M. Gekhtman, M. Shapiro and A. Vainshtein, ‘Cluster algebras and Poisson geometry’, *Moscow Mathematical Journal* 3, no. 3 (2003) 899-934.
- [31] D. Happel. *Triangulated categories in the representation theory of finite-dimensional algebras* London Mathematical Society Lecture Notes Series, 119 (Cambridge University Press, Cambridge, 1988).
- [32] V. Kac and P. Cheung, *Quantum Calculus* (Springer-Verlag, New York, 2001).
- [33] M. Kashiwara, ‘On crystal bases of the q -analogue of universal enveloping algebras’, *Duke Mathematical Journal* 63, no. 2 (1991) 465-516.
- [34] B. Keller, ‘Quiver mutation in Java’, *Java applet available at the author’s webpage*.
- [35] B. Keller, ‘Cluster algebras, quiver representations and triangulated categories’, *Preprint arXiv:0807.1960* (2008).
- [36] Y. Kimura, ‘Quantum Unipotent Subgroup and dual canonical basis’, *Preprint arXiv:1010.4242* (2010).
- [37] M. Kontsevich and Y. Soibelman ‘Stability structures, motivic Donaldson-Thomas invariants and cluster transformations’, *arXiv:0811.2435* (2008).
- [38] P. Lampe, ‘A quantum cluster algebra of Kronecker type and the dual canonical basis’, *Preprint arXiv:1002.2762* (2010).
- [39] B. Leclerc, ‘Dual canonical bases, quantum shuffles and q -characters’, *Mathematische Zeitschrift* 246, no. 4 (2004) 691-732.
- [40] S. Levendorkii and Y. Soibelman, ‘Algebras of Functions on Compact Quantum Groups, Schubert Cells and Quantum Tori’, *Communications in Mathematical Physics* 139, no. 1 (1991) 141-170.
- [41] G. Lusztig, ‘Quivers, perverse sheaves, and quantized enveloping algebras’, *Journal of the American Mathematical Society* 4, no. 2 (1991) 365-421.
- [42] G. Lusztig, *Introduction to quantum groups* Progress in Mathematics 110 (Birkhäuser, Boston, 1993).
- [43] G. Musiker and J. Propp, ‘Combinatorial interpretations for rank-two cluster algebras of affine type’, *Electronic Journal of Combinatorics* 14, no. 1 (2007) R15.
- [44] M. Reineke, ‘Multiplicative Properties of Dual Canonical Bases of Quantum Groups’, *Journal of Algebra* 211, no. 1 (1999) 134-149.
- [45] I. Reiten, ‘Dynkin diagrams and the representation theory of algebras’, *Notices of the American Mathematical Society* 44, no. 5 (1997) 546-556.

- [46] C.M. Ringel, 'The preprojective algebra of a quiver', *Algebras and modules, II (Geiranger)* CMS Conference Proceedings, 24 (American Mathematical Society, Providence, 1998), 467-480.
- [47] D. Rohleder, *Cluster-Kategorien vom Typ \mathbb{A}_n* , Bonn: Diplomarbeit am Mathematischen Institut (2007).
- [48] M. Rosso, 'Quantum groups and quantum shuffles', *Inventiones Mathematicae* 133, no. 2 (1998) 399-416.
- [49] D. Rupel, 'On Quantum Analogue of The Caldero-Chapoton Formula', *Preprint arXiv:1003.2652* (2010).
- [50] The On-Line Encyclopedia of Integer Sequences, 'Sequence A000111', published electronically at <http://oeis.org> (2004).
- [51] R.P. Stanley, 'Enumerative Combinatorics. Volume I', *Cambridge University Press* 2nd edition (2000).