

On the general solution of the Heideman-Hogan family of recurrences

Andrew N.W. Hone* and Chloe Ward

August 13, 2018

Abstract

We consider a family of nonlinear rational recurrences of odd order which was introduced by Heideman and Hogan. All of these recurrences have the Laurent property, implying that for a particular choice of initial data (all initial values set to 1) they generate an integer sequence. For these particular sequences, Heideman and Hogan gave a direct proof of integrality by showing that the terms of the sequence also satisfy a linear recurrence relation with constant coefficients. Here we present an analogous result for the general solution of each of these recurrences.

1 Introduction

The theory of integer sequences generated by linear recurrences has a long history in number theory, and finds many applications in areas such as coding and cryptography [6], but the case of nonlinear recurrences is much less well studied. For some time there has been considerable interest in rational recurrence relations of the form

$$x_{n+N} x_n = F(x_{n+1}, \dots, x_{n+N-1}), \quad (1.1)$$

where F is a polynomial in $N - 1$ variables, which surprisingly generate integer sequences. Several quadratic recurrences of this kind were found by

*School of Mathematics, Statistics and Actuarial Science, University of Kent, Canterbury CT2 7NF, U.K. E-mail: A.N.W.Hone@kent.ac.uk

Somos, and this inspired others to find new examples, as described in the articles by Gale [11]. An important early observation was that if (1.1) has the Laurent property, meaning that it generates Laurent polynomials in the initial values with integer coefficients, i.e.

$$x_n \in \mathbb{Z}[x_0^{\pm 1}, x_1^{\pm 1}, \dots, x_{N-1}^{\pm 1}] \quad \forall n \in \mathbb{Z},$$

then an integer sequence is generated automatically by choosing the initial values to be $x_0 = x_1 = \dots = x_{N-1} = 1$. Subsequently, as an offshoot of their development of cluster algebras, Fomin and Zelevinsky introduced the Caterpillar Lemma [7], which is a useful tool for proving the Laurent property for many recurrences of the form (1.1). In the special case where F is a sum of two monomials, Fordy and Marsh explained how such recurrences arise from cluster algebras associated with quivers that are periodic under cluster mutations [8], while for more general F a range of examples were found recently by Alman et al. [1], who considered mutation periodicity in the broader context of Laurent phenomenon (LP) algebras [17].

In this paper we are concerned with a particular family of nonlinear recurrences of odd order $N = 2k + 1$, given by

$$x_{n+2k+1} x_n = x_{n+2k} x_{n+1} + a(x_{n+k} + x_{n+k+1}), \quad (1.2)$$

where a is a non-zero parameter. This family was introduced in the case $a = 1$ by Heideman and Hogan [12], who proved that the sequence generated by (1.2) with the initial values $x_0 = x_1 = \dots = x_{2k} = 1$ consists entirely of integers. (By rescaling $x_n \rightarrow ax_n$, the parameter a can always be removed, but it will be useful to retain it here for bookkeeping purposes.) One way to see the integrality of this particular sequence is to show that (1.2) has the Laurent property, which was noted in [12] and proved in [13]; more recently, the family (1.2) was rediscovered in a search for period 1 seeds in LP algebras - see Theorem 3.10 in [1]. However, Heideman and Hogan's original proof of integrality was based on the following result.

Theorem 1.1. *The terms of the sequence generated by the recurrence (1.2) with initial values $x_j = 1$ for $j = 0, 1, \dots, 2k$ and $a = 1$ satisfy the linear relation*

$$x_{n+6k} - (2k^2 + 8k + 4)(x_{n+4k} - x_{n+2k}) - x_n = 0 \quad (1.3)$$

for all $n \in \mathbb{Z}$.

The integrality result in [12] is proved by starting from $x_j = 1$, $0 \leq j \leq 2k$, then determining the explicit form of the next $4k$ values x_j , $2k + 1 \leq j \leq 6k$ obtained by iterating (1.2) for $0 \leq n \leq 4k - 1$, where the value of x_{6k} is used to verify that (1.3) holds for $n = 0$, and finally showing by induction that if (1.3) is assumed to hold for all $n \geq 0$ then (1.2) also holds for all $n \geq 4k$. This particular sequence is also symmetric under reversal, in the sense that

$$x_{-n} = x_{n+2k} \quad \forall n \in \mathbb{Z}. \quad (1.4)$$

In that case, the efficacy of this inductive approach can be seen directly from an operator identity connecting the linear operator in (1.3) with the nonlinear equation (1.2), which can be rewritten in the form $\xi_n = 0$, where

$$\xi_n := \begin{vmatrix} x_n & x_{n+2k} \\ x_{n+1} & x_{n+2k+1} \end{vmatrix} - a(x_{n+k} + x_{n+k+1}). \quad (1.5)$$

Lemma 1.2. *Let \mathcal{S} denote the shift operator, such that $\mathcal{S} x_n = x_{n+1}$ for all n , and let*

$$\mathcal{L} = \mathcal{S}^{6k} - K(\mathcal{S}^{4k} - \mathcal{S}^{2k}) - 1,$$

where K is some fixed constant. Then

$$\mathcal{L} \xi_n = \mathcal{M}_n \cdot \mathcal{L} x_n, \quad (1.6)$$

where \mathcal{M}_n is the linear operator

$$\mathcal{M}_n = x_{n+6k} \mathcal{S}^{2k+1} - x_{n+6k+1} \mathcal{S}^{2k} - x_{n+2k} \mathcal{S} + x_{n+2k+1} - a(\mathcal{S}^{k+1} + \mathcal{S}^k).$$

The main result of this paper is the analogue of Theorem 1.1 for the case of arbitrary initial data.

Theorem 1.3. *The iterates of the recurrence (1.2) satisfy the linear relation*

$$x_{n+6k} - K(x_{n+4k} - x_{n+2k}) - x_n = 0 \quad (1.7)$$

for all $n \in \mathbb{Z}$, where

$$K = P^{(0)} + a P^{(1)} + a^2 P^{(2)}, \quad (1.8)$$

with

$$P^{(0)} = 1 + \frac{x_0}{x_{2k}} + \frac{x_{2k}}{x_0},$$

$$P^{(1)} = \left(1 + \frac{x_{2k}}{x_0}\right) \sum_{j=1}^k \frac{x_{j-1} + x_j}{x_{j+k-1}x_{j+k}} + \left(1 + \frac{x_0}{x_{2k}}\right) \sum_{j=1}^k \frac{x_{j+k-1} + x_{j+k}}{x_{j-1}x_j},$$

$$P^{(2)} = \frac{1}{x_k x_{2k}} + \sum_{j=0}^{k-1} \frac{1}{x_j} \left(\frac{1}{x_{j+k}} + \frac{1}{x_{j+k+1}} \right) + \sum_{\ell=1}^{k-1} \sum_{m=1}^{\ell} \frac{(x_{\ell} + x_{\ell+1})(x_{k+m-1} + x_{k+m})}{x_{k+\ell}x_{k+\ell+1}x_{m-1}x_m}.$$

In principle, it is possible to prove the above result directly by using the identity (1.6) in Lemma 1.2 and adapting the argument from [12]. To do so one should take $2k + 1$ initial values x_0, \dots, x_{2k} for the nonlinear recurrence (1.2), require the $4k$ vanishing conditions $\xi_0 = \xi_1 = \dots = \xi_{4k-1} = 0$ which fix $6k$ initial values $x_0, x_1, \dots, x_{6k-1}$ for the linear equation $\mathcal{L}x_n = 0$ together with the value of K , determined as

$$K = \frac{x_{6k} - x_0}{x_{4k} - x_{2k}}, \quad (1.9)$$

and then further verify that $\xi_j = 0$ for a total of $6k$ adjacent values of j (including the range $0 \leq j \leq 4k - 1$ already assumed); this implies that the corresponding solution of the initial value problem for $\mathcal{L}\xi_n = 0$ is the zero solution $\xi_n = 0$ for all n . Heideman and Hogan made this argument effective with the use of computer algebra, which they used (for arbitrary k) to calculate explicit expressions for the values of x_{2k+1}, \dots, x_{6k} corresponding to $x_0 = x_1 = \dots = x_{2k} = 1$, and hence to verify the value $K = 2k^2 + 8k + 4$ in (1.3) and other necessary identities; they also implicitly used the fact that this special sequence has the reversal symmetry (1.4) (although this fact was not stated in [12]), which means that once $\xi_j = 0$ holds for $0 \leq j \leq 4k - 1$ it automatically holds for $-2k + 1 \leq j \leq -1$ as well, so it is enough to verify in addition that $\xi_{4k} = 0$ in order to show that $\xi_n = 0$ for all n by induction. However, this argument is much harder to apply to the case of generic initial data, because the corresponding sequence need not have the symmetry (1.4), so here we prefer to adopt a different approach. Nevertheless, we are able to exploit the fact that the recurrence (1.2) is itself reversible in the sense of [18], making the proof below much simpler than it might be otherwise.

The result (1.3) can be restated as saying that the recurrence (1.2) is linearizable, with the coefficient K appearing in the linear relation (1.7) being a conserved quantity (this terminology is explained in more detail in the next section); the general solution for the case $k = 1$ was already covered in [14]. There are many other examples of nonlinear recurrences

that are linearizable, which arise in diverse contexts ranging from cluster algebras associated with affine A-type Dynkin diagrams [8, 9, 10, 16], to frieze relations [2], Q-systems for characters in representation theory [4], and period 1 seeds in LP algebras [1, 15]. In all these examples, the key to obtaining the linear recurrences is provided by certain determinantal identities for discrete Wronskians. The fact that (1.2) can be rewritten as the vanishing of the expression (1.5) involving a 2×2 determinant permits a linear relation to be derived in a straightforward way, although it turns out that this approach is insufficient to obtain the precise form of (1.7).

In the next section we provide the proof of Theorem 1.3, and in section 3 we present various corollaries, before making some conclusions.

2 Proof of the main theorem

Before proceeding with the proof, we give some discussion of terminology, and describe properties of (1.2) that will be useful later on. First of all, note that iterating the nonlinear recurrence is equivalent to iterating a birational map in dimension $2k + 1$, namely

$$\varphi : (x_0, x_1, \dots, x_{2k}) \mapsto \left(x_1, x_2, \dots, \frac{x_1 x_{2k} + a(x_{k+1} + x_k)}{x_0} \right). \quad (2.1)$$

If we always use this map to iterate then it is useful to regard the terms in the sequence $(x_n)_{n \in \mathbb{Z}}$ as rational functions (in fact, Laurent polynomials, but we will not need this) in the initial coordinates x_0, x_1, \dots, x_{2k} and a , obtained by the pullback of φ (or its inverse) applied to these variables, so that

$$(\varphi^*)^n x_0 = x_n \quad \forall n \in \mathbb{Z},$$

with $(\varphi^{-1})^* = (\varphi^*)^{-1}$. We say that a non-constant function $F(x_0, x_1, \dots, x_{2k})$ is a conserved quantity, or first integral, for the map φ if it is invariant under pullback, i.e. $\varphi^* F = F \cdot \varphi = F$, and we say that it is a p -invariant if it is periodic with period p , i.e. $(\varphi^*)^p F = F$.

From Theorem 3.10 in [1], the map can be factored as $\varphi = \rho \cdot \mu$, where ρ is a cyclic permutation of the coordinates and μ is a mutation in an LP algebra, but more interesting for our purposes is the fact that it is a reversible map (it has the discrete analogue of time-reversal symmetry [18]), meaning that it is conjugate to its own inverse.

Lemma 2.1. *The map φ satisfies*

$$\varphi = \sigma \cdot \varphi^{-1} \cdot \sigma,$$

where the reversing symmetry σ is the involution

$$\sigma : (x_0, x_1, \dots, x_{2k}) \mapsto (x_{2k}, x_{2k-1}, \dots, x_0).$$

Reversibility means that the reversing symmetry σ can be extended to the level of the whole sequence (x_n) by pullback, so that it acts according to

$$\sigma^* x_n = x_{2k-n} \quad \forall n \in \mathbb{Z}. \quad (2.2)$$

In order to obtain linear relations for the terms of the sequence, it will be convenient to introduce the 3×3 discrete Wronskian matrix

$$\Psi_n := \begin{pmatrix} x_n & x_{n+2k} & x_{n+4k} \\ x_{n+1} & x_{n+2k+1} & x_{n+4k+1} \\ x_{n+2} & x_{n+2k+2} & x_{n+4k+2} \end{pmatrix}, \quad (2.3)$$

which has 2×2 minors of the form appearing in (1.5).

Proposition 2.2. *The determinant*

$$\delta_n := \det \Psi_n$$

is a k -invariant for the map φ .

Proof: Using Dodgson condensation [5] (also known as the Desnanot-Jacobi identity) to expand the 3×3 determinant in terms of its 2×2 connected minors yields

$$\begin{aligned} x_{n+2k+1} \delta_n &= \begin{vmatrix} \xi_n + a s_{n+k} & \xi_{n+2k} + a s_{n+3k} \\ \xi_{n+1} + a s_{n+k+1} & \xi_{n+2k+1} + a s_{n+3k+1} \end{vmatrix} \\ &= L_n + a^2 \begin{vmatrix} s_{n+k} & s_{n+3k} \\ s_{n+k+1} & s_{n+3k+1} \end{vmatrix}, \end{aligned}$$

where $s_n = x_n + x_{n+1}$, and the quantity L_n is a sum of homogeneous linear and quadratic terms in ξ_j for certain j . A direct calculation then shows that

$$x_{n+2k+1} x_{n+3k+1} (\delta_{n+k} - \delta_n) = x_{n+2k+1} L_{n+k} - x_{n+3k+1} L_n + \Delta_n,$$

where

$$\Delta_n = a^2 \left(s_{n+2k+1} \xi_{n+2k} + s_{n+2k} \xi_{n+2k+1} - s_{n+3k+1} \xi_{n+k} - s_{n+3k} \xi_{n+k+1} \right),$$

which clearly vanishes, along with L_n and L_{n+k} , if $\xi_j = 0$ for all j . Therefore $\delta_{n+k} = (\varphi^*)^k \delta_n = \delta_n$ for all n , as required. \square

Remark 2.3. Working in the ambient field of rational functions, that is $\mathbb{C}(x_0, x_1, \dots, x_{2k}, a)$, and using explicit expressions for the first few iterates (see below) it can be verified directly that δ_0 and all its shifts $\delta_1, \dots, \delta_{k-1}$ are non-zero rational functions (actually, Laurent polynomials), e.g.

$$\delta_0 = \delta_{-2k} = \begin{vmatrix} x_{-2k} & x_0 & x_{2k} \\ x_{-2k+1} & x_1 & x_{2k+1} \\ x_{-2k+2} & x_2 & x_{2k+2} \end{vmatrix}$$

can be calculated from the formulae in Lemma 2.7, and by periodicity none of the shifts $(\varphi^*)^n \delta_0$ can be identically zero (as a rational function).

Corollary 2.4. The determinant of the 4×4 discrete Wronskian matrix

$$\hat{\Psi}_n := \begin{pmatrix} x_n & x_{n+2k} & x_{n+4k} & x_{n+6k} \\ x_{n+1} & x_{n+2k+1} & x_{n+4k+1} & x_{n+6k+1} \\ x_{n+2} & x_{n+2k+2} & x_{n+4k+2} & x_{n+6k+2} \\ x_{n+3} & x_{n+2k+3} & x_{n+4k+3} & x_{n+6k+3} \end{pmatrix}$$

is zero.

Proof of Corollary: Using Dodgson condensation once more to expand the 4×4 determinant in terms of its 3×3 connected minors, which are shifts of the determinant of (2.3), gives

$$\det \hat{\psi}_n \begin{vmatrix} x_{n+2k+1} & x_{n+4k+1} \\ x_{n+2k+2} & x_{n+4k+2} \end{vmatrix} = \begin{vmatrix} \delta_n & \delta_{n+2k} \\ \delta_{n+1} & \delta_{n+2k+1} \end{vmatrix} = \begin{vmatrix} \delta_n & \delta_n \\ \delta_{n+1} & \delta_{n+1} \end{vmatrix} = 0$$

by Proposition 2.2. □

The above results are almost, but not quite, sufficient to produce the linear relation in Theorem 1.3.

Theorem 2.5. The iterates of the nonlinear recurrence (1.2) satisfy the linear recurrence

$$x_{n+6k} - K^{(1)} x_{n+4k} + K^{(2)} x_{n+2k} - x_n = 0, \quad (2.4)$$

where $K^{(1)}, K^{(2)}$ are conserved quantities with

$$K^{(2)} = \sigma^* K^{(1)}, \quad (2.5)$$

as well as the linear recurrence

$$x_{n+3} - \gamma_n x_{n+2} + \beta_n x_{n+1} - \alpha_n x_n = 0, \quad (2.6)$$

where α_n is a k -invariant and β_n, γ_n are $2k$ -invariants.

Proof: An element of the kernel of $\hat{\Psi}_n$ is given by a column vector $\mathbf{v}_n = (-K^{(3)}, K^{(2)}, -K^{(1)}, 1)^T$, where the last entry has been scaled to 1 (which is valid since δ_n is non-vanishing by Remark 2.3), and a priori the other entries $K^{(j)}$ depend on n . The first three rows of the equation $\hat{\Psi}_n \mathbf{v}_n = \mathbf{0}$ give a linear system for the $K^{(j)}$, and by Cramer's rule the solution is

$$K^{(1)} = \frac{1}{\delta_n} \begin{vmatrix} x_n & x_{n+2k} & x_{n+6k} \\ x_{n+1} & x_{n+2k+1} & x_{n+6k+1} \\ x_{n+2} & x_{n+2k+2} & x_{n+6k+2} \end{vmatrix}, \quad K^{(2)} = \frac{1}{\delta_n} \begin{vmatrix} x_n & x_{n+4k} & x_{n+6k} \\ x_{n+1} & x_{n+4k+1} & x_{n+6k+1} \\ x_{n+2} & x_{n+4k+2} & x_{n+6k+2} \end{vmatrix}$$

and $K^{(3)} = \delta_n^{-1} \delta_{n+2k} = 1$. The last three rows of $\hat{\Psi}_n \mathbf{v}_n = \mathbf{0}$ give the same linear system with all indices shifted by 1, implying that $K^{(1)}$ and $K^{(2)}$ are independent of n . Now applying σ^* to (2.4), replacing $n \rightarrow -n - 4k$ and adding the result back to the original relation produces

$$(\sigma^* K^{(2)} - K^{(1)}) x_{n+4k} + (K^{(2)} - \sigma^* K^{(1)}) x_{n+2k} = 0$$

for all n , hence (2.5) must hold. The same argument applied to the kernel of the transpose matrix $\hat{\Psi}_n^T$ yields the relation (2.6) where

$$\beta_n = \frac{1}{\delta_n} \begin{vmatrix} x_n & x_{n+2} & x_{n+3} \\ x_{n+2k} & x_{n+2k+2} & x_{n+2k+3} \\ x_{n+4k} & x_{n+4k+2} & x_{n+4k+3} \end{vmatrix}, \quad \gamma_n = \frac{1}{\delta_n} \begin{vmatrix} x_n & x_{n+1} & x_{n+3} \\ x_{n+2k} & x_{n+2k+1} & x_{n+2k+3} \\ x_{n+4k} & x_{n+4k+1} & x_{n+4k+3} \end{vmatrix}$$

are $2k$ -invariants and $\alpha_n = \delta_n^{-1} \delta_{n+1}$ is a k -invariant. \square

By considering the monodromy of the linear equation (2.6) with periodic coefficients, the coefficients in (2.4) for can be written in terms of $\alpha_j, \beta_j, \gamma_j$. In terms of the matrix Ψ_n , the system (2.6) implies that

$$\Psi_{n+1} = \mathbf{L}_n \Psi_n, \quad \mathbf{L}_n = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \alpha_n & -\beta_n & \gamma_n \end{pmatrix}, \quad \mathbf{L}_n^{-1} = \frac{1}{\alpha_n} \begin{pmatrix} \beta_n & -\gamma_n & 1 \\ \alpha_n & 0 & 0 \\ 0 & \alpha_n & 0 \end{pmatrix},$$

so that

$$\Psi_{n+2k} = \mathbf{M}_n \Psi_n, \quad \mathbf{M}_n = \mathbf{L}_{n+2k-1} \mathbf{L}_{n+2k-2} \cdots \mathbf{L}_n,$$

while on the other hand

$$\Psi_{n+2k} = \Psi_n \tilde{\mathbf{L}}, \quad \tilde{\mathbf{L}} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & -K^{(2)} \\ 0 & 1 & K^{(1)} \end{pmatrix}, \quad \tilde{\mathbf{L}}^{-1} = \begin{pmatrix} K^{(2)} & 1 & 0 \\ -K^{(1)} & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}.$$

Then noting that $\text{tr} \tilde{\mathbf{L}} = \text{tr} \mathbf{M}_n$ and $\text{tr} \tilde{\mathbf{L}}^{-1} = \text{tr} \mathbf{M}_n^{-1}$, together with the observation that $\prod_{j=1}^k \alpha_j = 1$, we have the following.

Proposition 2.6. *The conserved quantities $K^{(1)}$ and $K^{(2)}$ can be written as polynomials in $\alpha_j, \beta_j, \gamma_j$, given by*

$$K^{(1)} = \text{tr } \mathbf{L}_{n+2k-1} \mathbf{L}_{n+2k-2} \cdots \mathbf{L}_n, \quad K^{(2)} = \text{tr } \mathbf{L}_n^{-1} \mathbf{L}_{n+1}^{-1} \cdots \mathbf{L}_{n+2k-1}^{-1}.$$

According to the result we are aiming for, Theorem 1.3, we expect to find $K^{(1)} = K^{(2)} = K$, a Laurent polynomial in x_0, x_1, \dots, x_{2k} , which by (2.5) must be invariant under the reversal symmetry. However, none of the formulae for $K^{(1)}, K^{(2)}$ obtained so far make this coincidence manifest, and none of them immediately yield a Laurent polynomial. Indeed, the preceding result is somewhat mysterious, since direct calculations for small values of k reveal that $\alpha_j, \beta_j, \gamma_j$ are not Laurent polynomials themselves. In order to prove the main result, we calculate explicit formulae for $2k$ iterates on either side of the initial data, which allows us to obtain K as a quadratic polynomial in a , by using (1.9) with suitably shifted indices.

Lemma 2.7. *The first $2k$ terms obtained by iterating (1.2) forwards from the initial values x_0, x_1, \dots, x_{2k} are given by*

$$\begin{aligned} x_{2k+j} &= x_0^{-1} x_j x_{2k} + a F_{2k+j}^{(1)}, \\ x_{3k+j} &= x_0^{-1} x_{k+j} x_{2k} + a F_{3k+j}^{(1)} + a^2 F_{3k+j}^{(2)}, \quad 1 \leq j \leq k, \end{aligned}$$

where the coefficients of the linear and quadratic terms in a are specified by

$$\begin{aligned} F_{2k+j}^{(1)} &= x_j \sum_{\ell=1}^j (x_{\ell-1} x_\ell)^{-1} (x_{k+\ell-1} + x_{k+\ell}), \\ F_{3k+j}^{(1)} &= x_0^{-1} x_{k+j} x_{2k} \sum_{\ell=1}^j (x_{k+\ell-1} x_{k+\ell})^{-1} (x_{\ell-1} + x_\ell) + x_k^{-1} x_{k+j} F_{3k}^{(1)}, \\ F_{3k+j}^{(2)} &= x_{k+j} \sum_{\ell=1}^j (x_{k+\ell-1} x_{k+\ell})^{-1} \left(F_{2k+\ell-1}^{(1)} + F_{2k+\ell}^{(1)} \right), \end{aligned}$$

for the same range of the index j , with $F_{2k}^{(1)} = 0$. The first $2k$ terms obtained by iterating (1.2) backwards from the same initial values are

$$\begin{aligned} x_{-j} &= x_{2k}^{-1} x_{2k-j} x_0 + a F_{-j}^{(1)}, \\ x_{-k-j} &= x_{2k}^{-1} x_{k-j} x_0 + a F_{-k-j}^{(1)} + a^2 F_{-k-j}^{(2)}, \quad 1 \leq j \leq k, \end{aligned}$$

where, for the same range of j values,

$$F_{-j}^{(1)} = \sigma^* F_{2k+j}^{(1)}, \quad F_{-k-j}^{(1)} = \sigma^* F_{3k+j}^{(1)}, \quad F_{-k-j}^{(2)} = \sigma^* F_{3k+j}^{(2)}.$$

Proof: The first $2k+1$ iterations of (1.2), either forwards or backwards, only require multiplication and addition of previous terms, as well as division by one of x_0, x_1, \dots, x_{2k} , so for the division there is no need to consider any cancellations between numerator and denominator (which are required for the Laurent property to hold at subsequent steps). It is plain to see that the first k terms produced by iterating are linear in a , while the next k terms are quadratic. Expanding x_{2k+j} in a and substituting into (1.2) with $n = j - 1$ it is straightforward to obtain the leading order term recursively, while the coefficient of the linear term satisfies the recursion

$$x_j^{-1}F_{2k+j}^{(1)} - x_{j-1}^{-1}F_{2k+j-1}^{(1)} = (x_{j-1}x_j)^{-1}(x_{k+j-1} + x_{k+j}), \quad 1 \leq j \leq k,$$

which can be summed telescopically (with $F_{2k}^{(1)} = 0$) to obtain the above formula for $F_{2k+j}^{(1)}$. Similarly, expanding in a for the next k iterations, the leading order term is found immediately, while for the term linear in a the recursion is

$$x_{k+j}^{-1}F_{3k+j}^{(1)} - x_{k+j-1}^{-1}F_{3k+j-1}^{(1)} = (x_{k+j-1}x_{k+j}x_0)^{-1}x_{2k}(x_{j-1} + x_j), \quad 1 \leq j \leq k.$$

which immediately yields the above expression for $F_{3k+j}^{(1)}$; and for the quadratic term the formula for the coefficient is found by solving the recursion

$$x_{k+j}^{-1}F_{3k+j}^{(2)} - x_{k+j-1}^{-1}F_{3k+j-1}^{(2)} = (x_{k+j-1}x_{k+j})^{-1}\left(F_{2k+j-1}^{(1)} + F_{2k+j}^{(1)}\right), \quad 1 \leq j \leq k.$$

Similarly, iterating $2k$ steps backwards produces the images of the forward iterates under the action of the reversing map, so the formulae for x_{-j} and x_{-k-j} follow by direct application of σ^* . \square

Proposition 2.8. *If K is defined by*

$$K = \frac{x_{4k} - x_{-2k}}{x_{2k} - x_0}, \quad (2.7)$$

where x_{4k} and x_{-2k} are given as Laurent polynomials in x_0, x_1, \dots, x_{2k} and a according to Lemma 2.7, then it is given explicitly by (1.8).

Proof: The formula (1.8) is readily checked at each order in a . At leading order this is trivial, while at order a and a^2 the identities

$$(x_{2k} - x_0)P^{(j)} = F_{4k}^{(j)} - F_{-2k}^{(j)}$$

are seen to hold for $j = 1, 2$. \square

Theorem 2.9. *The quantity K in (1.8) is a first integral for the map φ .*

Proof: To verify that K is a conserved quantity, observe that

$$\varphi^* K = \frac{x_{4k+1} - x_{-2k+1}}{x_{2k+1} - x_1}$$

from (2.7), and this is equal to K if and only if $\mathcal{L}x_{-2k+1} = 0$, where \mathcal{L} is the operator in Lemma 1.2. Now $\mathcal{L}x_{-2k}$ vanishes by (2.7), so from the identity

$$\begin{aligned} x_{2k}\mathcal{L}x_{-2k+1} - x_{2k+1}\mathcal{L}x_{-2k} - \xi_{2k} + K\xi_0 &= a(x_{3k} + x_{3k+1}) - Ka(x_k + x_{k+1}) \\ &\quad + x_{2k+1}x_{-2k} - x_{2k}x_{-2k+1} \end{aligned}$$

we see that it is sufficient to check that the right-hand above is zero. Substitution of the explicit expressions from Lemma 2.7 yields a cubic polynomial in a ; the order zero term clearly vanishes, while at order one, two and three it is straightforward to verify the identities

$$\begin{aligned} (x_k + x_{k+1})\left(x_0^{-1}x_{2k} - P^{(0)}\right) + x_0^{-1}x_1x_{2k}F_{-2k}^{(1)} + x_{2k}^{-1}x_0^2F_{2k+1}^{(1)} - x_{2k}F_{-2k+1}^{(1)} &= 0, \\ F_{3k}^{(1)} + F_{3k+1}^{(1)} + x_0^{-1}x_1x_{2k}F_{-2k}^{(2)} + F_{2k+1}^{(1)}F_{-2k}^{(1)} - x_{2k}F_{-2k+1}^{(2)} - (x_k + x_{k+1})P^{(1)} &= 0, \\ F_{3k+1}^{(2)} + (x_k + x_{k+1})\left(x_0^{-1}F_{-2k}^{(2)} - P^{(2)}\right) &= 0. \end{aligned}$$

This shows that $\varphi^*K = K$, and completes the proof of Theorem 1.3. \square

3 General solution and other corollaries

The result of Theorem 1.3 has many implications for the recurrence (1.2). Theorem 1.1 is just a special case: one finds the value $K = 2k^2 + 8k + 4$ by substituting $a = x_0 = x_1 = \dots = x_{2k} = 1$ into (1.8). Also, note that we have not made use of the Laurent property in the proof, yet by the formulae in Lemma 2.7 we see that $x_{-2k}, x_{-2k+1}, \dots, x_{4k}$ provide $6k$ initial data for the linear recurrence $\mathcal{L}x_n = 0$, and these are all Laurent polynomials in x_0, x_1, \dots, x_{2k} with coefficients in $\mathbb{Z}[a]$, as is K given by (1.8), so we arrive at the following.

Corollary 3.1. *The nonlinear recurrence (1.2) has the Laurent property, i.e. $x_n \in \mathbb{Z}[x_0^{\pm 1}, x_1^{\pm 1}, \dots, x_{2k}^{\pm 1}, a] \forall n \in \mathbb{Z}$.*

In addition to the homogeneous linear recurrences in Theorem 2.5, various inhomogeneous linear recurrences now follow.

Corollary 3.2. *The iterates of (1.2) satisfy the linear recurrences*

$$x_{n+4k} - (K - 1)x_{n+2k} + x_n = \nu_n, \quad (3.1)$$

where ν_n is a $2k$ -invariant,

$$\sum_{j=0}^{2k-1} x_{n+4k+j} - (K - 1)x_{n+2k+j} + x_{n+j} = K', \quad (3.2)$$

where

$$K' = \nu_0 + \nu_1 + \dots + \nu_{2k-1} \quad (3.3)$$

is a conserved quantity, and

$$x_{n+2} + \eta_n x_{n+1} + \zeta_n x_n = \epsilon_n, \quad (3.4)$$

where ϵ_n, ζ_n and η_n are all $2k$ -invariants.

Proof of Corollary: The operator \mathcal{L} can be factorized as

$$\mathcal{L} = (\mathcal{S}^{2k} - 1) \left(\mathcal{S}^{4k} - (K - 1)\mathcal{S}^{2k} + 1 \right) = (\mathcal{S} - 1) \left(\sum_{j=0}^{2k-1} \mathcal{S}^j \right) \left(\mathcal{S}^{4k} - (K - 1)\mathcal{S}^{2k} + 1 \right),$$

which means that (1.7) can be “integrated” in two different ways to yield (3.1) and (3.2). By shifting and summing $2k$ copies of (3.1), the conserved quantity K' is given as a symmetric function of the $2k$ independent shifts of ν_n according to (3.3).

The equation (1.7) also implies that, for all n , the vector $(1, -K, K, -1)^T$ belongs to the kernel of the matrix

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ x_n & x_{n+2k} & x_{n+4k} & x_{n+6k} \\ x_{n+1} & x_{n+2k+1} & x_{n+4k+1} & x_{n+6k+1} \\ x_{n+2} & x_{n+2k+2} & x_{n+4k+2} & x_{n+6k+2} \end{pmatrix}$$

so by writing a vector in the kernel of its transpose as the row vector $(\epsilon_n, -\zeta_n, \eta_n, -1)$, the relation (3.4) follows, and the same argument as in the proof of Theorem 2.5 shows that $\epsilon_n, \zeta_n, \eta_n$ are invariant under shifting $n \rightarrow n + 2k$. \square

The fact that the iterates of (1.2) satisfy a linear relation with constant coefficients means that they can be written explicitly in terms of the roots of the associated characteristic polynomial. Due to the particular form of (1.7), the general solution can also be written using either trigonometric functions or Chebyshev polynomials (with the latter form of the solution for $k = 1$ being included in the results of [14]). In order to do this, we introduce quantities θ and t such that

$$t = \frac{K-1}{2} = \cos \Theta, \quad \Theta = 2k\theta,$$

and recall that the Chebyshev polynomials of the first and second kinds are defined by

$$T_n(t) = \cos(n\Theta), \quad U_{n-1}(t) = \frac{\sin(n\Theta)}{\sin \Theta}$$

respectively, so that $T_0 = U_0 = 1$, $T_1 = T_{-1} = t$, $U_1 = 2t$, $U_{-1} = 0$.

Corollary 3.3. *The general solution of (1.2) can be written in the form*

$$x_n = a_n + b_n \cos(n\theta + \phi_n),$$

where a_n, b_n, ϕ_n are all periodic in n with periodic $2k$, or as

$$x_n = q_j + r_j T_m(t) + s_j U_m(t), \quad m = \left\lfloor \frac{n}{2k} \right\rfloor,$$

where $j = n \bmod 2k$ and for $j = 0, 1, \dots, 2k-1$ the coefficients are

$$\begin{pmatrix} q_j \\ r_j \\ s_j \end{pmatrix} = \frac{1}{2t(1-t)} \begin{pmatrix} t & -2t^2 & t \\ -1 & 2t & 1-2t \\ 1-t & 0 & t-1 \end{pmatrix} \begin{pmatrix} x_{2k+j} \\ x_j \\ x_{-2k+j} \end{pmatrix}.$$

4 Conclusions

We have proved that all sequences generated by the nonlinear recurrence (1.2) satisfy a linear relation, which was left as an open problem in [12]. A key feature in the proof was to use the reversibility property in Lemma 2.1. In fact, all the cluster maps obtained from period 1 quivers in [8], and many of the recurrences considered in [1], are also reversible with the same sort of reversing symmetry, which means that the above approach can be applied

in those cases too, and may prove useful for finding explicit formulae for conserved quantities (when they exist).

One question that remains open is whether there is any natural Poisson structure which is preserved by the map (2.1), since Poisson structures (or presymplectic structures) arise naturally in the context of cluster algebras, but whether there is something similar for LP algebras in general is an open question. In fact we expect that there is a Poisson bracket (albeit a rather degenerate one, of rank two) for a combination of two reasons: first, the map should have many conserved quantities in addition to K and K' given by (3.3), since any symmetric function of the shifts of a $2k$ -invariant is conserved; and second, φ has the logarithmic volume form

$$\omega = \frac{1}{x_0 x_1 \dots x_{2k}} dx_0 \wedge dx_1 \wedge \dots \wedge dx_{2k}$$

which is anti-invariant, in the sense that $\varphi^* \omega = -\omega$; so if there are at least $2k - 2$ independent conserved quantities, then the corresponding co-volume can be contracted with their differentials to construct a Poisson bracket, by one of the results in [3].

Acknowledgments. Some of these results first appeared in the PhD thesis [19], which was supported by EPSRC studentship EP/P50421X/1. ANWH is supported by EPSRC fellowship EP/M004333/1.

References

- [1] J. Alman, C. Cuenca and J. Huang, Laurent phenomenon sequences, *J. Algebr. Comb.* **43** (2016) 589–633.
- [2] I. Assem and C. Reutenauer and D. Smith, Friezes, *Adv. Math.* **225** (2010) 3134–3165.
- [3] G.B. Byrnes, F.Haggar and G.R.W. Quispel, Sufficient conditions for dynamical systems to have pre-symplectic or pre-implectic structures, *Physica A* **272** (1999) 99–129.
- [4] P. Di Francesco and R. Kedem, Q-Systems, Heaps, Paths and Cluster Positivity, *Comm. Math. Phys.* **293** (2010) 727–802.
- [5] C.L. Dodgson, Condensation of determinants, *Proc. R. Soc. Lond.* **15** (1866) 150–155.

- [6] G. Everest, A. van der Poorten, I. Shparlinski and T. Ward, *Recurrence Sequences*, AMS Mathematical Surveys and Monographs, vol. **104**, Amer. Math. Soc., Providence, RI, 2003.
- [7] S. Fomin and A. Zelevinsky, The Laurent Phenomenon, *Advances in Applied Mathematics* **28** (2002) 119–144.
- [8] A.P. Fordy and R.J. Marsh, Cluster Mutation-Periodic Quivers and Associated Laurent Sequences, *J. Algebr. Comb.* **34** (2009) 19–66.
- [9] A.P. Fordy, Mutation-Periodic Quivers, Integrable Maps and Associated Poisson Algebras, *Philos. Trans. R. Soc. Lond. Ser. A*, **369**, (2010) 1264–1279.
- [10] A.P. Fordy and A.N.W. Hone, Discrete integrable systems and Poisson algebras from cluster maps, *Comm. Math. Phys.* **325** (2014) 527–584.
- [11] D. Gale, The strange and surprising saga of the Somos sequences, *Mathematical Intelligencer* **13** (1) (1991) 40–42; Somos sequence update, *Mathematical Intelligencer* **13** (4) (1991) 49–50.
- [12] P. Heideman and E. Hogan, A New Family of Somos-like Recurrences, *Elec. J. Comb.* **15** (2008) #R54, 8pp.
- [13] E. Hogan, Experimental mathematics applied to the study of non-linear recurrences, PhD thesis, Rutgers University, 2011.
- [14] A.N.W. Hone, Nonlinear recurrence sequences and Laurent polynomials *Number Theory and Polynomials* (ed. J.McKee and C. Smyth), *LMS Lecture Note Series* **352** (2008) 188–210.
- [15] A.N.W. Hone and C. Ward, A family of linearizable recurrences with the Laurent property, *Bull. Lond. Math. Soc.* **46** (2014) 503–516.
- [16] B. Keller and S. Scherotzke, Linear recurrence relations for cluster variables of affine quivers, *Adv. Math.* **228** (2011) 1842–1862
- [17] T. Lam and P. Pylyavskyy, Laurent phenomenon algebras, *Cambridge Journal of Mathematics* **4** (2016) 121–162.
- [18] J.S.W. Lamb and J.A.G. Roberts, Time-reversal symmetry in dynamical systems: a survey, *Physica D* **112** (1998) 1–39.
- [19] C. Ward, Discrete integrability and nonlinear recurrences with the Laurent property, PhD thesis, University of Kent, 2013.