

# Kent Academic Repository

## Full text document (pdf)

### Citation for published version

Mansfield, Elizabeth L. and Rojo-Echeburua, Ana (2019) Moving Frames and Noether's Finite Difference Conservation Laws II. Transactions of Mathematics and its applications . (In press)

### DOI

### Link to record in KAR

<https://kar.kent.ac.uk/74250/>

### Document Version

Author's Accepted Manuscript

#### Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

#### Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

#### Enquiries

For any further enquiries regarding the licence status of this document, please contact:

[researchsupport@kent.ac.uk](mailto:researchsupport@kent.ac.uk)

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

## Moving Frames and Noether's Finite Difference Conservation Laws II.

E. L. MANSFIELD<sup>†</sup>, A. ROJO-ECHEBURÚA<sup>‡</sup>

SMSAS, University of Kent, Canterbury, CT2 7FS, UK

[Received on ; revised on ]

In this second part of the paper, we consider finite difference Lagrangians which are invariant under linear and projective actions of  $SL(2)$ , and the linear equi-affine action which preserves area in the plane.

We first find the generating invariants, and then use the results of the first part of the paper to write the Euler–Lagrange difference equations and Noether's difference conservation laws for any invariant Lagrangian, in terms of the invariants and a difference moving frame. We then give the details of the final integration step, assuming the Euler Lagrange equations have been solved for the invariants. This last step relies on understanding the Adjoint action of the Lie group on its Lie algebra. We also use methods to integrate Lie group invariant difference equations developed in Part I.

Effectively, for all three actions, we show that solutions to the Euler–Lagrange equations, in terms of the original dependent variables, share a common structure for the whole set of Lagrangians invariant under each given group action, once the invariants are known as functions on the lattice.

*Keywords:* Noether's Theorem, Finite Difference, Discrete Moving Frames

### Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>The linear action of <math>SL(2)</math> in the plane</b>	<b>6</b>
2.1	The Adjoint action	6
2.2	The discrete frame, the generating invariants and their syzygies	6
2.2.1	The generating discrete invariants	6
2.2.2	The generating differential invariants	7
2.3	The Euler–Lagrange equations and conservation laws	8
2.4	The general solution	10
<b>3</b>	<b>The <math>SA(2) = SL(2) \ltimes \mathbb{R}^2</math> linear action</b>	<b>12</b>
3.1	The Adjoint action	13
3.2	The discrete frame, the generating invariants and their syzygies	13
3.3	The Euler–Lagrange equations and the conservation laws.	15
3.4	The general solution	17
<b>4</b>	<b>The <math>SL(2)</math> projective action</b>	<b>18</b>
4.1	The Adjoint action	18

<sup>†</sup>Corresponding author. Email: e.l.mansfield@kent.ac.uk

<sup>‡</sup>Email: arer2@kent.ac.uk

4.2 The discrete frame, the generating invariants and their syzygies . . . . . 19  
 4.3 The generating discrete invariants . . . . . 19  
 4.4 The generating differential invariants . . . . . 20  
 4.5 The Euler–Lagrange equations and the conservation laws . . . . . 21  
 4.6 The general solution . . . . . 23

**5 Conclusions** **24**

**1. Introduction**

This is a continuation of *Moving Frames and Noether’s Finite Difference Conservation Laws I* Mansfield, Rojo-Echeburúa, Hydon & Peng (2019) where now we consider Lie group actions of the special linear group  $SL(2)$  and  $SL(2) \times \mathbb{R}^2$ . The Lie group  $SL(2)$  is the set of  $2 \times 2$  real (or complex) matrices with determinant equal to unity. Its typical element is written as

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad ad - bc = 1. \tag{1.1}$$

Its Lie algebra, the set of  $2 \times 2$  real (or complex) matrices with zero trace, is denoted  $\mathfrak{sl}(2)$ . Part I of this paper developed all the necessary theory and considered simpler solvable groups; here we show some additional techniques needed for semi-simple groups. Smooth variational problems with an  $SL(2)$  and  $SL(2) \times \mathbb{R}^2$  symmetry were considered using moving frame techniques in Gonçalves & Mansfield (2012), Gonçalves & Mansfield (2016) and Mansfield (2010). Here we show the finite difference analogue for these variational problems.

We first recall the notation, definitions and the main Theorems developed in the first part of this paper (Mansfield, Rojo-Echeburúa, Hydon & Peng (2019)), which we will need; citations to the literature on standard notions from the discrete calculus of variations and on moving frames can be found in the Introduction to Part 1.

We consider the dependent variables to take values in  $U \subset \mathbb{R}^q$  with coordinates  $\mathbf{u} = (u^1, \dots, u^q)$ . We use the notation  $u_j^\alpha = u^\alpha(n + j)$  for  $\alpha = 1, \dots, q$  and  $n, j \in \mathbb{Z}$ . The (forward) shift operator  $S$  acts on functions of  $n$  as follows:

$$S : n \mapsto n + 1, \quad S : f(n) \mapsto f(n + 1),$$

for all functions  $f$  whose domain includes  $n$  and  $n + 1$ . In particular,

$$S : u_j^\alpha \mapsto u_{j+1}^\alpha$$

on any domain where both of these quantities are defined. The forward difference operator is  $S - \text{id}$ , where  $\text{id}$  is the identity operator:

$$\text{id} : n \mapsto n, \quad \text{id} : f(n) \mapsto f(n), \quad \text{id} : u_j^\alpha \mapsto u_j^\alpha.$$

The shift  $S$  is an operator on  $P_n^{(-\infty, \infty)}(U)$  where  $P_n^{(J_0, J)}(U) \simeq U \times \dots \times U$  ( $J - J_0 + 1$  copies) with coordinates  $z = (\mathbf{u}_{J_0}, \dots, \mathbf{u}_J)$ , where  $J_0 \leq 0$  and  $J \geq 0$ . We denote the  $j^{\text{th}}$  power of  $S$  by  $S_j$ , so that  $\mathbf{u}_j = S_j \mathbf{u}_0$  for each  $j \in \mathbb{Z}$ .

We will consider actions that are assumed to be free and regular on a manifold  $M$ . Therefore, there exists a cross section  $\mathcal{K} \subset M$  that is transverse to the orbits  $\mathcal{O}(z)$  and, for each  $z \in M$ , the set  $\mathcal{K} \cap \mathcal{O}(z)$  has just one element, the projection of  $z$  onto  $\mathcal{K}$ . Using the cross-section  $\mathcal{K}$ , a moving frame for the group action on a neighbourhood  $\mathcal{U} \subset M$  of  $z$  can be defined as follows.

DEFINITION 1.1 (Moving Frame) Given a smooth Lie group action  $G \times M \rightarrow M$ , a moving frame is an equivariant map  $\rho : \mathcal{U} \subset M \rightarrow G$ . Here  $\mathcal{U}$  is called the domain of the frame.

A left equivariant map satisfies  $\rho(g \cdot z) = g\rho(z)$ , and a right equivariant map satisfies  $\rho(g \cdot z) = \rho(z)g^{-1}$ . In order to find the frame, let the cross-section  $\mathcal{K}$  be given by a system of equations  $\psi_r(z) = 0$ , for  $r = 1, \dots, R$ , where  $R$  is the dimension of the group  $G$ . One then solves the so-called normalization equations,

$$\psi_r(g \cdot z) = 0, \quad r = 1, \dots, R, \quad (1.2)$$

for  $g$  as a function of  $z$ . The solution is the group element  $g = \rho(z)$  that maps  $z$  to its projection on  $\mathcal{K}$ . In other words, the frame  $\rho$  satisfies

$$\psi_r(\rho(z) \cdot z) = 0, \quad r = 1, \dots, R.$$

LEMMA 1.1 (Normalized Invariants) Given a left or right action  $G \times M \rightarrow M$  and a right frame  $\rho$ , then  $\iota(z) = \rho(z) \cdot z$ , for  $z$  in the domain of the frame  $\rho$ , is invariant under the group action.

DEFINITION 1.2 The normalized invariants are the components of  $\iota(z)$ .

THEOREM 1.3 (Replacement Rule) If  $F(z)$  is an invariant of the given action  $G \times M \rightarrow M$  for a right moving frame  $\rho$  on  $M$ , then  $F(z) = F(\iota(z))$ .

DEFINITION 1.4 (Invariantization Operator) Given a right moving frame  $\rho$ , the map  $z \mapsto \iota(z) = \rho(z) \cdot z$  is called the *invariantization operator*. This operator extends to functions as  $f(z) \mapsto f(\iota(z))$ , and  $f(\iota(z))$  is called the *invariantization* of  $f$ .

If  $z$  has components  $z^\alpha$ , let  $\iota(z^\alpha)$  denote the  $\alpha^{\text{th}}$  component of  $\iota(z)$ .

A discrete moving frame is a sequence of moving frames  $(\rho_k)$ ,  $k = 1, \dots, N$  with a nontrivial intersection of domains which, locally, are uniquely determined by the cross-section  $\mathcal{K} = (\mathcal{K}_1, \dots, \mathcal{K}_N)$  to the group orbit through  $z \in M^N$ , where  $M^N$  is the Cartesian product manifold. For a right discrete frame, we define the invariants

$$I_{k,j} := \rho_k(z) \cdot z_j. \quad (1.3)$$

Suppose that  $M$  is  $q$ -dimensional. Therefore  $z_j$  has components  $z_j^1, \dots, z_j^q$ , and the  $q$  components of  $I_{k,j}$  are the invariants

$$I_{k,j}^\alpha := \rho_k(z) \cdot z_j^\alpha, \quad \alpha = 1, \dots, q. \quad (1.4)$$

We will denote the invariantization operator with respect to the frame  $\rho_k(z)$  by  $\iota_k$ , so that

$$I_{k,j} = \iota_k(z_j), \quad I_{k,j}^\alpha = \iota_k(z_j^\alpha).$$

DEFINITION 1.5 (Discrete Maurer–Cartan invariants) Given a right discrete moving frame  $\rho$ , the *right discrete Maurer–Cartan group elements* are

$$K_k = \rho_{k+1}\rho_k^{-1} \quad (1.5)$$

where defined.

We call the components of the Maurer–Cartan elements the *Maurer–Cartan invariants*.

A difference moving frame is a natural discrete moving frame that is adapted to difference equations by prolongation conditions.

**THEOREM 1.6** Given a right difference moving frame  $\rho$ , the set of all invariants is generated by the set of components of  $K_0 = \rho_1 \rho_0^{-1}$  and  $I_{0,0} = \rho_0(z) \cdot z_0$ .

As  $K_0$  is invariant, by (1.3) we have that

$$K_0 = \iota_0(\rho_1), \quad (1.6)$$

where  $\iota_0$  denotes invariantization with respect to the frame  $\rho_0$ .

Given any smooth path  $t \mapsto z(t)$  in the space  $\mathcal{M} = M^N$ , consider the induced group action on the path and its tangent. We extend the group action to the dummy variable  $t$  trivially, so that  $t$  is invariant. The action is extended to the first-order jet space of  $\mathcal{M}$  as follows:

$$g \cdot \frac{dz(t)}{dt} = \frac{d(g \cdot z(t))}{dt}.$$

If the action is free and regular on  $\mathcal{M}$ , it will remain so on the jet space and we may use the same frame to find the first-order differential invariants

$$I_{k,j;t}(t) := \rho_k(z(t)) \cdot \frac{dz_j(t)}{dt}. \quad (1.7)$$

**DEFINITION 1.7 (Curvature Matrix)** The curvature matrix  $N_k$  is given by

$$N_k = \left( \frac{d}{dt} \rho_k \right) \rho_k^{-1} \quad (1.8)$$

when  $\rho_k$  is in matrix form.

It can be seen that for a right frame,  $N_k$  is an invariant matrix that involves the first order differential invariants. The above derivation applies to all discrete moving frames. For a difference frame the follow syzygy holds

$$\frac{d}{dt} K_0 = (SN_0)K_0 - K_0 N_0. \quad (1.9)$$

As  $N_0$  is invariant, from (1.3) we have that

$$N_0 = \iota_0 \left( \frac{d}{dt} \rho_0 \right). \quad (1.10)$$

In order to calculate the invariantized variation of the Euler–Lagrange equations, we may use the differential–difference syzygy

$$\frac{d}{dt} \boldsymbol{\kappa} = \mathcal{H} \boldsymbol{\sigma}, \quad (1.11)$$

where  $\boldsymbol{\kappa}$  is a vector of generating invariants,  $\mathcal{H}$  is a linear difference operator with coefficients that are functions of  $\boldsymbol{\kappa}$  and its shifts, and  $\boldsymbol{\sigma}$  is a vector of generating first order differential invariants of the form (1.7). Note that (1.11) comes from rearranging components in (1.9).

DEFINITION 1.8 Given a linear difference operator  $\mathcal{H} = c_j S_j$ , the adjoint operator  $\mathcal{H}^*$  is defined by

$$\mathcal{H}^*(F) = S_{-j}(c_j F)$$

and the associated boundary term  $A_{\mathcal{H}}$  is defined by

$$F \mathcal{H}(G) - \mathcal{H}^*(F)G = (S - \text{id})(A_{\mathcal{H}}(F, G)),$$

for all appropriate expressions  $F$  and  $G$ .

Now suppose we are given a group action  $G \times M \rightarrow M$  and that we have found a difference frame for this action.

THEOREM 1.9 (Invariant Euler–Lagrange Equations) (Theorem 5.2 in Mansfield, Rojo-Echeburúa, Hydon & Peng (2019)). Let  $\mathcal{L}$  be a Lagrangian functional whose invariant Lagrangian is given in terms of the generating invariants as

$$\mathcal{L} = \sum L(n, \boldsymbol{\kappa}_0, \dots, \boldsymbol{\kappa}_{J_1}),$$

and suppose that the differential–difference syzygies are

$$\frac{d\boldsymbol{\kappa}}{dt} = \mathcal{H} \boldsymbol{\sigma}.$$

Then (with  $\cdot$  denoting the sum over all components)

$$E_{\mathbf{u}}(L) \cdot \mathbf{u}'_0 = (\mathcal{H}^* E_{\boldsymbol{\kappa}}(L)) \cdot \boldsymbol{\sigma}, \quad (1.12)$$

where  $E_{\boldsymbol{\kappa}}(L)$  is the difference Euler operator with respect to  $\boldsymbol{\kappa}$ . Consequently, the invariantization of the original Euler–Lagrange equations is

$$\iota_0(E_{\mathbf{u}}(L)) = \mathcal{H}^* E_{\boldsymbol{\kappa}}(L). \quad (1.13)$$

Consequently, the original Euler–Lagrange equations, in invariant form, are equivalent to

$$\mathcal{H}^* E_{\boldsymbol{\kappa}}(L) = 0.$$

THEOREM 1.10 (Theorem 7.1 in Mansfield, Rojo-Echeburúa, Hydon & Peng (2019)) Suppose that the conditions of Theorem 1.9 hold. Write

$$A_{\mathcal{H}} = \mathcal{C}_{\alpha}^j S_j(\boldsymbol{\sigma}^{\alpha}),$$

where each  $\mathcal{C}_{\alpha}^j$  depends only on  $n$ ,  $\boldsymbol{\kappa}$  and its shifts. Let  $\Phi^{\alpha}(\mathbf{u}_0)$  be the row of the matrix of characteristics corresponding to the dependent variable  $u_0^{\alpha}$  and denote its invariantization by  $\Phi_0^{\alpha}(I) = \Phi^{\alpha}(\rho_0 \cdot \mathbf{u}_0)$ . Then the  $R$  conservation laws in row vector form amount to

$$\mathcal{C}_{\alpha}^j S_j \{ \Phi_0^{\alpha}(I) \mathcal{A}d(\rho_0) \} = 0. \quad (1.14)$$

That is, to obtain the conservation laws, it is sufficient to make the replacement

$$\boldsymbol{\sigma}^{\alpha} \mapsto \{ \Phi^{\alpha}(g \cdot \mathbf{u}_0) \mathcal{A}d(g) \} \Big|_{g=\rho_0}. \quad (1.15)$$

in  $A_{\mathcal{H}}$ .

## 2. The linear action of $SL(2)$ in the plane

We consider the action of  $SL(2)$  on the prolongation space  $P_n^{(0,0)}(\mathbb{R}^2)$ , which has coordinates  $(x_0, y_0)$ . This action is given by

$$\begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \mapsto \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} \tilde{x}_0 \\ \tilde{y}_0 \end{pmatrix}, \quad ad - bc = 1. \quad (2.1)$$

### 2.1 The Adjoint action

For our calculations we need the adjoint representation of  $SL(2)$  relative to this group action. The infinitesimal vector fields are

$$\mathbf{v}_a = x\partial_x - y\partial_y, \quad \mathbf{v}_b = y\partial_x, \quad \mathbf{v}_c = x\partial_y.$$

We have that the induced action on these are

$$\begin{pmatrix} \tilde{\mathbf{v}}_a & \tilde{\mathbf{v}}_b & \tilde{\mathbf{v}}_c \end{pmatrix} = \begin{pmatrix} \mathbf{v}_a & \mathbf{v}_b & \mathbf{v}_c \end{pmatrix} \mathcal{A}d(g)^{-1}$$

where

$$\mathcal{A}d(g) = \begin{matrix} & a & b & c \\ \begin{matrix} a \\ b \\ c \end{matrix} & \begin{pmatrix} ad+bc & -ac & bd \\ -2ab & a^2 & -b^2 \\ 2cd & -c^2 & d^2 \end{pmatrix} \end{matrix}. \quad (2.2)$$

### 2.2 The discrete frame, the generating invariants and their syzygies

Taking the normalisation equations  $\tilde{x}_0 = 1$ ,  $\tilde{x}_1 = \tilde{y}_0 = 0$  and solving for  $a, b$  and  $c$ , we define the moving frame

$$\rho_0(x_0, y_0, x_1, y_1) = \begin{pmatrix} \frac{y_1}{\tau} & -\frac{x_1}{\tau} \\ -y_0 & x_0 \end{pmatrix} \in SL(2)$$

where we have set  $\tau = x_0y_1 - x_1y_0$ . Then  $\rho_k = S_k\rho_0$  gives the discrete moving frame  $(\rho_k)$ .

#### 2.2.1 The generating discrete invariants. The Maurer–Cartan matrix is

$$K_0 = \iota_0(\rho_1) = \begin{pmatrix} \kappa & \frac{1}{\tau} \\ -\tau & 0 \end{pmatrix} \quad (2.3)$$

where we have set  $\kappa = \frac{x_0y_2 - x_2y_0}{x_1y_2 - x_2y_1}$ . Note that  $\tau = \iota_0(y_1)$  and  $\kappa = \frac{\iota_0(y_2)}{S(\iota_0(y_1))}$  are invariant, by the equivariance of the frame.

By the general theory of discrete moving frames, the algebra of invariants is generated by  $\tau$ ,  $\kappa$  and their shifts.

2.2.2 *The generating differential invariants.* We now consider  $x_j = x_j(t)$ ,  $y_j = y_j(t)$  and we define some first order differential invariants by setting

$$I_{k,j;t}^x := \rho_k \cdot x'_j \quad \text{and} \quad I_{k,j;t}^y := \rho_k \cdot y'_j, \quad (2.4)$$

where  $x'_j = \frac{d}{dt}x_j(t)$  and  $y'_j = \frac{d}{dt}y_j(t)$ . We set the notation

$$\sigma^x := I_{0,0;t}^x \quad \text{and} \quad \sigma^y := I_{0,0;t}^y. \quad (2.5)$$

LEMMA 2.1 For all  $k, j$ , both  $I_{k,j;t}^x(t)$  and  $I_{k,j;t}^y(t)$  may be written in terms of  $\sigma^x$ ,  $\sigma^y$ ,  $\kappa$ ,  $\tau$  and their shifts.

*Proof.* First note that  $I_{j,j;t}^x(t) = S_j \sigma^x$ ,  $I_{j,j;t}^y(t) = S_j \sigma^y$ . We have next for each  $k > j$  that

$$I_{k,j;t}^x(t) = \rho_k \cdot x'_j = \rho_k \rho_{k-1}^{-1} \rho_{k-1} \rho_{k-2}^{-1} \rho_{k-2} \cdots \rho_j^{-1} \rho_j \cdot x'_j = (S_k K_0) \cdots (S_j K_0) S_j \sigma^x$$

while similar calculations hold for for  $k < j$  and for  $I_{k,j;t}^y(t)$ .  $\square$  For our calculations, we need to know  $I_{0,2;t}^x(t)$ ,  $I_{0,1;t}^x(t)$  and  $I_{0,1;t}^y(t)$  explicitly. We have

$$\begin{aligned} \begin{pmatrix} I_{0,1;t}^x(t) \\ I_{0,1;t}^y(t) \end{pmatrix} &= \rho_0 \begin{pmatrix} x'_1 \\ y'_1 \end{pmatrix} \\ &= \rho_0 \rho_1^{-1} \rho_1 \begin{pmatrix} x'_1 \\ y'_1 \end{pmatrix} \\ &= K_0^{-1} \begin{pmatrix} S \sigma^x \\ S \sigma^y \end{pmatrix} \\ &= \begin{pmatrix} -\frac{S \sigma^y}{\tau} \\ \tau S \sigma^x + \kappa S \sigma^y \end{pmatrix} \end{aligned} \quad (2.6)$$

while a similar calculation yields, setting  $\tau_j = S_j \tau$  and  $\kappa_j = S_j \kappa$ ,

$$\begin{aligned} \begin{pmatrix} I_{0,2;t}^x(t) \\ I_{0,2;t}^y(t) \end{pmatrix} &= K_0^{-1} (S K_0^{-1}) \begin{pmatrix} S_2 \sigma^x \\ S_2 \sigma^y \end{pmatrix} \\ &= \begin{pmatrix} -\frac{\tau_1}{\tau} & -\frac{\kappa_1}{\tau} \\ \kappa \tau_1 & \kappa \kappa_1 - \frac{\tau}{\tau_1} \end{pmatrix} \begin{pmatrix} S_2 \sigma^x \\ S_2 \sigma^y \end{pmatrix}. \end{aligned} \quad (2.7)$$

We now define

$$N_0 = \iota_0 \left( \frac{d}{dt} \rho_0 \right) = \begin{pmatrix} -\sigma^x & -\frac{I_{0,1;t}^x(t)}{\tau} \\ -\sigma^y & \sigma^x \end{pmatrix} \in \mathfrak{sl}(2). \quad (2.8)$$

From (1.9) we may calculate the differential-difference syzygy. Equating components in (1.9) and simplifying we obtain

$$\begin{aligned} \frac{d}{dt} \kappa &= \kappa(\text{id} - S) \sigma^x + \left( \frac{1}{\tau} - \frac{\tau}{\tau_1^2} S_2 \right) \sigma^y, \\ \frac{d}{dt} \tau &= \tau(S + \text{id}) \sigma^x + \kappa S \sigma^y \end{aligned} \quad (2.9)$$



so that

$$\frac{d}{dt} \begin{pmatrix} \kappa \\ \tau \end{pmatrix} = \mathcal{H} \begin{pmatrix} \sigma^x \\ \sigma^y \end{pmatrix}$$

where

$$\mathcal{H} = \begin{pmatrix} \kappa(\text{id} - S) & \frac{1}{\tau} - \frac{\tau}{\tau_1^2} S_2 \\ \tau(\text{id} + S) & \kappa S \end{pmatrix}. \quad (2.10)$$

### 2.3 The Euler–Lagrange equations and conservation laws

We are now in a position to obtain the Euler–Lagrange equations and conservation laws for a Lagrangian of the form

$$\mathcal{L}[x, y] = \sum L(\tau, \tau_1, \dots, \tau_{j_1}, \kappa, \kappa_1, \dots, \kappa_{j_2}).$$

Using Theorem 1.9, we have that the Euler–Lagrange system is  $0 = \mathcal{H}^* (E_\kappa(L) E_\tau(L))^T$  which is written explicitly as

$$\begin{aligned} 0 &= (\text{id} - S_{-1}) \kappa E_\kappa(L) + (\text{id} + S_{-1}) \tau E_\tau(L), \\ 0 &= -S_{-2} \left( \frac{\tau}{\tau_1^2} E_\kappa(L) \right) + \frac{1}{\tau} E_\kappa(L) + S_{-1} (\kappa E_\tau(L)). \end{aligned} \quad (2.11)$$

We recall that if  $\mathcal{H} = \sum_{k=0}^m c_k S_k$  then  $\mathcal{H}^* = \sum_{k=0}^m (S_{-k} c_k) S_{-k}$ . Further, we recall the formula

$$F \mathcal{H}(G) - \mathcal{H}^*(F)G = (S - \text{id}) A_{\mathcal{H}}(F, G)$$

where

$$A_{\mathcal{H}}(F, G) = \sum_{k=1}^m \left( \sum_{j=0}^{k-1} S_j \right) (S_{-k} (c_k F) G)$$

and where the identity

$$(S_k - \text{id}) = (S - \text{id}) \sum_{j=0}^{k-1} S_j$$

has been used.

To obtain the conservation laws we need only the boundary terms arising from  $E(L) \mathcal{H} (\sigma^x \ \sigma^y)^T - \mathcal{H}^*(E(L)) (\sigma^x \ \sigma^y)^T$ , which we record here. They are  $(S - \text{id}) A_{\mathcal{H}}$  where

$$\begin{aligned} A_{\mathcal{H}} &= \mathcal{C}_0^x \sigma^x + \mathcal{C}_0^y \sigma^y + \mathcal{C}_1^y S \sigma^y \\ &= [-S_{-1} (\kappa E_\kappa(L)) + S_{-1} (\tau E_\tau(L))] \sigma^x \\ &\quad + \left[ S_{-1} (\kappa E_\tau(L)) - S_{-2} \left( \frac{\tau}{\tau_1^2} E_\kappa(L) \right) \right] \sigma^y \\ &\quad - S_{-1} \left( \frac{\tau}{\tau_1^2} E_\kappa(L) \right) S \sigma^y, \end{aligned} \quad (2.12)$$

where this defines  $\mathcal{C}_0^x$ ,  $\mathcal{C}_0^y$  and  $\mathcal{C}_1^y$ .

To find the conservation laws from  $A_{\mathcal{H}}$ , we first calculate the invariantized form of the matrix of infinitesimals restricted to the variables  $x_0$  and  $y_0$

$$\Phi_0(I) = \begin{matrix} & & a & b & c \\ \begin{matrix} x_0 \\ y_0 \end{matrix} & & \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \end{matrix}$$

and then the replacement required by Theorem 1.10 is given by

$$S_k \sigma^x \mapsto (1 \ 0 \ 0) S_k \mathcal{A}d(\rho_0)$$

and

$$S_k \sigma^y \mapsto (0 \ 0 \ 1) S_k \mathcal{A}d(\rho_0).$$

Since  $S \mathcal{A}d(\rho_0) = \mathcal{A}d(K_0) \mathcal{A}d(\rho_0)$ , after collecting terms and simplifying we obtain the Noether's Conservation Laws in the form

$$\begin{aligned} \mathbf{k} &= [\mathcal{C}_0^x (1 \ 0 \ 0) + \mathcal{C}_0^y (0 \ 0 \ 1) + \mathcal{C}_1^y (0 \ 0 \ 1) \mathcal{A}d(K_0)] \mathcal{A}d(\rho_0) \\ &= V(I) \mathcal{A}d(\rho_0) \end{aligned} \tag{2.13}$$

where

$$\mathcal{A}d(\rho_0) = \begin{pmatrix} \frac{x_0 y_1 + x_1 y_0}{\tau} & \frac{y_0 y_1}{\tau} & -\frac{x_0 x_1}{\tau} \\ 2 \frac{x_1 y_1}{\tau^2} & \frac{y_1^2}{\tau^2} & -\frac{x_1^2}{\tau^2} \\ -2x_0 y_0 & -y_0^2 & x_0^2 \end{pmatrix}$$

and

$$\mathcal{A}d(K_0) = \begin{pmatrix} -1 & \kappa \tau & 0 \\ -2 \frac{\kappa}{\tau} & \kappa^2 & -\frac{1}{\tau^2} \\ 0 & -\tau^2 & 0 \end{pmatrix}$$

and where  $\mathcal{C}_0^x$ ,  $\mathcal{C}_0^y$  and  $\mathcal{C}_1^y$  are defined in Equation (2.12), the vector  $\mathbf{k} = (k_1, k_2, k_3)$  is a vector of constants and where this equation defines  $V(I) = (V_0^1, V_0^2, V_0^3)^T$ . Explicitly, the vector of invariants  $V(I)$  is of the form

$$V(I) = S_{-1} \left( \begin{matrix} \tau E_{\tau}(L) - \kappa E_{\kappa}(L) & E_{\kappa}(L) & \kappa E_{\tau}(L) - S_{-1} \left( \frac{\tau}{\tau_1^2} E_{\kappa}(L) \right) \end{matrix} \right).$$

We note that once the Euler–Lagrange equations have been solved for the sequences  $(\kappa_k)$  and  $(\tau_k)$ , then  $V(I)$  is known, so that (2.13) can be considered as an algebraic equation for  $\rho_0$ . This will be the focus of the next section.

Recall that from  $(S - \text{id})(V(I), \mathcal{A}d(\rho_0)) = 0$  we obtain the discrete Euler–Lagrange equations in the form  $SV(I), \mathcal{A}d(\rho_1 \rho_0^{-1}) = V(I)$  which yields the equations

$$\begin{pmatrix} V_0^1 & V_0^2 & V_0^3 \end{pmatrix} = \begin{pmatrix} V_1^1 & V_1^2 & V_1^3 \end{pmatrix} \begin{pmatrix} -1 & \kappa\tau & 0 \\ -2\frac{\kappa}{\tau} & \kappa^2 & -\frac{1}{\tau^2} \\ 0 & -\tau^2 & 0 \end{pmatrix}. \quad (2.14)$$

#### 2.4 The general solution

If we can solve for the discrete frame  $(\rho_k)$  we then have from the general theory that

$$\begin{pmatrix} x_k \\ y_k \end{pmatrix} = \rho_k^{-1} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} d_k & -b_k \\ -c_k & a_k \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} d_k \\ -c_k \end{pmatrix}$$

since the normalisation equations for  $\rho_k$  are  $\rho_k \cdot (x_k, y_k)^T = (1, 0)^T$ .

**THEOREM 2.1** Given a solution  $(\kappa_k), (\tau_k)$  to the Euler–Lagrange equations, so that the vector of invariants  $S_k V(I) = (V_k^1, V_k^2, V_k^3)^T$  appearing in the conservation laws are known and satisfy  $V_k^2 \neq 0$  for all  $k$ , (2.13), and that with the three constants  $\mathbf{k} = (k_1, k_2, k_3)^T$  satisfying  $k_3(k_1^2 + 4k_2k_3) \neq 0$  are given, then the general solution to the Euler–Lagrange equations, in terms of  $(x_k, y_k)$  is

$$\begin{pmatrix} x_k \\ y_k \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} Q \begin{pmatrix} \prod_{l=0}^k \zeta_l \lambda_{1,l} & 0 \\ 0 & \prod_{l=0}^j k \zeta_l \lambda_{2,l} \end{pmatrix} Q^{-1} \begin{pmatrix} c_0 \\ d_0 \end{pmatrix}$$

where here,  $c_0$  and  $d_0$  are two further arbitrary constants of integration,

$$Q = \begin{pmatrix} k_1 - \sqrt{k_1^2 + 4k_2k_3} & k_1 + \sqrt{k_1^2 + 4k_2k_3} \\ 2k_3 & 2k_3 \end{pmatrix}, \quad (2.15)$$

and where

$$\lambda_{1,l} = V_l^1 - \sqrt{k_1^2 + 4k_2k_3}, \quad \lambda_{2,l} = V_l^1 + \sqrt{k_1^2 + 4k_2k_3}, \quad \zeta_l = -\frac{\tau_l}{2V_l^2}. \quad (2.16)$$

*Proof.* If we set

$$\rho_0 = \begin{pmatrix} a_0 & b_0 \\ c_0 & d_0 \end{pmatrix}, \quad a_0 d_0 - b_0 c_0 = 1 \quad (2.17)$$

and write (2.13) in the form  $\mathbf{k} \mathcal{A}d(\rho_0)^{-1} = V(I)$  as three equations for  $\{a_0, b_0, c_0, d_0\}$ , we obtain

$$\begin{aligned} (a_0 d_0 + b_0 c_0) k_1 + 2b_0 d_0 k_2 - 2a_0 c_0 k_3 &= V_0^1, \\ c_0 d_0 k_1 + d_0^2 k_2 - c_0^2 k_3 &= V_0^2, \\ -a_0 b_0 k_1 - b_0^2 k_2 + a_0^2 k_3 &= V_0^3. \end{aligned} \quad (2.18)$$

Computing a Gröebner basis associated to these equations, together with the equation  $a_0d_0 - b_0c_0 = 1$ , using the lexicographic ordering  $k_3 < k_2 < k_1 < c_0 < b_0 < a_0$ , we obtain

$$k_1^2 + 4k_2k_3 - (V_0^1)^2 - 4V_0^2V_0^3 = 0, \quad (2.19a)$$

$$k_3c_0^2 - k_1c_0d_0 - k_2d_0^2 + V_0^2 = 0, \quad (2.19b)$$

$$2b_0V_0^2 - 2c_0k_3 + (k_1 - V_0^1)d_0 = 0, \quad (2.19c)$$

$$2a_0V_0^2 - c_0(k_1 + V_0^1) - 2k_2d_0 = 0. \quad (2.19d)$$

We note that (2.19a) is a first integral of the Euler–Lagrange equations, (2.19b) is a conic equation for  $(c_0, d_0)$  while (2.19c) and (2.19d) are linear for  $(a_0, b_0)$  in terms of  $(c_0, d_0)$ .

We have

$$\rho_1 = \begin{pmatrix} \kappa & \frac{1}{\tau} \\ -\tau & 0 \end{pmatrix} \rho_0$$

where  $\rho_1 = S\rho_0$ . Hence

$$c_1 = -\tau a_0, \quad \text{and} \quad d_1 = -\tau b_0.$$

Back-substituting for  $a_0$  and  $b_0$  from (2.19c) and (2.19d) yields, assuming  $V_0^2 \neq 0$

$$\begin{pmatrix} c_1 \\ d_1 \end{pmatrix} = \frac{-\tau}{2V_0^2} \left( V_0^1 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} k_1 & 2k_2 \\ 2k_3 & -k_1 \end{pmatrix} \right) \begin{pmatrix} c_0 \\ d_0 \end{pmatrix}. \quad (2.20)$$

Now, setting

$$\underline{c}_0 = \begin{pmatrix} c_0 \\ d_0 \end{pmatrix}, \quad \underline{c}_1 = \begin{pmatrix} c_1 \\ d_1 \end{pmatrix}, \quad \zeta_0 = \frac{-\tau}{2V_0^2} \quad \text{and} \quad X_0 = V_0^1 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} k_1 & 2k_2 \\ 2k_3 & -k_1 \end{pmatrix}$$

equation (2.20) can be written as

$$\underline{c}_1 = \zeta_0 X_0 \underline{c}_0. \quad (2.21)$$

Diagonalising  $X_0$  we obtain  $\Lambda_0$  diagonal such that

$$\Lambda_0 = Q^{-1} X_0 Q = \begin{pmatrix} \lambda_0^1 & 0 \\ 0 & \lambda_0^2 \end{pmatrix}$$

where

$$\lambda_0^1 = V_0^1 - \sqrt{k_1^2 + 4k_2k_3} \quad \text{and} \quad \lambda_0^2 = V_0^1 + \sqrt{k_1^2 + 4k_2k_3}$$

and

$$Q = \begin{pmatrix} k_1 - \sqrt{k_1^2 + 4k_2k_3} & k_1 + \sqrt{k_1^2 + 4k_2k_3} \\ 2k_3 & 2k_3 \end{pmatrix}. \quad (2.22)$$

Since  $Q$  is a constant matrix, it is now simple to solve the recurrence relation. From (2.21), supposing  $k_3 \sqrt{k_1^2 + 4k_2k_3} \neq 0$  so  $Q^{-1}$  exists, we obtain

$$\underline{c}_{k+1} = Q \begin{pmatrix} \prod_{l=0}^k \zeta_l \lambda_{1,l} & 0 \\ 0 & \prod_{l=0}^k \zeta_l \lambda_{2,l} \end{pmatrix} Q^{-1} \underline{c}_0$$

where here,  $\underline{c}_0$  is the initial data. Finally from the normalisation equations

$$\begin{aligned} \begin{pmatrix} x_k \\ y_k \end{pmatrix} &= \rho_k^{-1} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} d_k & -b_k \\ -c_k & a_k \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} d_k \\ -c_k \end{pmatrix} \\ &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} c_k \\ d_k \end{pmatrix} \end{aligned}$$

and the result follows as required.  $\square$

REMARK 2.1 The proof of the Theorem makes no use of Equation (2.19b). Here we note that it is consistent with the second component of (2.14). We have that (2.19b) can be written as

$$\begin{pmatrix} c_0 & d_0 \end{pmatrix} \begin{pmatrix} k_3 & -\frac{k_1}{2} \\ -\frac{k_1}{2} & k_2 \end{pmatrix} \begin{pmatrix} c_0 \\ d_0 \end{pmatrix} = -V_0^2$$

and therefore

$$\begin{pmatrix} c_1 & d_1 \end{pmatrix} \begin{pmatrix} k_3 & -\frac{k_1}{2} \\ -\frac{k_1}{2} & k_2 \end{pmatrix} \begin{pmatrix} c_1 \\ d_1 \end{pmatrix} = -V_1^2. \quad (2.23)$$

Substituting (2.23) into (2.20) yields after simplification the equation

$$V_1^2 = -\tau^2 V_0^3$$

as claimed.

### 3. The $SA(2) = SL(2) \times \mathbb{R}^2$ linear action

We write the general element of the equi-affine group,  $SA(2) = SL(2) \times \mathbb{R}^2$ , as  $(g, \alpha, \beta)$  where  $g \in SL(2)$  as in Equation (1.1), and  $\alpha, \beta \in \mathbb{R}$ . We then consider the equi-affine group action on  $P_n^{(0,0)}(\mathbb{R}^2)$  with coordinates  $(x_0, y_0)$  given by

$$(g, \alpha, \beta) \cdot (x_0, y_0) = (\tilde{x}_0, \tilde{y}_0) = (ax_0 + by_0 + \alpha, cx_0 + dy_0 + \beta), \quad ad - bc = 1.$$

The standard representation of this group is given by

$$(g, \alpha, \beta) \mapsto \begin{pmatrix} a & b & \alpha \\ c & d & \beta \\ 0 & 0 & 1 \end{pmatrix}.$$

### 3.1 The Adjoint action

The infinitesimals vector fields are of the form

$$\mathbf{v}_a = x\partial_x - y\partial_y, \quad \mathbf{v}_b = y\partial_x, \quad \mathbf{v}_c = x\partial_y, \quad \mathbf{v}_\alpha = \partial_x, \quad \mathbf{v}_\beta = \partial_y.$$

We have that the induced action on these vector fields is

$$(\tilde{\mathbf{v}}_a \quad \tilde{\mathbf{v}}_b \quad \tilde{\mathbf{v}}_c \quad \tilde{\mathbf{v}}_\alpha \quad \tilde{\mathbf{v}}_\beta) = (\mathbf{v}_a \quad \mathbf{v}_b \quad \mathbf{v}_c \quad \mathbf{v}_\alpha \quad \mathbf{v}_\beta) \mathcal{A}d(g, \alpha, \beta)^{-1}$$

where

$$\mathcal{A}d(g, \alpha, \beta) = \begin{matrix} a & b & c & \alpha & \beta \\ \begin{pmatrix} a & b & c & \alpha & \beta \\ ad+bc & -ac & bd & 0 & 0 \\ -2ab & a^2 & -b^2 & 0 & 0 \\ 2cd & -c^2 & d^2 & 0 & 0 \\ -\alpha(ad+bc)+2ab\beta & a(c\alpha-a\beta) & b(b\beta-d\alpha) & a & b \\ \beta(ad+bc)-2cd\alpha & c(c\alpha-a\beta) & d(b\beta-d\alpha) & c & d \end{pmatrix} \end{matrix}. \quad (3.1)$$

REMARK 3.1 We note that (3.1) can be written as

$$\mathcal{A}d(g, \alpha, \beta) = \begin{pmatrix} \text{Id}_3 & 0 \\ \alpha \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} + \beta \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix} & \text{Id}_2 \end{pmatrix} \begin{pmatrix} \mathcal{A}d(g) & 0 \\ 0 & g \end{pmatrix} \quad (3.2)$$

where  $\text{Id}_2$  and  $\text{Id}_3$  are the  $2 \times 2$  and  $3 \times 3$  identity matrices respectively.

### 3.2 The discrete frame, the generating invariants and their syzygies

We define a moving frame  $\rho_0$  given by the normalisation equations

$$(g, \alpha, \beta) \cdot (x_0, y_0) = (0, 0), \quad (g, \alpha, \beta) \cdot (x_1, y_1) = (1, 0), \quad (g, \alpha, \beta) \cdot (x_2, y_2) = (0, *)$$

where  $*$  is to be left free. Solving for the group parameters  $a, b, c, d, \alpha$  and  $\beta$  we obtain the following standard matrix representation of the moving frame

$$\rho_0 = \begin{pmatrix} \frac{y_2 - y_0}{\kappa} & \frac{x_0 - x_2}{\kappa} & \frac{x_2 y_0 - x_0 y_2}{\kappa} \\ y_0 - y_1 & x_1 - x_0 & x_0 y_1 - x_1 y_0 \\ 0 & 0 & 1 \end{pmatrix},$$

where

$$\kappa = (y_1 - y_2)x_0 + (y_2 - y_0)x_1 + (y_0 - y_1)x_2$$

is an invariant. Indeed,

$$\kappa = \rho_0 \cdot y_2.$$

We define the discrete moving frame to be  $(\rho_k)$  where

$$\rho_k = S_k \rho_0.$$

The Maurer–Cartan matrix in the standard representation is

$$K_0 = \iota_0(\rho_1) = \begin{pmatrix} \tau & \frac{1+\tau}{\kappa} & -\tau \\ -\kappa & -1 & \kappa \\ 0 & 0 & 1 \end{pmatrix} \quad (3.3)$$

where  $\kappa = \rho_0 \cdot y_2$  is given above, and

$$\tau = \frac{x_0(y_1 - y_3) + x_1(y_3 - y_0) + x_3(y_0 - y_1)}{x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)} = \frac{\rho_0 \cdot y_3}{\kappa_1}$$

where we have used the Replacement Rule, Theorem 1.3, and where  $\kappa_k = S_k \kappa$ . By the general theory of discrete moving frames the algebra of invariants is generated by  $\tau$ ,  $\kappa$  and their shifts. We note that one could take  $\rho_0 \cdot y_3$  and  $\kappa$  to be the generators, but the resulting formulae in the sequel are no simpler.

We obtain

$$N_0 = \iota_0 \left( \frac{d}{dt} \rho_0 \right) = \begin{pmatrix} \sigma^x - I_{0,1;t}^x(t) & \frac{\sigma^x - I_{0,2;t}^x(t)}{\kappa} & -\sigma^x \\ \sigma^y - I_{0,1;t}^y(t) & I_{0,1;t}^x(t) - \sigma^x & -\sigma^y \\ 0 & 0 & 0 \end{pmatrix} \quad (3.4)$$

where we have set  $\sigma^x := I_{0,0;t}^x(t)$  and  $\sigma^y := I_{0,0;t}^y(t)$ .

To obtain  $\rho_0 \cdot x'_j = I_{0,j;t}^x(t)$ ,  $\rho_0 \cdot y'_j = I_{0,j;t}^y(t)$ ,  $j = 1, 2$  in terms of  $\sigma^x$ ,  $\sigma^y$ ,  $\tau$ ,  $\kappa$  and their shifts, we have, since the translation part of the group plays no role in the action on the derivatives,

$$\begin{pmatrix} \rho_0 \cdot x'_1 \\ \rho_0 \cdot y'_1 \\ 0 \end{pmatrix} = \rho_0 \begin{pmatrix} x'_1 \\ y'_1 \\ 0 \end{pmatrix} = \rho_0 \rho_1^{-1} \rho_1 \begin{pmatrix} x'_1 \\ y'_1 \\ 0 \end{pmatrix} = K_0^{-1} \begin{pmatrix} S\sigma^x \\ S\sigma^y \\ 0 \end{pmatrix}$$

and similarly

$$\rho_0 \begin{pmatrix} x'_2 \\ y'_2 \\ 0 \end{pmatrix} = K_0^{-1} (S K_0^{-1}) \begin{pmatrix} S_2 \sigma^x \\ S_2 \sigma^y \\ 0 \end{pmatrix}.$$

Finally from (1.9) and the relations above, we have the differential-difference syzygy

$$\frac{d}{dt} \begin{pmatrix} \tau \\ \kappa \end{pmatrix} = \mathcal{H} \begin{pmatrix} \sigma^x \\ \sigma^y \end{pmatrix}, \quad (3.5)$$

where  $\mathcal{H}$  is a difference operator depending only on the generating difference invariants  $\tau$ ,  $\kappa$  and their shifts, and which has the form

$$\mathcal{H} = \begin{pmatrix} \mathcal{H}_{11} & \mathcal{H}_{12} \\ \mathcal{H}_{21} & \mathcal{H}_{22} \end{pmatrix}$$

where

$$\begin{aligned}
\mathcal{H}_{11} &= -\tau + \left(1 + \frac{\kappa}{\kappa_1}(1 + \tau)\right) S + \tau S_2 - \frac{\kappa}{\kappa_1^2} [\kappa_2(1 + \tau_1) - \kappa_1] S_3, \\
\mathcal{H}_{12} &= -\frac{1 + \tau}{\kappa} + \frac{\tau(1 + \tau_1)}{\kappa_1} S_2 - \frac{\kappa}{\kappa_1^2 \kappa_2} [\kappa_2 \tau_2(1 + \tau_1) - \kappa_1(1 + \tau_2)] S_3, \\
\mathcal{H}_{21} &= -\kappa - \kappa S + (\tau \kappa_1 - \kappa) S_2, \\
\mathcal{H}_{22} &= -1 - (1 + \tau) S + \left(\tau \tau_1 - \frac{\kappa(1 + \tau_1)}{\kappa_1}\right) S_2.
\end{aligned} \tag{3.6}$$

### 3.3 The Euler–Lagrange equations and the conservation laws.

We consider a Lagrangian of the form  $L(\tau, \dots, \tau_{J_1}, \kappa, \dots, \kappa_{J_2})$ . Then by Theorem 1.9 we have that the Euler–Lagrange equations are

$$\begin{aligned}
0 &= \mathcal{H}_{11}^* E_\tau(L) + \mathcal{H}_{21}^* E_\kappa(L), \\
0 &= \mathcal{H}_{12}^* E_\tau(L) + \mathcal{H}_{22}^* E_\kappa(L)
\end{aligned} \tag{3.7}$$

where the  $\mathcal{H}_{ij}$  are given in Equation (3.6).

The boundary terms contributing to the conservation laws are

$$\begin{aligned}
A_{\mathcal{H}} &= A_{\mathcal{H}_{11}}(E_\tau(L), \sigma^x) + A_{\mathcal{H}_{21}}(E_\kappa(L), \sigma^x) + \\
&\quad + A_{\mathcal{H}_{12}}(E_\tau(L), \sigma^y) + A_{\mathcal{H}_{22}}(E_\kappa(L), \sigma^y) \\
&= \sum_{k=0}^2 \mathcal{C}_k^x S_k \sigma^x + \mathcal{C}_k^y S_k \sigma^y
\end{aligned} \tag{3.8}$$

where this defines the  $\mathcal{C}_k^x, \mathcal{C}_k^y$ . Explicitly

$$\begin{aligned}
\mathcal{C}_0^x &= S_{-1} \left(1 + \frac{\kappa}{\kappa_1}(1 + \tau)\right) E_\tau(L) + S_{-2} (\tau E_\tau(L)) + S_{-3} \left(-\frac{\kappa}{\kappa_1^2} (\kappa_2(1 + \tau_1) - \kappa_1)\right) E_\tau(L) \\
&\quad + S_{-1} (-\kappa E_\kappa(L)) + S_{-2} ((\tau \kappa_1 - \kappa) E_\kappa(L)), \\
\mathcal{C}_1^x &= S_{-1} (\tau E_\tau(L)) + S_{-2} \left(-\frac{\kappa}{\kappa_1^2} (\kappa_2(1 + \tau_1) - \kappa_1)\right) E_\tau(L) + S_{-1} ((\tau \kappa_1 - \kappa) E_\kappa(L)), \\
\mathcal{C}_2^x &= S_{-1} \left(-\frac{\kappa}{\kappa_1^2} (\kappa_2(1 + \tau_1) - \kappa_1)\right) E_\tau(L), \\
\mathcal{C}_0^y &= S_{-2} \left(\frac{\tau(1 + \tau_1)}{\kappa_1} E_\tau(L)\right) + S_{-3} \left(-\frac{\kappa}{\kappa_1^2 \kappa_2} (\kappa_2 \tau_2(1 - \tau_1) - \kappa_1(1 + \tau_2))\right) E_\tau(L) \\
&\quad - S_{-1} (1 + \tau) E_\kappa(L) + S_{-2} \left(\tau \tau_1 - \frac{\kappa(1 - \tau_1)}{\kappa_1}\right) E_\kappa(L), \\
\mathcal{C}_1^y &= S_{-1} \left(\frac{\tau(1 + \tau_1)}{\kappa_1} E_\tau(L)\right) + S_{-2} \left(-\frac{\kappa}{\kappa_1^2 \kappa_2} (\kappa_2 \tau_2(1 - \tau_1) - \kappa_1(1 + \tau_2))\right) E_\tau(L) \\
&\quad + S_{-1} \left(\tau \tau_1 - \frac{\kappa(1 - \tau_1)}{\kappa_1}\right) E_\kappa(L), \\
\mathcal{C}_2^y &= S_{-1} \left(-\frac{\kappa}{\kappa_1^2 \kappa_2} (\kappa_2 \tau_2(1 - \tau_1) - \kappa_1(1 + \tau_2))\right) E_\tau(L).
\end{aligned} \tag{3.9}$$



To obtain the conservation laws we need the invariantized form of the matrix of infinitesimals restricted to the variables  $x_0$  and  $y_0$

$$\Phi_0(I) = \begin{array}{c} x_0 \\ y_0 \end{array} \begin{array}{ccccc} a & b & c & \alpha & \beta \\ \left( \begin{array}{ccccc} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \end{array}$$

and then the replacements required to obtain the conservation laws from  $A_{\mathcal{H}}$  are

$$S_k \sigma^x \mapsto (0 \ 0 \ 0 \ 1 \ 0) S_k \mathcal{A}d(\rho_0), \quad S_k \sigma^y \mapsto (0 \ 0 \ 0 \ 0 \ 1) S_k \mathcal{A}d(\rho_0).$$

Hence, the conservation laws are given by  $(S - \text{id})A = 0$  where

$$\begin{aligned} A &= \left[ \begin{array}{ccccc} 0 & 0 & 0 & 1 & 0 \end{array} \right] (\mathcal{C}_0^x + \mathcal{C}_1^x \mathcal{A}d(K_0) + \mathcal{C}_2^x \mathcal{A}d(K_0(SK_0))) \\ &\quad + \left[ \begin{array}{ccccc} 0 & 0 & 0 & 0 & 1 \end{array} \right] (\mathcal{C}_0^y + \mathcal{C}_1^y \mathcal{A}d(K_0) + \mathcal{C}_2^y \mathcal{A}d((SK_0)K_0)) \mathcal{A}d(\rho_0) \\ &= V(I) \mathcal{A}d(\rho_0) \end{aligned} \quad (3.10)$$

where

$$\mathcal{A}d(K_0) = \begin{pmatrix} \text{Id}_3 & 0 \\ \begin{pmatrix} \tau & -\kappa & 0 \\ \kappa & 0 & -\tau \end{pmatrix} & \text{Id}_2 \end{pmatrix} \begin{pmatrix} -2\tau & \tau\kappa & -\frac{1+\tau}{\kappa} & 0 & 0 \\ -\frac{2\tau(1+\tau)}{\kappa} & \tau^2 & -\frac{(1+\tau)^2}{\kappa^2} & 0 & 0 \\ 2\kappa & -\kappa^2 & 1 & 0 & 0 \\ 0 & 0 & 0 & \tau & \frac{1+\tau}{\kappa} \\ 0 & 0 & 0 & -\kappa & -1 \end{pmatrix}$$

and where this defines the vector of invariants,  $V(I) = (V_0^1, V_0^2, V_0^3, V_0^4, V_0^5)^T$  and where the  $\mathcal{C}_j^x, \mathcal{C}_j^y$  are defined in Equation (3.8) and (3.9).

We can thus write the conservation laws in the form

$$\mathbf{k} = V(I) \mathcal{A}d(\rho_0) \quad (3.11)$$

where  $\mathbf{k} = (k_1, k_2, k_3, k_4, k_5)$  is a vector of constants and where

$$\mathcal{A}d(\rho_0) = \begin{pmatrix} \text{Id}_3 & 0 \\ \begin{pmatrix} \frac{x_0 y_2 - x_2 y_0}{\kappa} & x_1 y_0 - x_0 y_1 & 0 \\ x_0 y_1 - x_1 y_0 & 0 & \frac{x_2 y_0 - x_0 y_2}{\kappa} \end{pmatrix} & \text{Id}_2 \end{pmatrix} \begin{pmatrix} (\mathcal{A}d(g))|_{\rho_0} & 0 \\ 0 & g|_{\rho_0} \end{pmatrix}$$

where

$$\mathcal{A}d(g)|_{\rho_0} = \begin{pmatrix} \frac{(2y_0 - y_1 - y_2)x_0 - (x_1 + x_2)y_0 + y_2x_1 + x_2y_1}{\kappa} & \frac{(y_0 - y_2)(y_0 - y_1)}{\kappa} & \frac{(x_0 - x_2)(x_1 - x_0)}{\kappa} \\ \frac{2(y_0 - y_2)(x_0 - x_2)}{\kappa^2} & \frac{(y_0 - y_2)^2}{\kappa^2} & -\frac{(x_0 - x_2)^2}{\kappa^2} \\ 2(y_0 - y_1)(x_1 - x_0) & -(y_0 - y_1)^2 & (x_0 - x_1)^2 \end{pmatrix},$$

and

$$g|_{\rho_0} = \begin{pmatrix} \frac{y_2 - y_0}{\kappa} & \frac{x_0 - x_2}{\kappa} \\ y_0 - y_1 & x_1 - x_0 \end{pmatrix}.$$

We will show in the next section that a first integral of the Euler–Lagrange equations is given by

$$k_1k_4k_5 + k_2k_5^2 - k_3k_4^2 = V_0^1V_0^4V_0^5 + V_0^2(V_0^5)^2 - V_0^3(V_0^4)^2. \quad (3.12)$$

### 3.4 The general solution

We now show how to obtain the solution to the Euler–Lagrange equations in terms of the original variables, given the vector of invariants and the constants in the conservation laws, (3.11).

**THEOREM 3.1** Suppose a solution  $(\tau_k), (\kappa_k)$  to the Euler–Lagrange equations (3.7), is given, so that the vector of invariants  $(S_k V(I))$  appearing in the conservation laws (3.10) is known, and that  $V_0^4 V_0^5 \neq 0$ . Suppose further that a vector of constants  $\mathbf{k} = (k_1, k_2, k_3, k_4, k_5)$  satisfying  $k_4 k_5 \neq 0$  is given. Then the general solution to the Euler–Lagrange equations, in terms of  $(x_k, y_k)$  is given by

$$\begin{pmatrix} x_k \\ y_k \\ 1 \end{pmatrix} = \rho_k^{-1} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} -\alpha_k d_k + \beta_k b_k \\ \alpha_k c_k - \beta_k c_k \\ 1 \end{pmatrix}. \quad (3.13)$$

where, setting  $\mu := k_1k_4k_5 + k_2k_5^2 - k_3k_4^2$ ,

$$\begin{aligned} a_0 &= -\frac{V_0^5}{V_0^4}c_0 + \frac{k_4}{V_0^4} \\ b_0 &= -\frac{V_0^5k_5}{V_0^4k_4}c_0 + \frac{k_4k_5 - V_0^4V_0^5}{V_0^4k_4} \\ d_0 &= \frac{k_5}{k_4}c_0 + \frac{V_0^4}{k_4} \\ \alpha_0 &= \frac{\mu V_0^5}{(V_0^4k_4)^2}c_0^2 + \frac{(k_2k_5^2 + k_3k_4^2 + \mu)V_0^4(V_0^5)^2 - 2\mu k_4k_5V_0^5}{(V_0^4k_4)^2V_0^5k_5}c_0 \\ &\quad + \frac{1}{(V_0^4k_4)^2V_0^5k_5} \left( k_2k_5(V_0^4V_0^5)^2 - (k_2k_5^2 + k_3k_4^2 + \mu)V_0^4V_0^5k_4 + k_4^2k_5(V_0^3(V_0^4)^2 + \mu) \right) \\ \beta_0 &= -\frac{\mu}{k_4^2V_0^4}c_0^2 - \frac{V_0^4(k_2k_5^2 + k_3k_4^2 + \mu)}{k_4^2k_5V_0^4}c_0 + \frac{k_4^2V_0^2 - k_2(V_0^4)^2}{k_4^2V_0^4} \end{aligned} \quad (3.14)$$

and where

$$c_k = \prod_{l=0}^{k-1} \left( \frac{\kappa_l V_l^5}{V_l^4} - 1 \right) c_0 - \sum_{l=0}^{k-1} \prod_{m=l+1}^{k-1} \left( \frac{\kappa_l V_m^5}{V_m^4} - 1 \right) \frac{k_4 \kappa_l}{V_l^4} \quad (3.15)$$

where in this last equation,  $c_0$  is the initial datum, or constant of integration.

*Proof.* If we can solve for the discrete frame  $(\rho_k)$

$$\rho_k = \begin{pmatrix} a_k & b_k & \alpha_k \\ c_k & d_k & \beta_k \\ 0 & 0 & 1 \end{pmatrix},$$

then we have by the normalisation equations, that Equation (3.13) holds. We consider (3.11) as five equations for  $\{a_0, b_0, c_0, d_0, \alpha_0, \beta_0\}$ , which when written out in detail are

$$\begin{aligned} 0 &= (a_0 d_0 + b_0 c_0) k_1 + 2b_0 d_0 k_2 - 2a_0 c_0 k_3 + (b \beta_0 + d_0 \alpha_0) k_4 - (a_0 \beta_0 + c_0 \alpha_0) - V_0^1, \\ 0 &= -c_0^2 k_3 + c_0 k_1 d_0 - c_0 k_5 \beta_0 + k_2 d_0^2 + k_4 d_0 k_2 - V_0^2, \\ 0 &= a_0^2 k_3 - a_0 b_0 k_1 + a_0 k_5 \alpha_0 - b_0^2 k_2 - b_0 k_4 \alpha_0 - V_0^3, \\ 0 &= -c_0 k_5 + k_4 d_0 - V_0^4, \\ 0 &= a_0 k_5 - b_0 k_4 - V_0^5. \end{aligned}$$

Computing a Gröbner basis associated to these equations with the lexicographic ordering  $k_2 < k_1 < a_0 < b_0 < d_0 < \beta_0 < \alpha_0$ , we obtain the first integral noted in Equation (3.12), and the expressions for  $a_0, b_0, d_0, \alpha_0$  and  $\beta_0$  in terms of  $c_0$  given in Equations (3.14), provided  $V_0^4, V_0^5, k_4$  and  $k_5$  are all non zero.

We have  $\rho_1 = K_0 \rho_0$  so that we have a recurrence equation for  $(c_k)$ , specifically,

$$c_1 = -\kappa a_0 - c_0 = \left( \frac{\kappa V_0^5}{V_0^4} - 1 \right) c_0 - \frac{k_4 \kappa}{V_0^4}$$

where we have back substituted for  $a_0$  from (3.14). This is linear and can be easily solved to obtain the expression for  $c_k$  given in Equation (3.15). Substituting this into the shifts of (3.14) yields  $(a_k), (b_k), (d_k), (\alpha_k)$  and  $(\beta_k)$  and substituting these into (3.13) yields the desired result.  $\square$

#### 4. The $SL(2)$ projective action

In this example, we show some techniques for calculating the recurrence relations when the action is nonlinear. We detail the calculations for a class of one-dimensional  $SL(2)$  Lagrangians, which are invariant under the projective action of  $SL(2)$ . This is defined by

$$\tilde{x}_0 = g \cdot x_0 = \frac{ax_0 + b}{cx_0 + d}, \quad ad - bc = 1. \quad (4.1)$$

##### 4.1 The Adjoint action

The infinitesimal vector fields are

$$\mathbf{v}_a = 2x\partial_x, \quad \mathbf{v}_b = \partial_x, \quad \mathbf{v}_c = -x^2\partial_x.$$

We have that the induced action on these are

$$(\tilde{\mathbf{v}}_a \quad \tilde{\mathbf{v}}_b \quad \tilde{\mathbf{v}}_c) = (\mathbf{v}_a \quad \mathbf{v}_b \quad \mathbf{v}_c) \mathcal{A}d(g)^{-1}$$

where

$$\mathcal{A}d(g) = \begin{matrix} & a & b & c \\ \begin{matrix} a \\ b \\ c \end{matrix} & \begin{pmatrix} ad+bc & -ac & bd \\ -2ab & a^2 & -b^2 \\ 2cd & -c^2 & d^2 \end{pmatrix} \end{matrix} \quad (4.2)$$

which matches with (2.2) as expected.

#### 4.2 The discrete frame, the generating invariants and their syzygies

We choose the normalisation equations

$$\tilde{x}_0 = \frac{1}{2}, \quad \tilde{x}_1 = 0, \quad \tilde{x}_2 = -\frac{1}{2}$$

which we can solve, together with  $ad - bc - 1$  for  $a, b, c$  and  $d$  to find the frame

$$\rho_0 = \frac{\sqrt{x_0 - x_2}}{\sqrt{(x_0 - x_1)(x_1 - x_2)}} \begin{pmatrix} \frac{1}{2} & -\frac{x_1}{2} \\ \frac{x_2 - 2x_1 + x_0}{x_0 - x_2} & \frac{x_0x_1 - 2x_0x_2 + x_1x_2}{x_0 - x_2} \end{pmatrix} \quad (4.3)$$

and we take

$$\rho_k = S_k \rho_0.$$

#### 4.3 The generating discrete invariants

The famous, historical invariant for this action, given four points, is the cross ratio,

$$\kappa = \frac{(x_0 - x_1)(x_2 - x_3)}{(x_0 - x_3)(x_2 - x_1)}. \quad (4.4)$$

By the Replacement Rule, Replacement Rule, Theorem 1.3, we have that

$$\kappa(x_0, x_1, x_2, x_3) = \kappa(\rho_0 \cdot x_0, \rho_0 \cdot x_1, \rho_0 \cdot x_2, \rho_0 \cdot x_3) = \kappa\left(\frac{1}{2}, 0, -\frac{1}{2}, I_{0,3}^x\right)$$

or

$$\kappa = \frac{1 + 2I_{0,3}^x}{1 - 2I_{0,3}^x}.$$

The Maurer–Cartan matrix is then,

$$K_0 = \iota_0(\rho_1) = \sqrt{\frac{\kappa - 1}{4\kappa}} \begin{pmatrix} 1 & \frac{1}{2} \\ -\frac{6\kappa + 2}{\kappa - 1} & 1 \end{pmatrix}. \quad (4.5)$$

By the general theory of moving frames, the discrete invariants are generated by  $\kappa$  and its shifts.

We now show how the recurrence relations may be obtained for this non-linear action.

#### 4.4 The generating differential invariants

We now consider  $x_j = x_j(t)$  where  $t$  is an invariant parameter. The induced action on these is given by

$$g \cdot x'_j = \frac{d}{dt} g \cdot x_j = \frac{x'_j}{(cx_j + d)^2}$$

and hence we have for

$$\rho_k = \begin{pmatrix} a_k & b_k \\ c_k & d_k \end{pmatrix}$$

that

$$\rho_k \cdot x'_j = \frac{x'_j}{(c_k x_j + d_k)^2}.$$

We define

$$\sigma_j^x := \rho_0 \cdot x_{j,t} = \frac{x'_j}{(c_0 x_j + d_0)^2} = \frac{x'_j (x_1 - x_0)}{(x_0 - x_2)(x_0 - x_1)}$$

where  $c_0$  and  $d_0$  are given in Equation (4.3). In terms of the  $\sigma_j^x$ , it is straightforward to show

$$N_0 = \iota_0 \left( \frac{d}{dt} \rho_0 \right) = \begin{pmatrix} \frac{1}{2} \sigma_1^x - \frac{1}{2} \sigma_0^x & -\sigma_1^x \\ 2\sigma_0^x - 4\sigma_1^x + 2\sigma_1^x & -\frac{1}{2} \sigma_1^x + \frac{1}{2} \sigma_0^x \end{pmatrix}.$$

We now obtain the recurrence relations for the  $\sigma_j^x$ . First observe that since  $\rho_k \cdot x'_j$  is an invariant, we have for all  $k$  and  $j$  that

$$\rho_k \cdot x'_j = \frac{\tilde{x}'_j}{(\tilde{c}_k \tilde{x}_j + \tilde{d}_k)^2} = \frac{\tilde{x}'_j}{(\tilde{c}_k \tilde{x}_j + \tilde{d}_k)^2} \Big|_{\rho_0} = \frac{\rho_0 \cdot x'_j}{(\tilde{c}_k|_{\rho_0} \rho_0 \cdot x_j + \tilde{d}_k|_{\rho_0})^2}.$$

Now the frames  $\rho_k$  are equivariant, and so  $\tilde{\rho}_k = \rho_k g^{-1}$ . Hence

$$\begin{pmatrix} \tilde{a}_k & \tilde{b}_k \\ \tilde{c}_k & \tilde{d}_k \end{pmatrix} \Big|_{\rho_0} = \rho_k \rho_0^{-1} = \rho_k \rho_{m-1}^{-1} \cdots \rho_1 \rho_0^{-1} = (S^{m-1} K_0) \cdots K_0.$$

In particular, we have

$$S\sigma_0^x = \rho_1 \cdot x'_1 = \frac{\rho_0 \cdot x'_1}{((K_0)_{2,1} \rho_0 \cdot x_1 + (K_0)_{2,2})^2} = \frac{4\kappa}{\kappa - 1} \sigma_1^x \quad (4.6)$$

since  $\rho_0 \cdot x_1 = 0$  and  $\rho_0 \cdot x'_1 = \sigma_1^x$ . Next,

$$S_2\sigma_0^x = \rho_2 \cdot x'_2 = \frac{\rho_0 \cdot x'_2}{(((SK_0)K_0)_{2,1} \rho_0 \cdot x_2 + ((SK_0)K_0)_{2,2})^2} = \frac{\kappa_1(\kappa - 1)}{(\kappa_1 - 1)\kappa} \sigma_2^x \quad (4.7)$$

noting that by the normalisation equations,  $\rho_0 \cdot x_1 = 0$  and  $\rho_0 \cdot x_2 = -1/2$ . Similarly, one can prove that

$$S\sigma_1^x = \frac{\kappa - 1}{4\kappa} \sigma_2^x. \quad (4.8)$$

We are now ready to calculate the differential difference syzygy. Calculating (1.9) and equating components and using the syzygies (4.6), (4.6) and (4.8), we obtain

$$\begin{aligned} \frac{d}{dt} \kappa &= \frac{\kappa(\kappa-1)\kappa_1(\kappa_2-1)}{\kappa_2(\kappa_1-1)} S_3 \sigma_0^x + \frac{\kappa(\kappa_1-1)}{\kappa_1} S_2 \sigma_0^x - (\kappa-1) S \sigma_0^x - \kappa(\kappa-1) \sigma_0^x \\ &= \mathcal{H} \sigma_0^x \end{aligned} \quad (4.9)$$

where this defines the linear difference operator  $\mathcal{H}$ .

#### 4.5 The Euler–Lagrange equations and the conservation laws

Given a Lagrangian of the form

$$\mathcal{L}[x] = \sum L(\kappa, \kappa_1, \dots, \kappa_J)$$

we have from Theorem 1.9 that the Euler–Lagrange equation is

$$0 = \mathcal{H}^*(E_\kappa(L))$$

where  $\mathcal{H}$  is given in Equation (4.9). Set  $\mathcal{H} = \alpha S_3 + \beta S_2 + \gamma S + \delta$  where this defines  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$ , specifically,

$$\begin{aligned} \alpha &= \frac{\kappa(\kappa-1)\kappa_1(\kappa_2-1)}{\kappa_2(\kappa_1-1)}, \\ \beta &= \frac{\kappa(\kappa_1-1)}{\kappa_1}, \\ \gamma &= -(\kappa-1), \\ \delta &= -\kappa(\kappa-1). \end{aligned} \quad (4.10)$$

Then the Euler–Lagrange equation is

$$0 = \mathcal{H}^*(E_\kappa(L)) = S_{-3}(\alpha E_\kappa(L)) + S_{-2}(\beta E_\kappa(L)) + S_{-1}(\gamma E_\kappa(L)) + \delta E_\kappa(L).$$

To obtain the conservation law, we need the matrix of infinitesimals, which is

$$\Phi_0 = x_0 \begin{pmatrix} a & b & c \\ 2x_0 & 1 & -x_0^2 \end{pmatrix}$$

and so

$$\Phi_0(I) = x_0 \begin{pmatrix} a & b & c \\ 1 & 1 & -\frac{1}{4} \end{pmatrix}.$$

Recall the relation

$$E_\kappa(L) \mathcal{H} \sigma_0^x = \mathcal{H}^*(E_\kappa(L)) \sigma_0^x + (S - \text{id})(A_{\mathcal{H}}(E_\kappa(L), \sigma_0^x))$$

where for any suitable arguments  $F$  and  $G$ ,

$$\begin{aligned} A_{\mathcal{H}}(F, G) &= (S_{-3}(\gamma F) + S_{-2}(\beta F) + S_{-1}(\alpha F)) G \\ &\quad + (S_{-1}(\beta F) + S_{-2}(\alpha F)) S(G) + S_{-1}(\alpha F) S_2(G). \end{aligned} \quad (4.11)$$

Hence by Theorem 1.10, the conservation law is

$$\begin{aligned} \mathbf{k} = & (\mathbb{S}_{-1}(\gamma\mathbf{E}_\kappa(L)) + \mathbb{S}_{-2}(\beta\mathbf{E}_\kappa(L)) + \mathbb{S}_{-1}(\alpha\mathbf{E}_\kappa(L))) \Phi_0(I) \mathcal{A}d(\rho_0) \\ & + (\mathbb{S}_{-3}(\beta\mathbf{E}_\kappa(L)) + \mathbb{S}_{-2}(\alpha\mathbf{E}_\kappa(L))) \Phi_0(I) (\mathbb{S} \mathcal{A}d(\rho_0)) \\ & + \mathbb{S}_{-1}(\alpha\mathbf{E}_\kappa(L)) \Phi_0(I) (\mathbb{S}_2 \mathcal{A}d(\rho_0)). \end{aligned} \quad (4.12)$$

Using

$$\begin{aligned} \mathbb{S} \mathcal{A}d(\rho_0) &= \mathcal{A}d(\rho_1) = \mathcal{A}d(K_0) \mathcal{A}d(\rho_0), \\ \mathbb{S}_2 \mathcal{A}d(\rho_0) &= \mathcal{A}d(\mathbb{S}(K_0)) \mathcal{A}d(K_0) \mathcal{A}d(\rho_0) \end{aligned}$$

and collecting terms, we arrive at the conservation laws in the form

$$\mathbf{k} = V(I) \mathcal{A}d(\rho_0) \quad (4.13)$$

where this defines the vector  $V(I) = (V_0^1, V_0^2, V_0^3)^T$  and where

$$\mathcal{A}d(\rho_0) = \begin{pmatrix} \frac{x_1^2 - x_0 x_2}{(x_0 - x_1)(x_1 - x_2)} & \frac{2x_1 - x_2 - x_0}{2(x_0 - x_1)(x_1 - x_2)} & \frac{x_1(2x_0 x_2 - x_1(x_0 + x_2))}{2(x_0 - x_1)(x_1 - x_2)} \\ \frac{(x_0 - x_2)x_1}{2(x_0 - x_1)(x_1 - x_2)} & \frac{x_0 - x_2}{4(x_0 - x_1)(x_1 - x_2)} & \frac{x_1^2(x_2 - x_0)}{4(x_0 - x_1)(x_1 - x_2)} \\ \frac{2(x_2 - 2x_1 + x_0)((x_1 - 2x_2)x_0 + x_1 x_2)}{(x_0 - x_2)(x_0 - x_1)(x_1 - x_2)} & -\frac{(x_2 - 2x_1 + x_0)^2}{(x_0 - x_2)(x_0 - x_1)(x_1 - x_2)} & \frac{((x_1 - 2x_2)x_0 + x_1 x_2)^2}{(x_0 - x_2)(x_0 - x_1)(x_1 - x_2)} \end{pmatrix}.$$

Explicitly,  $V(I)$  is given by

$$\begin{aligned} V(I) = & \begin{pmatrix} 1 & 1 & \frac{1}{4} \end{pmatrix} \{ (\mathbb{S}_{-1}(\gamma\mathbf{E}_\kappa(L)) + \mathbb{S}_{-2}(\beta\mathbf{E}_\kappa(L)) + \mathbb{S}_{-1}(\alpha\mathbf{E}_\kappa(L))) \\ & + (\mathbb{S}_{-3}(\beta\mathbf{E}_\kappa(L)) + \mathbb{S}_{-2}(\alpha\mathbf{E}_\kappa(L))) \mathcal{A}d(K_0) \\ & + \mathbb{S}_{-1}(\alpha\mathbf{E}_\kappa(L)) \mathcal{A}d(\mathbb{S}(K_0)) \mathcal{A}d(K_0) \} \end{aligned}$$

where

$$\mathcal{A}d(K_0) = \begin{pmatrix} \frac{\kappa + 1}{2\kappa} & \frac{3\kappa + 1}{2\kappa} & \frac{\kappa - 1}{8\kappa} \\ \frac{-\kappa + 1}{4\kappa} & \frac{\kappa - 1}{4\kappa} & \frac{-\kappa + 1}{16\kappa} \\ -\frac{3\kappa + 1}{\kappa} & -\frac{(3\kappa + 1)^2}{(\kappa - 1)\kappa} & \frac{\kappa - 1}{4\kappa} \end{pmatrix}$$

and where

$$\mathcal{A}d(\mathbb{S}(K_0)) \mathcal{A}d(K_0) = \frac{1}{\kappa \kappa_1} \begin{pmatrix} \frac{(1 - \kappa_1)\kappa + \kappa_1 + 1}{2} & \frac{(1 - 3\kappa_1)\kappa^2 + 2(1 - \kappa_1)\kappa + 1}{2(\kappa - 1)} & \frac{(\kappa_1 + 1)(1 - \kappa)}{8} \\ \frac{(\kappa_1 - 1)(\kappa + 1)}{4} & \frac{(\kappa_1 - 1)(\kappa + 1)^2}{4(\kappa - 1)} & \frac{(\kappa_1 + 1)(1 - \kappa)}{16} \\ \frac{(3\kappa - 1)\kappa_1^2 + 2(\kappa - 1)\kappa_1 - \kappa - 1}{\kappa_1 - 1} & -\frac{(\kappa_1(3\kappa - 1) - \kappa - 1)^2}{(\kappa_1 - 1)(\kappa - 1)} & \frac{(\kappa - 1)(\kappa_1 + 1)^2}{4(\kappa_1 - 1)} \end{pmatrix}.$$

#### 4.6 The general solution

If we can solve for the discrete frame  $(\rho_k)$  then we have

$$x_k = \rho_k^{-1} \cdot \frac{1}{2} = \frac{d_k - 2b_k}{2a_k - c_k} \quad (4.14)$$

since  $\rho_k \cdot x_k = \frac{1}{2}$  is the normalisation equation.

The Adjoint representation of the Lie group  $G$  is, in this case, precisely that of the linear action discussed in §2, and so we may make use of the simplification of the algebraic equations for the group parameters stemming from the conservation laws,  $\mathbf{k} = V(I)\mathcal{A}d(\rho_0)$ , that we described there. However, the Maurer–Cartan matrix is different, and so the recurrence relations needed to complete the solution, differ. Nevertheless, we again find that the remaining recurrence relations are diagonalisable, and are therefore easily solved.

We again have equations (2.19a)–(2.19d), where now the  $V_0^i$  are those of equation (4.13); we rewrite these here for convenience,

$$k_1^2 + 4k_2k_3 - (V_0^1)^2 - 4V_0^2V_0^3 = 0, \quad (4.15a)$$

$$k_3c_0^2 - k_1c_0d_0 - k_2d_0^2 + V_0^2 = 0, \quad (4.15b)$$

$$2b_0V_0^2 - 2c_0k_3 + (k_1 - V_0^1)d_0 = 0, \quad (4.15c)$$

$$2a_0V_0^2 - c_0(k_1 + V_0^1) - 2k_2d_0 = 0. \quad (4.15d)$$

The recurrence relation is  $\rho_1 = K_0\rho_0$  or

$$\begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} = \sqrt{\frac{\kappa-1}{4\kappa}} \begin{pmatrix} 1 & \frac{1}{2} \\ -\frac{6\kappa+2}{\kappa-1} & 1 \end{pmatrix} \begin{pmatrix} a_0 & b_0 \\ c_0 & d_0 \end{pmatrix}$$

leading to the equations

$$c_1 = \sqrt{\frac{\kappa-1}{4\kappa}} \left( -\frac{6\kappa+2}{\kappa-1}a_0 + c_0 \right), \quad d_1 = -\sqrt{\frac{\kappa-1}{4\kappa}} \left( -\frac{6\kappa+2}{\kappa-1}b_0 + d_0 \right).$$

Using these to eliminate  $a_0$  and  $b_0$  from (4.15c) and (4.15d), leads to the linear system,

$$\begin{pmatrix} c_1 \\ d_1 \end{pmatrix} = Q\Lambda_0Q^{-1} \begin{pmatrix} c_0 \\ d_0 \end{pmatrix}$$

where  $Q$  is a constant matrix and  $\Lambda_0$  is diagonal. Indeed, setting

$$\mu = \sqrt{k_1^2 + 4k_2k_3}$$

we have

$$Q = \frac{1}{2\mu} \begin{pmatrix} \mu + k_1 & \mu - k_1 \\ 2k_3 & -2k_3 \end{pmatrix}.$$



Further,  $\Lambda_0 = \text{diag}(\lambda_0^1, \lambda_0^2)$  where

$$\lambda_0^1 = \frac{1}{2\sqrt{\kappa-1}\sqrt{\kappa}V_0^2} \left( -(3\kappa+1)(\mu+V_0^1) + (\kappa-1)V_0^2 \right),$$

$$\lambda_0^2 = \frac{1}{2\sqrt{\kappa-1}\sqrt{\kappa}V_0^2} \left( (3\kappa+1)(\mu-V_0^1) + (\kappa-1)V_0^2 \right).$$

We have then that

$$\begin{pmatrix} c_k \\ d_k \end{pmatrix} = Q \begin{pmatrix} \prod_{l=0}^{k-1} \lambda_l^1 & 0 \\ 0 & \prod_{l=0}^{k-1} \lambda_l^2 \end{pmatrix} Q^{-1} \begin{pmatrix} c_0 \\ d_0 \end{pmatrix}$$

where in this last,  $c_0$  and  $d_0$  are the initial data and  $\lambda_k^l = S_k \lambda_0^l$ .

Substituting these into the  $k^{\text{th}}$  shifts of (4.15c) and (4.15d), specifically,

$$2b_k V_k^2 - 2c_k k_3 + (k_1 - V_k^1) d_k = 0, \quad (4.16a)$$

$$2a_k V_k^2 - c_k (k_1 + V_k^1) - 2k_2 d_k = 0 \quad (4.16b)$$

yields the expressions for  $a_k$  and  $b_k$  needed to obtain, finally,  $x_k$  given in (4.14).

## 5. Conclusions

In these two papers we have introduced difference moving frames and applications to variational problems. We have also shown how to use these frames to solve Lie group invariant difference (recurrence) relations. We have considered relatively simple, solvable Lie group actions and some  $SL(2)$  actions. Open problems include the efficient use of difference frames for numerical approximations which preserve symmetry.

## Acknowledgements

The authors would like to thank the SMSAS at the University of Kent and the EPSRC (grant EP/M506540/1) for funding this research.

## References

- Gonçalves, T. M. N. & Mansfield, E. L. (2012), ‘On moving frames and Noether’s conservation laws’, *Studies in Applied Mathematics* **128**(1), 1–29.
- Gonçalves, T. M. N. & Mansfield, E. L. (2016), ‘Moving frames and Noether’s conservation laws – the general case’, *Forum of mathematics Sigma* (4), 55.
- Mansfield, E. L. (2010), *A practical guide to the invariant calculus*, Vol. 26 of *Cambridge Monographs on Applied and Computational Mathematics*, Cambridge University Press.
- Mansfield, E. L., Rojo-Echeburúa, A., Hydon, P. E. & Peng, L. (2019), ‘Moving frames and Noether’s finite difference conservation laws I.’, *Transactions of Mathematics and its Applications*, in press.