

Kent Academic Repository

Full text document (pdf)

Citation for published version

Gope, K. and Prabhudesai, V.S. and Mason, Nigel and Krishnakumar, E. (2015) Dissociative electron attachment to chlorine dioxide probed by velocity slice imaging. *Journal of Physics: Conference Series*, 635 (7). 072035. ISSN 1742-6588.

DOI

<https://doi.org/10.1088/1742-6596%2F635%2F7%2F072035>

Link to record in KAR

<https://kar.kent.ac.uk/74681/>

Document Version

Publisher pdf

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

PAPER • OPEN ACCESS

Dissociative electron attachment to chlorine dioxide probed by velocity slice im-aging

To cite this article: Krishnendu Gope *et al* 2015 *J. Phys.: Conf. Ser.* **635** 072035

View the [article online](#) for updates and enhancements.

Related content

- [Water disinfection agents and disinfection by-products](#)
J Ilavský, D Barloková, O. Kapusta *et al.*
- [Momentum imaging of dissociative electron attachment in biologically relevant molecules](#)
Marvin Weyland, Xueguang Ren, Thomas Pflüger *et al.*
- [Isotope effect in dissociative electron attachment to H₂ at 4 eV](#)
J Fedor, J Koišek and R Janeková



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Dissociative electron attachment to chlorine dioxide probed by velocity slice imaging

Krishnendu Gope^{* 1}, Vaibhav S Prabhudesai^{*}, Nigel Mason[†] and E. Krishnakumar^{* 2}

^{*} Dept. Nuclear and Atomic Physics, Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India

[†] Dept. of Physical Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK

Synopsis We report the study of dissociative electron attachment (DEA) to chlorine dioxide (OCIO) in the energy range of 1 to 10 eV using velocity slice imaging technique. DEA to OCIO shows Cl^- , O^- , O_2^- , and OCl^- as fragment channels at various electron energies. Based on the momentum images of these ions we unravel the molecular dynamics leading to DEA.

Chlorine dioxide (OCIO) is an important molecule in atmospheric chemistry as a major source of atomic chlorine in the stratosphere as it decomposes photo-chemically forming O_2 as the other product. Dissociative electron attachment (DEA) to OCIO can also lead to atomic chlorine formation.

DEA to OCIO has been studied in the past with reports of absolute cross-section measurements [1]. However, the dynamics that leads to DEA has not been reported so far. Here we report the study of DEA dynamics for OCIO using velocity slice imaging technique.

We observed Cl^- , O^- , O_2^- , and OCl^- ions from the DEA measurements on OCIO formed at various electron energies. The OCl^- being the most dominant channel was found to peak around 0.7 eV, whereas Cl^- and O_2^- signals peaked around 0.7 eV, 4 eV, 8 eV. Out of these the peak at 0.7 eV was found to be the dominant peak. The O^- ion yield was found to peak at 1.2 eV, 4 eV, 8 eV.

We carried out the momentum imaging of various fragment ions across different resonances using velocity slice imaging [2]. The VSI spectrometer was modified with longer flight tube and larger (75 mm diameter) phosphor screen based position sensitive detector [3]. For each of the ion fragments, we obtained the kinetic energy and angular distribution data. The velocity slice images obtained for Cl^- channel at 8 eV is shown in Fig. 1 as an example. For Cl^- channel, with very low energy threshold (-3.37 eV) [1], it was expected to carry good amount

of kinetic energy from the 8 eV resonance. However, the low kinetic energy observed in Cl^- channel indicates an interesting many body fragmentation dynamics behind the DEA process. The angular distribution also shows a forward-backward asymmetry.

In this poster we will describe the dynamics involved in DEA to OCIO based on the kinetic energies and angular distributions measured using velocity slice images for all the fragments observed at various resonances.




Figure 1. Momentum image of OCl^- , O^- and Cl^- from OCIO at 0.7eV, 4eV and 8eV electron energy respectively.

References

- [1] Gilbert Senn et al. 1999 *J. Phys. B At. Mol. Opt. Phys.* **32** 3615
- [2] Dhananjay Nandi, Vaibhav S. Prabhudesai, E. Krishnakumar and A. Chatterjee 2005 *Rev. Sci. Instrum.* **76** 053107
- [3] Ewelina Szyman´ska, Vaibhav S. Prabhudesai, Nigel J. Mason and E. Krishnakumar 2013 *Phys. Chem. Chem. Phys.* **15** 998

¹ E-mail: krishnendu@tifr.res.in

² E-mail: ekumar@tifr.res.in

