



# Kent Academic Repository

**Gope, K., Prabhudesai, V.S., Mason, Nigel and Krishnakumar, E. (2015)**  
*Dissociative electron attachment to molecular chlorine.* *Journal of Physics: Conference Series*, 635 (7). ISSN 1742-6588.

## Downloaded from

<https://kar.kent.ac.uk/74682/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1088/1742-6596/635/7/072024>

## This document version

Publisher pdf

## DOI for this version

## Licence for this version

CC BY (Attribution)

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

## Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

PAPER • OPEN ACCESS

## Dissociative electron attachment to molecular chlorine

To cite this article: Krishnendu Gope *et al* 2015 *J. Phys.: Conf. Ser.* **635** 072024

View the [article online](#) for updates and enhancements.

### Related content

- [Time-resolved detection of atomic chlorine by coherent anti-stokes Raman spectroscopy](#)  
Wang Lixin, Kang Jin, Huang Nantang et al.
- [XeF<sub>2</sub> gas-assisted focused-electron-beam-induced etching of GaAs with 30 nm resolution](#)  
A Ganczarzyk, M Geller and A Lorke
- [A dissociative electron attachment cross-section estimator](#)  
James J Munro, Stephen Harrison, Milton M Fujimoto et al.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

## Dissociative electron attachment to molecular chlorine

Krishnendu Gope<sup>\*1</sup>, Vaibhav S Prabhudesai<sup>\*</sup>, Nigel Mason<sup>†</sup> and E. Krishnakumar<sup>\*2</sup>

<sup>\*</sup> Dept. Nuclear and Atomic Physics, Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India

<sup>†</sup> Dept. of Physical Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK

**Synopsis** We report the study of dissociative electron attachment to molecular chlorine in the energy range of 1 to 20 eV using velocity slice imaging technique. Based on the angular distributions at various electron energies, we identify the partial waves involved in the resonance formation that lead to DEA.

Velocity slice imaging (VSI) is a well-established technique to investigate the molecular dynamics involved in dissociative electron attachment (DEA) to simple molecules [1]. This technique enables one to measure the kinetic energy and angular distribution of the fragment ions produced in the DEA process. The kinetic energy distribution of the fragment ions points to the kinematics of the process leading to DEA. On the other hand the angular distribution of the fragment negative ions with respect to the incoming electron beam indicates the symmetry of the negative ion resonant states involved in the DEA process. This information in combination with the kinetic energy distribution unravels the dynamics of the DEA process.


Due to their high electron affinity, the halogen molecules form a class of interesting diatomic molecules. These molecules are known to show more than one resonances in the DEA channel with very little threshold for the formation of halide anion. This leads to the substantial kinetic energy release in the fragmentation process in DEA.

Particularly for chlorine, the DEA studies are important from the plasma processing point of view. Chlorine is known to affect the properties of electrical plasmas considerably and hence the efficiency of technologies utilizing the plasma, such as those generated in either pure chlorine or mixtures of chlorine with other gases. Chlorine being a highly oxidizing agent much attention has been paid to the investigation of its properties.

Dissociative electron attachment to chlorine has been extensively studied in the past with reports of measurements of absolute cross-section [2] and angular distribution in the limited angular range [3]. However the kinetic energy distribution and the angular distribution for the entire 0-360° angular range have not been reported so far. Here we report

the study of DEA using velocity slice imaging at various resonances observed in the DEA to chlorine molecules [1]. The VSI spectrometer was modified with longer flight tube and larger (75mm diameter) phosphor screen based position sensitive detector [4].

In our measurements we measured the ion yield curve up to 90eV electron energy and we found that the chloride anion yield show three resonances at 0eV, 2.5eV and 5.6eV which is consistent with earlier measurements. The velocity slice images obtained at various electron energies across the resonances are shown in Fig. 1. In this poster we will describe the dynamics involved in DEA to chlorine molecule based on the kinetic energies and angular distributions obtained using these velocity slice images.



**Figure 1.** Image of Cl<sup>-</sup> from Cl<sub>2</sub> at 2.5eV, 4eV and 6eV electron energy.

### References

- [1] Dhananjay Nandi, Vaibhav S. Prabhudesai, E. Krishnakumar and A. Chatterjee 2005 *Rev. Sci. Instrum.* **76** 053107
- [2] M. V. Kurepa *et al.* 1978 *J. Phys. B: Atom. Mol. Phys.* **11** 3719
- [3] R. Azria *et al.* 1982 *J. Phys. B: At. Mol. Phys.* **15** L569
- [4] Ewelina Szyman'ska, Vaibhav S. Prabhudesai, Nigel J. Mason and E. Krishnakumar 2013 *Phys. Chem. Chem. Phys.* **15** 998

<sup>1</sup> E-mail: [krishnendu@tifr.res.in](mailto:krishnendu@tifr.res.in)

<sup>2</sup> E-mail: [ekumar@tifr.res.in](mailto:ekumar@tifr.res.in)

