



Kent Academic Repository

Mathie, Alistair, Clarke, Catherine E, Ranatunga, Kishani M and Veale, Emma L. (2003) *What are the roles of the many different types of potassium channel expressed in cerebellar granule cells?* Review of: *Cerebellum* by UNSPECIFIED. *Cerebellum* (London, England), 2 (1). pp. 11-25. ISSN 1473-4222.

Downloaded from

<https://kar.kent.ac.uk/75933/> The University of Kent's Academic Repository KAR

The version of record is available from

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

What are the roles of the many different types of potassium channel expressed in cerebellar granule cells?

Alistair Mathie, Catherine E Clarke, Kishani M Ranatunga, Emma L Veale

Biophysics Section, Blackett Laboratory, Department of Biological Sciences, Imperial College of Science Technology and Medicine, London, UK

Potassium (K) channels have a key role in the regulation of neuronal excitability. Over a hundred different subunits encoding distinct K channel subtypes have been identified so far. A major challenge is to relate these many different channel subunits to the functional K currents observed in native neurons. In this review, we have concentrated on cerebellar granule neurons (CGNs). We have considered each of the three principal super families of K channels in turn, namely, the six transmembrane domain, voltage-gated super family, the two transmembrane domain, inward-rectifier super family and the four transmembrane domain, leak channel super family. For each super family, we have identified the subunits that are expressed in CGNs and related the properties of these expressed channel subunits to the functional currents seen in electrophysiological recordings from these neurons. In some cases, there are strong molecular candidates for proteins underlying observed currents. In other cases the correlation is less clear. We show that at least 26 potassium channel α subunits are moderately or strongly expressed in CGNs. Nevertheless, a good empirical model of CGN function has been obtained with just six distinct K conductances. The transient K_A current in CGNs, seems due to expression of $K_v4.2$ channels or $K_v4.2/4.3$ heteromers, while the K_{Ca} current is due to expression of large-conductance slo channels. The G-protein activated K_{IR} current is probably due to heteromeric expression of $K_{IR}3.1$ and $K_{IR}3.2$. Perhaps $K_{IR}2.2$ subunits underlie the K_{IR} current when it is constitutively active. The leak conductance can be attributed to TASK-1 and or TASK-3 channels. With less certainty, the I_{K-slow} current may be due to expression of one or more members of the KCNQ or EAG family. Lastly, the delayed-rectifier K_v current has as many as six different potential contributors from the extensive K_v family of α subunits. Since many of these subunits are highly regulated by neurotransmitters, physiological regulators and, often, auxiliary subunits, the resulting electrical properties of CGNs may be highly dynamic and subject to constant fine-tuning.

Keywords:

Potassium channels, cerebellar granule cells/neurons, K_v , K_{IR} , two pore domain K channels, weaver mice

Mathie A, Clarke CE, Ranatunga KM, Veale EL.

What are the roles of the many different types of potassium channel expressed in cerebellar granule cells? *Cerebellum* 2003; 2: 11–25

Introduction

Potassium (K) channels play an important role in a number of different aspects of the electrical responses of the nervous system. K channel activity determines neuronal action potential frequency, shapes the neuronal action potential waveform and controls the strength of synaptic contacts between neurons.¹ Furthermore, K channels have a role in setting (or contributing to) the neuronal resting membrane potential and in regulating the excitability of individual neurons. Since the first

molecular cloning of K channel components in the late 1980s, well over a hundred different proteins—subunits of distinct types of K channels—have been identified and this number is still growing.² There are a number of good reviews that discuss particular aspects of K channels such as their structure, distribution and functional properties.^{2–14}

A major focus of our work concerns the physiological role of K channels in the CNS. We concentrate, largely on the cerebellum. However, even simply to describe the localization and function of K channels in the cerebellum is a daunting task. In their authoritative 1999 review, Coetzee et al.² described the tissue expression of 35 different principal (α) K channel subunits in the cerebellum. Since then, the number has grown to well over 40. In this review, then, we will concentrate primarily on cerebellar granule neurons (CGNs). We will describe the K currents seen, physiologically in native granule cells and correlate these data with information available from localization and distribution studies of the many K channel subunits expressed in these cells. By combining this information and building as complete

Received 8 March 2002; Revised 23 April 2002; Accepted 23 April 2002

Correspondence:

Alistair Mathie, Biophysics Section, Blackett Laboratory, Department of Biological Sciences, Imperial College of Science Technology and Medicine, Prince Consort Road, London SW7 2BW, UK.
Tel: +44 20 7594 7691. Fax: +44 20 7589 0191.
E mail: a.mathie@ic.ac.uk

a picture as possible of the channels and currents present in CGNs, we hope to provide a description of how multiple functioning K channels act together to regulate the excitability of these important neurons. While interesting in their own right, CGNs are often considered to be model CNS neurons, perhaps because of their abundance and comparative ease of isolation and identification. So, in addition, the information provided should give some idea of the level of complexity of K channels in other neurons in the CNS.

Cerebellar granule cells

Granule cells of the cerebellum are small glutamatergic interneurons that are by far the most numerous cell type in the brain. It is estimated that they number somewhere between 10^{10} and 10^{11} .¹⁵ CGNs receive inputs from mossy fibers or from the axons of brush cells.¹⁶ Granule cell axonal projections form the parallel fibers of the cerebellum and terminate on the dendrites of Purkinje neurons. It has been estimated that a single Purkinje neuron can receive inputs from 80,000 CGNs. Golgi cells, stellate cells and basket cells also receive inputs from granule cell parallel fibers.

Because of their abundance and their many interactions with other cell types in the cerebellum, CGNs play an important transduction role in cerebellar function. This is exemplified in the *weaver* mutation in mice (probably a mutation of an inward-rectifier K channel, K_{IR} 3.2; see below) where most CGNs fail to differentiate and then die, causing severe disruption of the cerebellar network and severe motor dysfunction.^{17,18} In normal, mature, CGNs, firing is most often fast and repetitive,¹⁹ but theta-frequency spike bursting can also be observed, which may play an important role in determining synchronization, rhythmicity and learning in the cerebellum.²⁰ Both the complement and the activity of granule cell potassium channels will profoundly determine the firing properties of these cells.

Potassium channel superfamilies

From the sequences of known potassium channel α subunits, it is clear that three major superfamilies of K channels can be delineated (Figure 1). They are the six transmembrane domain channels which are voltage- and/or calcium-gated; the two transmembrane domain inward-rectifiers and the four transmembrane, two-pore domain leak potassium channels.

The six transmembrane domain super family is made up of a number of different families, such as the voltage-gated K_V family of channels ($K_V1.x - K_V9.x$) which are thought to underlie voltage-gated delayed-rectifier and A-type K channels. KCNQ and EAG channels are also in this super family. These are low-threshold voltage-gated currents which are G-protein regulated and which often have a non-inactivating component open near the resting membrane potential of the cell. Functionally, in neurons, these channels underlie currents such as the M current. Finally in this super family are the slo and SK families of K channel which underlie functionally observed calcium-activated K currents. The two transmembrane domain super family of inward-rectifier K channels (K_{IR} 1.x – 7.x) underlie strong inward-rectifier currents, inward-rectifier currents activated by G-protein coupled receptors and, together with sulphonylurea receptors, ATP-sensitive K channels. The more recently described four transmembrane, two-pore domain K channels (such as TWIK-1, TASK-1 and TREK-1) are thought to underlie leak currents open at all potentials and are expressed heterologously throughout the CNS.

Six transmembrane domain K channels

Functional studies of voltage-gated K currents

In functional studies, the voltage-gated K current of mature CGNs is usually divided, quite straightforwardly, into two components: a transient component (or A-type current) which activates then inactivates rapidly




Figure 1
Potassium channel superfamilies.




Figure 2
 K_V and K_A currents in rat cerebellar granule cells. Currents were evoked by progressive 10 mV depolarizing steps for 85 ms from a holding potential of -70 mV following a 250 ms prepulse to either -140 mV (left hand side) or -50 mV (right hand side). The currents on the left-hand side following a hyperpolarizing prepulse have both a transient and a sustained component while those on the right-hand side following a depolarizing prepulse have only a sustained component.

following depolarization and a sustained current (or delayed-rectifier) which shows little inactivation over the time course of a normal depolarizing voltage step (see Figure 2).

These two components of voltage-gated outward current have been well characterized by a number of different groups since their first description in the late 1980s, using whole-cell voltage clamp recording methods from both CGNs in tissue culture and acutely isolated cells in cerebellar slices.^{21–29}

The sustained current activates with a time constant of about 10 ms at test potentials more depolarized than around -50 mV and shows no inactivation over a period of at least 150 ms.^{23,26} It is moderately susceptible to block by tetraethylammonium ions (TEA), being blocked by around 60% by 5 mM TEA,²⁷ 82% by 10 mM TEA³⁰ and completely blocked by 20 mM TEA.^{23,26} The sustained current, in some instances, can be further separated into a calcium-independent component and a calcium-dependent component²³ (also see below).

Removal of inactivation of the transient current, I_{A_s} , by use of a hyperpolarizing prepulse, followed by step depolarizations to potentials positive to -60 mV, reveals a current that peaks in 1–2 ms then inactivates with a time constant of around 10 ms.²³ This current is comparatively insensitive to TEA. 4-Aminopyridine (4-AP, 1–5 mM) reduces both the transient and sustained component of current^{23,27} consistent with its lack of selectivity between most members of the K_V family.^{2,10}

K_V channel distribution

In a simple world, CGNs would express two K_V channel subunits, one with the characteristics of the sustained component of current and the other with the character-

istics of the transient current. Unfortunately, the true picture is considerably more complicated. α -subunits from each of the four major classes of K_V channels ($K_V1.x - K_V4.x$) are strongly expressed in CGNs (Table 1).

K_V1 channels

CGNs have been shown to have strong immunoreactivity for $K_V1.1, 1.3, 1.5$ & 1.6 channels.³¹ In the absence of accessory subunits, all four of these clones give functional currents that resemble delayed-rectifier type, slow-inactivating currents. Chung et al.³¹ also found low immunoreactivity for $K_V1.2$ and no staining for $K_V1.4$. These results correlate reasonably well with earlier labeling studies (see Table 1 and refs 32–35), however earlier studies have found no labeling for $K_V1.6$ in CGNs.^{32,36,37} In a separate study, good $K_V1.1$ staining was seen in cell bodies of adult granule cells, however expressed channel numbers were seen to decline in aged rats.³⁸ This study, again, found much weaker labeling for $K_V1.2$ in granule cell bodies and that this didn't change with age. It is of interest that in other regions of the cerebellum, notably basket cell termini there is strong expression of $K_V1.2$ channel subunits.^{39,40} This suggests a role of these subunits in the regulation of transmitter release, functioning either as homomers or even heteromers with other K_V1 subunits.

The K_V1 family is often associated with regulatory β subunits which alter their functional properties from those expected of homomer channels. For example, $K_V\beta3.1$ is highly expressed in cerebellum and converts $K_V1.5$ from a non-inactivating delayed-rectifier type current to an inactivating current.⁴¹ Similarly $K_V1.1$ and $K_V\beta1$ are expressed in CGNs.^{35,42} Again $K_V\beta1$ is known to convert $K_V1.1$ from a non-inactivating to an inactivating current if the two subunits are co-expressed.⁴³ It is worth noting however that CGNs express both $K_V\beta1$ and $K_V\beta2$ at comparatively low levels compared with other regions of the brain.⁴⁴ The labelling for $K_V\beta1$ is much less intense than that seen in Purkinje cells.⁴⁵

Mutations in $K_V1.1$ channels give rise to episodic ataxia (or familial periodic cerebellar ataxia, a dominantly inherited human cerebellar disease) and to myokymia syndrome. These are associated with non-functional or malfunctioning $K_V1.1$ channels^{46,47} perhaps because of the normal role of this subunit in regulating granule cell excitability. $K_V1.1$ subunits are often expressed at synaptic terminals and dendrites and such a distribution is also seen in CGNs. A reduction in $K_V1.1$ channel activity will lead, therefore, to a prolongation of action potential duration.

K_V2 channels

$K_V2.1$ and $K_V2.2$ channels have distinct non-overlapping distributions in the mammalian brain. Both channels, when expressed as homomers, give rise to

Table 1Expression of K_v channel α subunits in cerebellar granule cells

Channel	Synonym	Species	Age	RNA/protein	Level	References
K _v 1.1	RCK1	rat mouse	P7/ adult	RNA/protein	moderate/ abundant	31, 32, 34, 35, 38, 39 (but see 37)
K _v 1.2	RCK5	rat mouse	young/ adult	RNA/protein	low	31, 32, 33, 34, 38, 39, 145
K _v 1.3	RCK3	rat	adult	RNA/protein	moderate	31, 32, 34
K _v 1.4	RCK4	rat	adult	RNA/protein	none	31, 32
K _v 1.5	RCK7	rat	adult	RNA/protein	moderate	31, 32
K _v 1.6	RCK2	rat	adult	RNA/protein	none	32, 36, 37 (but see 31)
K _v 2.1	DRK1	rat	adult	RNA	low	32, 36, 48
K _v 2.2	CDRK	rat	adult	RNA	abundant	48
K _v 3.1	Raw2	rat mouse	adult	RNA/protein	abundant	32, 36, 50, 51, 55
K _v 3.2	Raw1	rat	adult	RNA	low	32
K _v 3.3		mouse	adult	RNA	none	50, 144
K _v 3.4	Raw3	rat	adult	RNA/protein	low	32, 34
K _v 4.1		rat	adult	RNA	none	57
K _v 4.2		rat mouse	P3–P10/ adult	RNA	highly abundant	32, 53, 56, 57
K _v 4.3		rat	adult	RNA	abundant	56, 57
K _v 5.1	IK8	rat	adult	RNA	none	36, 37
K _v 6.1	K13	rat	adult	RNA	moderate/ abundant	36, 37
K _v 8.1		rat hamster	adult	RNA	abundant	60, 61
K _v 9.1		mouse	adult	RNA	abundant	63 (see also 64)
K _v 9.2		mouse	adult	RNA	abundant	63
K _v 9.3		mouse	adult	RNA	in brain	64

Note: see text for information on KCNQ, EAG, slo and SK channel α -subunit expression in CGNs.

delayed-rectifier currents with slow inactivation and moderate sensitivity to block by TEA. K_v2.2 expression is high in CGNs⁴⁸ while low levels of K_v2.1 are seen.^{32,36,48} Expression levels of K_v2.2 can be enhanced up to six-fold by mK_v β 4 with no change in channel properties. mK_v β 4 is also expressed in CGNs.⁴⁹ Interestingly K_v5.1, 6.1, 6.2, 8.1, 9.1, 9.2, & 9.3 give no functional currents in expression systems but can form heteromers with K_v2.x channels changing their properties² (see below).

K_v3 channels

K_v3.1 channels are expressed in the somata of CGNs, along the entire length of their axons (the parallel fibers) and in presynaptic boutons, however there is low or no expression in dendrites and glomeruli.^{32,36,50–53} In cerebellar slices, application of TEA at a concentration that preferentially blocks K_v3.1 channels resulted in parallel fiber action potential broadening and an increase in excitatory post-synaptic currents at granule cell–Purk-

inje cell synapses.⁵⁴ Expression of the K_v3.3b channel is highly enriched in the brain, particularly in the cerebellum, where its expression is confined to Purkinje cells and deep cerebellar nuclei with little expression in CGNs. There are also detectable levels of K_v3.2 and 3.4 found in CGNs.^{32,34} Expressed K_v3 channels have extremely rapid kinetics of activation and deactivation and a high threshold of activation (more positive than –20 mV). The deactivation of K_v3 channels is at least ten-fold faster than all K_v1 channels. As a result, K_v3 channels are thought to underlie the fast after-hyperpolarization required for high frequency firing. They are highly sensitive to block by TEA.¹⁰

K_v3.1 and K_v3.3 knock-out animals exist, which on their own show little phenotype.⁵⁵ However, severe ataxia is seen when both channels are mutated in the same animals. This is thought to result from a combination of a lack of K_v3.1 in CGNs and a lack of K_v3.3 in Purkinje cells.⁵⁵ Surprisingly, there is no gross alteration in cerebellar morphology associated with the double mutant.

K_V4 channels

CGNs express K_V4.2 and 4.3 but not K_V4.1 channels.^{32,36,53,56,57} Both K_V4.2 and 4.3 give rise to transient A-type potassium conductances in expression systems. A number of studies have suggested that K_V4 subunits are the major component of transient I_A currents in the mammalian CNS.^{56,57} K_V4 channels can be regulated by a family of K channel interacting proteins (KChIPs⁵⁸). These act to increase total K_V4 current, accelerate recovery from inactivation and moderately slow inactivation. They are calcium-binding proteins some of which (KChIP1 and KChIP3) are predominantly expressed in neurons. Another member of this family KChIP4a, acts to remove inactivation of K_V4 currents.⁵⁹ Both KChIP1 and KChIP4a are expressed in CGNs⁵⁹ and can act in combination to modify K_V4 channel properties.

K_V5-9 subunits

There are several K_V subunits (K_V5.1, 6.1, 6.2, 8.1, 9.1, 9.2 and 9.3) that do not form functional channels when expressed on their own but do seem to form heteromers with K_V2.x channels giving them different functional properties from homomeric K_V2.x channels.²

In CGNs, the more important of these subunits seem to be K_V8.1, K_V9.1 and K_V9.2. K_V8.1 is known to be present in the granule cell layer of the cerebellum.^{60,61} At high levels of expression it abolishes the function of other K_V channels (K_V2.1, 2.2 and 3.4) via association with these subunits through its N terminal domain. At lower levels it modifies the activity of K_V2.1 and 2.2. The major effect of the co-expression of K_V8.1 with K_V2.2 in COS cells concerns the inactivation process. The voltage-dependence of K_V2.2 is shifted toward negative potentials, and its rate of inactivation is considerably slowed down. Clearly then, any K_V2.2/K_V8.1 heteromers present in CGNs would have kinetic properties different from K_V2.2 homomers.⁶²

There are also high levels of K_V9.1 and 9.2 expression in cerebellar granule cells.⁶³ K_V9.1 and 9.2 cause minimal changes in inactivation of K_V2.2 channels but reduce the current amplitude and shift the voltage-dependence of inactivation of the channel to more negative values. K_V9.3 channels are expressed in the CNS, but their expression levels and functional importance in CGNs is not known.⁶⁴

K_V5.1, K_V6.1 and K_V6.2 are known to regulate K_V2.1 functional properties.^{63,65} However, K_V5.1 is not expressed in the cerebellum to any degree.^{36,37} K_V6.1 is highly expressed in CGNs^{36,37} but the presence or absence of K_V6.2 and the functional importance of K_V6.1 remains to be established.

Correlation of functional currents and expressed K_V subunits

Considering homomers of the primary subunits alone, a

plausible case can be made for seven different voltage-gated K_V channels (1.1, 1.3, 1.5, 2.2, 3.1, 4.2 and 4.3) as being major contributors to the whole-cell voltage-gated K current in CGNs. If heteromers can be formed, or if regulatory subunits have a prominent role, the situation is even more complicated. How can we reconcile these data with the two primary components of current, transient and sustained, observed in functional studies from CGNs?

The best correlative evidence currently available concerns the transient component of current. Shibata et al.³⁵ have constructed dominant negative cDNA for the K_V4.2 subunit. When this is expressed in CGNs it results in a large reduction in A current in these cells. This provides strong evidence that the native A current is carried through K_V4.2 channels, however an interaction with K_V4.3 channels cannot be ruled out. The functional properties of K_V4.2 channels in expression systems are broadly similar (but not identical) to those seen of native A currents in the CGNs.^{26,35} Such differences may be accounted for by the multiple regulatory actions of KChIPs on K_V4 channels in CGNs.⁵⁹ Nevertheless, either K_V4.3 homomeric channels or heteromers of K_V4.2 and 4.3 channels may contribute to the A current. K_V1.4 subunits which give A currents in expression systems, are not expressed in CGNs, however co-expression of K_V1.1³⁵ with K_Vβ1⁴² could give rise to a functional A current in CGNs as could co-expression of K_Vβ3.1 with K_V1.5 (see above). Any of these possibilities could explain the residual 25% of transient current seen in CGNs treated with dominant negative K_V4.2.³⁵

The delayed-rectifier component is much more difficult to assign to a particular K_V subunit. Since loss of either K_V1.1 or 3.1 in CGNs can lead to the generation of episodic ataxia, this would suggest that both these subunits may be functionally important. The whole-cell granule cell sustained current, however, is much less sensitive to block by TEA (60% at 5 mM) when compared with K_V1.1 and K_V3 channels which have, respectively, IC₅₀s for block of 0.5 and around 0.2 mM. By contrast, K_V1.3 and 1.5 are rather too insensitive to TEA block to be candidates for the sustained current. This leaves K_V1.6 and K_V2.2, which have IC₅₀s of around 2–7 mM and 8 mM respectively for TEA block.^{2,10} This criterion, taken together with the distribution studies (Table 1 and above), would suggest that K_V2.2 is the most likely candidate to underlie the sustained voltage-gated current in CGNs since there is some doubt as to whether K_V1.6 is expressed.

Of course the whole-cell sustained current in CGNs may easily comprise of more than one expressed α subunit. Furthermore, the existence of splice variants and the potential formation of heteromers between different α subunits or between α and β subunits adds complexity. All K_V channels have consensus sequence sites for serine/threonine phosphorylation, while at least some can be phosphorylated on tyrosine residues (see for example refs 66, 67). These phosphorylation events

can markedly alter channel properties (see ref. 10), which would make identification even more difficult. Finally, it is likely that different K_V channels are expressed in different regions of CGNs. For example, $K_V1.1$ and 3.1 expression may be predominantly localized to presynaptic terminals and parallel fibers.

EAG and KCNQ channels

The EAG and KCNQ family of K channels encode voltage-gated potassium channels that are open close to the resting membrane potential of neurons.^{2,12,13} Both EAG currents and KCNQ currents can be inhibited following the activation of certain G-protein coupled receptors such as muscarinic acetylcholine receptors.

EAG family

The different members of the EAG family of K channels have quite distinct kinetic and pharmacological properties when studied in expression systems.¹² They can be divided into three subfamilies (eag 1–2; elk 1–3 (eag-like) and erg 1–3 (eag-related)) on the basis of their distinct structural and functional properties. One noticeable feature, however, is that all eight known members of the EAG family give a large component of non-inactivating current near the membrane potential.⁶⁸ Interestingly, like K_V channels, they may form heteromers within subfamilies.

In situ hybridization studies in CGNs reveals very high levels of eag1, but no detectable eag2; moderate levels of erg1 but no erg2 or erg3; and moderate to high levels of elk2, but no elk1 or elk3.^{68,69} Interestingly then, there is no overlap of expression of members of the same EAG subfamily in CGNs, therefore there is probably no heteromer formation between different EAG channel subunits.

In general, roles for EAG potassium channels in the mammalian CNS remain to be found.⁶⁸ However Hoshi et al.⁷⁰ have suggested that the non-inactivating current, $I_{K_{SO}}$, seen in CGNs (ref. 71 and below) may be due to co-expression of eag1 with a novel auxiliary subunit KCR1 (see also refs 72, 73).

KCNQ family

KCNQ channels in neurons are also thought to encode for non-inactivating voltage-gated currents with a low threshold for activation. For example, KCNQ2/KCNQ3 heteromers have been suggested to underlie M currents in a number of different neurons.^{13,74} Each of these KCNQ subunits when expressed alone results in much less M-like current than when the two are co-expressed.⁷⁴ Saganich et al.⁶⁸ showed very high levels of KCNQ2 expression, yet no or very little KCNQ3 expression in CGNs (see also ref. 75). In many neurons expression of KCNQ2 and KCNQ3 overlaps, so it is surprising that this does not seem to occur in CGNs.

There is little expression of KCNQ1 channels in the brain,² while little or no KCNQ4 or KCNQ5 expression is detected in CGNs.^{76–78} So KCNQ2 is the only detectable KCNQ subunit identified so far in CGNs.

A recent study has shown that CGNs in cerebellar slices possess a novel slow K conductance, which in many respects resembles the M current seen in other neurons (see Figure 3). This current was resistant to 4-AP (4 mM) and TEA (20 mM) but reduced by external Ba^{2+} or internal Cs^+ ions.²⁰ It is possible that this current results from the expression of KCNQ2 and/or one or more of the EAG family of subunits expressed in CGNs.

Ca-activated K channels

Ca-activated K channels can be divided into three broad groups. The large conductance maxi-K channels (or BK channels) corresponding to the slo family of K channels subunits; the small-conductance or SK channels (corresponding to SK1–3) and the intermediate conductance K channels which probably correspond to SK4 (also referred to as $I_{K_{Ca1}}$) channels.




Figure 3

A slow outward K current (I_{K-slow}) in cerebellar granule cells. (A) I_{K-slow} was generated by a step from -80 mV to $+30$ mV in the presence of 20 mM TEA and 4 mM 4-AP. It was reversibly inhibited by Ba^{2+} (1 mM). (B) Activation of I_{K-slow} by a series of 1 sec, 10 mV depolarizing steps from -80 mV. Adapted from D'Angelo et al. (2001), ref. 20, with permission. Copyright 2001, the Society for Neuroscience.

BK/slo channels

Functional studies in CGNs have shown the presence of a Ca-activated K conductance. Originally this was seen in single-channel recordings where a large conductance channel was observed with a single channel conductance of 110–185 pS.⁷⁹ This would suggest that the channel is a member of the BK class of Ca-activated K channels. BK channels in CGNs are activated following metabotropic glutamate receptor activation⁷⁹ via ryanodine receptors coupled to L-type Ca channels.⁸⁰ A corresponding calcium-dependent whole-cell K current has been observed in granule cells in cerebellar slices²⁶ and it is believed that BK channels participate in granule cell repolarization following an action potential.²⁰ BK channels may also stabilize repetitive firing by enhancing fast after hyperpolarizing potential and Na channel de-inactivation.²⁰

It is generally envisaged that such BK channels result from expression of a member of the slo family of K channel subunits.^{81,82} However expression levels of slo are rather low in CGNs when compared with other regions of the brain, including cerebellar Purkinje cells.^{83–85}

SK channels

There are only moderate levels of SK2 expression in CGNs, while no SK1 or SK3 could be detected by *in situ* hybridization.⁸⁶ SK4 (or IK_{Ca1}) channels are thought to be predominantly expressed in non-excitable tissue.⁸⁷ Furthermore, there is no effect of apamin on whole-cell K currents in CGNs, which suggests that SK channels may not have a major functional role in these cells.²⁰

Two Transmembrane Domain K channels

Inward-rectifier currents in granule cells

A G-protein activated inwardly-rectifying K current is evident from studies in cultured CGNs from mouse.^{88–92} The current is activated by GTP- γ -S, baclofen, somatostatin and trans-APCD and blocked by Ba^{2+} , Cs^+ and treatment with *pertussis* toxin. A study in mouse CGNs *in situ* showed that the inward-rectifier current changed its properties through the time-course of cell development⁹³ (see Figure 4). The majority of cells in the internal granule cell layer, 10–15 days after birth, displayed only a G-protein activated inwardly-rectifying current similar to that seen in the cultured cells. However, a constitutively active inward current was observed in the remainder of cells. CGNs at a more advanced developmental stage, 20–21 days after birth, displayed, for the most part, this latter current. Both currents share similar pharmacological properties, but they exhibit different biophysical characteristics. The G-

protein activated current displays mostly fast activation with no inactivation. The fast activation appears almost voltage-independent and instantaneous, similar to observations in cultured CGNs.^{88,89} Meanwhile, the constitutively active current has slower activation and a steeper voltage-dependence. Note, also, that there is no evidence for any I_h -like current in cerebellar granule cells.²⁰

Distribution and localization of K_{IR} subunits

In terms of the G-protein activated $K_{IR}3$ channels, a variety of different methods have revealed the presence of $K_{IR}3.1$, 3.2 and 3.3 subunits and lower levels of $K_{IR}3.4$ subunits in CGNs of both rat and mouse^{94–100} (see Table 2). Co-immunoprecipitation studies show direct physical association of $K_{IR}3.1$ and $K_{IR}3.2$ in COS cells¹⁰¹ and mouse brain tissue⁹⁵ suggesting that they exist and function as heteromers.

There are high levels of the strong inward-rectifier $K_{IR}2.2$ in CGNs but no detectable levels of $K_{IR}2.3$ or 2.4 subunits^{96,102,103} (see Table 2). It is not clear from the literature whether $K_{IR}2.1$ subunits are present in CGNs.^{96,102} For ATP-sensitive K channels, *in situ* hybridization also reveals weak mRNA signal for $K_{IR}6.2$ (but no detectable $K_{IR}6.1$) and low levels of its associated subunit SUR1 in CGNs.^{104,105}

A number of K_{IR} channel subunits ($K_{IR}1.1$, 4.1, 4.2 and 7.1) are not expressed in CNS neurons.⁹⁷ Furthermore, $K_{IR}5.1$, although expressed in the CNS, cannot function as a homomeric channel, as it seems to only form functional heteromers with $K_{IR}4.1$ or 4.2.¹⁰⁶




Figure 4
 K_{IR} currents in mouse cerebellar granule cells. Recordings were performed with patch pipettes containing GTP- γ -S. Traces were taken at the beginning (control) and 6 min (GTP- γ -S) after the beginning of the recordings. The voltage-clamp protocol is illustrated at the bottom. Cell 1 was recorded at P12 and is an immature postmigratory neuron. It has no constitutively active current but has an inward-rectifier current activated by GTP- γ -S. Cell 2 was recorded at P22 and is a mature postmigratory neuron. It has a constitutively active inward-rectifier current which is not enhanced further by GTP- γ -S. Adapted from Rossi et al. (1998), ref. 93, with permission. Copyright 1998, the Society for Neuroscience.

Table 2.
Expression of K_{IR} channel α subunits in cerebellar granule cells

Channel	Synonym	Species	Age	RNA/protein	Level	References
$K_{IR}1.1$	ROMK	rat		RNA	not in brain	97
$K_{IR}2.1$	IRK1	rat	adult	RNA/protein	low	96, 99
		mouse				(but see 102)
$K_{IR}2.2$	IRK2	rat	P21-adult	RNA	abundant	96, 102
		mouse				
$K_{IR}2.3$	IRK3	rat	E13-adult	RNA	none	96, 102
		mouse				
$K_{IR}2.4$	IRK4	rat	adult	RNA	none	103
$K_{IR}3.1$	GIRK1	rat	E15-adult	RNA/protein	moderate	95, 96, 97, 99
		mouse				(but see 146)
$K_{IR}3.2$	GIRK2	rat	E13-adult	RNA/protein	abundant	94, 95, 96, 97, 98
		mouse				100, 146
$K_{IR}3.3$	GIRK3	rat	E15-adult	RNA	abundant	96, 97, 146
		mouse				
$K_{IR}3.4$	GIRK4	rat	adult	RNA/protein	none	96, 100
$K_{IR}4.1$	BIR10	rat		RNA	not in brain	97
$K_{IR}4.2$		rat		RNA	not in brain	97
$K_{IR}5.1$	BIR9	rat		RNA	low in brain	97 (see 106)
$K_{IR}6.1$	uKATP-1	rat	adult	RNA/protein	none	105
$K_{IR}6.2$	BIR IKATP	rat	adult	RNA	low	104
		mouse				
$K_{IR}7.1$		rat		RNA	not in brain	97

Functional properties of K_{IR} subunits

$K_{IR}3.2$ homomultimers in *Xenopus* oocytes display G-protein activated, K selective currents which are enhanced by carbachol, acetylcholine and $G\beta\gamma$ ^{88,101,107} $K_{IR}3.1$ homomers in *Xenopus* oocytes produce similar currents but with slower activation kinetics.¹⁰⁷ Currents elicited by the presence of both $K_{IR}3.1$ and $K_{IR}3.2$ are also enhanced by G-protein coupled receptor activation.¹⁰¹ They exhibit slow kinetics, attributed to the presence of $K_{IR}3.1$.¹⁰⁷ Similar currents are obtained by engineered $K_{IR}3.1/K_{IR}3.2$ dimers.¹⁰⁷ It is thought likely that, *in vivo*, these channels form as heteromers.

Correlation of the channel subtypes present with functional properties

The activation of the inwardly-rectifying currents by GTP- γ -S and various GABA_B and muscarinic agonists in CGNs, identify it as composed of $K_{IR}3.x$ channels. The slower activation in response to hyperpolarization suggests the presence of the $K_{IR}3.1$ subunit. Considered together with the distribution and localization studies and the *weaver* mutation below, it is most likely that the dominant inward-rectifier K current in CGNs is due to the expression of $K_{IR}3.1$ and $K_{IR}3.2$ heteromers. There may also be a contribution of $K_{IR}3.3$ subunits. The developmental change from G-protein activated to constitutive, as well as from generally fast to slower activa-

tion kinetics in CGNs *in situ* is not understood. It may be that $K_{IR}2.2$ channels underlie the constitutively active current, since CGNs express this subunit in abundance (Table 2). At present, no information is available on developmental expression of $K_{IR}2.2$.

The *weaver* mutation

The *weaver* mutation of mouse (*wv*) arose spontaneously in C7BL/6 mice over 30 years ago and was identified by, and named after, its instability of gait. Homozygous (*wv/wv*) mice exhibit mild locomotor hyperactivity, male sterility, and severe ataxia within two weeks of birth. The cerebellum is smaller in size with the loss of CGNs occurring within one week of birth. Rakic & Sidman¹⁷ observed that *wv* granule cell precursors are normal at birth, but display defects through differentiation. They fail to extend neurites or migrate along glial fibres to form bipolar processes.

The *wv* gene was mapped to mouse chromosome 16, a region associated with Down's Syndrome in humans,¹⁰⁸ and sequencing revealed a missense mutation in a region coding for $K_{IR}3.2$.¹⁸ The mutation, G156S (nucleotide G935A), resides in the 'H5' or 'p' region that is highly conserved amongst K channels and structurally forms the selectivity filter of the channel.¹⁰⁹ The gene is autosomal recessive,¹⁰⁷ and heterozygous mice (*wv/+*) show milder abnormalities than homozy-

gous ones. However, it should be noted that $K_{IR3.2}$ knock-out mice ($-/-$), whilst still more susceptible to induced seizures than wild-type (WT), remain morphologically similar to WT and exhibit milder cerebellar abnormalities than *wv*. This would suggest that the *wv* phenotype may not simply be due to the loss of functional $K_{IR3.2}$ channels.

wvK_{IR3.2} channels expressed in cell lines display altered currents to $WT K_{IR3.2}$ (see Figure 5). The cells themselves, whether CHO cells or *Xenopus* oocytes, show decreased survival.¹⁰¹ The currents become constitutively active and non-selective for K^+ .^{88,101,107} Differing results have emerged from studies on cultured CGNs. Some studies observe a basal, non-selective current that depolarizes the cell's resting potential,^{88,107} analogous to the constitutively active current observed in *wvK_{IR3.2}*-containing channels in oocytes. This current can be blocked by the cation channel blockers MK-801 and QX-314, which are also able to enhance cell viability and neurite outgrowth.⁸⁸ However, other studies reveal a profound reduction in G-protein activated currents¹⁰⁷ or no inwardly-rectifying currents with or without G-protein activation.^{91,92} Slices of *wv* mouse cerebellum display altered architecture with no discernible internal granule cell layer. In slices, pre-migratory *wv*CGNs do not exhibit any G-protein activated inward-rectifier currents. Putative CGNs that have escaped degeneration express a constitutively active inwardly-rectifying current with reduced selectivity to K that is blocked by QX-314. The kinetics are slower, compatible with *wvK_{IR3.1}* homomultimers. However there is no net increase in leakage current.⁹³

With such a variety of results and no explanation for the differences, no clear hypothesis of mechanism emerges. However, the dominant effect is of a depolar-

ized granule cell resting potential, whether this occurs by a gain of basal Na conductance or a loss of agonist-activated K conductance. This could lead to cell death or failure to drive cell differentiation and motility via the activation of voltage-gated Ca^{2+} channels and NMDA receptors. Alternatively *wvK_{IR3.2}* may act less directly, altering the expression of other genes or the mutation in $K_{IR3.2}$ may be merely closely associated with the *wv* gene and be a coincidental finding.⁹²

Four transmembrane domain/two-pore domain K channels

Leak currents in granule cells

CGNs possess a non-inactivating outward potassium current that is active at all membrane potentials. This current, termed IK_{SO} (for standing-outward current), which has a major role in regulating the excitability of these cells, is openly rectifying and is inhibited by the activation of muscarinic acetylcholine receptors^{71,110,111} (see Figure 6). IK_{SO} is insensitive to the classical K channel blockers tetraethylammonium and 4-aminopyridine, but can be blocked by Ba^{2+} , small extracellular acidification¹¹² and the endocannabinoid, anandamide.¹¹³ It is enhanced by the volatile anaesthetic agent halothane.¹¹³ Inhibition of this current by muscarinic receptor activation or extracellular acidification increases granule cell excitability^{71,112} (see Figure 7). CGNs have also been shown to possess a background K channel that can be activated by arachidonic acid and other polyunsaturated fatty acids.¹¹⁴ Activation of this current by arachidonic acid reduces granule cell excitability.¹¹⁴




Figure 5 Reduction of K_{IR} currents in mouse cerebellar granule cells from *weaver* mice. Examples of GTP- γ -S-activated currents recorded from wild-type (A), *weaver* (B), and Girk2 null (C) mutant mice are shown. GTP- γ -S-activated currents are markedly reduced in granule cells isolated from *weaver* and Girk2 null mutant mice. Currents were elicited by voltage steps from 130 to +10 mV from a holding potential of 0 mV. The GTP- γ -S-activated current was obtained by subtracting the current recorded at the beginning of the whole-cell recording from the current recorded after GTP- γ -S activation. (D) Current-voltage plots show the strong rectification for +/+, +/-, and +/ww granule cells. (E) Bar graph shows the mean amplitude of the GTP- γ -S-activated currents divided by the cell capacitance for +/+, +/ww, ww/ww, +/-, and -/- granule cells. Adapted from Slesinger et al. (1997), ref. 90, with permission. Copyright 1997, the National Academy of Sciences (USA).




Figure 6
Muscarinic ACh receptor activation inhibits $I_{K_{SO}}$. Granule cells were held at -20 mV and stepped to -60 mV for 400 ms. This protocol was repeated once every 5 seconds. The upper panel shows currents recorded in the absence and presence of muscarine ($10 \mu\text{M}$). The time course for muscarinic inhibition of $I_{K_{SO}}$ is shown in the lower panel. $I_{K_{SO}}$ was measured at -20 mV after the CGNs had been at this potential for 4.6 seconds. Adapted from Boyd & Mathie (2002), ref. 110.

Two pore domain potassium channels and localization in CGNs

Recently a group of K selective pore-forming subunits with four transmembrane domains and two pore domains per subunit have been discovered. Named two pore domain potassium (2PK) channels, due to this unique feature, it is these channels that are thought to underlie the “leak” potassium currents that have been recorded in CGNs. To date, 14 members of the mammalian 2PK family have been described.^{14,115,116} These subunits are thought to form together as dimers to provide the four P domains required to produce a functional channel.¹⁴ They can be grouped, loosely, into a number of different classes on the basis of structural and functional properties. TWIK-1 (tandem of P domains in a weak-inward-rectifier K channel) was the first to be cloned¹¹⁷ and all subsequent channels have been named in relation to this one. A separate nomenclature refers to this channel as KCNK-1.¹⁴




Figure 7
 $I_{K_{SO}}$ blockade increases granule cell excitability. In current clamp recording, injection of 100 pA of current depolarized the cerebellar granule cell but did not evoke action-potential firing (A). Application of $10 \mu\text{M}$ muscarine blocked $I_{K_{SO}}$ and increased granule cell excitability such that the same amount of current injection now depolarized the cell sufficiently to evoke action-potential firing (B, C). Adapted from Watkins & Mathie (1996), ref. 71.

TWIK-1 and TWIK-2^{118,119} as their names suggest, are thought to be weak inward-rectifier K channels but there is some doubt about whether they can form functional channels independently of auxiliary subunits (e.g. ref. 14). A number of 2PK channels have been shown to be sensitive to small acidifications of the external solution and have been termed TWIK-related acid-sensitive K channels (TASK). TASK-1, 3 & 5 have rather different structural and functional characteristics to TASK-2 and TASK-4.^{120–129} Closely related to TASK-2 are the TWIK-related alkaline pH activated K channels (TALK-1 and TALK-2) with TALK-2 being identical to TASK-4.¹³⁰ In addition, TWIK related K channels (TREK-1 and TREK-2) and the TWIK-related arachidonic acid sensitive K channel (TRAAK) are potentiated by arachidonic acid, volatile anaesthetics, heat and mechanical stress.^{116,131–133} Tandem pore domain halothane inhibited K channels (THIK-1 and THIK-2) have been cloned¹³⁴ and a background current in mouse cerebellar Purkinje cells with many properties in common with this family of 2PK channels has recently been described.¹³⁵ Finally, a 2PK clone, KCNK-7 does

not form functional channels when heterologously expressed alone or in combination.¹³⁶ THIK-2 and TASK-5 have also proved to be non-functional so far.

No fewer than seven of these subunits have been found in CGNs (see Table 3). High levels of TASK-1, TASK-3, TREK-2, TWIK-1^{128,137–139} and THIK-2¹³⁴ were found, whilst lower, but measurable levels of TREK-1, TRAAK^{137,140,141} were also seen. The TREK-2 isoform in CGNs has very recently been shown to be TREK-2c.¹⁴¹ In addition, TWIK-2 and KCNK-7 subunits are expressed in the cerebellum.^{119,138}

The molecular identity of leak currents in CGNs

All of the functional properties of $I_{K_{SO}}$ correlate well with those of the 2PK channel TASK-1. Both $I_{K_{SO}}$ currents and currents through TASK-1 channels in expression systems are open rectifiers which are insensitive to TEA and 4-AP, blocked by Ba^{2+} , muscarinic receptor activation, anandamide and external acidification, enhanced by halothane and not enhanced by arachidonic acid. This has led to the suggestion that TASK-1 channels underlie $I_{K_{SO}}$.^{112,113,139} However, it is extremely difficult to distinguish the functional properties of TASK-1 channels from those of the more recently cloned TASK-3 channels at the whole-cell level, so it is possible either that TASK-3 currents contribute to $I_{K_{SO}}$ ¹⁴² or that heterodimers could be formed between TASK-1 and TASK-3.¹⁴³ Similarly, the arachidonic

acid enhanced current in CGNs has been attributed to TREK-1 channels¹¹⁴ but recent expression studies might suggest a more likely correlate is the TREK-2c channel which is highly expressed in CGNs.¹⁴¹

Conclusion

A surprisingly large number of K channels are expressed in CGNs (and, it seems likely, every other neuron in the brain too). More precisely, there is evidence for strong or moderate expression levels of at least 26 K channel α subunits: $K_v1.1$, $K_v1.3$, $K_v1.5$, $K_v2.2$, $K_v3.1$, $K_v4.2$, $K_v4.3$, $K_v6.1$, $K_v8.1$, $K_v9.1$, $K_v9.2$, *eag1*, *erg1*, *elk2*, *KCNQ2*, *slo*, *SK2*, $K_{IR}2.2$, $K_{IR}3.1$, $K_{IR}3.2$, $K_{IR}3.3$, TASK-1, TASK-3, TREK-2c, THIK-2 and TWIK-1. Furthermore, there is evidence for detectable levels of expression of a number of other α subunits while several regulatory β subunits are also expressed. It is not at all clear why CGNs need so many different K channels.

If we consider granule cells from a functional perspective, D'Angelo et al.²⁰ have shown that it is quite possible to obtain a good empirical model of CGN function through the use of only five distinct K conductances, a leak conductance, three Na conductances and a Ca conductance. The five K conductances are a delayed rectifier conductance (K_v), a transient A-type conductance (K_A), a Ca-activated conductance (K_{Ca}) an inward-rectifier conductance (K_{IR}) and a slowly activating conduc-

Table 3
Expression of 2PK channel subunits in cerebellar granule cells

Channel	Synonym	Species	Age	RNA/protein	Level	References
TASK-1	KCNK-3	rat human	P7/ adult	RNA/protein	moderate/ abundant	112, 128, 137 139, 147, 148
TASK-2	KCNK-5	human	adult	RNA	none in cerebellum	138 (but see 149)
TASK-3	KCNK-9 KT3.2	rat	adult	RNA	abundant	127, 128, 137 139
TASK-4	KCNK-17 TALK-2	human	adult	RNA	not in brain	130
TASK-5	KT3.3	rat	adult	RNA	none	128
TALK-1	KCNK-16	human	adult		not in brain	130
TREK-1	KCNK-2	rat human	adult	RNA	low	137, 140, 141
TREK-2	KCNK-10	rat	adult	RNA	abundant	137, 141
TRAAK	KCNK-4	rat	adult	RNA	low	137
TWIK-1	KCNK-1	rat	adult	RNA	abundant	137, 139
TWIK-2	KCNK-6	human	adult	RNA	low in cerebellum	119, 138
KCNK-7		human	adult	RNA	abundant in cerebellum	138
THIK-1	KCNK-13	rat	adult	RNA	none	134
THIK-2	KCNK-12	rat	adult	RNA	abundant	134

tance (K_{slow}). There is also a requirement for a leak conductance (I_{leak}). In fast repetitive firing mode, K_V and K_{Ca} account for most of the active K current during the spike and fast AHP. K_A , K_{IR} and K_{slow} are active during the interspike interval, K_A acting to delay spike activation. Bursting behaviour is revealed when K_{Ca} is reduced (such as by TEA) or by G-protein coupled receptor action.

As we have shown above, in some cases, it is possible to make a good correlation between a functional current and an expressed channel. The transient K_A current for example, seems to be predominantly due to $K_V4.2$ channels or $K_V4.2/4.3$ heteromers, while the K_{Ca} current is likely due to expression of large-conductance slo channels. Similarly the G-protein activated K_{IR} current is probably due to heteromeric expression of $K_{\text{IR}3.1}$ and $K_{\text{IR}3.2}$, while the constitutively active component of K_{IR} may be due to $K_{\text{IR}2.2}$ expression. The leak conductance can be attributed to TASK-1 and/or TASK-3 channels. With less certainty, the K_{slow} current identified by D'Angelo et al.²⁰ may result from the expression of KCNQ2 and/or one or more of the EAG family of channels, however a possible small contribution of KCNQ3 or KCNQ5 channels cannot be ignored. The delayed-rectifier, K_V current is least well accounted for with five or more possible K_V subunits as potential contributors.

Measuring the whole-cell CGN current may well be important in determining granule cell excitability, but at the level of individual dendrites or release sites, CGNs may tailor the complement of K channel subunits present to suit particular roles. For example, the biophysical properties of $K_V1.1$ and $K_V3.1$ channels may be best suited to regulating transmitter release from granule cell terminals,² while TASK-1/TASK-3 channels on the soma may fundamentally control granule cell excitability. The function of all these K channels can be regulated by co-expression of regulatory subunits (e.g. $K_V2.2$ with $K_V8.1$ or $K_V1.1$ with $K_V\beta1$), by the action of neurotransmitters such as acetylcholine (e.g. $K_{\text{IR}3.1/3.2}$, TASK-1/TASK-3, eag1) or by other physiological regulators. Therefore the complement of functioning K channels determining the electrical properties of granule cells at any given moment may be highly dynamic and subject to constant fine-tuning.

Acknowledgements

Our work on potassium channels in cerebellar granule cells was supported by the MRC and the Wellcome Trust. Catherine Clarke is supported by a BBSRC studentship.

References

- Hille B. Ion channels of excitable membranes. Sunderland, MA: Sinauer, 2001.
- Coetzee WA, Amarillo Y, Chiu J, Chow A, Lau D, McCormack T, Moreno H, Nadal MS, Ozaita A, Pountney D, Saganich M, Vega-Saenz de Miera E, Rudy B. Molecular diversity of K^+ channels. *Ann NY Acad Sci* 1999; 868: 233–285.
- Pongs O. Molecular biology of voltage-dependent potassium channels. *Physiol Rev* 1992; 72: S69–S88.
- Chandy KG, Gutman GA. Voltage-gated potassium channel genes. In: North RA, editor. Handbook of receptors and channels: ligand and voltage-gated ion channels. Boca Raton: CRC Press, 1995: 1–71.
- Dolly O, Parcej DN. Molecular properties of voltage-gated K^+ channels. *J Bioenerg Biomembr* 1996; 28: 231–253.
- Jan LY, Jan YN. Voltage-gated and inwardly rectifying potassium channels. *J Physiol* 1997; 505: 267–282.
- Nichols CG, Lopatin AN. Inwardly rectifying potassium channels. *Ann Rev Physiol* 1997; 59: 171–191.
- Sanguinetti MC, Spector PS. Potassium channelopathies. *Neuropharmacol* 1997; 36: 755–762.
- Aguilar-Bryan L, Clement JP, Gonzalez G, Kunjilwar K, Babenko A, Bryan J. Toward understanding the assembly and structure of K_{ATP} channels. *Physiol Rev* 1998; 78: 227–245.
- Mathie A, Woollorton JRA, Watkins CS. Voltage-activated potassium channels in mammalian neurons and their block by novel pharmacological agents. *Gen Pharmacol* 1998; 30: 13–24.
- Bezanilla F. The voltage sensor in voltage-dependent ion channels. *Physiol Rev* 2000; 80: 555–592.
- Bauer CK, Schwarz JR. Physiology of EAG channels. *J Membrane Biol* 2001; 182: 1–15.
- Robbins J. KCNQ potassium channels: physiology, pathophysiology and pharmacology. *Pharmacol Therapeut* 2001; 90: 1–19.
- Goldstein SAN, Bockenhauer D, O'Kelly I, Zilberberg N. Potassium leak channels and the KCNK family of two-P-domain subunits. *Nature Rev Neurosci* 2001; 2: 175–184.
- Herrup K, Kuemerle B. The compartmentalization of the cerebellum. *Ann Rev Neurosci* 1997; 20: 61–90.
- Voogd J, Glickstein M. The anatomy of the cerebellum. *Trends in Neurosci* 1998; 21: 370–375.
- Rakic P, Sidman RL. Sequence of developmental abnormalities leading to granule cell deficit in cerebellar cortex of weaver mutant mice. *J Comp Neurol* 1973; 152: 103–132.
- Patil N, Cox DR, Bhat D, Faham M, Myers RM, Peterson AS. A potassium channel mutation in weaver mice implicates membrane excitability in granule cell differentiation. *Nature Genetics* 1995; 11: 126–129.
- Gabbiani F, Midtgaard J, Knopfel T. Synaptic integration in a model of cerebellar granule cells. *J Neurophysiol* 1994; 72: 999–1009.
- D'Angelo E, Nieuws T, Maffei A, Armano S, Rossi P, Taglietti V, Fontana A, Naldi G. Theta-frequency bursting and resonance in cerebellar granule cells: experimental evidence and modeling of a slow K^+ -dependent mechanism. *J Neurosci* 2001; 21: 759–770.
- Hockberger PE, Tseng H-Y, Connor JA. Immunocytochemical and electrophysiological differentiation of rat cerebellar granule cells in explant cultures. *J Neurosci* 1987; 7: 1370–1383.
- Robello M, Carignani C, Marchetti C. A transient voltage-dependent outward current in cultured cerebellar granule neurons. *Biosci Rep* 1989; 9: 451–457.
- Cull-Candy SG, Marshall CG, Ogden D. Voltage-activated membrane currents in rat cerebellar granule neurons. *J Physiol* 1989; 414: 179–199.
- Jalonen T, Johansson S, Holopainen I, Oja SS, Arhem P. Single-channel and whole-cell currents in rat cerebellar granule cells. *Brain Res* 1990; 535: 33–38.
- Carignani C, Robello M, Marchetti C, Maga L. A transient outward current dependent on external calcium in rat cerebellar granule cells. *J Membrane Biol* 1991; 122: 259–265.
- Bardoni R, Belluzzi O. Kinetic study and numerical reconstruction of A-type current in granule cells of rat cerebellar slices. *J Neurophysiol* 1993; 69: 2222–2231.

27. Watkins CS, Mathie A. Modulation of the gating of the transient outward potassium current of rat isolated cerebellar granule neurons by lanthanum. *Pflügers Archiv/Eur J Physiol* 1994; 428: 209–216.
28. Zegarramorán O, Moran O. Properties of the transient potassium currents in cerebellar granule cells. *Exp Brain Res* 1994; 98: 298–304.
29. Jones G, Boyd DF, Yeung SY, Mathie A. Inhibition of delayed rectifier K⁺ conductance in cultured rat cerebellar granule neurons by activation of calcium-permeable AMPA receptors. *Eur J Neurosci* 2000; 12: 935–944.
30. Watkins CS, Mathie A. Effect on K⁺ currents in rat cerebellar granule neurons of a membrane-permeable analogue of the calcium chelator BAPTA. *Brit J Pharmacol* 1996; 118: 1773–1778.
31. Chung YH, Shin CM, Kim MJ, Lee BK, Cha CI. Immunohistochemical study on the distribution of six members of the Kv1 channel subunits in the rat cerebellum. *Brain Res* 2001; 895: 173–177.
32. Kues WA, Wunder F. Heterogeneous expression patterns of mammalian potassium channel genes in developing and adult rat brain. *Eur J Neurosci* 1992; 4: 1296–1308.
33. Sheng M, Tsaur ML, Jan YN, Jan LY. Contrasting subcellular localization of the Kv_v1.2 K⁺ channel subunit in different neurons of rat brain. *J Neurosci* 1994; 14: 2408–2417.
34. Veh RW, Lichtinghagen R, Sewing S, Wunder F, Grumbach IM, Pongs O. Immunohistochemical localization of 5 members of the Kv_v1 channel subunits—contrasting subcellular locations and neuron-specific co-localizations in rat brain. *Eur J Neurosci* 1995; 7: 2189–2205.
35. Shibata R, Nakahira K, Shibasaki K, Wakazono Y, Imoto K, Ikenaka K. A-type K⁺ current mediated by the Kv4 channel regulates the generation of action potential in developing cerebellar granule cells. *J Neurosci* 2000; 20: 4145–4155.
36. Drewe JA, Verma S, Frech G, Joho RH. Distinct spatial and temporal expression patterns of K⁺ channel messenger RNAs from different subfamilies. *J Neurosci* 1992; 12: 538–548.
37. VermaKurvari S, Border B, Joho RH. Regional and cellular expression patterns of four K⁺ channel mRNAs in the adult rat brain. *Mol Brain Res* 1997; 46: 54–62.
38. Chung YH, Shin CM, Kim MJ, Lee BK, Cha CI. Age-related changes in the distribution of Kv_v1.1 and Kv_v1.2 channel subunits in the rat cerebellum. *Brain Res* 2001; 897: 193–198.
39. Wang H, Kunkel DD, Schwartzkroin PA, Tempel BL. Localization of Kv_v1.1 and Kv_v1.2 K-channel proteins, to synaptic terminals, somata and dendrites in the mouse brain. *J Neurosci* 1994; 14: 4588–4599.
40. Southan AP, Robertson B. Patch-clamp recordings from cerebellar basket cell bodies and their presynaptic terminals reveal an asymmetric distribution of voltage-gated potassium channels. *J Neurosci* 1998; 18: 948–955.
41. Leicher T, Bähring R, Isbrandt D, Pongs O. Coexpression of the KCNA3B gene product with Kv1.5 leads to a novel A-type potassium. *J Biol Chem* 1998; 273: 35095–35101.
42. Downen M, Belkowsky S, Knowles H, Cardillo M, Prystowsky MB. Developmental expression of voltage-gated potassium channel beta subunits. *Dev Brain Res* 1999; 117: 71–80.
43. Rettig J, Heinemann SH, Wunder F, Lorra C, Wittka R, Dolly O, Pongs O. Inactivation properties of voltage-gated K⁺ channels altered by the presence of beta-subunit. *Nature* 1994; 369: 289–294.
44. Rhodes KJ, Monaghan MM, Barreuzeta NX, Nawoschik S, BekeleArcuri Z, Matos MF, Nakahira K, Schechter LE, Trimmer JS. Voltage-gated K⁺ channel beta subunits: expression and distribution of Kv beta 1 and Kv beta 2 in adult rat brain. *J Neurosci* 1996; 16: 4846–4860.
45. Butler DM, Ono JK, Chang T, McCaman RE, Barish ME. Mouse brain potassium channel beta 1 subunit mRNA. Cloning and distribution during development. *J Neurobiol* 1998; 34: 135–150.
46. Browne DL, Gancher ST, Nutt JG, Brunt ER, Smith EA, Kramer P, Litt M. Episodic ataxia/myokymia syndrome is associated with a point mutation in the human potassium channel gene, KCNA1. *Nature Genetics* 1994; 8: 136–140.
47. Adelman JP, Bond CT, Pessia M, Maylie J. Episodic ataxia results from voltage-dependent potassium channels with altered functions. *Neuron* 1995; 15: 1449–1454.
48. Hwang PM, Glatt CE, Brecht DS, Yellen G, Snyder SH. A novel K⁺ channel with unique localizations in mammalian brain—molecular-cloning and characterization. *Neuron* 1992; 8: 473–481.
49. Fink M, Duprat F, Lesage F, Heurteaux C, Romey G, Barhanin J, Lazdunski M. A new K⁺ channel beta subunit to specifically enhance Kv2.2 (CDRK) expression. *J Biol Chem* 1996; 271: 26341–26348.
50. Sekirnjak C, Martone ME, Weiser M, Deerinck T, Bueno E, Rudy B, Ellisman M. Subcellular localization of the K⁺ channel subunit Kv3.1b in selected rat CNS neurons. *Brain Res* 1997; 766: 173–187.
51. Perney TM, Marshall J, Martin KA. Expression of the messenger RNAs for the Kv_v3.1 potassium channel gene in the adult and developing rat brain. *J Neurophysiol* 1992; 68: 756–766.
52. Weiser M, Demiera EVS, Kentros C, Moreno H, Franzen L, Hillman D, Baker H, Rudy B. Differential expression of shaw-related K⁺ channels in the rat central nervous system. *J Neurosci* 1994; 14: 949–972.
53. Shibata R, Wakazono Y, Nakahira K, Trimmer JS, Ikenaka K. Expression of Kv3.1 and Kv4.2 genes in developing cerebellar granule cells. *Dev Neurosci* 1999; 21: 87–93.
54. Sabatini BL, Regehr WG. Control of neurotransmitter release by presynaptic waveform at the granule cell to Purkinje cell synapse. *J Neurosci* 1997; 17: 3425–3435.
55. Espinosa F, McMahon A, Chan E, Wang S, Ho CS, Heintz N, Joho RH. Alcohol hypersensitivity, increased locomotion, and spontaneous myoclonus in mice lacking the potassium channels Kv3.1 and Kv3.3. *J Neurosci* 2001; 21: 6657–6665.
56. Tsaur ML, Chou CC, Shih YH, Wang HL. Cloning, expression and CNS distribution of Kv4.3, an A-type K⁺ channel alpha subunit. *FEBS Lett* 1997; 400: 215–220.
57. Serodio P, Rudy B. Differential expression of Kv4 K⁺ channel subunits mediating subthreshold transient K⁺ (A-type) currents in rat brain. *J Neurophysiol* 1998; 79: 1081–1091.
58. An WF, Bowly MR, Bett M, Cao J, Ling HP, Mendoza G, Hinson JW, Mattsson KI, Strassle BW, Trimmer JS, Rhodes KJ. Modulation of A-type potassium channels by a family of calcium sensors. *Nature* 2000; 403: 553–556.
59. Holmqvist MH, Cao J, Hernandez-Pineda R, Jacobson MD, Carroll KI, Sung MA, Betty M, Ge P, Gilbride KJ, Brown ME, Jurman ME, Lawson D, Silos-Santiago I, Xie Y, Covarrubias M, Rhodes KJ, Distefano PS, An WF. Elimination of fast inactivation in Kv_v4 A-type potassium channels by an auxiliary subunit domain. *Proc Natl Acad Sci USA* 2002; 99: 1035–1040.
60. Hugnot JP, Salinas M, Lesage F, Guillemare E, deWeille J, Heurteaux C, Mattei MG, Lazdunski M. Kv8.1, a new neuronal potassium channel subunit with specific inhibitory properties towards shab and shaw channels. *EMBO J* 1996; 15: 3322–3331.
61. Castellano A, Chiara MD, Mellstrom B, Lopez-Barneo J. Identification and functional characterization of a K⁺ channel alpha-subunit with regulatory properties specific to brain. *J Neurosci* 1997; 17: 4652–4661.
62. Salinas M, deWeille J, Guillemare E, Lazdunski M, Hugnot JP. Modes of regulation of shab K⁺ channel activity by the Kv8.1 subunit. *J Biol Chem* 1997; 272: 8774–8780.
63. Salinas M, Duprat F, Heurteaux C, Hugnot JP, Lazdunski M. New modulatory alpha subunits for mammalian shab K⁺ channels. *J Biol Chem* 1997; 272: 24371–24379.
64. Stocker M, Kerschensteiner D. Cloning and tissue distribution of two new potassium channel alpha-subunits from rat brain. *Biochem Biophys Res Commun* 1998; 248: 927–934.
65. Kramer JW, Post MA, Brown AM, Kirsch GE. Modulation of potassium channel gating by coexpression of Kv2.1 with regulatory Kv5.1 or Kv6.1 alpha-subunits. *Am J Physiol: Cell Physiol* 1998; 43: C1501–C1510.
66. Huang XY, Morielli AD, Peralta EG. Tyrosine kinase-dependent suppression of a potassium channel by the G protein-coupled m1 muscarinic acetylcholine receptor. *Cell* 1993; 75: 1145–1156.
67. Jonas EA, Kaczmarek LK. Regulation of potassium channels by protein kinases. *Curr Opin Neurobiol* 1996; 6: 318–323.
68. Saganich MJ, Machado E, Rudy B. Differential expression of genes encoding subthreshold-operating voltage-gated K⁺ channels in brain. *J Neurosci* 2001; 21: 4609–4624.
69. Ludwig J, Weseloh R, Karschin C, Liu Q, Netzer R, Engeland B, Stansfeld C, Pongs O. Cloning and functional expression of rat eag2, a new member of the ether-a-go-go family of potassium

- channels and comparison of its distribution with that of eag1. *Mol Cell Neurosci* 2000; 16: 59–70.
70. Hoshi N, Takahashi H, Shahidullah M, Yokoyama S, Higashida H. KCRI, a membrane protein that facilitates functional expression of non-inactivating K⁺ currents associates with rat eag voltage-dependent K⁺ channels. *J Biol Chem* 1998; 273: 23080–23085.
 71. Watkins CS, Mathie A. A non-inactivating K⁺ current sensitive to muscarinic receptor activation in rat cultured cerebellar granule neurons. *J Physiol* 1996; 491:401–412.
 72. Marrion NV. Does r-eag contribute to the M current? *Trends in Neurosci* 1997; 20: 243–244.
 73. Mathie A, Watkins CS. Is EAG the answer to the M current? *Trends in Neurosci* 1997; 20: 14.
 74. Wang HS, Pan ZM, Shi WM, Brown BS, Wymore RS, Cohen IS, Dixon JE, McKinnon D. KCNQ2 and KCNQ3 potassium channel subunits: molecular correlates of the M-channel. *Science* 1998; 282: 1890–1893.
 75. Tinel N, Lauritzen I, Chouabe C, Lazdunski M, Borsotto M. The KCNQ2 potassium channel splice variants, function and developmental expression. *Brain localization and comparison with KCNQ3*. *FEBS Lett* 1998; 438: 171–176.
 76. Kharkovets T, Hardelin J-P, Safieddine S, Schweizer M, El-Amraoui A, Petit C, Jentsch TJ. KCNQ4 a K channel mutated in a form of dominant deafness, is expressed in the inner ear and the central auditory pathway. *Proc Natl Acad Sci USA* 2000; 97: 4333–4338.
 77. Lerche C, Scherer CR, Seeböhm G, Derst C, Wei AD, Busch AE, Steinmeyer K. Molecular cloning and functional expression of KCNQ5, a potassium channel subunit that may contribute to neuronal M current diversity. *J Biol Chem* 2000; 275: 22395–22400.
 78. Schroeder BC, Hechenberger M, Weinreich F, Kubisch C, Jentsch TJ. KCNQ5, a novel potassium channel broadly expressed in brain, mediates M-type currents. *J Biol Chem* 2000; 275: 24089–24095.
 79. Fagni L, Bossu JL, Bockaert J. Activation of a large conductance Ca²⁺ dependent K⁺ channel by stimulation of glutamate phosphoinositide-coupled receptors in cultured cerebellar granule cells. *Eur J Neurosci* 1991; 3: 778–789.
 80. Chavis P, Ango F, Michel JM, Bockaert J, Fagni L. Modulation of big K⁺ channel activity by ryanodine receptors and L-type Ca²⁺ channels in neurons. *Eur J Neurosci* 1998; 10: 2322–2327.
 81. Adelman JP, Shen KZ, Kavanaugh MP, Warren RA, Wu YN, Lagrutta A, Bond CT, North RA. Calcium-activated potassium channels expressed from cloned complementary DNAs. *Neuron* 1992; 9: 209–216.
 82. TsengCrank J, Foster CD, Krause JD, Mertz R, Godinot N, Dichiaro TJ, Reinhart PH. Cloning, expression and distribution of functionally distinct Ca²⁺-activated K⁺ channel isoforms from human brain. *Neuron* 1994; 13: 1315–1330.
 83. Knaus HG, Schwarzer C, Koch ROA, Eberhart A, Kaczorowski GJ, Glossmann H, Wunder F, Pongs O, Garcia ML, Sperk G. Distribution of high-conductance Ca²⁺-activated K⁺ channels in rat brain: targeting to axons and nerve terminals. *J Neurosci* 1996; 16: 955–963.
 84. Chang CP, Dworetzky SI, Wang JC, Goldstein ME. Differential expression of the alpha and beta subunits of the large-conductance calcium-activated potassium channel. Implication for channel diversity. *Mol Brain Res* 1997; 45: 33–40.
 85. Müller YL, Reitsstetter R, Yool AJ. Regulation of Ca²⁺-dependent K⁺ channel expression in rat cerebellum during postnatal development. *J Neurosci* 1998; 18: 16–25.
 86. Stocker M, Pedarzani P. Differential distribution of three Ca²⁺-activated K⁺ channel subunits, SK1, SK2, and SK3, in the adult rat central nervous system. *Mol Cell Neurosci* 2000; 15: 476–493.
 87. Joiner WJ, Wang L-Y, Tang MD, Kaczmarek LK. hSK4, a member of a novel subfamily of calcium-activated potassium channels. *Proc Natl Acad Sci USA* 1997; 94: 11013–11018.
 88. Kofuji P, Hofer M, Millen KJ, Millonig JH, Davidson N, Lester HA, Hatten ME. Functional analysis of the weaver mutant GIRK2 K⁺ channel and rescue of weaver granule cells. *Neuron* 1996; 16: 941–952.
 89. Surmeier JD, Mermelstein PG, Goldowitz D. The weaver mutation of GIRK2 results in a loss of inwardly-rectifying K⁺ current in cerebellar granule cells. *Proc Natl Acad Sci USA* 1996; 93: 11191–11195.
 90. Slesinger PA, Stoffel M, Jan YN, Jan LY. Defective GABA_B receptor-activated inwardly-rectifying K⁺ currents in cerebellar granule cells isolated from weaver and Girk2 null mutant mice. *Proc Natl Acad Sci USA* 1997; 94: 12210–12217.
 91. Liesi P, Stewart RR, Wright JM. Involvement of GIRK2 in post-natal development of the weaver cerebellum. *J Neurosci Res* 2000; 60: 164–173.
 92. Mjaatvedt AE, Cabin DE, Cole SE, Long LJ, Breitweiser GE, Reeves RH. Assessment of a mutation in the H5 domain of GIRK2 as a candidate for the weaver mutation. *Genome Res* 1995; 5: 453–463.
 93. Rossi P, De Filippi G, Armano S, Taglietti V, D'Angelo E. The weaver mutation causes a loss of inward-rectifier current regulation in premigratory granule cell of the mouse cerebellum. *J Neurosci* 1998; 18: 3537–3547.
 94. Dixon AK, Gubituz AK, Ashford MLJ, Richardson PJ, Freeman TC. Distribution of the mRNA encoding the inwardly rectifying K⁺ channel, BIR1 in rat tissues. *FEBS Lett* 1995; 374: 135–140.
 95. Liao YJ, Jan YN, Jan LY. Heteromultimerization of G-protein inwardly rectifying K⁺ channel proteins GIRK1 and GIRK2 and their altered expression in weaver brain. *J Neurosci* 1996; 16: 7137–7150.
 96. Karschin C, Dissmann E, Stuhmer W, Karschin A. IRK(1-3) and GIRK(1-4) inwardly rectifying K⁺ channel mRNAs are differentially expressed in the adult rat brain. *J Neurosci* 1996; 16: 3559–3570.
 97. Karschin C, Karschin A. Ontogeny of gene expression of Kir channel subunits in the rat. *Mol Cell Neurosci* 1997; 10: 131–148.
 98. Lauritzen I, De Weille JD, Adelbrecht C, Lesage F, Murer MG, Raisman-Vozari R, Lazdunski M. Comparative expression of the inward-rectifier K⁺ channel GIRK2 in the cerebellum of normal and weaver mutant mice. *Brain Res* 1997; 753: 8–17.
 99. Miyashita T, Kubo Y. Localization and developmental changes of the expression of two inward rectifying K⁺-channel proteins in the rat brain. *Brain Res* 1997; 750: 251–263.
 100. Murer G, Adelbrecht C, Lauritzen I, Lesage F, Lazdunski M, Agid Y, Raisman-Vorazi R. An immunocytochemical study on the distribution of two G-protein-gated inward rectifier potassium channels (GIRK2 and GIRK4) in the adult rat brain. *Neurosci* 1997; 80: 345–357.
 101. Navarro B, Kennedy ME, Velimirovic B, Bhat D, Peterson AS, Clapham DE. Nonselective and Gbg-insensitive weaver K⁺ channels. *Science* 1996; 272: 1950–1953.
 102. Horio Y, Morishige K, Takahashi N, Kurachi Y. Differential distribution of classical inwardly rectifying potassium channel mRNAs in the brain: Comparison of IRK2 with IRK1 and IRK3. *FEBS Lett* 1996; 379: 239–243.
 103. Topert C, Döring F, Wischmeyer E, Karschin C, Brockhaus J, Ballanyi K, Derst C, Karschin A. Kir2.4: A novel K⁺ inward rectifier channel associated with motoneurons of cranial nerve nuclei. *J Neurosci* 1998; 18: 4096–4105.
 104. Karschin C, Ecker C, Ashcroft FM, Karschin A. Overlapping distribution of KATP channel-forming Kir6.2 subunit and the sulphonylurea receptor SUR1 in rodent brain. *FEBS Lett* 1997; 401: 59–64.
 105. Thomzig A, Wenzel M, Karschin C, Eaton MJ, Skatchkov SN, Karschin A, Veh RW. Kir6.1 is the principal pore-forming subunit of astrocyte but not neuronal plasma membrane K-ATP channels. *Mol Cell Neurosci* 2001; 18: 671–690.
 106. Pessia M, Imbrici P, D'Adamo C, Salvatore L, Tucker SJ. Differential pH sensitivity of Kir4.1 and Kir4.2 potassium channels and their modulation by heteropolymerisation with Kir5.1. *J Physiol* 2001; 532: 359–367.
 107. Slesinger PA, Patil N, Liao YJ, Jan YN, Jan LY, Cox DR. Functional effects of the mouse weaver mutation on G-protein-gated inwardly rectifying K⁺ channels. *Neuron* 1996; 16: 321–331.
 108. Reeves RH, Crowley MR, Lorenzon N, Pavan WJ, Sweeney RJ, Goldowitz D. The mouse neurological mutant weaver maps within the region of chromosome-16 that is homologous to human chromosome-21. *Genomics* 1989; 5: 522–526.
 109. Doyle DA, Cabral JM, Pfuetzner RA, Kuo A, Gulbis JM, Cohen SL, Chiat BT, MacKinnon R. The structure of the potassium channel: molecular basis of K⁺ conduction and selectivity. *Science* 1998; 280: 69–77.

110. Boyd DF, Mathie A. Inhibition of the potassium current IK_{SO} , in cerebellar granule cells, by the inhibitors of MEK1 activation, PD 98059 and U 0126. *Neuropharmacol* 2002; 42: 221–228.
111. Boyd DF, Millar JA, Watkins CS, Mathie A. The role of Ca^{2+} stores in the muscarinic inhibition of the K^+ current IK_{SO} in neonatal rat cerebellar granule cells. *J Physiol* 2000; 529: 321–331.
112. Millar JA, Barratt L, Southan AP, Page KM, Fyffe REW, Robertson B, Mathie A. A functional role for the two-pore domain potassium channel, TASK-1, in cerebellar granule neurons. *Proc Natl Acad Sci USA* 2000; 97: 3614–3618
113. Maingret F, Patel AJ, Lazdunski M, Honore E. The endocannabinoid anandamide is a direct and selective blocker of the background K^+ channel TASK-1. *EMBO J* 2001; 20: 47–54.
114. Lauritzen I, Blondeau N, Heurteaux C, Widmann C, Romey G, Lazdunski M. Polyunsaturated fatty acids are potent neuroprotectors. *EMBO J* 2000; 19: 1784–1793
115. Lesage F, Lazdunski M. Molecular and functional properties of two-pore-domain potassium channels. *Am J Physiol Renal Physiol* 2000; 279: F793–F801.
116. Patel AJ, Honore E. Properties and modulation of mammalian 2P domain K^+ channels. *Trends Neurosci* 2001; 24: 339–346.
117. Lesage F, Guillemare E, Fink M, Duprat F, Lazdunski M, Romey G, Barhanin J. TWIK-1, a ubiquitous human weakly inward rectifying K^+ channel with a novel structure. *EMBO J* 1996; 15: 1004–1011.
118. Chavez RA, Gray AT, Zhao BB, Kindler CH, Mazurek MJ, Mehta Y, Forsayeth JR, Yost CS. TWIK-2, a new weak inward rectifying member of the tandem pore domain potassium channel family. *J Biol Chem* 1999; 274: 7887–7892.
119. Patel AJ, Maingret F, Magnone V, Fosset M, Lazdunski M, Honore E. TWIK-2, an inactivating 2P domain K^+ channel. *J Biol Chem* 2000; 275: 28722–28730
120. Reyes R, Duprat F, Lesage F, Fink M, Salinas M, Farman N, Lazdunski M. Cloning and expression of a novel pH-sensitive two pore domain K^+ channel from human kidney. *J Biol Chem* 1998; 273: 30863–30869.
121. Kim Y, Bang H, Kim D. TASK-3, a new member of the tandem pore K^+ channel family. *J Biol Chem* 2000; 275: 9340–9347.
122. Rajan S, Wischmeyer E, Liu GX, Preisig-Muller R, Daut J, Karschin A, Derst C. TASK-3, a novel tandem pore domain acid-sensitive K^+ channel. *J Biol Chem* 2000; 275: 16650–16657.
123. Meadows HJ, Randall AD. Functional characterization of human TASK-3, an acid-sensitive two-pore domain potassium channel. *Neuropharmacol* 2001; 40: 551–559.
124. Decher N, Maier M, Dittrich W, Gassenhuber J, Bruggemann A, Busch AE, Steinmeyer K. Characterization of TASK-4, a novel member of the pH-sensitive, two-pore domain potassium channel family. *FEBS Lett* 2001; 492: 84–89
125. Ashmole I, Goodwin PA, Stanfield PR. TASK-5 a novel member of the tandem pore K^+ channel family. *Pflugers Archiv* 2001; 442: 828–833.
126. Kim D, Gnatenco C. TASK-5, a new member of the tandem-pore K^+ channel family. *Biochem Biophys Res Commun* 2001; 284: 923–930.
127. Vega-Saenz De Miera E, Lau DHP, Zhadina M, Pountney D, Coetzee WA, Rudy B. KT3.2 and KT3.3, two novel human two-pore K^+ channels closely related to TASK-1. *J Neurophysiol* 2001; 86: 130–142.
128. Karschin C, Wischmeyer E, Preisig-Muller R, Rajan S, Derst C, Grzeschik K-H, Daut J, Karschin A. Expression pattern in brain of TASK-1, TASK-3 and a tandem pore domain K^+ channel subunit, TASK-5, associated with the central auditory nervous system. *Mol Cell Neurosci* 2001; 18: 632–648.
129. Duprat F, Lesage F, Fink M, Reyes R, Heurteaux C, Lazdunski M. TASK, a human background K^+ channel to sense external pH variations near physiological pH. *EMBO J* 1997; 16: 5464–5471.
130. Girard C, Duprat F, Terrenoire C, Tinel N, Fosset M, Romey G, Lazdunski M, Lesage F. Genomic and functional characteristics of novel human pancreatic 2P domain K^+ channels. *Biochem Biophys Res Commun* 2001; 282: 249–256.
131. Fink M, Duprat F, Lesage F, Reyes R, Romey G, Heurteaux C, Lazdunski M. Cloning, functional expression and brain localization of a novel unconventional outward rectifier K^+ channel. *EMBO J* 1996; 15: 6854–6862.
132. Fink M, Lesage F, Duprat F, Heurteaux C, Reyes R, Fosset M, Lazdunski M. A neuronal two P domain K^+ channel stimulated by arachidonic acid and polyunsaturated fatty acids. *EMBO J* 1998; 17: 3297–3308.
133. Bang H, Kim Y, Kim D. TREK-2, a new member of the mechanosensitive tandem-pore K^+ channel family. *J Biol Chem* 2000; 275:17412–17419
134. Rajan S, Wischmeyer E, Karschin C, Preisig-Muller R, Grzeschik KH, Daut J, Karschin A, Derst C. THIK-1 and THIK-2, a novel subfamily of tandem pore domain K^+ channels. *J Biol Chem* 2001; 276: 7302–7311.
135. Bushell T, Clarke C, Mathie A, Robertson B. Pharmacological characterization of a non-inactivating outward current observed in mouse cerebellar Purkinje neurones. *Br J Pharmacol* 2002; 135: 705–712.
136. Salinas M, Reyes R, Lesage F, Fosset M, Heurteaux C, Romey G, Lazdunski M. Cloning of a new mouse two-P domain channel subunit and a human homologue with a unique pore structure. *J Biol Chem* 1999; 274: 11751–11760.
137. Talley EM, Solorzano G, Lei Q, Kim D, Bayliss DA. CNS distribution of members of the two-pore-domain (KCNK) potassium channel family. *J Neurosci* 2001; 21: 7491–7505.
138. Medhurst AD, Rennie G, Chapman CG, Meadows H, Duckworth MD, Kelsell RE, Gloger II, Pangalos MN. Distribution analysis of human two pore domain potassium channels in tissues of the central nervous system and periphery. *Brain Res Mol Brain Res* 2001; 86: 101–114
139. Brickley SG, Revilla V, Cull-Candy SG, Wisden W, Farrant M. Adaptive regulation of neuronal excitability by a voltage-independent potassium conductance. *Nature* 2001; 409: 88–92.
140. Hervieu GJ, Cluderay JE, Gray CW, Green PJ, Ranson JL, Randall AD, Meadows HJ. Distribution and expression of TREK-1, a two-pore-domain potassium channel, in the adult rat CNS. *Neurosci* 2001; 103: 899–919.
141. Gu W, Schlichthorl G, Hirsch JR, Engels H, Karschin C, Karschin A, Derst C, Steinlein OK, Daut J. Expression pattern and functional characteristics of two novel splice variants of the two-pore domain potassium channel TREK-2. *J Physiol* 2002; 539:657–668.
142. Han J, Truell J, Gnatenco C, Kim D. Several tandem-pore K^+ channels contribute to the background current in cerebellar granule neurons. *Biophys J* 2002; 82: 636a.
143. Czirjak G, Enyedi P. Formation of functional heterodimers between the TASK-1 and TASK-3 two-pore domain potassium channel subunits. *J Biol Chem* 2002; 277: 5426–5432.
144. GoldmanWohl DS, Chan E, Baird D, Heintz N. $K_v3.3B$ — A novel shaw type potassium channel expressed in terminally differentiated cerebellar Purkinje cells and deep cerebellar nuclei. *J Neurosci* 1994; 14: 511–522.
145. McNamara NMC, Averill S, Wilkin GP, Dolly JO, Priestley JV. Ultrastructural localization of a voltage-gated K^+ channel alpha subunit ($K_v1.2$) in the rat cerebellum. *Eur J Neurosci* 1996; 8: 688–699.
146. Shu-Cheng C. Ehrhard P, Goldowitz D, Smeyne RJ. Developmental expression of the GIRK family of inward rectifying potassium channels: implications for abnormalities in the *weaver* mutant mouse. *Brain Res* 1997; 778: 251–264.
147. Kindler CH, Pietruck C, Yost CS, Sampson ER, Gray AT. Localization of the tandem pore domain K^+ channel TASK-1 in the rat central nervous system. *Brain Res Mol Brain Res* 2000; 80: 99–108
148. Talley EM, Lei QB, Sirois JE, Bayliss DA. TASK-1, a two-pore domain K^+ channel, is modulated by multiple neurotransmitters in motoneurons. *Neuron* 2000; 25: 399–410.
149. Gabriel A, Abdallah M, Yost CS, Winegar BD, Kindler CH. Localization of the tandem pore domain K^+ channel KCNK5 (TASK-2) in the rat central nervous system. *Mol Brain Res* 2002; 98:153–163.