

# Kent Academic Repository

## Full text document (pdf)

### Citation for published version

Cazenave, Marine (2020) Hip joint loading differences between *Paranthropus robustus* and *Australopithecus africanus* revealed by the cortical bone distribution at the femoral neck. In: European Society for the study of Human Evolution 2020. (Unpublished)

### DOI

### Link to record in KAR

<https://kar.kent.ac.uk/84673/>

### Document Version

Publisher pdf

#### Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

#### Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

#### Enquiries

For any further enquiries regarding the licence status of this document, please contact:

[researchsupport@kent.ac.uk](mailto:researchsupport@kent.ac.uk)

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

# Hip joint loading differences between *Paranthropus robustus* and *Australopithecus africanus* revealed by the cortical bone distribution at the femoral neck

Marine Cazenave<sup>1</sup>, Amélie Beaudet<sup>2,3</sup>

1 - Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, UK · 2 - School of Geography, Archaeology and Environmental Studies, University of the Witwatersrand, Johannesburg, South Africa · 3 - Department of Anatomy, University of Pretoria, South Africa.

Given the ability of the cortical and trabecular tissues to respond to the loading environment, the reconstruction of the locomotor repertoire of australopiths (*Australopithecus* and *Paranthropus*) has increasingly integrated information from the inner structure of the appendicular skeleton [1-4]. There has been a particular focus on the internal structure of the hip joint because of its critical role in locomotion. The asymmetrical distribution of cortical bone in the femoral neck, with thinner superior (S) and thicker inferior (I) cortex, has been shown to characterize bipedal humans, reflecting the stereotypical loading of the hip joint. In australopiths, the combination of a human-like S/I asymmetry at the base of neck and a less asymmetric distribution at mid-neck has been interpreted as a commitment to terrestrial bipedalism associated to a slightly altered gait kinematics compared to *Homo* [1,3]. A recent X-ray microtomographic ( $\mu$ CT) analysis of cortical bone thickness distribution across the femoral neck of four *P. robustus* specimens has identified three additional features: (i) a human-like trend of lateral-to-medial decrease in S/I asymmetry; (ii) an accentuated contrast between the relatively thicker anterior and the thinner posterior walls, and (iii) a marked lateral-to-medial thinning of both cortices. However, it is still unknown whether, and to what extent, these features also characterize *Australopithecus*. Given previous evidence for some trabecular variation between *Au. africanus* and *P. robustus* in different skeletal elements [4,5], we predict that a refined assessment of the *Au. africanus* femoral neck cortical distribution has the potential to reveal gait-related differences with respect to *P. robustus*.

Using  $\mu$ CT data (resolution 30  $\mu$ m), we assessed the cortical bone distribution in two proximal femora from Sterkfontein Caves, South Africa: StW 479 from Member 4 attributed to *Au. africanus* and StW 311 from Member 5, originally attributed to *Au. africanus*, but recently suggested to represent either early *Homo* or *P. robustus* [4]. Both fossils are housed at the Evolutionary Studies Institute and were scanned at the Paleosciences Centre of the Univ. Witwatersrand (South Africa). Due to differences in preservation, we extracted a set of slices at regular intervals from the base (lateral) of the neck to 70% of the total neck length in StW 479 and one at mid-neck of StW 311. We compared their functional signals to the published evidence from the *P. robustus* specimen SK 82 from Swartkrans (South Africa) and 25 extant humans and 8 extant chimpanzees [3].

Our refined S/I measurements of StW 479 support previous CT-based estimates [1]. StW 479 shares with SK 82 a S/I asymmetry in cortical thickness at the base of neck that decreases latero-medially. These features, also observed in extant humans, are thought to reflect habitual bipedal locomotion and the action of the gluteal abductor muscles [1,3]. However, while SK 82 has a less asymmetric S/I ratio than in humans, especially at mid-neck, StW 479 shows a human-like distribution along the neck. In addition, relatively thicker posterior and thinner anterior walls are identified in StW 479, while an opposite pattern is observed in SK 82 [3]. Such differences suggest that the locomotor related-biomechanical environment of the hip joint differs between *Au. africanus* and *P. robustus*, supporting locomotor diversity within the early hominin record. StW 311 shows similarities with StW 479 in cortical bone distribution at mid-neck and differs from the condition in SK 82. Since its trabecular bone distribution pattern at the head is similar to the nonhuman ape condition and differs from the *Au. africanus* pattern [4], further investigation of the trabecular network across the StW 311 neck are needed to help clarify its loading conditions and taxonomic attribution.

K. Jakata (Johannesburg) and B. Zipfel (Johannesburg) for access to the  $\mu$ CT record of the fossil materials; J. Dumoncel (Toulouse) for the development of a custom-written program in the MATLAB v. 8.1 (MathWorks, 2013) software to automatically measure the cortical thickness; MC is funded by the Fyssen Foundation and AB is funded by the University of the Witwatersrand.

**References:** [1] Ruff, C.B., Higgins, R., 2013. Femoral neck structure and function in early hominins. *American Journal of Physical Anthropology* 150, 512-525. [2] Kivell, T.L., Davenport, R., Hublin, J.-J., Thackeray, J.F., Skinner, M.M., 2018. Trabecular architecture and joint loading of the proximal humerus in extant hominoids, *Ateles*, and *Australopithecus africanus*. *American Journal of Physical Anthropology* 167, 348-365. [3] Cazenave, M., Braga, J., Oetlér, A., Pickering, T.R., Heaton, J.L., Nakatsukasa, M., Thackeray, J.F., de Beer, F., Hoffman, J., Dumoncel, J., Macchiarelli, R., 2019. Cortical bone distribution in the femoral neck of *Paranthropus robustus*. *Journal of Human Evolution* 135, 102666. [4] Georgiou, L., Dunmore, C.J., Bardo, A., Buck, L.T., Hublin, J.-J., Pahr, D.H., Stratford, D., Synek, A., Kivell, T.L., Skinner, M.M., 2020. Evidence for habitual climbing in a middle Pleistocene hominin in South Africa. *Proceedings of the National Academy of Sciences USA*. <https://doi.org/10.1073/pnas.1914481117>. [5] Su, A., Carlson, K.J., 2017. Comparative analysis of trabecular bone structure and orientation in South African hominin tali. *Journal of Human Evolution* 106, 1-18.