

Kent Academic Repository

Full text document (pdf)

Citation for published version

Brylka, Asteria and Wolke, Dieter and Ludyga, Sebastian and Bilgin, Ayten and Spiegler, Juliane and Trower, Hayley and Gkiouleka, Anna and Gerber, Markus and Brand, Serge and Grob, Alexander and Weber, Peter and Heinonen, Kati and Kajantie, Eero and Räikkönen, Katri and Lemola, Sakari (2021) Physical Activity, Mental Health, and Well-Being in Very Pre-Term and Term Born Adolescents:

DOI

<https://doi.org/10.3390/ijerph18041735>

Link to record in KAR

<https://kar.kent.ac.uk/86582/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

Physical Activity, Mental Health, and Well-being in Very Preterm and Term Born Adolescents. An Individual Participant Data Meta-analysis of two Accelerometry Studies

Asteria Brylka¹, Dieter Wolke¹, Sebastian Ludyga², Ayten Bilgin^{1, 15}, Juliane Spiegler^{1,9}, Hayley Trower¹, Anna Gkiouleka¹, Markus Gerber², Serge Brand^{2,3,4,5,6}, Alexander Grob⁷, Peter Weber⁸, Kati Heinonen¹⁰, Eero Kajantie^{11,12,13}, Katri Räikkönen¹⁰, & Sakari Lemola^{1,14}

¹ *Department of Psychology, University of Warwick, Coventry, United Kingdom*

² *Department of Sport, Exercise and Health, Sport Sciences Section, University of Basel, Basel, Switzerland*

³ *Psychiatric Clinics, Center for Affective, Stress, and Sleep Disorders, University of Basel, Basel, Switzerland*

⁴ *Sleep Disorders Research Center, Kermanshah University of Medical Sciences (KUMS), Kermanshah, Iran*

⁵ *Substance Abuse Prevention Research Center, Health Institute, Kermanshah University of Medical Sciences (KUMS), Kermanshah, Iran*

⁶ *School of Medicine, Tehran University of Medical Sciences (TUMS), Tehran, Iran*

⁷ *Department of Psychology, University of Basel, Basel, Switzerland*

⁸ *University Children's Hospital Basel, Division of Neuropediatrics and Developmental Medicine, Basel, Switzerland*

⁹ *University of Lübeck, Department of Paediatrics, Lübeck, Germany*

¹⁰ *Department of Psychology & Logopedics, University of Helsinki, Helsinki, Finland*

¹¹ *National Institute of Health and Welfare, Helsinki, Finland*

¹² *PEDEGO Research Unit, Medical Research Center Oulu, Oulu University Hospital and University of Oulu Finland*

¹³ *Children's Hospital, Helsinki University Hospital and the University of Helsinki, Helsinki, Finland*

¹⁴ *Bielefeld University, Department of Psychology, Bielefeld, Germany*

¹⁵ *School of Psychology, University of Kent, Canterbury, United Kingdom*

Correspondence to:

Sakari Lemola, PhD

Department of Psychology,

Bielefeld University,

Bielefeld, Germany

sakari.lemola@uni-bielefeld.de

Abstract: This study examined whether physical activity is associated with better mental health and well-being among very preterm (≤ 32 weeks) and term born (≥ 37 weeks) adolescents alike or whether the associations are stronger in either of the groups. Physical activity was measured with accelerometry in children born very preterm and at term in two cohorts, the Basel Study of Preterm Children (BSPC; 40 adolescents born ≤ 32 weeks of gestation and 59 term born controls aged 12.3 years) and the Millennium Cohort Study (MCS; 45 adolescents born ≤ 32 weeks of gestation and 3,137 term born controls aged 14.2 years on average). In both cohorts, emotional and behavioural problems were mother-reported using the Strengths and Difficulties Questionnaire. Subjective well-being was self-reported using the Kidscreen-52 Questionnaire in the BSPC and single items in the MCS. Hierarchical regressions with ‘preterm status x physical activity’-interaction effects were subjected to individual participant data (IPD) meta-analysis. IPD meta-analysis showed that higher levels of physical activity were associated with lower levels of peer problems and higher levels of psychological well-being, better self-perception/body image, and school related well-being. Overall, the effect-sizes were small and the associations did not differ significantly between very preterm and term born adolescents. Future research may examine the mechanisms behind effects of physical activity on mental health and wellbeing in adolescence as well as which type of physical activity might be most beneficial in term and preterm born children.

Keywords: physical activity; mental health; well-being; preterm birth; adolescence; accelerometry.

Abbreviations: Basel Study of Preterm Children (BSPC); Millennium Cohort Study (MCS); Individual Participant Data (IPD) Meta-Analysis.

Children and adults born very preterm or with very low birth weight have an increased risk of neurodevelopmental and mental health problems (Aarnoudse-Moens, Weisglas-Kuperus, van Goudoever & Oosterlaan, 2009; Wolke, Baumann, Strauss, Johnson & Marlow, 2015). During adolescence and young adulthood, those born very preterm are often described as being more socially withdrawn and having problems with peers (Eryigit-Madzwamuse, Strauss, Baumann, Bartmann & Wolke, 2015). While prematurity increases the risk of adverse outcomes, many very preterm born children and adolescents do not have any symptoms of mental health problems, showing resilience in the face of complications and adversity (Wolke, 2018). However, research on resilience factors among very preterm children and adolescents is rather scarce. Existing research has identified sensitive parenting in childhood (Wolke, Jaekel, Hall & Baumann, 2013) as a potential resilience factor for academic achievement among very preterm children (i.e. in very preterm children sensitive parenting was more strongly associated with academic outcomes than in term born children). By contrast, a recent study reported that very preterm children may be more strongly affected by risk factors for mental health such as family dysfunction, parental mental health problems, and peer victimisation while not benefitting as much from potential resilience factors such as number of close friends or sensitive parental care compared to term born children (Van Lieshout et al., 2018).

Physical activity has been suggested to promote mental health and well-being among children and adolescents in the general population (e.g. Andersen et al., 2017; Biddle & Asare, 2011; Lubans et al., 2016). Further, it has been linked to increased happiness (Booker, Skew, Kelly & Sacker, 2015), positive affect (Wen et al., 2018), improved facial emotion recognition (Brand et al., 2019), and decreased levels of anxiety, depressive symptoms, and socio-emotional difficulties (Booker, Skew, Kelly & Sacker, 2015; Kirkcaldy, Shephard & Siefen, 2002). In particular, it has been suggested that physical activity plays an important

role for adolescents in building a favourable self-image and self-esteem, and in decreasing social inhibition and anxiety by evoking positive social feedback and recognition from peers (Liu, Wu, & Ming, 2015; Kirkcaldy, Shephard & Siefen, 2002).

Very preterm children tend to have more physical and motor limitations (Evensen et al., in press; Spittle et al., 2018) and show reduced pulmonary function, aerobic capacity, and muscle strength (Takken et al., 2010). Despite this, no difference in physical activity levels has been reported during adolescence between very preterm and term born children (Dahan-Oliel, Mazer & Majnemer, 2012; Nordvall-Lassen et al., 2018; Spiegler, Mendonça & Wolke, 2019), although lower levels of physical activity have been observed in young adults (Kaseva et al., 2012; Tikanmäki et al., 2017). However, research is scarce on whether physical activity has similar benefits for mental health and well-being among very preterm children and adolescents as it has among their term born peers (Dahan-Oliel et al. 2014). As very preterm children are often reported to show higher levels of emotional and peer problems, increased physical activity may particularly benefit their self-image and recognition from peers and decrease social inhibition.

The objective of the current study was to examine three competing explanatory models regarding how physical activity may impact on very preterm born adolescents' mental health and well-being (Figure 1). The first model proposes that physical activity protects very preterm and term born adolescents alike from lower mental health and well-being; the second model proposes that physical activity is a resiliency factor that particularly benefits very preterm adolescents; while the third model proposes that very preterm adolescents benefit less from physical activity than those born at term (Van Lieshout et al., 2018). While the first model involves a main effect (i.e. physical activity showing the same association in both groups), the other two models describe moderation effects involving a stronger (the 2nd model) or weaker (the 3rd model) positive association among very preterm adolescents in

comparison to term born adolescents. These models are tested using physical activity objectively assessed with accelerometry, mother-reported behavioral and emotional problems and child-reported well-being in two samples of very preterm born adolescents and term born controls. Physical activity was assessed with accelerometry to avoid the common-method bias, which may inflate associations when both predictor and outcome are reported by the same person (Podsakoff, MacKenzie, Lee & Podsakoff, 2003).




Figure 1.

Three competing hypothetical models of the impact of physical activity in very preterm and termborn adolescents. A) Universal Protective Factor, B) Group-specific Resiliency Factor, C) Less Benefit for at Risk Group.

Methods

Participants

Basel Study of Preterm Children (BSPC). The BSPC is a cohort study of very preterm children and term born controls. The very preterm sub-sample was recruited from an initial cohort of 260 prematurely born children, who were treated at the University Children’s Hospital of Basel, Switzerland, between June 2001 and December 2006, and for whom gestational age was retrieved from hospital records (Maurer et al., 2016). Cross-sectional data used for the current study come from the third wave of the BSPC collected between February 2016 and October 2017, when the children were eleven to fifteen-year old (average age = 12.2 years, $SD= 1.1$). In total, 40 children and adolescents born very preterm (≤ 32 weeks and 0 days of gestation) had sufficient days of valid accelerometry recordings to be included (i.e.,

at least 4 days with a wear time of at least 70% including one weekend day). The control subsample included 59 term born participants, who were originally recruited through official birth notifications (≥ 37 weeks and 0 days of gestation; also including post-terms) as an age and sex matched comparison group in the first wave of the BSPC when children were 8 years old on average (Maurer et al., 2016).

Millennium Cohort Study (MCS). The Millennium Cohort Study (MCS) is a representative longitudinal study of 18,818 infants born in the UK (Connelly & Platt, 2014). A random sample was drawn from Child Benefit registers of infants born in the UK, between September 2000 and January 2002, and who were living in the UK at the age of 9 months. Parents were interviewed for the first time when the children were aged 9 months (survey 1), and again at 14 years (survey 6). Validity of parent-reported gestational age at 9 months after birth has been reported by Poulsen and colleagues (Poulsen et al., 2011). Cross-sectional data at wave 6 were used for the current study, when participants were 14.2 years old ($SD=0.3$). Data of twins and triplets were included in the analysis. Detailed analysis of sociodemographic data and a flow chart of study participation of different gestational age groups have been published before (Spiegler, Mendoca & Wolke, 2019). In total, 10,337 MCS members had been invited for the accelerometry protocol. After excluding 6,258 participants with insufficient accelerometer data (i.e. participants with a wear time of 70% or less on one of two accelerometry assessment days), and 292 with missing data on gestational age or gestational age ranging from 32+1 to 36+6 weeks, the study sample included 45 very preterm born adolescents (≤ 32 weeks and 0 days of gestation) and 3,137 term born participants (≥ 37 weeks and 0 days of gestation; also including post-terms). Very preterm children were not more likely to drop out of the analysis compared to their term born counterparts (participation rate among very preterm adolescents was 17.8% compared to 18.7% among those born at term; $\chi^2(1)=0.13$, $P=0.72$).

Sample characteristics of very preterm and term born adolescents are shown in Table 1 for both cohorts.

Table 1.

Sample Characteristics of the very preterm and term born adolescents and reliability indices for study scales.

	<i>Basel Study of Preterm Children</i>				<i>Millennium Cohort Study</i>				Std. pooled mean difference ¹
	Very preterm <i>n=40</i> <i>N(%) / M(SD)</i>	Control <i>n=59</i> <i>N(%) / M(SD)</i>	Very preterm vs. Control <i>p</i>	Cronbach's α	Very preterm <i>n=45</i> <i>N(%) / M(SD)</i>	Control <i>n=3137</i> <i>N(%) / M(SD)</i>	Very preterm vs. Control <i>p</i>	Cronbach's α	
Gestational age (weeks)	30.2 (2.1)	39.5 (1.5)	<i>p</i> < .001	-	29.5 (2.4)	39.6 (1.3)	<i>p</i> < .001	-	
Age at assessment (years)	12.3 (1.0)	12.1 (1.1)	<i>p</i> = .233	-	14.3 (0.4)	14.2 (0.3)	<i>p</i> = .130	-	
Sex (n Male)	21 (51.2)	31 (50.8)	<i>p</i> = .968	-	22 (48.9)	1438 (45.8)	<i>p</i> = .684	-	
Neurosensory impairment (n)	2 (4.9)	1 (1.6)	<i>p</i> = .563	-	2 (4.4)	56 (1.8)	<i>p</i> = .197	-	
Ethnic minority group (n)	10 (24.4)	7 (11.5)	<i>p</i> = .086	-	11 (24.4)	408 (13.0)	<i>p</i> = .024	-	
Parental tertiary education (n)	14 (34.1)	44 (72.1)	<i>p</i> < .001	-	16 (35.6)	1028 (32.8)	<i>p</i> = .693	-	
Physical activity(minutes/day)	162 (50)	172 (58)	<i>p</i> = .400	-	119 (59)	129 (73)	<i>p</i> = .334	-	
Mental health (mother-reported):									
Total Behavioral/Emotional Difficulties	7.34 (4.90)	7.36 (5.68)	<i>p</i> = .987	.83	8.20 (5.95)	7.12 (5.40)	<i>p</i> = .187	.71	.129
Emotional symptoms	1.82 (1.85)	1.36 (1.67)	<i>p</i> = .206	.70	2.36 (2.49)	1.84 (2.02)	<i>p</i> = .088	.72	.259*
Conduct problems	1.25 (1.28)	1.64 (1.88)	<i>p</i> = .259	.59	1.26 (1.64)	1.12 (1.38)	<i>p</i> = .488	.62	-.037
Hyperactivity/inattention	2.95 (2.23)	2.81 (2.47)	<i>p</i> = .776	.80	2.69 (2.20)	2.60 (2.25)	<i>p</i> = .793	.76	.047
Peer problems	1.33 (1.42)	1.55 (1.75)	<i>p</i> = .499	.57	1.89 (1.75)	1.56 (1.73)	<i>p</i> = .219	.63	.058
Well-being (child-reported):									
Overall well-being	126.08 (13.39)	125.75 (12.24)	<i>p</i> = .900	.88	33.64 (7.64)	33.42 (6.48)	<i>p</i> = .815	-	.031
Psychological well-being	20.43 (2.48)	20.34 (2.19)	<i>p</i> = .853	.81	5.73 (1.53)	5.59 (1.40)	<i>p</i> = .483	-	.079
Self-perception	17.37 (2.98)	16.95 (2.71)	<i>p</i> = .471	.69	5.16 (1.54)	4.80 (1.55)	<i>p</i> = .132	-	.203
Peer relations	19.68 (3.05)	18.90 (3.36)	<i>p</i> = .240	.85	5.67 (1.55)	5.97 (1.28)	<i>p</i> = .118	-	-.017
School related well-being	19.40 (2.70)	19.03 (3.24)	<i>p</i> = .558	.82	11.02 (3.23)	11.00 (2.46)	<i>p</i> = .956	-	.048

¹ Pooled mean difference estimate from the random effects meta-analysis (See Supplemental Table 1)**p* = .033

Ethical considerations

Ethical approval from the respective ethic committees (BSPC: Ethikkommission Nordwest- und Zentralschweiz, reference number 122/11; MCS: London - Hampstead Research Ethics Committee, REC reference 14/LO/0868) were obtained and participants gave written informed consent.

Measures

Physical activity. In the BSPC, physical activity was measured over seven consecutive days with triaxial accelerometers (ActiGraph LLC; Pensacola, FL, USA) worn on the wrist of the non-dominant hand throughout the day. ActiLife software 6.13.3 (ActiGraph LLC) was used for the initialisation of accelerometers, download, and processing of collected data. Prior to analyses, wear time validation was performed with the Troiano and colleagues' (2008) algorithm implemented in the program that defines non-wear time by an interval of 60 or more consecutive minutes of zero activity intensity counts, with allowance for one to two minutes of counts between zero and 100. Recorded days with a non-wear time of 30% or higher were excluded from the analyses. A dataset was considered valid, if data covered at least four days after controlling for non-wear time, and if one of these days was a valid weekend day (see Barreira et al., 2015). Physical activity on valid days was coded into number of minutes of sedentary, light, moderate, and vigorous physical activity using counts per minute cut-points, which have been validated in children (Mattocks et al. 2007). Cut-off points for moderate and vigorous physical activity were 3600 counts/min and 6200 counts/min, respectively. The level of objective physical activity per week was calculated by summing minutes of vigorous and moderate physical activity performed by the participants divided by the number of days of accelerometry data collection and multiplied by 7. To use physical activity in cross-cohort analysis, the variable was z-standardized.

In the MCS, objective physical activity was measured using GENEActiv Original accelerometer devices worn on the wrist of the non-dominant hand on one day during the week and one weekend day calculating Euclidean Norm Minus One (ENMO). Total number of minutes are indicated summing up 1-minute epochs with ENMO>100mg (Heywood, 2018). This variable representing the total number of minutes of moderate to vigorous physical activity was highly correlated with other definitions of moderate-vigorous physical activity (e.g. it correlated at $r=0.85$ ($p < 0.001$) with moderate to vigorous physical activity defined as bouts of 5-minutes windows with 80% of time spent in moderate-vigorous physical activity). To calculate a weekly average, weekday physical activity was multiplied by 5 and weekend physical activity was multiplied by 2 before summing the two values. To use physical activity in cross-cohort analysis, the variable was z-standardized.

Behavioral and emotional problems. Mothers reported on behavioral and emotional problems in both the BSPC and the MCS using 20 items of the Strengths & Difficulties Questionnaire (SDQ) (Goodman, 1997). The response scale for each item ranged from zero (*not true*) to two (*certainly true*). The overall score and the sub-scores for the four domains of emotional symptoms (five items), conduct problems (five items), hyperactivity and inattention (five items), and peer problems (five items) were used in analyses. Higher scores correspond to more severe behavioral and emotional problems. Reliability coefficients (α) are shown in Table 1.

Positive well-being. Positive well-being was self-reported with the Kidscreen-52 (Ravens-Sieberer et al., 2005) in the Basel Study of Preterm Children (BSPC), and with single items in the Millennium Cohort Study (MCS). Facets of positive well-being measured in both studies included *psychological well-being* (Kidscreen-52-example item: “Have you been satisfied with your life?”, MCS-single item: “How do you feel about your life as a whole?”), *self-perception* (Kidscreen-52-example item: “Did you worry about your

appearance?", MCS-single item: "How do you feel about the way you look?"), *peer relations* (Kidscreen-52-example item: "Could you rely on your friends?", MCS-single item: "How do you feel about your friends?"), *school related well-being* (Kidscreen-52-example item: "Have you been happy at school?", MCS-single item: "How do you feel about the school you go to?"). The overall well-being scale involved all Kidscreen52 items in the BSPC and a mean score of the MCS-single item scales. The Kidscreen-52 included 5-point Likert scales, while the MCS-single items included a 7-point Likert scale (1='completely happy' to 7='not at all happy'); all scales were coded such that higher values denoted higher levels of well-being.

Covariates. The following covariates were included in the analysis: participants' age at the assessment of the outcomes, sex, neurosensory impairment, ethnic minority group membership, and parental education. Neurosensory impairment was defined as parent-reported severe visual impairments, hearing deficits, or motor impairments (including cerebral palsy), or cognitive ability scores of at least 3 SDs below the population average. Cognitive ability scores were assessed with the Wechsler Intelligence Scale for Children-IV in BSPC, at an average age of 10 years (during the 2nd study wave of the BSPC), and with the British Ability Scales (Picture Similarity and Pattern Construction) in the MCS, at the age of 5 years. Ethnic minority group membership was measured with adolescents' first language, which was used as a proxy (0 = German; 1 = other) in the BSPC, and with parent-reported ethnic minority status (0= white British; 1 = other) in the MCS. Parental education was defined by the highest educational level of either parent (0 = no parent had tertiary education; 1 = either the mother or the father had tertiary education).

Statistical Analysis

Preliminary analyses were conducted separately for both cohorts and involved comparing physical activity, behavioral and emotional problems, and well-being between very preterm and term born adolescents as well as examining Pearson's correlations between physical activity and covariates. This was followed by random-effects meta-analysis to compare pooled means of physical activity, behavioral and emotional problems, and well-being of very preterm and term born adolescents across the two cohorts.

As the main analysis, two-step individual participant data (IPD) meta-analysis was conducted. In the first stage, following the procedure of Aiken and West (1991), hierarchical regression analyses were conducted with physical activity, preterm status, and their interaction term (*physical activity X preterm status*) as predictors added to the regression model in consecutive steps. All analyses were controlled for sex, age at assessment, neurosensory impairment, ethnic minority group membership, and parental education. Outcome variables were mother-reported behavioral and emotional problems or child-reported well-being. Outcome variables and physical activity were z-standardized prior to analysis to allow for comparisons between the two cohorts and unstandardized regression coefficients are displayed. These analyses were conducted using SPSS (version 24; IBM Corporation, Armonk NY, USA).

In the second stage, two separate sets of meta-analyses were conducted for main effects of physical activity (i.e. regression coefficients for physical activity before interaction terms were added to the hierarchical regression models) and interaction effects (*physical activity X preterm status*). Pooled effects and 95% confidence intervals (CIs) were computed using the random-effects method with DerSimonian and Laird technique (DerSimonian & Laird, 1986). In all meta-analyses, between-study heterogeneity was tested using the Cochran's Q statistic and quantified by the I^2 value. Low heterogeneity was defined as an I^2 value of 0% to 25%, moderate heterogeneity as an I^2 of 25% to 75%, and high heterogeneity

as an I^2 of 75% to 100%. Two-stage random-effects IPD meta-analysis was conducted using the 'ipdmetan' command (Fisher, 2015) in STATA (version 15.0; StataCorp, College Station TX, USA). We applied Bonferroni correction to account for multiple comparisons ($p=.05/10=.005$).

We further conducted two sensitivity analyses. In the first sensitivity analysis, we repeated all analyses excluding those who were twins/triplets (as including these participants this could have resulted in skewed findings). In the second sensitivity analysis, we used propensity score matching using the 'psmatch2' command in STATA in the MCS cohort, where there was a large difference in the number of participants between the VPT and FT groups, and repeated the regression analysis. Using propensity score matching has advantages including that it provides a more precise treatment effect by creating a balanced data set (Rosenbaum & Rubin, 1983)

Results

Descriptive Statistics

Participants' characteristics are presented in Table 1. In the BSPC sample, very preterm adolescents were less likely to have parents with tertiary education than term born adolescents. Moreover, very preterm adolescents were more likely to belong to ethnic minority groups in the MCS sample. Pooled mean differences based on random effects meta-analysis revealed significantly higher levels of emotional problems in very preterm compared to term born children (Std. mean difference = 0.26, $P = 0.03$; supplemental table 1). However, no further significant differences in mean levels of behavioural and emotional problems, well-being, and physical activity between the groups were found.

In the BSPC sample, physical activity correlated significantly with age ($r = -.42, p < .001$) but not with sex, neurosensory impairment, ethnic minority group membership, and parental education (all p -values > 0.20). In the MCS sample, physical activity was associated with sex ($r = -.12, p < .001$), ethnic group ($r = -.06, p = .002$), and parental education ($r = -.06, p < .001$), indicating lower physical activity among girls, participants with ethnic minority background, and with parents with lower educational levels, but was not correlated with age.

Individual Participant Data (IPD) Meta-Analysis

Table 2 displays the findings of the IPD meta-analysis. In the IPD meta-analysis of the main effects, physical activity was associated with lower levels of emotional symptoms ($\beta = -.048$; 95% CI = $-.081 - -.014$) and peer problems ($\beta = -.086$; 95% CI = $-.120 - -.053$) but higher levels of hyperactivity/inattention ($\beta = .066$; 95% CI = $.016 - .116$). Physical activity was also associated with higher levels of overall well-being ($\beta = .057$; 95% CI = $.022 - .091$), psychological well-being ($\beta = .053$; 95% CI = $.019-.087$), more favourable self-perception ($\beta = .061$; 95% CI = $.028 - .094$), and better school related well-being ($\beta = .056$; 95% CI = $.022 - .090$). Using Bonferroni correction ($p = .05/10 = .005$), the associations between physical activity and hyperactivity/inattention and emotional problems are non-significant. No statistically significant heterogeneity existed between the study cohorts in the analyses of the main effects (p -values $> .14$). Findings remained the same in the sensitivity analysis repeating the analysis without the twin/triplet participants (i.e., all formerly significant associations remained significant at the Bonferroni corrected level of $p < 0.005$; Supplementary Table 1).

Table 2.

Meta-analysis of associations of physical activity as well as physical activity x preterm status interactions with behavioral/ emotional difficulties and well-being.

	Data Points	β	<i>p</i>	95% CI Lower Bound	95% CI Upper Bound	Cochran <i>Q</i> Test	<i>I</i> ²	Test for Heterogeneity (<i>P</i>)
Main Effects of Physical Activity								
Total Behavioral/ Emotional Difficulties	2	-.017	.819	-.164	.130	1.90	47.3	.16
Emotional symptoms	2	-.048	.005	-.081	-.014	0.76	0.5	.47
Conduct problems	2	.007	.938	-.173	.187	2.20	54.5	.13
Hyperactivity/Inattention	2	.066	.010	.016	.116	1.06	5.4	.30
Peer Problems	2	-.086	<.001	-.120	-.053	0.63	0.0	.42
Overall Well-Being	2	.057	.001	.022	.091	0.17	0.0	.67
Psychological	2	.053	.002	.019	.087	0.24	0.0	.62
Self-perception	2	.061	<.001	.028	.094	0.32	0.0	.57
Peer relations	2	.014	.440	-.021	.049	0.34	0.0	.55
School related well-being	2	.056	.001	.022	.090	0.13	0.1	.77
Interaction between Preterm status and Physical Activity								
Total Behavioral/ Emotional Difficulties	2	-.244	.084	-.521	.033	1.07	6.5	.30
Emotional symptoms	2	-.157	.353	-.426	.112	0.22	0.0	.64
Conduct problems	2	-.127	.366	-.404	.149	0.20	0.0	.65
Hyperactivity/Inattention	2	-.281	.046	-.557	-.005	0.94	0.0	.33
Peer Problems	2	-.182	.183	-.451	.086	0.16	0.0	.69
Overall Well-Being	2	.070	.842	-.621	.762	5.89	83.0	.02
Psychological	2	-.098	.730	-.651	.456	4.05	75.3	.04
Self-perception	2	-.058	.837	-.604	.489	3.77	73.5	.05
Peer relations	2	.052	.713	-.225	.328	0.15	0.1	.79
School related well-being	2	.030	.896	-.416	.475	2.61	61.8	.10

Note. Dependent variables and physical activity were z-standardized before analyses; unstandardized regression coefficients are displayed.

In the IPD meta-analysis of interaction effects the only significant association was between *physical activity x preterm status* interaction and hyperactivity/inattention ($\beta = -.281$; 95% CI = $-.557 - -.005$), which became nonsignificant after correction for multiple comparisons. Apart from this association, there were no significant *physical activity x preterm status* interactions with any of the outcome variables. There was significant heterogeneity between the two samples in general and psychological well-being, which were not explained by female sex, minority ethnicity and university level parental education (*P*s $>.05$). For both domains, the heterogeneity was indicative of increased well-being among

very preterm adolescents with higher physical activity in the BSPC sample, but of an inverse relationship for the MCS.

The findings of the hierarchical regression analyses for the two cohorts (i.e. the first stage of the IPD meta-analysis) are displayed in Table 3. Findings regarding the physical activity x preterm status interactions remained similar when using propensity score matching for the MCS cohort (data not shown).

Table 3.

Hierarchical regressions of child- and mother-reported outcomes (z-standardized).

	<i>Basel Study of Preterm Children</i>						<i>Millennium Cohort Study</i>					
	<i>n=40 Very preterms ($\leq 32^{nd}$ Gest. week); n=59 Term borns ($\geq 37^{th}$ Gest. week)</i>						<i>n=45 Very preterms ($\leq 32^{nd}$ Gest. week); n=3137 Term borns ($\geq 37^{th}$ Gest. week)</i>					
	Physical activity¹			Preterm \times PA			Physical activity¹			Preterm \times PA		
	β	SE	p	β	SE	p	β	SE	p	β	SE	p
Mental health (mother-rated SDQ):												
Overall behavioral and emotional difficulties	-.167	.111	0.136	-.394	0.212	0.066	-.012	.016	0.459	-.109	.175	0.530
Emotional symptoms	-.123	.109	0.261	-.225	0.211	0.289	-.045	.017	0.009	-.096	.178	0.590
Conduct problems	-.166	.119	0.168	-.194	0.231	0.402	.012	.016	0.44	-.067	.169	0.690
Hyperactivity-inattention	-.052	.117	0.656	-.427	0.224	0.061	.069	.017	<0.001	-.150	.175	0.391
Peer problems	-.171	.108	0.119	-.240	0.210	0.255	-.084	.017	<0.001	-.130	.178	0.465
Well-being (child-rated):												
Overall well-being (scales combined)	.105	.117	0.378	.444	0.226	0.053	.055	.017	0.002	-.262	.182	0.151
Psychological	.105	.110	0.341	.201	0.215	0.352	.051	.017	0.003	-.364	.180	0.044
Self-perception	.125	.116	0.282	.238	0.226	0.296	.058	.017	0.001	-.320	.177	0.072
Peer relations	-.049	.109	0.654	.091	0.213	0.668	.016	.018	0.384	.016	.187	0.931
School related well-being	.024	.111	0.828	.270	0.215	0.212	.057	.018	0.001	-.185	.182	0.309

Note. Dependent variables and physical activity were z-standardized before analyses; unstandardized regression coefficients are displayed.

All models control preterm birth status, sex, age, sensory or motor impairments, ethnic minority status, and parental education.

¹ Coefficients for physical activity from hierarchical regression models before the interaction term is entered to the model.

Discussion

IPD meta-analysis for the two cohorts revealed that higher levels of physical activity were associated with lower levels of peer problems reported by mothers. Similarly, higher levels of physical activity were associated with higher overall well-being, psychological and school-related well-being and better self-perception and body image as reported by adolescents themselves. The association of physical activity with behavioral and emotional

problems and well-being was not found to be different between very preterm born and term born adolescents. Thus, consistent with the first of the three competing explanatory models, physical activity is similarly associated with behavioral and emotional problems and well-being among all adolescents. Very preterm adolescents were neither more vulnerable nor more susceptible to the effects of physical activity. However, the observed effect sizes were small.

It has been suggested that physical activity may have a positive effect on mental health and well-being among adolescents, as it may evoke positive social feedback and recognition from peers, which in turn improves self-image and decreases social inhibition and anxiety (Kirkcaldy, Shephard & Siefen, 2002). Very preterm adolescents and adults are more often socially withdrawn and anxious (Eryigit-Madzwamuse, Strauss, Baumann, Bartmann & Wolke, 2015; Pyhälä et al., 2017) as well as socially excluded and bullied by peers (Wolke, Baumann, Strauss, Johnson & Marlow, 2015). Activities that decrease social anxiety and enhance peer recognition, such as physical activity, were expected to particularly improve mental health and well-being in very preterm adolescents. However, the findings indicate that physical activity was associated with improved self-perception and decreased peer problems among both very preterm and term born adolescents.

Finally, the findings are consistent with studies indicating that during adolescence, those born very preterm show similar levels of physical activity as their term born counterparts (Dahan-Oliel, Mazer & Majnemer, 2012; Nordvall-Lassen et al., 2018; Spiegler, Mendonça & Wolke, 2019), while differences may develop later in young adulthood when those born preterm at very low birth weight (<1500 g) report over 50% less physical activity than term born peers, measured by questionnaire, while no differences were seen when measured by accelerometry (Kaseva et al., 2012; Tikanmäki et al., 2017).

The following limitations of the study need to be mentioned. **First**, preliminary analyses only showed significant mean differences between very preterm and term born adolescents regarding emotional problems, while no further significant differences in behavioral and emotional problems and well-being were found in both cohorts. However, there is a large body of evidence suggesting that differences regarding attention difficulties and peer problems exist as well (Aarnoudse-Moens et al., 2009; Johnson & Wolke, 2013; Pyhälä et al., 2017). It is possible that the main analyses were affected as the cohorts included very preterm children with less attention difficulties and peer problems than expected. Relatedly, the two samples differ in how they were recruited and how representative they are for the entire population of very preterm and term born children. While it is conceivable that generally analyses based on the Millennium Cohort Study may lead to more generalizable conclusions, this may not necessarily transfer to analyses related to the relatively small subsample of very preterm children. **Second**, while IPD meta-analysis of two cohorts was used to increase statistical power, the overall number of very preterm adolescents studied was still relatively small and power to detect interaction effects was limited (observed power for the strongest interaction effect, i.e. regarding hyperactivity and inattention, was 0.72 before Bonferroni correction). This is in contrast to the much larger number of term born adolescents particularly in the MCS, which allowed examining the main effects of physical activity with ample statistical power. **Third**, the two cohort studies used different accelerometry devices, different definitions for valid accelerometry recordings, and different scales to measure wellbeing. Relatedly, the accelerometers in both studies were worn on the wrist, while studies on physical activity often use waist-worn devices (e.g. Barreira et al., 2015; Troiano et al., 2008). Further, while MCS recruited all participants early in life, the BSPPC followed a case-control approach matching very preterm children with control children according to age and sex at the first study wave when the children were 8 years old. These

differences may have increased heterogeneity of the findings between the two studies. **Fourth**, only a relatively small percentage of the original birth cohorts had valid accelerometry readings in adolescence. While in MCS the drop out was not selective for very preterm birth, a dropout analysis from birth to follow-up was not feasible for the BSPC because the comparison sample was recruited at the first follow-up time point at 8 years of age on average. **Fifth**, while the use of objective measures of physical activity may be regarded as a strength of the study, it may also involve a limitation. Accelerometry is better able to measure the number of steps, while it is not sensitive to other types of physical activity such as cycling or static exercise (see Martikainen et al 2012 for a discussion). It is possible that some types of exercise have more positive effect on well-being among adolescents than others, as it has been suggested for learning to play golf vs. playing soccer (Schulze, 2019). Particularly highly competitive team sports might also impose a threat to self-esteem and well-being for some adolescents. The current study does not inform us regarding the actual type of physical activity that has the greatest potential to positively affect mental health and well-being among very preterm and term born adolescents. **Finally**, causality cannot be inferred due to the observational cross-sectional design.

Conclusion

In conclusion, this study suggests that physical activity is associated with improved mental health and well-being for all adolescents, whether born full term or very preterm, although the effect size is small. It remains a task for future studies to determine whether distinct types and aspects of physical activity have a greater influence among very preterm or term born adolescents.

Author Contributions: A Brylka contributed to the data curation, formal analysis, and writing the initial manuscript.

D W contributed to the conceptualisation, funding acquisition, methodology, supervision, and reviewing/editing of the manuscript.

S Ludyga contributed to the formal analysis, resources, and reviewing/editing of the manuscript.

A Bilgin contributed to formal analysis and reviewing/editing of the manuscript

J S contributed to the methodology, formal analysis, and reviewing/editing of the manuscript.

H T contributed to data curation and reviewing/editing of the manuscript.

A G contributed to funding acquisition and reviewing/editing of the manuscript.

M G contributed to the formal analysis, resources, and reviewing/editing of the manuscript.

S B contributed to funding acquisition and reviewing/editing of the manuscript.

A G contributed to funding acquisition and reviewing/editing of the manuscript.

P W contributed to funding acquisition and reviewing/editing of the manuscript.

K H contributed to investigation of the literature and reviewing/editing of the manuscript.

E K contributed to investigation of the literature, funding acquisition, and reviewing/editing of the manuscript.

K R contributed to investigation of the literature, funding acquisition, and reviewing/editing of the manuscript.

S Lemola contributed to the conceptualisation, funding acquisition, methodology, data curation, investigation, and reviewing/editing of the manuscript.

All authors read and approved the final manuscript.

Funding: The research was funded by the New Opportunities for Research Funding Agency Co-operation in Europe (NORFACE), Dynamics of Inequality Across the Life-course

Program (DIAL; Grant number: 462-16-040). The Basel Study of Preterm Children was funded by the Swiss National Science Foundation (Projects: Sleep, cognitive, and socio-emotional development in preterm children during middle and late childhood, Grant number: 143962; Socio-emotional development and mental health of pre-term children: The role of HPA axis function, sleep, neuroplasticity, and physical exercise during the transition to adolescence, Grant number: 159362). The funding sources had no involvement in study design, collection, analysis, and interpretation of the data, the writing of the manuscript, and the submission process.

Acknowledgements: We thank The Center for Longitudinal Studies, UCL Institute of Education for the use of these data and the UK Data Archive and UK Data Service for making them available. However, they bear no responsibility for the analysis or interpretation of these data. We thank all participants of the Basel Study of Preterm Children and of the Millennium Cohort Study.

Conflicts of Interest: None

References

- Aarnoudse-Moens, C. S. H., Weisglas-Kuperus, N., van Goudoever, J. B., & Oosterlaan, J. (2009). Meta-analysis of neurobehavioral outcomes in very preterm and/or very low birth weight children. *Pediatrics*, *124*, 717–728.
- Aiken, L. S., & West, S. G. (1991). *Multiple regression: Testing and interpreting interactions*. Newbury Park, CA: Sage.
- Andersen, J. R., Natvig, G. K., Aadland, E., Moe, V. F., Kolotkin, R. L., Anderssen, S. A., & Resaland, G. K. (2017). Associations between health-related quality of life, cardiorespiratory fitness, muscle strength, physical activity and waist circumference in 10-year-old children: the ASK study. *Quality of life research*, *26*(12), 3421-3428.
- Barreira, T. V., Schuna, J. M., Tudor-Locke, C., Chaput, J. P., Church, T. S., Fogelholm, M., ... & Maher, C. (2015). Reliability of accelerometer-determined physical activity and sedentary behavior in school-aged children: a 12-country study. *International journal of obesity supplements*, *5*(S2), S29.
- Biddle, S. J., & Asare, M. (2011). Physical activity and mental health in children and adolescents: a review of reviews. *British journal of sports medicine*, *45*(11), 886-895.
- Booker, C. L., Skew, A. J., Kelly, Y. J., & Sacker, A. (2015). Media use, sports participation, and well-being in adolescence: Cross-sectional findings from the UK household Longitudinal Study. *American Journal of Public Health*, *105*, 173–179.
- Brand, S., Gerber, M., Colledge, F., Holsboer-Trachsler, E., Pühse, U., & Ludyga, S. (2019). Acute Exercise and Emotion Recognition in Young Adolescents. *Journal of Sport and Exercise Psychology*, (00), 1-8.

- Cerrillo-Urbina, A. J., García-Hermoso, A., Sánchez-López, M., Pardo-Guijarro, M. J., Santos Gómez, J. L., & Martínez-Vizcaíno, V. (2015). The effects of physical exercise in children with attention deficit hyperactivity disorder: A systematic review and meta-analysis of randomized control trials. *Child: care, health and development*, *41*(6), 779-788.
- Connelly, R. & Platt, L. (2014) Cohort profile: UK Millennium Cohort Study (MCS). *International Journal of Epidemiology*, *43*, 1719-1725.
- Dahan-Oliel, N., Mazer, B., & Majnemer, A. (2012). Preterm birth and leisure participation: A synthesis of the literature. *Research in Developmental Disabilities*, *33*, 1211–1220.
- Dahan-Oliel, N., Mazer, B., Maltais, D. B., Riley, P., Nadeau, L. & Majnemer, A. (2014) Child and environmental factors associated with leisure participation in adolescents born extremely preterm. *Early Human Development*, *90*, 665-672.
- DerSimonian R, Laird N. (1986). Meta-analysis in clinical trials. *Controlled Clinical Trials*, *7*(3):177–188.
- Eryigit-Madzwamuse, S., Strauss, V., Baumann, N., Bartmann, P., & Wolke, D. (2015). Personality of adults who were born very preterm. *Archives of Disease in Childhood-Fetal and Neonatal Edition*, *100*(6), F524-F529.
- Evensen KAI, Ustad T, Tikanmäki M, Haaramo P, Kajantie E. Long-term motor outcomes of very preterm and/or very low birth weight individuals without cerebral palsy: A review of the current evidence. *Semin Neonatol*. In press.
- Fisher DJ. (2015). Two-stage Individual Participant Data Meta-analysis and Generalized Forest Plots. *The Stata Journal*, *15*(2):369-396.

- Galán, I., Boix, R., Medrano, M. J., Ramos, P., Rivera, F., Pastor-Barriuso, R., & Moreno, C. (2013). Physical activity and self-reported health status among adolescents: a cross-sectional population-based study. *BMJ open*, 3(5), e002644.
- Goodman R. (1997). The strengths and difficulties Questionnaire: a research note. *Journal of Child Psychology and Psychiatry* 38, 581e6.
- Heywood J. (2018). *Millennium Cohort Study. Physical Activity: Accelerometer dataset MCS6(2015). User Guide*. UCL; Centre for Longitudinal Studies. Retrieved from http://doc.ukdataservice.ac.uk/doc/8156/mrdoc/pdf/mcs6_2018_accelerometer.pdf. Accessed July 2019.
- Johnson, S., & Wolke, D. (2013). Behavioural outcomes and psychopathology during adolescence. *Early Human Development*, 89(4), 199-207.
doi:<http://dx.doi.org/10.1016/j.earlhumdev.2013.01.014>
- Kaseva N, Wehkalampi K, Strang-Karlsson S, Salonen M, Pesonen AK, Räikkönen K, Tammelin T, Hovi P, Lahti J, Heinonen K, Järvenpää AL, Andersson S, Eriksson JG, Kajantie E. Lower conditioning leisure-time physical activity in young adults born preterm at very low birth weight. *PLoS One* 2012;7:e32430.
- Kajantie, E., Strang-Karlsson, S., Hovi, P., Räikkönen, K., Pesonen, A. K., Heinonen, K., ... & Andersson, S. (2010). Adults born at very low birth weight exercise less than their peers born at term. *The Journal of Pediatrics*, 157(4), 610-616.
- Kirkcaldy, B. D., Shephard, R. J., & Siefen, R. G. (2002). The relationship between physical activity and self-image and problem behaviour among adolescents. *Social Psychiatry and Psychiatric Epidemiology*, 37, 544–550.

- Liu, M., Wu, L., & Ming, Q. (2015). How does physical activity intervention improve self-esteem and self-concept in children and adolescents? Evidence from a meta-analysis. *PloS one*, 10(8), e0134804.
- Lubans, D., Richards, J., Hillman, C., Faulkner, G., Beauchamp, M., Nilsson, M., Kelly, P., Smith, J., Raine, L. and Biddle, S. (2016). Physical Activity for Cognitive and Mental Health in Youth: A Systematic Review of Mechanisms. *Pediatrics*, 138(3):e20161642
- Martikainen, S., Pesonen, A. K., Lahti, J., Heinonen, K., Tammelin, T., Kajantie, E., ... & Rääkkönen, K. (2012). Physical activity and psychiatric problems in children. *The Journal of Pediatrics*, 161(1), 160-162.
- Mattocks, C., Leary, S., Ness, A., Deere, K., Saunders, J., Tilling, K., Kirkby, J., Blair, S.N. and Riddoch, C. (2007). Calibration of an accelerometer during free-living activities in children. *International Journal of Pediatric Obesity*, 2(4), 218-226.
- Maurer, N., Perkinson-Gloor, N., Stalder, T., Hagmann-von Arx, P., Brand, S., Holsboer-Trachsler, E., Wellmann, S., Grob, A., Weber, P. and Lemola, S. (2016). Salivary and hair glucocorticoids and sleep in very preterm children during school age. *Psychoneuroendocrinology*, 72, 166–174.
- Nordvall-Lassen, M., Hegaard, H. K., Obel, C., Lindhard, M. S., Hedegaard, M., & Henriksen, T. B. (2018). Leisure time physical activity in 9-to 11-year-old children born moderately preterm: a cohort study. *BMC pediatrics*, 18(1), 163.
- Podsakoff, P. M., MacKenzie, S. B., Lee, J. Y., & Podsakoff, N. P. (2003). Common method biases in behavioral research: A critical review of the literature and recommended remedies. *Journal of applied psychology*, 88(5), 879.

- Poulsen, G., Kurinczuk, J. J., Wolke, D., Boyle, E. M., Field, D., Alfirevic, Z., & Quigley, M. A. (2011). Accurate reporting of expected delivery date by mothers 9 months after birth. *Journal of Clinical Epidemiology*, *64*(12), 1444-1450.
doi:10.1016/j.jclinepi.2011.03.007
- Pyhälä, R., Wolford, E., Kautiainen, H., Andersson, S., Bartmann, P., Baumann, N., . . . Rääkkönen, K. (2017). Self-Reported Mental Health Problems Among Adults Born Preterm: A Meta-Analysis. *Pediatrics*. doi:10.1542/peds.2016-2690
- Ravens-Sieberer, U., Gosch, A., Rajmil, L., Erhart, M., Bruil, J., Duer, W., Auquier, P., Power, M., Abel, T., Czemy, L. and Mazur, J. (2005). KIDSCREEN-52 quality-of-life measure for children and adolescents. *Expert review of pharmacoeconomics & outcomes research*, *5*(3), 353-364.
- Rosenbaum, P.R. & Rubin D.B. (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika*, *70* (1), 41-55.
- Schulze, C. (2019). Effect of playing golf on children's mental health. *Mental Health & Prevention*, *13*, 31-34.
- Spiegler, J., Mendonca, M., & Wolke, D. (2019). Prospective Study of Physical Activity of Preterm Born Children from Age 5 to 14 Years. *The Journal of pediatrics*, *208*, 66-73.
- Spiegler, J., Eves, R., Mendonça, M., & Wolke, D. (2019). Association of physical activity and cardiorespiratory function or BMI and body composition in preterm-born individuals: a systematic review. *Acta Paediatrica*, *108*(7), 1205-1214.
- Spittle, A. J., Cameron, K., Doyle, L. W., Cheong, J. L., & Victorian Infant Collaborative Study Group. (2018). Motor Impairment Trends in Extremely Preterm Children: 1991–2005. *Pediatrics*, *141*(4), e20173410.

- Takken, T., van Haastert, I. C., Scholman, W. L., Block, A. J., Eijssermans, M. J. C., & Vries, L. S. (2010). Aerobic exercise capacity and its relationship with pulmonary function, muscle strength, physical activity and motor competence in preterm born children: The PREMAFIT-pilot study. In D. C. Lieberman (Ed.), *Aerobic Exercise and Athletic Performance: Types, Duration and Health Benefits*, 293–306. London, United Kingdom: Nova Science Publishers Inc.
- Tikanmäki M, Kaseva N, Tammelin T, Sipola-Leppänen M, Matinolli HM, Eriksson JG, Järvelin MR, Vääräsmäki M, Kajantie E. Leisure-time physical activity in young adults born preterm. *J Pediatr* 2017;189:135-142.
- Troiano, R. P., Berrigan, D., Dodd, K. W., Masse, L. C., Tilert, T., & McDowell, M. (2008). Physical activity in the United States measured by accelerometer. *Medicine & Science in Sports & Exercise*, 40(1), 181-188.
- Van Lieshout, R. J., Ferro, M. A., Schmidt, L. A., Boyle, M. H., Saigal, S., Morrison, K. M., & Mathewson, K. J. (2018). Trajectories of psychopathology in extremely low birth weight survivors from early adolescence to adulthood: a 20-year longitudinal study. *Journal of Child Psychology and Psychiatry*, 59(11), 1192-1200.
- Wen, C. K. F., Liao, Y., Maher, J. P., Huh, J., Belcher, B. R., Dzubur, E., & Dunton, G. F. (2018). Relationships among affective states, physical activity, and sedentary behavior in children: Moderation by perceived stress. *Health Psychology*, 37(10), 904.
- Wolke, D. (2018). Preterm birth: high vulnerability and no resiliency? Reflections on van Lieshout et al.(2018). *Journal of Child Psychology and Psychiatry*, 59(11), 1201-1204.
- Wolke, D., Baumann, N., Strauss, V., Johnson, S., & Marlow, N. (2015). Bullying of preterm children and emotional problems at school age: cross-culturally invariant effects. *The Journal of pediatrics*, 166(6), 1417-1422.

Wolke, D., Jaekel, J., Hall, J., & Baumann, N. (2013). Effects of sensitive parenting on the academic resilience of very preterm and very low birth weight adolescents. *Journal of Adolescent Health, 53*(5), 642-647.

Supplementary Table 1.

Meta-analysis of associations of physical activity as well as physical activity x preterm status interactions with behavioral/ emotional difficulties and well-being without twins/triplets.

	Data Points	β	p	95% CI Lower Bound	95% CI Upper Bound	Cochran Q Test	I ²	Test for Heterogeneity (P)
Main Effects of Physical Activity								
Total Behavioral/ Emotional Difficulties	2	-.011	.501	-.045	.022	.01	.000	.943
Emotional symptoms	2	-.043	.013	-.077	-.009	.01	.000	.908
Conduct problems	2	.010	.542	-.022	.042	.08	.000	.772
Hyperactivity/Inattention	2	.071	<.001	.038	.105	.05	.000	.818
Peer Problems	2	-.083	<.001	-.117	-.049	.06	.000	.808
Overall Well-Being	2	.054	.002	.019	.088	.27	.000	.602
Psychological	2	.049	.004	.015	.083	.42	.000	.517
Self-perception	2	.060	.001	.026	.093	.000	.000	.978
Peer relations	2	.013	.474	-.022	.048	.50	.000	.480
School related well-being	2	.056	.001	.022	.091	.18	.000	.670
Interaction between Preterm status and Physical Activity								
Total Behavioral/ Emotional Difficulties	2	-.129	.470	-.480	.221	.30	.000	.582
Emotional symptoms	2	-.068	.705	-.419	.283	.000	.000	.948
Conduct problems	2	-.169	.436	-.594	.256	1.28	22.0	.258
Hyperactivity/Inattention	2	-.174	.344	-.533	.186	.85	.000	.358
Peer Problems	2	-.086	.636	-.444	.271	0.02	.000	.895
Overall Well-Being	2	.111	.678	-.413	.635	1.92	48.0	.165
Psychological	2	-.118	.567	-.522	.286	1.30	23.1	.254
Self-perception	2	-.074	.706	-.458	.310	1.12	10.4	.291
Peer relations	2	.057	.758	-.307	.421	.15	.000	.695
School related well-being	2	.067	.721	-.300	.434	.43	.000	.512

Please note that we applied Bonferroni correction ($p=0.05/10=0.005$) to account for the influence of multiple comparisons

