



# Kent Academic Repository

Hambly, Karen, Mundy, Emma, Poomsalood, Somruthai and Stephens, D. (2018) *Return to running following knee osteochondral repair using an anti-gravity treadmill (P03-581)*. In: *Knee Surgery, Sports Traumatology, Arthroscopy. Abstracts of the 18th ESSKA Congress, 9 – 12 May 2018, Glasgow, UK*. 26 (Supp 1). S174. Springer

## Downloaded from

<https://kar.kent.ac.uk/67001/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1007/s00167-018-4867-9>

## This document version

Author's Accepted Manuscript

## DOI for this version

## Licence for this version

UNSPECIFIED

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

## Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

# Return to running following knee osteochondral repair using an anti-gravity treadmill

Authors, K. HAMBLY<sup>1</sup>, S. POOMSALOOD<sup>1</sup>, E. MUNDY<sup>1</sup> and D. STEPHENS<sup>1</sup>

<sup>1</sup>School of Sport and Exercise Sciences, University of Kent, Medway, United Kingdom

## INTRODUCTION

Anti-gravity treadmills are being increasingly used after knee surgery to reduce ground reaction forces during walking and running.<sup>1,2</sup>

Antigravity treadmills have been shown to be safe and feasible to use in early rehabilitation following total knee replacement.<sup>1</sup> However, there are no studies on the use of antigravity treadmills in a knee osteochondral population despite their increasing inclusion in rehabilitation guidelines.

## AIM

The purpose of this study was to assess the impact of an anti-gravity treadmill return to running programme on self-efficacy and subjective knee function following knee osteochondral surgery.

## METHOD

Two otherwise healthy female endurance runners who had undergone knee osteochondral surgery were recruited.

Patient A - 39 year old 9 months post-Bone Marrow Aspirate Concentrate (BMAC)<sup>3,4</sup> for a left knee femoral cartilage grade 3-4 defect 3 cm<sup>2</sup>.

Patient B - 54 year old 11 weeks post-surgery for a partial lateral meniscectomy and chondroplasty.

An anti-gravity treadmill (Figure 1) was used to manipulate loading during a graduated phased return to running (Table 1).

Self-efficacy was evaluated using the Self-Efficacy for Rehabilitation outcomes scale (SER)<sup>5</sup> and the Knee Self-Efficacy Scale (K-SES).<sup>6</sup> Subjective knee function was evaluated using the Knee Injury and Osteoarthritis Outcome Score (KOOS).<sup>7</sup>

## METHOD



FIGURE 1. Anti-gravity treadmill

Week	Percentage Body Weight (%)	Running Speed (km/hr)	Running Time (Mins)	Rating of Perceived Exertion (RPE)
1	30	6.7	5	7
2	30	7.2	10	7
3 Session 1	40	7.6	10	8
3 Session 2	40	7.7	15	9
4 Session 1	50	7.5	15	9.5
4 Session 2	50	8.0	20	11
5 Session 1	60	8.3	20	11.5
5 Session 2	60	8.0	25	11.5
6 Session 1	70	7.5	25	11
6 Session 2	70	7.1	30	11.5
7	80	8.0	30 mins alternating 5 mins running & 5 mins walking	11
8	80	7.5	30 mins alternating 5 mins running & 5 mins walking	10

TABLE 1. Example anti-gravity treadmill rehabilitation programme

## RESULTS




FIGURE 2a. Patient A Self-efficacy scores across the anti-gravity treadmill programme on a scale of 0-10 where a higher score indicates a greater level of self-efficacy.




FIGURE 2b. Patient A KOOS subscale scores across the anti-gravity treadmill programme on a scale of 0-100 where a higher score indicates a greater level of function.




FIGURE 3a. Patient B Self-efficacy scores across the anti-gravity treadmill programme on a scale of 0-10 where a higher score indicates a greater level of self-efficacy.




FIGURE 3b. Patient B KOOS subscale scores across the anti-gravity treadmill programme on a scale of 0-100 where a higher score indicates a greater level of function.

### Summary: Patient A

- SER increased 57%
- K-SES present increased 89%
- K-SES future increased 65%
- KOOS Sport/Rec subscale showed a clinically important improvement<sup>8</sup>

### Summary: Patient B

- SER increased 18%
- K-SES present increased 33%
- K-SES future increased 33%
- KOOS Sport/Rec and QoL subscales showed clinically important improvements<sup>8</sup>

## CONCLUSIONS

The programmes resulted in improved knee and rehabilitation self-efficacy and subjective knee function following osteochondral repair of the knee.

These case reports illustrate the importance of considering self-efficacy in individualising rehabilitation after knee osteochondral surgery and highlights the potential role for anti-gravity treadmills in enhancing self-efficacy and subjective knee function in preparation for a return to sport.

## REFERENCES

1. Bugbee WD, Pulido PA, Goldberg T, D'Lima DD. Use of an anti-gravity treadmill for early postoperative rehabilitation after total knee replacement: A pilot study to determine safety and feasibility. *Am J Orthop (Belle Mead NJ)*. 2016;45(4):E167-73.
2. Eastlack RK, Hargens AR, Groppo ER, Steinbach GC, White KK, Pedowitz RA. Lower body positive-pressure exercise after knee surgery. *Clin Orthop Rel Res*. 2005;431:213-19.
3. Chahla J, Dean CS, Moatshe G, Pascual-Garrido C, Serra Cruz R, LaPrade RF. Concentrated bone marrow aspirate for the treatment of chondral injuries and osteoarthritis of the knee: A systematic review of outcomes. *Orthop J Sports Med*. 2016;4(1):DOI: 10.1177/2325967115625481.
4. Gobbi A, Karnatzikos G, Scotti C, Mahajan V, Mazzucco L, Grigolo B. One-step cartilage repair with bone marrow aspirate concentrated cells and collagen matrix in full-thickness knee cartilage lesions: Results at 2-year follow-up. *Cartilage*. 2011;2(3):286-99.
5. Waldrop D, Lightsey O, Ethington C, Woemmel C, Coke A. Self-efficacy, optimism, health competence and recovery from orthopaedic surgery. *J Cousins Psychol*. 2001;48:233-8.
6. Thomee P, Wahrborg P, Borjesson M, Thomee R, Eriksson BI, Karlsson J. A new instrument for measuring self-efficacy in patients with an anterior cruciate ligament injury. *Scand J Med Sci Sports*. 2006;16(3):181-7.
7. Roos EM, Engelhart L, Ranstam J, Anderson AF, Irrgang JJ, Marx RG, et al. ICRS recommendation document: Patient-reported outcome instruments for use in patients with articular cartilage defects. *Cartilage*. 2011;2(2):122-36.
8. Howard J, Whale Conley CE, Mattacola CG, Lattermann C. Minimal clinically important differences for the Knee Injury Osteoarthritis Outcomes Score (KOOS) among knee cartilage patients. *International Cartilage Repair Society 13th Congress; Sorrento, Italy 2016*.

## CONTACT INFORMATION

Dr Karen Hamby, School of Sport and Exercise Sciences, Medway Building, University of Kent, Chatham Maritime, Kent ME4 4AG, UK

[K.Hamby@kent.ac.uk](mailto:K.Hamby@kent.ac.uk)

@karenhamby