



Kent Academic Repository

Neufuss, Johanna, Robbins, Martha M., Baeumer, Jana, Humle, Tatyana and Kivell, Tracy L. (2018) *Gait characteristics of vertical climbing in mountain gorillas and chimpanzees*. *Journal of Zoology*, 306 (2). pp. 129-138. ISSN 0952-8369.

Downloaded from

<https://kar.kent.ac.uk/67175/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1111/jzo.12577>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Gait characteristics of vertical climbing in mountain gorillas and chimpanzees

Johanna Neufuss¹, Martha M. Robbins², Jana Baeumer², Tatyana Humle³, Tracy L. Kivell^{1,4}

¹Animal Postcranial Evolution (APE) Laboratory, Skeletal Biology Research Centre, School of Anthropology & Conservation, University of Kent, Canterbury, UK

²Department of Primatology, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

³Durrell Institute of Conservation and Ecology, School of Anthropology & Conservation, University of Kent, Canterbury, UK

⁴Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

***Corresponding author**

Johanna Neufuss. Animal Postcranial Evolution Laboratory, Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, CT2 7NR, UK. email: jn259@kentforlife.net

Short title: vertical climbing in gorillas and chimpanzees

Number of text pages: 18; **Figures:** 2; **Tables:** 7

Grant sponsorship: This work was supported by the University of Kent 50th anniversary scholarship (to JN), the European Research Council Starting Grant #336301 (to TLK and JN) and the Max Planck Society (to MR, JB, and TLK). We declare we have no competing interests.

Abstract

Biomechanical analyses of arboreal locomotion in great apes in their natural environment are scarce and thus attempts to correlate behavioural and habitat differences with variations in morphology are limited. The aim of this study was to investigate the gait characteristics of vertical climbing in mountain gorillas (*Gorilla beringei beringei*) and chimpanzees (*Pan troglodytes*) in a natural environment to assess differences in the climbing styles that may relate to variation in body size. We investigated temporal variables (i.e., cycle duration, duty factors, and stride frequency) and footfall sequences (i.e., diagonal vs. lateral sequence gaits) during vertical climbing (both ascent and descent) in 11 wild mountain gorillas and compared these data to those of eight semi-free-ranging chimpanzees, using video records *ad libitum*. Comparisons of temporal gait parameters revealed that large-bodied mountain gorillas exhibited a longer cycle duration, lower stride frequency and generally a higher duty factor than smaller-bodied chimpanzees. While both apes were similarly versatile in their vertical climbing performance in the natural environment, mountain gorillas most often engaged in diagonal sequence/diagonal couplet gaits and chimpanzees most often used lateral sequence/diagonal couplet gaits. This study revealed that mountain gorillas adapt their climbing strategy to accommodate their large body mass in a similar manner previously found in captive western lowland gorillas, and that chimpanzees are less variable in their climbing strategy than has been documented in captive bonobos.

Key words: locomotion, gait parameters, biomechanics, apes, gorillas, chimpanzees

Introduction

Vertical climbing is essential to the locomotor and foraging strategies of great apes (e.g., Hunt, 1992a; Remis, 1995; Pilbeam, 2002; Robbins & McNeillage, 2003). Records of the frequency of vertical climbing in wild African apes vary depending on the species and population studied as well as on differences in habitat structure and resource availability (e.g., Tuttle & Watts, 1985; Doran, 1993, 1996; Remis, 1995; Crompton *et al.*, 2010). Most studies agree that mountain gorillas (*Gorilla beringei beringei*) are less arboreal than chimpanzees (*Pan troglodytes*) (Tuttle & Watts, 1985; Remis, 1998; Hunt, 2004; Crompton *et al.*, 2010; Hunt, 2016) and are thought to spend less than 1 % of total locomotor time engaging in vertical climbing (Tuttle & Watts, 1985). However, arboreal locomotor behaviors in mountain gorillas have to date only been broadly described (e.g., Schaller, 1963; Doran, 1996) and the frequency is likely underestimated (Crompton, 2016).

During arboreal locomotion, and particularly vertical climbing, primates face several biomechanical challenges. As substrate inclination increases, the difficulty in maintaining stability also increases, and these challenges are greatest on vertical supports (Cartmill, 1974; Cartmill, 1985; Preuschoft, 2002). When climbing up inclined substrates primates must use their limbs to overcome gravitational force (e.g., Hirasaki *et al.*, 1993, 2000; Preuschoft, 2002; Hanna *et al.*, 2017), while a shear force acts against the direction of movement (Preuschoft, 2002). The higher forces that are required to propel the body upwards, together with the shift in body weight towards the hind limbs, increases the risk of toppling backwards if an animal is not able to make secure substrate contact (e.g., Cartmill, 1974; Preuschoft & Witte, 1991; Preuschoft, 2002). Primates typically must use their hands/forelimb (to pull) and feet/hindlimb (to push) to generate greater propulsive forces as substrate incline increases (Hirasaki *et al.*, 1993; Hanna *et al.*, 2017), although smaller primates (<1 kg) show greater versatility in limb use (Hanna *et al.* 2017). In contrast,

when descending an inclined or vertical substrate, greater braking may be applied to resist acceleration (e.g., Cartmill, 1985; Preuschoft, 2002). Heavier animals will, typically, exhibit a prolonged support phase or higher duty factor (i.e., increased limb contact with the substrate) during vertical climbing to increase stability, enabling a more cautious movement (e.g., Cartmill, 1974; Cartmill & Milton, 1977). On vertical supports, primates overcome some of these biomechanical challenges by using powerful grasping and adapting their forelimb and hand posture (Hunt *et al.*, 1996; Hirasaki *et al.*, 2000; Isler, 2005; Nakano *et al.*, 2006). For example, recent work has found that mountain gorillas and chimpanzees generally use the same hand grips, power grips and diagonal power grips, as well as similar forelimb postures on vertical supports of a similar size, which is consistent with their general similarity in hard and soft tissue morphology of the hand and forelimb (Neufuss *et al.*, 2017).

Attempts to correlate variations in African ape morphology (e.g., hand or limb proportions, body mass) with behavioral and habitat differences are limited because kinematics (e.g., movement of body segments, gait parameters) of arboreal locomotion, such as vertical climbing, are scarce especially in the wild. A few studies have investigated the spatio-temporal parameters and gait characteristics of vertical climbing in apes and other primates, all of which have been conducted in captivity (Hirasaki, Kumakura & Matano, 1993, 2000; Isler 2002, 2003, 2005; Isler & Grueter, 2006; Schoonaert *et al.*, 2016). Within great apes, there is only one naturalistic study that compared the gait parameters of vertical climbing in rehabilitated and wild Sumatran orangutans to captive individuals (Isler & Thorpe, 2003). Captive chimpanzee climbing patterns and limb joint kinematics have been briefly described (Nakano *et al.*, 2006) but gait parameters of fore- and hindlimbs have not been examined. Current knowledge about the spatio-temporal gait characteristics of gorilla vertical climbing stems solely from a captive study of western lowland gorillas

(*Gorilla gorilla gorilla*) using a rope as locomotor support (Isler, 2002). Isler (2002) identified key differences in gorilla climbing performance associated with age and sex; vertical climbing in an adult male gorilla was characterised by higher duty factors, relatively shorter strides and more variable footfall patterns compared with adult female gorillas and bonobos. Isler (2002) interpreted these kinematic differences as evidence that vertical climbing on a rope was more challenging for adult male gorillas due to their larger body mass. However, the vertical climbing 'characteristics' in mountain gorillas and chimpanzees within a natural environment have never been investigated, and the potential differences in the climbing performance between these two apes that differ significantly in body size (e.g., Sarmiento, 1994; Smith & Jungers, 1997) are not yet known.

The aim of this study is to provide further insights into the arboreal locomotor strategies of mountain gorillas and chimpanzees by describing the temporal gait parameters and footfall sequences during vertical climbing (both ascent and descent) on differently-sized natural substrates. We predict that vertical climbing of large-bodied mountain gorillas will be characterized by longer cycle durations, higher duty factors, lower stride frequencies, a higher number of limbs used as support and less variable footfall patterns compared to smaller-bodied chimpanzees. More specifically, we hypothesize that mountain gorillas will adapt their climbing strategy to accommodate their large body mass in a similar manner to that documented in captive western lowland gorillas (females 71.0-97.5 kg; males 162.5-175.2 kg across *G. beringei*, *G. gorilla*, *G. graueri*; Smith & Jungers, 1997) and, likewise, vertical climbing of smaller-bodied chimpanzees will be similar to that of bonobos given their generally similar body size (females 33.2-45.8 kg; males 42.7-59.7 kg across *P. paniscus* and *P. t. troglodytes*, *P. t. schweinfurthii*, and *P. t. verus*; Smith & Jungers, 1997).

Materials and Methods

Species and study sites

Mountain gorillas (*Gorilla beringei beringei*) were observed in the Bwindi Impenetrable National Park (331 km²). Data were collected on one fully habituated group of gorillas (Kyagurilo) between October-December 2014 and March-July 2015 during two fruiting seasons.

Chimpanzee (*Pan troglodytes* ssp.) vertical climbing data were collected between August-September 2014 on two colonies of semi-free-ranging chimpanzees at the Chimfunshi Wildlife Orphanage Trust (CWO), Zambia.

Data collection

Vertical climbing for any given individual was divided into 'sequences' and 'limb cycles'. A 'sequence' was defined as a continued period of climbing behavior. A sequence started when the right hindlimb was initially placed in contact with the substrate and a sequence ended when the individual stopped vertical climbing. A sequence was generally composed of multiple limb cycles. A limb cycle was defined as the interval between touchdown of one limb and the subsequent touchdown of the same limb (i.e., right foot/ hand to right foot/hand).

Both apes were filmed *ad libitum* during vertical ascent and descent climbing using a high-definition HDR-CX240E video camera (Sony, Japan) at a frequency of 50 Hz. The mountain gorillas were observed for an average of 4hours/day. All gorilla climbing sequences were recorded at relatively close range (7 m to ~20 m) on a sample of 11 individuals (Table 1). The gorillas had the opportunity to climb on differently-sized substrates ranging from lianas (6-10 cm diameter) to extremely large tree trunks (>50 cm diameter) (for more details on substrate sizes see Neufuss

et al., 2017). A total of eight adult semi-free-ranging chimpanzees were filmed at relatively close range (~10 m) (Table 1). Although the chimpanzees climbed in a natural environment, the sanctuary is located in a miombo woodland, and thus, substrates were limited to tree trunks of different sizes (i.e., no data were collected on lianas).

Table 1 Studied individuals and number of sequences and limb cycles analyzed.

Species	Individual	Sex/Age	Total no. of climbing sequences	No. of hindlimb cycles for vertical ascent	No. of hindlimb cycles for vertical descent

<i>G. b. beringei</i>	JN	female/adult	11	17	6
	ST	female/adult	11	13	6
	KR	female/adult	9	18	-
	TN	female/ adult	5	10	-
	TW	female/adult	7	10	3
	MG	female/adult	8	2	8
	BY	female/adult	5	3	-
	TD	female/adult	2	2	-
	KA	male/blackback	4	7	-
	MK	male/silverback	6	12	-
	RC	male/silverback	2	3	-
	Total			70	97
<i>P. troglodytes ssp.</i>	RI	female/adult	2	-	6
	KB	female/adult	8	12	7
	MI	female/adult	11	9	15
	KY	female/adult	7	6	6
	JU	female/adult	3	-	8
	UN	female/adult	4	-	5
	TA	male/adult	1	-	4
	CO	male/adult	1	-	2
Total			37	27	53

Types of vertical climbing

Two types of vertical climbing have been previously described in relation to substrate size (Hunt *et al.*, 1996): (1) when climbing on smaller substrates, such a liana or thin

tree, flexion of the elbow helps to elevate the body ('flexed-elbow' vertical climbing); (2) on larger substrates, such as tree trunks, the elbow is typically extended throughout the motion cycle ('extended-elbow' vertical climbing). In both mountain gorillas and chimpanzees, smaller substrates evoke flexed-elbow climbing while a substrate diameter larger than 20 cm is likely to evoke extended-elbow climbing (Hunt *et al.*, 1996; Neufuss *et al.*, 2017).

Analysis of gait characteristics

The footfall sequence and gait parameters such as cycle duration, duty factor and stride frequency were determined by reviewing video sequences frame-by-frame using free motion software (Kinovea 0.8.15). **Cycle duration** (CD) is defined as the time between two initial contacts with the substrate (or 'touchdowns') by the same limb (e.g., Isler 2002, Isler & Thorpe, 2003). The relative duration of the stance phase, or **duty factor** (S), is the fraction of the cycle duration that a particular limb contacts the substrate (Hildebrand, 1966). **Stride frequency** is the number of strides per unit of time, or $1/CD$ (Schoonaert *et al.*, 2016). The limb cycles were classified as either symmetrical or asymmetrical gaits according to the timing of footfalls, following Hildebrand (1967). **Stride symmetry** was calculated as the percentage of cycle duration separating the time between touchdowns of the right and left hindlimbs (Hildebrand, 1966). A cycle was considered symmetrical if the opposing limb's touchdown occurred between 40 % and 60 % of the cycle duration, allowing comparisons with results of Isler (2002, 2003, 2005) and Schoonaert *et al.* (2016). A cycle that was outside of this range (i.e., <40 % or >60 %) was considered as asymmetrical.

Symmetrical cycles were then further classified as being either diagonal sequence (DS) or lateral sequence (LS) gaits. In a **DS gait**, hindlimb touchdown is followed by

the touchdown of the opposite forelimb (right hindlimb > left forelimb), whereas in a

LS **gait**

the	Gait sequence pattern	Definition
-----	-----------------------	------------

ipsilateral forelimb follows (right hindlimb > right forelimb) (Hildebrand, 1966). **Limb phase** (D, also called diagonality) is a quantification of the timing of the footfalls and defined as the duration of time between hindlimb touchdown and touchdown of the ipsilateral forelimb, expressed as a percentage of the stride cycle (Hildebrand, 1966; Cartmill et al., 2002). DS and LS strides can be further subdivided into five categories: diagonal couplets (DC), lateral couplets (LC), single foot (SF), pace and trot (Hildebrand, 1967) based on the relative timing of touchdown of the limbs (see Fig. 1; Table 2). Finally, we further investigated the average number of limbs supporting the animal's body during ascent and descent climbing (i.e., two- vs. three-limb support). Limb support was categorized as diagonal, lateral, tripedal or quadrupedal following Vilensky and Gankiewicz (1989) (Table 2).

Table 2 Definitions of gait sequence patterns^a and types of limb support^b.

diagonal couplets (DC)	diagonally opposite fore-and hindlimb touchdown at the same time
lateral couplets (LC)	the footfalls on the same body side are evenly spaced in time
single foot (SF)	the footfalls of all the limbs are evenly spaced
pace	the footfalls of the fore- and hindlimb on the same side are evenly spaced
trot	diagonally opposite limbs are evenly spaced in time
Types of limb support	Definition
diagonal pair	support by either combination of diagonal limbs
lateral pair	support by either pair of limbs on the same side
tripedal	support by any combination of three limbs
quadrupedal	support by all four limbs

^aGait sequence patterns defined according to Hildebrand (1967) and ^btype of limb support follow definitions by Vilensky and Gankiewicz (1989) and Isler (2002).

Statistics

All statistical analyses were performed using SPSS (IBM SPSS Statistics for Windows, Version 24.0). Gait parameters were analyzed using mean values for all cycles/individual, so that each individual was contributing only a single observation within each condition to ensure independence of data points. Given the small sample size for each species, statistical analyses are used here primarily to summarize the observed patterns rather than to explicitly test hypotheses. Differences in gait parameters between fore- and hindlimbs during vertical ascent vs. descent, including flexed- vs. extended-elbow climbing, were assessed with independent-samples t-tests. Following Isler (2002), significance of inter- and intraspecific (sex classes) differences in the gait parameters between gorillas and chimpanzees was tested using a one-way ANOVA and a *Scheffé's* post-hoc test ($\alpha=0.05$). The overall sample size was too small to allow more sophisticated statistical tests that would account for

species, sex, or type of vertical climbing etc. as fixed effects and the individual as a random effect. Therefore, results of these statistical analyses should be interpreted with caution.

Results

We recorded a total of 70 climbing sequences, containing N=120 limb cycles (ascent: 97; descent: 23), for 11 mountain gorillas and 37 climbing sequences, containing N=80 limb cycles (ascent: 27; descent: 53), for eight chimpanzees (Table 1).

Gait sequence patterns

Mountain gorillas used DS gaits more often (68%) than trot (18%) and LS gaits (14%) during both ascent and descent climbing (Table 3, Fig. 1). Most of these DS gaits were further classified as diagonal couplets, followed by lateral couplets and single foot. Lateral couplets and LS pace were not observed. Chimpanzees showed the opposite pattern, using LS gaits (58%) more often than DS gaits (23%) and trot (20%; Table 3; Fig. 1) during both ascent and descent climbing. Most of these strides were diagonal couplets and single foot gaits. LS pace was not observed in chimpanzees (Fig. 1).

In mountain gorillas, limb phase was significantly higher (mean: 0.62, SD: 0.08; $t(17)=3.59$, $p=0.002$) for most limb cycles (20 cycles: 50-59%, 30 cycles: 60-69%, 18 cycles: 80-89%) than in chimpanzees (mean: 0.46, SD: 0.12; 15 cycles: 20-29%, 21 cycles: 40-49%, 14 cycles: 30-39%). This difference in limb phase is consistent with the more frequent use of DS gaits in mountain gorillas and LS gaits in chimpanzees.

Table 3 Frequency of gait sequence patterns in mountain gorillas and chimpanzees.

	Diagonal Sequence Gaits						Lateral Sequence Gaits				
	DS	DC	SF	LC	Pace	Trot	LS	DC	SF	LC	Pace
Mountain gorillas	68%	42%	23%	24%	11%	18%	14%	59%	41%	-	-
Chimpanzees	23%	33%	39%	17%	11%	20%	58%	59%	39%	2%	-

DS:
diag

onal sequence, LS: lateral sequence, DC: diagonal couplet, SF: single foot, LC: lateral couplet. Note that the total number of strides was N=120 in gorillas and N=80 in chimpanzees. In gorillas, the number of DS gaits types was N=81, trot: N=22 and LS gaits: N=17. In chimpanzees, the number of DS gait types was N=18; trot: N=16 and LS gaits: N=46.




Fig. 1. Classification of footfall patterns during symmetrical strides of vertical climbing in mountain gorillas (black diamonds) and chimpanzees (grey circles) following Hildebrand (1967). The x-axis shows the duty factor, or relative duration of the stance phase in percent of total cycle duration. The y-axis shows the delay of the ipsilateral forelimb following hindlimb touchdown, as a percentage of total cycle duration. Both mountain gorillas and chimpanzees used lateral and diagonal sequence gaits.

Temporal gait parameters

Table 4 summarizes the mean cycle duration and the relative duration of the support phases (i.e., duty factor) of fore- and hindlimbs of mountain gorillas and chimpanzees. When comparisons could be made, there were no significant species differences in either variable in the gait parameters between fore- and hindlimbs during (1) ascent and descent climbing and (2) flexed-elbow and extended-elbow climbing (Table 5). Thus, all results were pooled for each species.

There were no intraspecific significant differences in forelimb or hindlimb cycle duration, stride frequency, or duty factor between males and females within both gorillas and chimpanzees (Table 6). Interspecifically, however, significant variation was found across all gait parameters (Fig. 2, Table 6). Overall, gorillas (i.e., sexes pooled) showed a significantly longer cycle duration (mean: 2.7, SD: 0.8; ANOVA, $F(3)=9.52$, $MSE=1.45$, $p=0.001$) than chimpanzees (mean: 1.6, SD: 0.3) (Fig. 2). Cycle duration was significantly longer in female gorillas (mean: 2.7, SD: 0.9) than in female (mean: 1.6, SD: 0.2) and male (mean: 1.3, SD: 0.4) chimpanzees (Table 6). In contrast, male gorillas had a significantly longer cycle duration (mean: 2.4, SD: 0.4) than male chimpanzees but not compared to female chimpanzees (Table 6).

Table 4 Cycle duration and duty factor of vertical ascent and descent in mountain gorillas and chimpanzees.

Species	Individual	Sex/ Age	No. of limb cycles (hind/fore)	Cycle duration Hindlimb (sec)	Cycle duration Forelimb (sec)	Duty factor Hindlimb (%)	Duty factor Forelimb (%)
<u>Vertical ascent</u>							
<u>extended-elbow climb</u>							
<i>G. b. beringei</i>	JN	Female adult	12/12	1.93 (0.20)	1.87 (0.32)	66 (0.04)	69 (0.04)
	ST	Female adult	13/11	2.76 (0.61)	2.49 (0.68)	70 (0.09)	69 (0.07)
	KR	Female adult	18/18	2.32 (0.54)	2.32 (0.52)	63 (0.04)	68 (0.05)
	TN	Female adult	10/10	2.04 (0.54)	2.45 (0.87)	71 (0.06)	72 (0.07)
	TW	Female adult	10/10	2.37 (0.38)	2.48 (0.58)	68 (0.09)	69 (0.06)
	BY	Female adult	3/3	2.22 (0.08)	2.10 (0.15)	68 (0.02)	71 (0.05)
	TD	Female adult	2/2	3.41 (0.53)	2.96 (0.74)	69 (0.09)	73 (0.08)
	KA	Male blackback	7/7	3.03 (0.53)	2.63 (0.16)	76 (0.06)	71 (0.05)
	MK	Male silverback	10/8	2.26 (0.71)	2.50 (1.65)	72 (0.04)	66 (0.05)
	RC	Male silverback	3/3	2.19 (0.35)	1.93 (0.13)	65 (0.05)	71 (0.03)
<i>P. troglodytes</i> ssp.	KB	Female adult	7/7	1.82 (0.38)	1.82 (0.21)	68 (0.05)	63 (0.07)
	MI	Female adult	9/8	1.73 (0.78)	1.73 (0.72)	65 (0.08)	59 (0.04)
	KY	Female adult	6/5	1.90 (0.47)	1.94 (0.67)	69 (0.07)	66 (0.09)
<u>flexed-elbow climb</u>							
<i>G. b. beringei</i>	JN	Female adult	5/5	2.35 (0.13)	2.35 (0.22)	64 (0.03)	64 (0.04)
	MG	Female adult	2/2	1.86 (0.14)	1.76 (0.21)	72 (0.03)	68 (0.00)
	MK	Male silverback	2/2	2.22 (0.30)	3.39 (0.72)	71 (0.03)	69 (0.01)
<u>Vertical descent</u>							
<u>extended-elbow climb</u>							
<i>G. b. beringei</i>	JN	Female adult	4/3	1.65 (0.24)	2.49 (1.40)	66 (0.03)	65 (0.05)
	ST	Female adult	6/6	2.62 (0.76)	4.53 (0.93)	71 (0.04)	76 (0.04)
	TW	Female adult	3/3	4.34 (3.02)	3.35 (1.91)	69 (0.07)	71 (0.08)
	MG	Female adult	8/8	2.40 (0.70)	3.50 (1.59)	66 (0.06)	65 (0.11)
<i>P. troglodytes</i> ssp.	RI	Female adult	6/6	1.43 (0.38)	1.56 (0.55)	61 (0.09)	61 (0.08)
	KB	Female adult	12/12	1.83 (0.71)	1.78 (0.43)	66 (0.11)	62 (0.08)
	MI	Female adult	15/15	1.40 (0.31)	1.48 (0.34)	64 (0.06)	67 (0.07)
	KY	Female adult	6/6	1.60 (0.32)	1.71 (0.33)	65 (0.04)	65 (0.08)
	JU	Female adult	8/6	1.38 (0.20)	1.40 (0.15)	62 (0.03)	63 (0.07)
	UN	Female adult	5/5	1.81 (0.33)	1.55 (1.60)	68 (0.02)	68 (0.03)
	TA	Male adult	4/4	1.34 (0.14)	1.45 (0.18)	61 (0.02)	58 (0.03)
	CO	Male adult	2/2	1.23 (0.00)	0.94 (0.04)	62 (0.17)	58 (0.05)
<u>flexed-elbow climb</u>							
<i>G. b. beringei</i>	JN	Female adult	2/2	2.42 (0.52)	4.75 (0.47)	73 (0.00)	71 (0.01)
<i>P. troglodytes</i> ssp.	JU	Female adult	2/2	1.28 (0.18)	1.22 (0.47)	68 (0.06)	66 (0.16)
	UN	Female adult	2/2	1.62 (0.30)	1.87 (0.08)	55 (0.04)	68 (0.02)

Table 5 Independent-samples t-test of gait parameters between fore- (RF) and hindlimbs (RH) during vertical ascent and descent.

RF vs. RH during vertical ascent			
Species	cycle duration	stride frequency	duty factor
Mountain gorillas (N=10)	t(22)=0.21, p=0.834	t(22)=0.16, p=0.871	t(24)=-0.25, p=0.8
Chimpanzees (N=3)	t(4)=-1.71, p=0.873	t(4)=0.00, p=1.00	t(4)=1.98, p=0.119

RF vs. RH during vertical descent			
Species	cycle duration	stride frequency	duty factor
Mountain gorillas (N=4)	t(8)=-1.4, p=0.194	t(8)=-5.78, p=0.571	t(8)=0.51, p=0.63
Chimpanzees (N=8)	t(19)=0.35, p=0.728	t(18)=-0.2, p=0.82	t(14)=-1.53, p=0.88

Table 6 Scheffé's post-hoc test of gait parameters in mountain gorillas and chimpanzees.

Intraspecific	cycle duration	stride frequency	duty factor
female vs. male gorillas (N=10)	p=1.000	p=0.999	p=0.566
female vs. male chimpanzees (N=8)	p= 0.888	p=0.287	p=0.811

Interspecific	cycle duration	stride frequency	duty factor
female gorillas vs. female chimpanzees	p<0.001	p=0.001	p=0.002
female gorillas vs. male chimpanzees	p=0.003	p<0.001	p<0.001
male gorillas vs. female chimpanzees	p=0.075	p=0.004	p=0.040
male gorillas vs. male chimpanzees	p=0.005	p=0.001	p=0.003

The duty factor was significantly higher in gorillas overall (mean: 69 %, SD: 3; ANOVA, $F(3)=5.71$, $MSE=53.16$, $p=0.009$) than in chimpanzees (mean: 63 %, SD: 3.6) (Fig. 2). The duty factor was significantly different between female gorillas (mean: 70 %, SD: 3) and female (mean: 63 %, SD: 3) and male (mean: 58 %, SD: 0) chimpanzees (Table 6). Similarly, male gorillas showed a significantly higher duty factor (mean: 69 %, SD: 2.9) than female and male chimpanzees (Table 6). In both apes, the duty factor was most frequently between 60-69 % (gorillas: 65 limb cycles; chimpanzees: 42 limb cycles), but in gorillas the duty factor was higher for more limb cycles (36 limb cycles: 70-79 %, 8 cycles: 80-89 %) than in chimpanzees (18 limb cycles: 70-79 %) (Fig. 2).

The stride frequency was significantly lower in gorillas overall (mean: 0.40, SD: 0.11; ANOVA, $F(3)=22.16$, $MSE=0.12$, $p<0.001$) compared to chimpanzees (mean: 0.65, SD: 0.13) (Fig. 2). Interspecific differences showed that female gorillas exhibited a significantly lower stride frequency (mean: 0.38, SD: 0.12) than female (mean: 0.62, SD: 0.14) and male chimpanzees (mean: 0.78, SD: 0.04). Male gorillas similarly had a significantly lower stride frequency (mean: 0.43, SD: 0.03) than female and male chimpanzees (Table 6).

The average number of supporting limbs was not significantly different between gorillas (ascent: 2.7, SD: 0.4; descent: 2.4, SD: 0.05) and chimpanzees (ascent: 2.6, SD: 0.3; descent: 2.4, SD: 0.1), with both apes using on average three limbs as support more frequently during vertical ascent than during descent (Table 7).




Fig. 2. Box-and-whisker plots of interspecific variation in forelimb cycle duration, duty factor and stride frequency between mountain gorillas and chimpanzees. These differences were significant.

Table 7 Limb support during vertical ascent and descent in mountain gorillas and chimpanzees.

Species	Individual	Sex/Age	Total no. of hindlimb cycles (100%)	D ^a	L ^b	T ^c	Q ^d	Mean no. of supp. limbs
<u>Vertical ascent</u>								
<i>Gorilla beringei b.</i>	JN	Female adult	17	3 (17.6%)	1 (5.9%)	12 (70.6%)	1 (5.9%)	2.8
	ST	Female adult	13	3 (23.1%)	4 (30.8%)	6 (46.2%)	0 (0.0%)	2.5
	KR	Female adult	18	3 (16.7%)	0 (0.0%)	15 (83.3%)	0 (0.0%)	2.8
	TN	Female adult	10	2 (20.0%)	1 (10.0%)	7 (70.0%)	0 (0.0%)	2.7
	TW	Female adult	10	0 (0.0%)	0 (0.0%)	60 (60%)	40 (40%)	3.4
	MG	Female adult	2	0 (0.0%)	0 (0.0%)	2 (100%)	0 (0.0%)	2.0
	TD	Female adult	2	2 (100%)	0 (0.0%)	0 (0%)	0 (0.0%)	2.0
	BY	Female adult	3	2 (66.7%)	0 (0.0%)	1 (33.3%)	0 (0.0%)	2.3
	KA	Male blackback	7	0 (0.0%)	0 (0.0%)	7 (0.0%)	0 (0.0%)	3.0
	MK	Male silverback	12	0 (0.0%)	2 (16.7%)	10 (83.3%)	0 (0.0%)	2.8
	RC	Male silverback	3	0 (0.0%)	0 (0.0%)	3 (100%)	0 (0.0%)	3.0
<i>Pan troglodytes ssp.</i>	KB	Female adult	7	2 (28.6%)	1 (14.3%)	4 (57.2%)	0 (0.0%)	2.6
	MI	Female adult	12	1 (8.3%)	5 (41.6%)	6 (50.0%)	0 (0.0%)	2.5
	KY	Female adult	6	0 (0.0%)	2 (33.3%)	4 (66%)	0 (0.0%)	2.7
<u>Vertical descent</u>								
<i>Gorilla beringei b.</i>	JN	Female adult	6	1 (16.7%)	2 (33.3%)	3 (50.0%)	0 (0.0%)	2.4
	ST	Female adult	6	4 (66.7%)	1 (16.7%)	1 (16.7%)	0 (0.0%)	2.4
	TW	Female adult	3	0 (0.0%)	2 (66.7%)	1 (33.3%)	0 (0%)	2.3
	MG	Female adult	8	5 (62.5%)	0 (0.0%)	3 (37.5%)	0 (0%)	2.4
<i>Pan troglodytes ssp.</i>	RI	Female adult	8	3 (37.5%)	2 (25%)	3 (37.5%)	0 (0%)	2.4
	KB	Female adult	7	0 (0%)	3 (42.8%)	4 (57.2%)	0 (0%)	2.6
	MI	Female adult	16	1 (6.3%)	5 (31.3%)	10 (62.5%)	0 (0%)	2.6
	KY	Female adult	5	0 (0%)	0 (0%)	5 (100%)	0 (0%)	3.0
	JU	Female adult	8	5 (62.5%)	0 (0%)	3 (37.5%)	0 (0%)	2.4
	UN	Female adult	5	2 (40%)	1 (20%)	2 (40%)	0 (0%)	2.4
	TA	Male adult	4	2 (50%)	2 (50%)	0 (0%)	0 (0%)	2
	CO	Male adult	2	0 (0%)	2 (100%)	0 (0%)	0 (0%)	2

Discussion

This study provides the first description of gait characteristics in mountain gorillas and chimpanzees during vertical climbing in a natural environment. These new data provide a broader comparative context to understand variation in primate locomotion in general, and particularly the biomechanical adaptations for vertical climbing in large-bodied primates. Our study found support for the hypothesis that, due to variation in body size, gait characteristics (i.e., temporal gait parameters, footfall sequences and limb support pattern) of vertical climbing differs between larger-bodied mountain gorillas and smaller-bodied chimpanzees. However, due to limited sample sizes in the analysis, we caution that any species differences (or intraspecific similarities) found in this study need to be validated on larger samples and perhaps on more varied natural substrates. Firstly, we found no intraspecific sex differences within the temporal gait parameters in mountain gorillas or chimpanzees (Table 6). Male and female gorillas showed similarly high duty factors when their hands contacted the substrate (female mean: 70 %; male mean: 69 %), suggesting that ascent and descent climbing as well as flexed-elbow and extended-elbow climbing are similarly mechanically challenging for both sexes, despite their large variation in body mass (e.g., Schultz, 1934; Sarmiento, 1994; Smith & Jungers, 1997). However, a larger sample size on male individuals is needed to clarify whether mountain gorillas differ more between the sexes during vertical climbing than is found in this study.

As predicted, mountain gorillas and chimpanzees showed striking differences in their temporal gait parameters, with gorillas having significantly longer cycle duration, higher duty factor and lower stride frequency than chimpanzees (Table 6). The higher duty factor of the mountain gorilla's forelimb indicates that the gorilla's hand is

held in contact with the substrate for longer compared to chimpanzees. Forceful hand grips and the use of variable thumb postures relative to substrate size are, therefore, likely particularly important in large-bodied mountain gorillas (Neufuss *et al.*, 2017), as they face greater biomechanical challenges during vertical climbing than smaller-bodied chimpanzees. The lower cycle duration and stride frequency combined with longer contact times, represents a more cautious and stable climbing strategy of mountain gorillas, which would increase safety when traversing unfamiliar or irregular substrates in an arboreal environment (Pontzer & Wrangham, 2004), as well as likely reduce energetic costs as found in other climbing primates (Isler, 2003; Isler & Thorpe, 2003; Hanna & Schmitt, 2011). Indeed, several studies have indicated that arboreal great apes display energy-saving adaptations in their locomotion to cope with their large body mass in an energetically challenging environment (Thorpe, Crompton & Alexander, 2007; Pontzer *et al.*, 2010). An interspecific comparison between western lowland gorillas and bonobos, as well as between juvenile and adult gorillas and orang-utans, supports both suggestions on climbing safety and energy-saving adaptations (Isler, 2002, 2005). Further investigation of the spatio-temporal gait parameters will show whether mountain gorillas also use relative long strides and climb at low speed similar to western lowland gorillas (Isler, 2005).

We also predicted that mountain gorillas would adapt their climbing strategy to accommodate their large body mass in a similar manner to that previously described in western lowland gorillas (Isler, 2002, 2003) and, likewise, that smaller-bodied chimpanzees would show a similar climbing style to that of bonobos (Isler 2002). This prediction was only partially supported.

Similar to mountain gorillas, Isler (2002) also found high mean duty factors in captive western lowland gorillas, although she attributed this in part to their difficulty in climbing up a vertical rope (Isler, 2002, 2003). Although all of the chimpanzees in our study exhibited comparatively lower mean duty factors like that of bonobos, Isler (2002) found that female and male bonobos differed considerably in their gait parameters during rope climbing, with females showing a shorter cycle duration and lower duty factor than males. Female bonobos could climb at a faster speed while the male bonobos would typically climb more slowly, combining long strides with a long cycle duration (Isler, 2002). However, a recent study of bonobos climbing a pole found similar gait parameters between the sexes (Schoonaert *et al.*, 2016), suggesting that a flexible rope may pose a greater locomotor challenge than climbing on a rigid support. This is in contrast to climbing in gorillas, where a compliant rope or liana might become stiffer in line of action while a gorilla climbs up, using their large body mass against the substrate. However, since the sample size for male chimpanzees in this study and for female bonobos in Schoonaert *et al.* (2016) were small, a more comprehensive analysis of the spatio-temporal gait parameters in a larger sample on different substrate types will clarify whether chimpanzees also show the same level of intraspecific variation reported in bonobos (Isler, 2002).

This study also found that the climbing performance of mountain gorillas is not less versatile than that of chimpanzees in a natural environment, but that gorilla individuals most often used a different footfall sequence pattern (diagonal sequence gaits) compared to chimpanzees (Fig. 2; Table 3). However, at a given duty factor, mountain gorillas used a significantly higher limb phase on vertical supports than that of chimpanzees, likely leading to an increase in the frequency of simultaneous footfalls of diagonally opposite limbs (diagonal sequence-diagonal couplet, DSDC) to improve the balancing abilities in the arboreal environment (e.g., Cartmill *et al.*, 2002,

2007; Stevens, 2006). By using DSDC gaits, the mountain gorilla hindlimb can touch down before the contralateral forelimb to secure a firm foothold on the substrate and both limbs can provide body support for part of the gait cycle. Changes in duty factor and gait sequence patterns result in a higher proportion of strides with support by more limbs at one time on substrates that challenge stability (Stevens, 2006). This holds true for our sample of mountain gorillas, which used mainly three limbs as body support during vertical ascent (Table 7). This is consistent with the climbing strategy documented in western lowland gorillas in captivity, in which they also used mainly three-limb support and mostly engaged in trot or DSDC gaits (Isler, 2002). However, our sample size on individuals descending substrates was comparatively small and thus, results on limb support pattern during vertical descent should be interpreted with caution (Table 7). More data on vertical descent climbing will clarify whether mountain gorillas also support their body mainly by three limbs as during vertical ascent.

In contrast, smaller-bodied chimpanzees used lateral sequence, and particularly diagonal couplet (LSDC) gaits most often, in which the hind foot touches down slightly later than the contralateral forelimb and the body is balanced on two diagonally opposite limbs. This is perhaps not surprising as it has long been acknowledged that either DS or LS gaits can be used in combination with diagonal limb couplet support patterns (e.g., Muybridge, 1887; Hildebrand, 1966, 1976). The diagonal couplet support enables primates to arrange the limbs as a widely splayed diagonal bipod and allows the center of mass to be contained within the base of support, reducing the risk of slipping and falling off the support during climbing (Cartmill *et al.*, 2002). Similar to our sample of chimpanzees, bonobos also used LS gaits more often than lowland gorillas during rope climbing but used two-limb, rather than three-limb, supports (Isler, 2002). Bonobo pole climbing showed a similar

pattern to our study with a more frequent use of LS gaits than DS gaits (Schoonaert *et al.*, 2016). However, the variation in footfall sequences between gorillas and chimpanzees suggests that hypotheses (i.e., limb interference avoiding: Hildebrand, 1980; stability: Cartmill *et al.*, 2002; energetic benefit: Griffin *et al.*, 2004) of primate-specific DS gait adaptations for arboreal locomotion require further investigation. A computer simulation of chimpanzee quadrupedal locomotion found a preference for LS/lateral couplet gaits (Sellers *et al.*, 2013). Furthermore, Stevens (2006) showed that primates could readily switch between DS and LS gaits on different arboreal supports likely because of differences in relative stance and swing phase durations. These results suggest that neither DS nor LS gaits offer a particular advantage for stability on horizontal (e.g., Shapiro & Raichlen, 2005; Stevens, 2006, 2008) and potentially vertical supports. This may reflect the need for most primates to use both arboreal and terrestrial substrates and thus, gait flexibility is the key feature of primate locomotion rather than the choice of a specific footfall sequence (Stevens, 2006; Higurashi, Hirasaki & Kumakura, 2009).

Conclusion

This quantitative analysis of temporal gait characteristics of mountain gorillas and chimpanzees in a natural environment showed that the climbing style can vary within each species. However, further field research and additional laboratory studies on more challenging (i.e., differently sized, less stable, irregular surface) supports are needed to clarify if captive studies are adequately representative for the patterns found in natural environments. Likewise, more work is needed to further characterize arboreal kinematic variability and gait choice among a wider range of primates (body proportions, body size) during vertical ascent and descent.

Acknowledgements

We thank the Uganda Wildlife Authority and the Ugandan National Council for Science and Technology for permission to conduct research in the Bwindi Impenetrable National Park, Uganda. We are grateful to the Institute for Tropical Forest Conservation (ITFC) in Bwindi for providing logistical support and to all the field assistants of ITFC who assisted in the project. We are also very thankful to the Chimfunshi Wildlife Orphanage Trust, Zambia, for giving us permission to study their sanctuary chimpanzees and particularly I. C. Mulenga for assistance with logistics. We thank D. Schmitt for his help with the gait analysis in this study. We also thank the Associate Editor and two anonymous reviewers for their helpful comments that greatly improved this manuscript. This research was supported by the European Research Council Starting Grant #336301 (to TLK and JN) and the Max Planck Society (to MR, JB, and TLK).

References

- Alexander, R.M., Jayes, A.S., Maloiy, G.M.O. & Wathuta, E.M. (1981). Allometry of the leg muscles of mammals. *J. Zool.* **194**, 539-552.
- Cartmill, M. (1974). Pads and claws in arboreal locomotion. In *Primate Locomotion*. 45-83. Jenkins, Jr F.A. (Ed.). New York: Academic Press.
- Cartmill, M. & Milton, K. (1977). The Lorisiform wrist joint and the evolution of "brachiating" adaptations in the Hominoidea. *Am. J. Phys. Anthropol.* **47**, 249-272.
- Cartmill, M. (1985). Climbing. In *Functional Vertebrate Morphology*. 73-88. Hildebrand, M., Bramble, D.M., Liem, K.F., Wake, D.B. (Eds). Cambridge, MA: Harvard University Press.
- Cartmill, M., Lemelin, P. & Schmitt, D. (2002). Support polygons and symmetrical gaits in mammals. *Zool. J. Linn. Soc.* **136**, 401-420.

Cartmill, M., Lemelin, P. & Schmitt, D. (2007). Understanding the adaptive value of diagonal-sequence gaits in primates: a comment on Shapiro and Raichlen, 2005. *Am. J. Phys. Anthropol.* **133**, 822-825.

Crompton, R.H., Sellers, W.I. & Thorpe, S.K.S. (2010). Arboreality, territoriality and bipedalism. *Phil. Trans. Soc. B.* **365**, 3301-3314.

Crompton, R.H. (2016). The hominins: a very conservative tribe? Last common ancestors, plasticity and ecomorphology in hominidae. *J. Anat.* **228**, 686-699.

Doran, D.M. (1996). Comparative positional behavior of the African apes. In *Great Ape Societies*. 213-224. McGrew, W.C., Marchant, L.F. & Nishida, T. (Eds.). Cambridge: Cambridge University Press.

Griffin, T.M., Main, R.P. & Farley, C.T. (2004). Biomechanics of quadrupedal walking: how do four-legged animals achieve inverted pendulum-like movements? *J. Exp. Biol.* **207**, 3545-3558.

Hanna, J.B. & Schmitt, D. (2011). Locomotor energetics in primates: gait mechanics and their relationship to the energetics of vertical and horizontal locomotion. *Am. J. Phys. Anthropol.* **145**, 43-54.

Hanna, J.B., Granatosky, M.C., Rana, P. & Schmitt, D. (2017). The evolution of vertical climbing in primates: Evidence from reaction forces. *J. Exp. Biol.* doi: 10.1242/jeb.157628

Higurashi, Y., Hirasaki, E. & Kumakura, H. (2009). Gaits of Japanese macaques (*Macaca fuscata*) on a horizontal ladder and arboreal stability. *Am. J. Phys. Anthropol.* **138**, 448-457.

- Hildebrand, M. (1966). Analysis of the symmetrical gaits of tetrapods. *Folio. Biotheor.* **6**, 9-22.
- Hildebrand, M. (1967). Symmetrical gaits of primates. *Am. J. Phys. Anthropol.* **26**, 119-130.
- Hildebrand, M. (1976). Analysis of tetrapod gaits: general consideration and symmetrical gaits. In *Neural control of locomotion*. 203-236. Herman, R.M., Grillner, S., Stein, P.S.G. & Stuart, D.C. (Eds.). New York: Plenum Press.
- Hildebrand, M. (1980). The adaptive significance of tetrapod gait selection. *Am. Zool.* **20**, 255-267.
- Hirasaki, E., Kumakura, H. & Matano, S. (1993). Kinesiological characteristics of vertical climbing in *Ateles geoffroyi* and *Macaca fuscata*. *Folia. Primatol.* **61**, 148-156.
- Hirasaki, E., Kumakura, H. & Matano, S. (2000). Biomechanical analysis of vertical climbing in the spider monkey and the Japanese Macaque. *Am. J. Phys. Anthropol.* **113**, 455-472.
- Hunt, K.D. (1991). Mechanical implications of chimpanzee positional behavior. *Am. J. Phys. Anthropol.* **86**, 521-536.
- Hunt, K.D. (1992). Positional behavior of *Pan troglodytes* in the Mahale Mountains and Gombe Stream National Parks, Tanzania. *Am. J. Phys. Anthropol.* **87**, 83-105.
- Hunt, K.D., Cant, J.G.H., Gebo, D.L., Rose, M.D. & Walker, S.E. (1996). Standardized descriptions of primate locomotor and postural modes. *Primates* **37**, 363-387.

Hunt, K.D. (2004). The special demands of great ape locomotion and posture. In *The evolution of thought: evolutionary origins of great ape intelligence*: 172-189. Russon, A.E., Begun, D.R. (Eds.). Cambridge: Cambridge University Press.

Hunt, K.D. (2016). Why are there apes? Evidence for the co-evolution of ape and monkey ecomorphology. *J. Anat.* **228**, 630-685.

Isler, K. (2002). Characteristics of vertical climbing in African apes. *Senckenbergiana Lethaea* **82**, 115-124.

Isler, K. (2003). Footfall patterns, stride length and speed of vertical climbing in spider monkeys (*Ateles fusciceps robustus*) and woolly monkeys (*Lagothrix lagotricha*). *Folia Primatol.* **75**, 133-149.

Isler, K. & Thorpe, S.K.S. (2003). Gait parameters in vertical climbing of captive, rehabilitant and wild Sumatran orang-utans (*Pongo pygmaeus abelii*). *J. Exp. Biol.* **206**, 4081-4096.

Isler, K. (2005). 3D-Kinematics of vertical climbing in hominoids. *Am. J. Phys. Anthropol.* **126**, 66-81.

Isler, K. & Grueter, C.C. (2006). Arboreal locomotion in wild black-and-white snub-nosed monkeys (*Rhinopithecus bieti*). *Folia Primatol* **77**, 195-211.

Muybridge, E. (1887). *Animals in motion*. Brown, L.S. (Ed.). (Reprinted in 1957). New York: Dover Publications.

Nakano, Y. (2002). The effects of substratum inclination on locomotor patterns in primates. *Z. Morphol. Anthropol.* **83**, 189-199.

Nakano, Y., Hirasaki, E. & Kumakura, H. (2006). Patterns of vertical climbing in primates. In *Human origins and environmental backgrounds*: 97-104. Ishida, H.,

Tuttle, R., Pickford, M., Ogihara, N. & Nakatsukasa, M. (Eds.). New York: Springer Publishing.

Neufuss, J., Robbins, M., Baeumer, J., Humle, T. & Kivell, T.K. (2017). Comparison of hand use and forelimb posture during vertical climbing in mountain gorillas and chimpanzees. *Am. J. Phys. Anthropol.* doi: 10.1002/ajpa.23303.

Pilbeam, D. (2002). Perspectives on the Miocene Hominoidea. In *The Primate Fossil Record*: 303-310. Hartwig, W.C. (Ed.). Cambridge: Cambridge University Press.

Pontzer, H. & Wrangham, R.W. (2004). Climbing and the daily energy cost of locomotion in wild chimpanzees: implications for hominoid locomotor evolution. *J. Hum. Evol.* **46**, 317-335.

Pontzer, H., Raichlen, D., Shumaker, R., Ocobock, C. & Wich, S. (2010). Metabolic adaptation for low energy throughput in orangutans. *Proc. Nat. Acad. Sci.* **107**, 14048-14052.

Preuschoft, H. & Witte, H. (1991). Biomechanical reasons for the evolution of hominid body shape In *Origine(s) de la bipédie chez les hominides*: 59–77. Coppens, Y. & Senut, B. (Eds). Paris: Editions du CNRS.

Preuschoft, H. (2002). What does “arboreal locomotion” mean exactly and what are the relationship between “climbing”, environment and morphology? *Z. Morphol. Anthropol.* **83**, 171-188.

Preuschoft, H. (2004). Mechanisms for the acquisition of habitual bipedality: are there biomechanical reasons for the acquisition of upright bipedal posture? *J. Anat.* **204**, 363-384.

Rawlings, B., Davila-Ross, M. & Boyson, S.T. (2014). Semi-wild chimpanzees open hard-shelled fruits differently across communities. *Anim Cogn* **17**, 891-899.

Remis, M.J. (1995). Effect of body size and social context on the arboreal activities of lowland gorillas in the Central African Republic. *Am. J. Phys. Anthropol.* **97**, 413-433.

Remis, M.J. (1998). The gorilla paradox. In *Primate locomotion*: 95-106. Strasser, E., Fleagle, J.G., Rosenberger, A.L. & McHenry, H.M. (Eds). New York: Springer Publishing.

Remis, M.J. (1999). Tree structure and sex differences in arboreality among western lowland gorillas (*Gorilla gorilla gorilla*) at Bai Hokou, Central African Republic. *Primates* **40**, 383-396.

Robbins, M.M. (2001). Variation in the social system of mountain gorillas: The male perspective. In *Mountain Gorillas: Three Decades of Research at Karisoke*: 29-58. Robbins, M.M., Sicotte, P. & Stewart, K.J. (Eds.). Cambridge: Cambridge University Press.

Robbins, M.M. & McNeilage, A. (2003). Home range and frugivory patterns of mountain gorillas in Bwindi impenetrable national park, Uganda. *Int. J. Primatol.* **24**, 467-490.

Sarmiento, E.E. (1994). Terrestrial traits in the hands and feet of gorillas. *American Museum Novitates* **3091**, 56.

Schoonaert, K., D'Août, K., Samuel, D., Talloen, W., Nauwelaerts, S., Kivell, T.L. & Aerts, P. (2016). Characteristics and spatio-temporal variables of climbing in bonobos (*Pan paniscus*). *Am. J. Primatol.* **78**, 1165-1177.

- Schultz, A.H. (1934). Some distinguishing characters of the mountain gorilla. *J. Mammal.* **15**, 51-61.
- Shapiro, L.J. & Raichlen, D.A. (2005). Lateral sequence walking in infant *Papio cynocephalus*: implications for the evolution of diagonal sequence walking in primates. *Am. J. Phys. Anthropol.* **126**, 205-213.
- Smith, R.J. & Jungers, W.L. (1997). Body mass in comparative primatology. *J. Hum. Evol.* **32**, 523-559.
- Stevens, N.J. (2006). Stability, limb coordination and substrate type: The ecorelevance of gait sequence pattern in primates. *J. Exp. Zool.* **305**, 953-963.
- Stevens, N.J. (2008). The effect of branch diameter on primate gait sequence pattern. *Am. J. Primatol.* **70**, 356-362.
- Thorpe, S.K.S., Crompton, R.H. & Alexander, R.McN. (2007). Orangutans utilise compliant branches to lower the energetic cost of locomotion. *Biol. Lett.* **3**, 253-256.
- Tuttle, R.H. & Watts, D.P. (1985). The positional behaviour and adaptive complexes of Pan gorilla. In *Primate Morphophysiology, Locomotor Analyzes, and Human Bipedalism*: 261-288. Kondo, S. (Ed.). Tokyo: University of Tokyo Press.
- Vielnsky, J.A. & Gankiewicz, E. (1989). Early development of locomotor behavior in vervet monkeys. *Am. J. Primatol.* **17**, 11-25.