



Kent Academic Repository

Taylor, Samantha L., Crawley-Snowdon, Harriet, Wagstaff, Jane L., Rowe, Michelle L., Shepherd, Mark, Williamson, Richard A. and Howard, Mark J. (2013) *Measuring protein reduction potentials using 15N HSQC NMR spectroscopy*. *Chemical Communications*, 49 (18). pp. 1847-1849. ISSN 1359-7345.

Downloaded from

<https://kar.kent.ac.uk/33499/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1039/c3cc38952a>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Measuring protein reduction potentials using ^{15}N HSQC NMR spectroscopy

Samantha L. Taylor^a, Harriet Crawley-Snowdon^a, Jane L. Wagstaff^a, Michelle L. Rowe^a, Mark Shepherd^a, Richard A. Williamson^{a,*} and Mark J. Howard^{a,*}

⁵ Received (in XXX, XXX) Xth XXXXXXXXX 20XX, Accepted Xth XXXXXXXXX 20XX

DOI: 10.1039/b000000x

NMR spectroscopy was used to measure reduction potentials of four redox proteins by following multiple ^{15}N HSQC protein resonances across a titration series using mixtures of oxidised and reduced glutathione. Results for PDI **a**, PDI **ab** and DsbA agree with the literature and our result for ERp18 confirms this protein as an oxidoreductase of comparable or greater reducing strength than PDI **a**.

In order for proteins to be stable and to conduct their catalytic functions efficiently, proteins must be correctly folded and the presence of disulphide bonds can ensure a protein folds correctly. A family of thiol-disulphide oxidoreductase proteins located within the cells catalyse the formation and rearrangement of disulphide bonds. Typically these enzymes contain an active site motif of CXXC and exist in either the oxidised or reduced state.¹

PDI (protein disulphide isomerase) is located within the endoplasmic reticulum and aids in the folding of disulphide bond-containing proteins via the formation (oxidation), rearrangement (isomerisation) and removal (reduction) of non-native disulphide bonds. Human PDI is a 57 kDa protein that contains four thioredoxin-like domains in the order **a**, **b**, **b'**, **a'** with an extended linker (**x**) between the **b'** and **a'** domains and an acidic 19-residue C-terminal tail (**c**). The two catalytic **a** and **a'** domains contain the conserved CGHC active site motif. The **b** and **b'** domains, although non-catalytic, are considered essential for full PDI function, with the **b** domain providing structural stability and the **b'** domain containing the primary ligand binding site.²⁻⁴ This study uses the **a** and **ab** domain fragments of human PDI (*ca.* 14 and 27 kDa in size, respectively). To date, the PDI family contains up to 21 members with differing domain architectures.¹ ERp18, at 18 kDa, is one of the smallest known members of the PDI family with the thioredoxin fold and a CGAC active site.^{5, 6} The fourth protein used in this study is *E. coli* DsbA, the archetypal disulphide oxidoreductase in bacteria.⁷

Determining redox potentials allows the identification of a protein's role *in vivo*; proteins with low redox potentials in the region of -270 mV, such as thioredoxin,⁸ will have reducing properties, while proteins with higher reduction potentials around -122 mV, such as DsbA, will have oxidising properties.⁹ Reduction potentials have typically been determined by monitoring reactions using tryptophan fluorescence¹⁰, or alternatively, using HPLC or radiolabelling. These methods operate by separating and quantifying the relative amounts of

reduced and oxidised forms of the protein. However this method could only be applied to proteins with one active site. Recently, the reduction potentials of human PDI were elegantly determined using chemical modification and mass spectrometry to show the influence of adjacent active sites on re-oxidation of PDI by the protein Ero1 α .¹¹ This approach allows the measurement of two active sites simultaneously using a differential alkylation approach followed by mass spectrometry.

Measurement of the reduction potential using ^{15}N HSQC has been achieved previously, which allowed monitoring of one active site cysteine in cytoplasmic desulfothioredoxin from *Desulfovibrio vulgaris*¹² and 1D ^1H NMR has been used to study reduction potentials in rubredoxin variants.¹³ Here we demonstrate a novel and much improved method with respect to using ^{15}N HSQC NMR data. The cooperative mechanism whereby the protein switches between oxidised and reduced states enables a choice of two resonances to be monitored that mirror the conversion from one oxidation state to the other. In addition, all resonances that significantly shift between oxidised and reduced states are measured and each resonance provides a single Hill equation fit and reduction potential. Fitting each resonance to an individual curve fit provides the opportunity to identify and remove systematic and data specific errors to increase the precision and accuracy of the overall average reduction potential determined. Errors can be overlooked if all data is treated as one within an individual Hill plot.

Proof of concept is demonstrated for proteins containing one active site, however, it could easily be expanded to obtain redox potentials of proteins with two or more active sites. This would require NMR assignments to identify resonances from amino acids that report on each active site. Assignment of resonances to individual amino acids is summarised in the ESI†. Conversely, for proteins with a single active site, no residue-specific assignments are necessary, as only changes in each resonance are required to monitor the oxidative status.

^{15}N enriched recombinant human PDI **ab** and ERp18 were expressed in *E. coli* and purified as previously described.^{2, 6} PDI **a** and DsbA were produced using protocols outlined in the ESI†.

To determine the reduction potential of PDI **ab**, PDI **a**, ERp18 and DsbA different ratios of [GSH]/[GSSG] were used where the total concentration ([GSH]+[GSSG]) equalled 5 mM. All NMR data was obtained at 298 K using a 14.1 T (600 MHz ^1H) Bruker Avance III NMR spectrometer equipped with a QCI-F cryoprobe.

All NMR samples were 330 μ L within a Shigemi NMR tube and contained either 0.065mM (PDI **a**) or 0.3 mM (PDI **ab**, ERp18, DsbA) protein in 20 mM sodium phosphate buffer (pH 7.3) containing 50 mM sodium chloride and 5% (v/v) deuterium oxide. All buffers had nitrogen gas bubbled through them for 30-60 min to ensure the removal of any dissolved oxygen and all proteins were found to reach redox equilibrium within 5 minutes of setting up each condition. All 15 N HSQC spectra were acquired with 2048 points in the direct F2 dimension (1 H) and 256 points in the F1 dimension (15 N). NMR data processing was completed using NMRpipe and example spectra are shown in Figure 1. NMR backbone assignments have previously been completed for all the proteins^{2, 6, 14} and were used to identify the peaks that shifted with oxidation status.




Fig. 1 15 N HSQC spectra of (a) PDI **a**, (b) PDI **ab**, (c) ERp18 and (d) DsbA. Fully oxidised spectra are shown in red, and fully reduced spectra are shown in black.

Peak heights and volumes were determined using Analysis v2.2¹⁵ for each oxidised and reduced peak in the spectrum. Peak volumes provided better curve fits for both the oxidised and reduced peaks. The fraction reduced was determined from this data and fitted in Kaleidagraph version 4.1 (Synergy Software) to the Hill equation to determine the equilibrium constant (K_{eq}) using Equation 1, where F_{red} is the fraction of reduced protein. Example data fits to the Hill equation are shown in Figure 2, where millimolar concentrations were used to generate this plot. Therefore, K_{eq} is solved from Equation 1 in millimolar units.

$$F_{red} = ([GSH]^2/[GSSG]) / (K_{eq} + [GSH]^2/[GSSG]) \quad (\text{Eq.1})$$

The reduction potential of each protein was then determined by the Nernst equation (Equation 2) where $T = 298$ K, $n = 2$, and the standard reduction potential of GSH ($E'_{0(GSH)} = -240$ mV). If R is defined as the molar gas constant, K_{eq} must be used with molar concentration units in Equation 2.

$$E'_0 = E'_{0(GSH)} - (RT/nF) \ln(K_{eq}) \quad (\text{Eq.2})$$

Both equations have been described previously with respect to mass spectrometry reduction potential determination¹¹ and are equally valid with our approach.




Fig. 2 Two fits shown using the Hill equation, PDI **a**, lysine 14 (green) with a redox potential of -144.43 ± 0.34 mV and DsbA serine 43 (red) with a redox potential of -124.63 ± 0.98 mV.

Errors were determined using the Levenberg-Marquardt R-value for the fit of the Hill equation and two controls were run; fully oxidised protein (in 5 mM GSSG) and fully reduced protein (in 5 mM dithiothreitol (DTT)). DTT was used as it is capable of fully reducing catalytic PDI domains. Individual plots are shown in the ESI†. Average reduction potentials and curve fit error values obtained by this method are summarised in Table 1.

Table 1. K_{eq} and mean reduction potential values (E'_0) obtained including number of resonances used in final K_{eq} and E'_0 analysis.

	PDI a	PDI ab	ERp18	DsbA
Number of resonances	18	12	6	6
K_{eq} (mM)	0.59	1.59	2.61	0.16
E'_0 (mV)	-144.4 ± 0.7	-156.8 ± 1.2	-161.5 ± 3.3	-125.6 ± 0.6

This study has measured the reduction potential at one active site of the protein, however, the reduction potentials found in this study are similar to those previously determined.^{1, 8, 11}

The use of multiple resonances to determine the reduction potential is extremely useful. For example, PDI **a**, which contains 120 amino acids and one CGHC active site, provided 15 N HSQC spectra with over 18 backbone amide cross peaks with significant chemical shift changes between the reduced and oxidised states. This provides a degree of data redundancy allowing weak or overlapped peaks to be removed from the analysis and a higher precision in the final value of the reduction potential determined. Across all moving resonances in PDI **a**, individual reduction potentials were calculated and ranged from -148.73 mV to -140.39 mV. Using only optimal data from resonances that were not overlapped or of low signal to noise enables a value of -144.4 ± 0.7 mV to be quoted. We also measured the reduction potential of PDI **a** and PDI **ab** as -144.6 ± 1.1 mV and -157.0 ± 2.1 mV respectively via single Hill plots that used all resonance data simultaneously; these plots are available in the ESI†. However, our approach provides smaller errors and hence greater precision because handling resonance data separately allows for the identification and exclusion of poor data.

The values for PDI **a** are consistent with values previously

published, where the reduction potential varies from -110 mV¹⁶ to -190 mV.¹⁷ PDI **ab** data is also consistent and confirms that the addition of the structural **b** domain does not significantly alter the reduction potential of the catalytic **a** domain. This data confirms our method is equally applicable for larger proteins as PDI **ab** is *ca.* 27 kDa. The molecular weight limit for this method will be in excess of 60 kDa when acquiring TROSY optimised ¹⁵N-¹H HSQC data. ERp18 provided a reduction potential of -161.5 ± 3.3 mV that suggests this protein is slightly more reducing than PDI **a**. ERp18 was previously reported to have a similar reduction potential to PDI **a**, but the value reported was considered of low precision.⁵ The value obtained for DsbA is also within the range of that determined by other approaches.^{18, 19}

When monitoring the redox changes in a protein, either peak height or volume could be used although peak volumes tended to provide better data fitting to the Hill equation and so was preferred. Our data reveals that reduction potentials for each protein could be determined by measuring peak volumes from only the reduced peak in the spectrum. Equally, the oxidised peak could be used to determine the fraction reduced, but care is required because NMR line broadening of the oxidised state resonance may complicate peak height or volume measurements. The effect of peak line broadening was demonstrated with Hill plots and associated reduction potentials obtained using only oxidised resonances for ERp18 and PDI **a** (data in ESI†). ERp18 is known to exhibit relaxation line broadening in the oxidised state⁶ and the ESI† data demonstrates these data providing poor Hill plot fits and values even when spectrum sensitivity is not an issue as shown in Figure 1c and from oxidised and reduced ERp18 spectra displayed in the assignment section of the ESI†. We suggest that to prevent inaccuracy, all peak volumes or heights should be measured from either the oxidised or reduced state and not a mixture of the two forms. However, line broadening effects that alter quantitation could be managed by adjusting peak volumes or heights for relaxation using methods similar to those recently described in the study of metabolites.²⁰

Overall, this method of measuring protein reduction potentials using ¹⁵N HSQC NMR is quick and convenient and has great potential to be utilised with proteins containing multiple active sites. This method can also probe the effect of physical and physiological conditions that influence the mechanistic process of disulphide bond biochemistry through different buffer systems, crowding agents and substrates. Whilst in this study the oxidation and reduction of the proteins were conducted using a [GSH]/[GSSG] mixture, this method could also be applied to alternative systems, for example [NADP]/[NADPH].

Notes and references

^a School of Bioscience, University of Kent, Canterbury, Kent, CT2 7NJ, UK. Fax: 44 (0) 1227763912; Tel: 44 (0) 1227 764000; E-mail: r.a.williamson@kent.ac.uk or m.j.howard@kent.ac.uk. We would like to thank The Wellcome Trust for supporting this project through Equipment Grant 091163/Z/10/Z (MJH and RAW) and Project Grant 093125/B/10/Z (MJH and RAW). Additional thanks to Dr. Begoña Heras (La Trobe University) and Prof. Mark Schembri (University of Queensland) for the production and supply of the *E. coli* DsbA clone. SLT and MJH would also like to thank Prof. John Baldock for funding SLT through the University of Kent PVC Research Fund.

† Electronic Supplementary Information (ESI) available: [Isotopic enrichment notes, Protein expression and purification, NMR assignment notes, reduction potential Hill Plots for PDI **a**, **ab**, ERp18 and DsbA, combined Hill plots of PDI **a** and **ab** and oxidised data Hill plots of PDI **a** and ERp18]. See DOI: 10.1039/b000000x/

1. F. Hatahet and L. W. Ruddock, *Antioxid Redox Signal*, 2009, **11**, 2807-2850.
2. L. J. Byrne, A. Sidhu, A. K. Wallis, L. W. Ruddock, R. B. Freedman, M. J. Howard and R. A. Williamson, *Biochem J*, 2009, **432**, 209-217.
3. V. D. Nguyen, K. Wallis, M. J. Howard, A. M. Haapalainen, K. E. Salo, M. J. Saaranen, A. Sidhu, R. K. Wierenga, R. B. Freedman, L. W. Ruddock and R. A. Williamson, *J Mol Biol*, 2008, **383**, 1144-1155.
4. A. Y. Denisov, P. Maattanen, C. Dabrowski, G. Kozlov, D. Y. Thomas and K. Gehring, *Febs J*, 2009, **276**, 1440-1449.
5. H. I. Alanen, R. A. Williamson, M. J. Howard, A.-K. Lappi, H. P. Jantti, S. M. Rautio, S. Kellokumpu and L. W. Ruddock, *J Biol Chem*, 2003, **278**, 28912-28920.
6. M. L. Rowe, L. W. Ruddock, G. Kelly, J. M. Schmidt, R. A. Williamson and M. J. Howard, *Biochemistry*, 2009, **48**, 4596-4606.
7. J. Couprie, M. L. Remerowski, A. Bailleul, M. Courcon, N. Gilles, E. Quemeneur and N. Jamin, *Protein Sci*, 1998, **7**, 2065-2080.
8. F. Aslund, K. D. Berndt and A. Holmgren, *J Biol Chem*, 1997, **272**, 30780-30786.
9. E. Mossner, M. Huber-Wunderlich, A. Rietsch, J. Beckwith, R. Glockshuber and F. Aslund, *J Biol Chem*, 1999, **274**, 25254-25259.
10. A. Holmgren, *Journal of Biological Chemistry*, 1979, **254**, 9627-9632.
11. J. E. Chambers, T. J. Tavender, O. B. Oka, S. Warwood, D. Knight and N. J. Bulleid, *J Biol Chem*, 2010, **285**, 29200-29207.
12. E. B. Garcin, O. Bornet, L. Elantak, N. Vita, L. Pieulle, F. Guerlesquin and C. Sebban-Kreuzer, *J Biol Chem*, 2012, **287**, 1688-1697.
13. I. J. Lin, E. B. Gebel, T. E. Machonkin, W. M. Westler and J. L. Markley, *Proc Natl Acad Sci U S A*, 2005, **102**, 14581-14586.
14. J. Kemmink, N. J. Darby, K. Dijkstra, R. M. Scheek and T. E. Creighton, *Protein Science*, 1995, **4**, 2587-2593.
15. W. F. Vranken, W. Boucher, T. J. Stevens, R. H. Fogh, A. Pajon, M. Llinas, E. L. Ulrich, J. L. Markley, J. Ionides and E. D. Laue, *Proteins*, 2005, **59**, 687-696.
16. H. C. Hawkins, M. de Nardi and R. B. Freedman, *Biochem J*, 1991, **275 (Pt 2)**, 341-348.
17. J. Lundstrom and A. Holmgren, *Biochemistry*, 1993, **32**, 6649-6655.
18. U. Grauschopf, J. R. Winther, P. Korber, T. Zander, P. Dallinger and J. C. Bardwell, *Cell*, 1995, **83**, 947-955.
19. R. P. Szajewski and G. M. Whitesides, *Journal of the American Chemical Society*, 1980, **102**, 2011-2026.
20. K. F. Hu, W. M. Westler and J. L. Markley, *Journal of the American Chemical Society*, 2011, **133**, 1662-1665.