



Kent Academic Repository

Hopker, James G., Passfield, Louis, Coleman, Damian A, Jobson, Simon A., Edwards, L. and Carter, Helen (2009) *The effects of training on gross efficiency in cycling: a review*. International journal of sports medicine, 30 (12). pp. 845-850. ISSN 1439-3964.

Downloaded from

<https://kar.kent.ac.uk/36044/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1055/s-0029-1237712>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

The Effects of Training on Gross Efficiency in Cycling: A Review

Authors

J. Hopker¹, L. Passfield¹, D. Coleman², S. Jobson¹, L. Edwards³, H. Carter⁴

Affiliations

¹ Centre for Sport Studies, University of Kent, England

² Department of Sports Science, Leisure and Tourism, Canterbury Christ Church University, Kent, England

³ Department of Physiology, Anatomy and Genetics, University of Oxford, England

⁴ Chelsea School, University of Brighton, England

Key words

- endurance training
- economy
- cycling
- delta efficiency

Abstract

There has been much debate in the recent scientific literature regarding the possible ability to increase gross efficiency in cycling via training. Using cross-sectional study designs, researchers have demonstrated no significant differences in gross efficiency between trained and untrained cyclists. Reviewing this literature provides evidence to suggest that methodological inadequacies may have played a crucial role in the conclusions drawn from the majority of these

studies. We present an overview of these studies and their relative shortcomings and conclude that in well-controlled and rigorously designed studies, training has a positive influence upon gross efficiency. Putative mechanisms for the increase in gross efficiency as a result of training include, muscle fibre type transformation, changes to muscle fibre shortening velocities and changes within the mitochondria. However, the specific mechanisms by which training improves gross efficiency and their impact on cycling performance remain to be determined.

Introduction

Gross efficiency is defined as the ratio of power output to energy expenditure and is a key determinant of endurance cycling performance [14, 17, 18, 36, 49, 53]. Despite this, previous studies investigating gross efficiency during cycling have found no differences between trained and untrained cyclists [6, 45, 50, 52]. The results of these studies supported the recently challenged assumption that training has no effect on a cyclist's gross efficiency [34]. These observations of non-significant differences between trained and untrained individuals using cross-sectional research designs have limited the extent of investigation into the mechanisms that could be responsible for changes in efficiency with training. Recently, using more rigorous experimental designs researchers have provided evidence that gross efficiency can increase with training and is higher in a trained population [33, 34, 60]. Nonetheless, one recent study documenting improvements in cycling efficiency in a Grand Tour Champion [19] has provoked much debate within the scientific literature [25, 47, 61]. Currently, researchers are beginning to identify the specific circumstances under which training can lead to an increase in gross efficiency during cycling.

Considered in full, the extant literature also provides some clues regarding the possible mechanisms and adaptations that are responsible. The purpose of this review is to provide: a summary of the different methods used to calculate efficiency, a discussion of the requirements for reliable measurement of efficiency during cycling, an overview of the published literature relating to acute and training-related effects on gross efficiency, and suggestions for future research.

Terminology

During steady-state cycle ergometry, efficiency has been extensively used to provide a convenient index of the effectiveness with which an individual can convert chemical energy into mechanical power [4, 23, 28, 42, 52, 66]. The most commonly used measure of efficiency is gross efficiency ($[\text{work accomplished}/\text{energy expended}] \times 100$). In calculating gross efficiency, the caloric equivalent of steady state $\dot{V}O_2$ and the respiratory exchange ratio (RER) are used to calculate energy expenditure. The term 'gross efficiency' is normally reported as a percentage of total energy expenditure [15].

accepted after revision
July 20, 2009

Bibliography

DOI 10.1055/s-0029-1237712
Published online:
November 25, 2009
Int J Sports Med 2009; 30:
845–850 © Georg Thieme
Verlag KG Stuttgart · New York
ISSN 0172-4622

Correspondence

Dr. James Hopker
Centre for Sport Studies
University of Kent
Chatham Maritime
Kent
ME4 4AG
Tel.: +44/1634/88 88 14
Fax: +44/1634/88 88 09
j.g.hopker@kent.ac.uk

Other indices of efficiency used in the literature include work, net and delta efficiency. The calculation of net efficiency requires the energy expenditure at rest to be measured (and assigned to the performance of internal work), which is then subtracted from the total amount of energy expended for the external work accomplished. Thus, net efficiency does not take into account the cost of moving the involved limbs. Gaesser and Brooks [23] suggested that for a valid measurement this cost should be taken into account. For example, the energetic cost of cycling on an ergometer, but with a work rate of 0W (i.e. unloaded pedalling) would need to be subtracted from the total energy expenditure for the measured work rate.

The calculation of work efficiency accounts for the additional cost of moving the legs, but this measurement of zero load in cycling can be problematic. Kautz and Neptune [40] have argued that zero load cycling does not provide a valid baseline for reference to a range of work intensities. Gaesser and Brooks [23] proposed a floating baseline measure for the physiological and external energy cost of exercise, i.e. delta efficiency (DE). Here energy expenditure at a lower work rate is subtracted from the energy expenditure at a higher work rate ($[\text{delta work accomplished}/\text{delta energy expended}] \times 100$). Coyle et al. [16] suggested that this is the most valid calculation of whole-body efficiency as it attempts to partial out the influence of unmeasured work. However, the linear relationship that is produced between energy cost and work rate does not necessarily mean that it is a valid measure (i.e. it is independent of the work rate used). For efficiency to increase with power output [12,23] energy expenditure must increase non-linearly due to the decreasing relative contribution of non-propulsive factors (e.g. basal metabolism and moving the legs). The use of DE may also be limited as both Moseley and Jeukendrup [49] and Hopker et al. [33] have shown it to have greater day-to-day variability than gross efficiency.

The issues involved in these measures have been extensively discussed by Gasser and Brooks [23], Stainsby et al. [64], Cavanagh and Kram [11] and Kautz and Neptune [40]. Due to the various criticisms of base-line subtraction methods outlined above, it is unrealistic to attempt to attribute a portion of the total body energy cost to muscle work during a whole body exercise such as cycling. Therefore we will focus on gross efficiency for the rest of this review.

The measurement of efficiency and other methodological issues



Steady state testing

In order to measure efficiency accurately all gas collection must take place under steady state exercise conditions, otherwise measured pulmonary $\dot{V}O_2$ may not adequately reflect muscle O_2 consumption [55]. At the onset of exercise or in the transition from one work rate to another, during light to moderate intensities, pulmonary $\dot{V}O_2$ increases in a mono-exponential manner to reach a steady state within 2–3 min [75]. Heavy exercise intensities, characterised by a sustained metabolic acidosis, have been found to result in a delayed steady state [74]. Consequently, the relationship between $\dot{V}O_2$ and power output presents a marked deviation from linearity at higher intensities when examined carefully [26, 78]. The exercise intensities eliciting this additional O_2 consumption, or $\dot{V}O_2$ slow component, should not be used to determine efficiency values. The calculated energy equivalent

for a given $\dot{V}O_2$ depends upon the equivalence of RER and muscle RQ. A decrease of 0.05 in RER reduces calculated energy expenditure by 1.3% typically increasing GE by 0.4%. Therefore as $\dot{V}CO_2$ affects RER it is important to ensure its stability prior to taking efficiency measurements. During steady state exercise resulting in minimal metabolic acidosis, $\dot{V}CO_2$ may take considerably longer (at least 4 min) to reach steady state [13,72]. For these reasons, long work stages (≥ 5 min) should be used when collecting gas data for the calculation of efficiency. Boning et al. [6] and Moseley et al. [50] used 3 min stage durations for their studies of cycling efficiency. Therefore they may have failed to allow sufficient time to achieve a steady state when investigating differences in efficiency between trained and untrained cyclists.

Exercise intensity

A particular criticism of past research is the low exercise intensities used for assessing efficiency during cycling. Researchers have previously sought to investigate optimal levels of energy expenditure during cycling using untrained participants [23,62,66,77]. As a result, it has only been possible to study responses to low work rates. These values are typically very low if inferences from such findings are made to trained cyclists who commonly race at much higher work rates.

Efficiency has been shown to increase with work rate [12,23]. It is thought that this is largely due to the unmeasured work (i.e. that required to sustain basal metabolism and body position on the bike) forming a smaller percentage of total energy expenditure at higher work rates [64]. However, more recently Moseley and Jeukendrup [49] have demonstrated that a plateau in gross efficiency occurs at the higher work rates ($>240W$) used by trained cyclists. Ideally, the work rate used for the determination of gross efficiency should encompass the functional range of the population of interest. However, when trained cyclists are considered this is potentially difficult due to the $\dot{V}O_2$ slow component associated with higher racing intensities.

Standardisation of other factors

Most authors agree that changes in muscle shortening velocity (i.e. pedal cadence) markedly affect efficiency. For a given power output, increasing cadence has been shown to decrease gross efficiency [12,28,45,63]. Therefore, cadence must be standardised when conducting repeated measurements of efficiency in the same individual. Changes in riding position may also influence the efficiency values obtained. Alterations in seat tube angle and saddle height have been shown to change gross efficiency [30,56]. Price and Donne [56] found an energetic optimum for combinations of seat tube angle and seat height (70° seat tube angle; 100% trochanteric height). Alterations in the muscle length-tension relationships (quadriceps versus hamstrings) and ankle patterns could account for differences in efficiency found with increased seat tube angles and heights. Thus, the dimensions of a cyclist's bicycle should be replicated when using a cycle ergometer and maintained during any subsequent tests.

High ambient temperatures (35.5 °C) have been shown to cause decreases in gross efficiency [31]. With hyperthermia the energy cost of the exercise may increase due to greater circulation, sweating and ventilation. This in turn may reduce efficiency as the work accomplished would remain unchanged. Therefore ambient temperature should be tightly controlled, especially

when repeating tests during different phases of a season under varied climatic conditions.

Finally, the control of cyclists' pre-testing regimen presents a further methodological issue. Racing cyclists often train and compete for several hours at intensities of 60% $\dot{V}O_{2\text{ peak}}$ and above [44]. A gradual $\dot{V}O_2$ drift is commonly observed during such prolonged exercise. This $\dot{V}O_2$ drift manifests as a reduction in gross efficiency and is apparent even in trained cyclists when exercising for 75 min at 60% $\dot{V}O_{2\text{ peak}}$ [54]. Passfield and Doust [54] further demonstrated that this acute reduction in gross efficiency was significantly correlated with a lower 5 min time-trial performance. Similarly, there is also some evidence to suggest that muscle damage from high intensity training might decrease efficiency during subsequent exercise performance [43,71].

Therefore, it is important to monitor training in the days prior to testing athletes for efficiency. The time course for restoration of gross efficiency after exercise has not been established. The ability to detect meaningful changes in cycling performance over time, and to study the effects of selected interventions, requires careful consideration of the validity and reliability of testing methods and equipment [3,35]. If reliability of repeated measurements is not ensured then there will be increased chances of falsely rejecting the null hypothesis in any research study. Therefore, it is critical that appropriate methods are employed which minimise the variability of within steady state $\dot{V}O_2$ and $\dot{V}CO_2$ data, exercise intensities utilised, ergometer set-up and participant pre-test preparation.

Chronic changes in gross efficiency with training

A question which has been addressed by several researchers is whether cycling efficiency can be increased by training. Previous cross-sectional studies have failed to find any differences in efficiency between trained and untrained cyclists [6,45,46,50,52]. However a number of these investigations can be criticized on the basis of their methods and their failure to address the risk of committing a type 2 statistical error. Specifically, a lack of statistical power in past research studies has meant an inability to detect significant differences between study populations. Other

confounding factors are the usage of short stage protocols and artificially imposed cadences. **Table 1** shows the main findings of research studies that have sought to investigate differences in efficiency between participants of varied cycling ability. To illustrate the likely chance of a type 2 statistical error we have calculated the effect size and statistical power using post hoc methods [5,69]. Effect size was calculated by the division of the mean difference between the two groups (μ_1 and μ_2) by the pooled standard deviation (SD) [69];

$$ES = (\mu_1 - \mu_2) / SD$$

δ (the value for non centrality) and achieved statistical power was then calculated using GPower software [22].

Gross efficiency during cycling has been reported to be in the range of 18–23% [16]. An improvement in efficiency implies an increase in mechanical power output for a given metabolic cost [53]. Using the extremes of the 18–23% range suggests that for the same rate of metabolic energy expenditure, an efficient cyclist would produce 28% more power than a less efficient cyclist ((23–18)/18=28%). Evidence of longitudinal changes in gross efficiency was provided by Coyle [19], who reported an 8.8% increase in the gross efficiency of a Grand Tour Champion over a 7-year-period. Unfortunately, it was not possible to tightly control all aspects of data collection in this case study. Consequently, this study has been the subject of repeated criticism for its design, method and analysis [25,47,61]. Sassi et al. [60] have reported seasonal changes (albeit not significant) in gross efficiency in a small group of competitive cyclists. Seasonal variation was not controlled in Coyle's [19] study and therefore could account for the majority of the improvement he reports. Using a longitudinal study design Hopker et al. [34] have demonstrated that competitive cyclists can increase their gross efficiency by as much as 5%. These increases in gross efficiency were significantly correlated with the volume and intensity of training completed. More specifically, high intensity training was most strongly related to increases in gross efficiency.

Even though the studies outlined above have demonstrated that gross efficiency can increase as a result of training, it is still unclear whether such chronic changes will actually impact on performance. Currently, the effect of training-induced increases in efficiency on cycling performance has not been assessed.

Table 1 Methods and findings of cross sectional research studies that have investigated differences in efficiency between participants of varied cycling ability. The table is restricted to studies which present enough data to estimate effect size and statistical power.

Author(s)	Sample Size	Stage Duration	Power Outputs used	Efficiency measure	Cadence imposed	Effect Size/ Statistical Power	Efficiency finding
Boning et al., (1984)	15 (9 trained cyclists; 6 untrained)	3 mins	50, 100, 200W	GE & NE	yes (40, 60, 70, 80, 100 rev.min ⁻¹)	0.87/0.33 at 200W	GE ($p < 0.05$) & NE (both 1% mean difference)
Nickleberry & Brooks (1996)	12 (6 competitive; 6 recreational)	4 mins	50–200W	GE & DE	yes (50 and 80 rev.min ⁻¹)	0.26/0.07 at 200W	no significant difference (>1% difference between groups)
Marsh et al., (2000)	31 (11 competitive cyclists; 10 trained runner; 10 non-cyclists)	5 mins	trained cyclists; 100, 150, 200W Untrained 75, 100, 150W	DE	yes (50, 65, 80, 95, 110 rev.min ⁻¹)	1.25/0.77 at 80 rev.min ⁻¹	no significant difference (~1% difference between groups at 80 & 95 rev.min ⁻¹)
Moseley et al., (2004)	69 trained cyclists (divided on $\dot{V}O_{2\text{ peak}}$, low, medium and high)	3 mins	95W increasing by 35W	GE & DE	yes (80–90 rev.min ⁻¹)	2.33/1.00 between low and high $\dot{V}O_{2\text{ peak}}$ groups	GE & DE no significant difference (GE 0.9% difference Med-High groups; DE 1.2% difference Low to High groups)
Hopker et al., (2007)	30 (14 trained cyclists; 16 recreational)	10 mins	150W, 50 & 60% W_{max}	GE	preferred cadence used	1.51–1.54/0.98 across intensities used	GE significantly higher in trained group (mean +1.4% $p < 0.05$)

Horowitz et al. [36] have shown that cyclists with a high gross efficiency (21.9%) were able to sustain a significantly higher power output (27W) during a 1-hour cycle time-trial performance than a group with a lower gross efficiency (20.4%). Jeukendrup et al. [39] used a mathematical modeling approach to predict that a 1% increase in a cyclist's gross efficiency would result in a 63-s improvement in 40 km time-trial time. It is therefore quite possible that the changes in gross efficiency following training described by Hopker et al. [34] and Sassi et al. [60] will afford a performance advantage.

Potential mechanisms for chronic changes in gross efficiency

▼ Muscle fibre type transformation

Previous work has demonstrated that there is a positive correlation between the percentage of type I fibres and gross efficiency [16,36]. Horowitz et al. [36] have shown that cyclists with a higher percentage of type I fibres also had a significantly higher efficiency and were able to maintain a 9% higher power output during a one-hour performance trial. Both studies largely provide correlation based evidence, and intervention studies are required to evaluate this relationship more thoroughly. Coyle et al. [15] reported a positive relationship between years of endurance training and percentage of type I fibres ($r=0.75$). Again, due to the cross-sectional study design, it was not possible to determine whether the percentage of type I fibres were a response to years of training, or that those cyclists with more type I fibres continue to train and race longer.

Jansson and Kaijser [38] compared metabolic responses in trained cyclists and untrained individuals exercising at 65% $\dot{V}O_{2peak}$. The trained cyclists had a significantly higher percentage of type I fibres (70% vs. 40%) and a greater gross efficiency (22% vs. 19%). Coyle [19] suggested that much of the 8% increase in efficiency observed in a Grand Tour Champion was the result of an increased percentage of type I muscle fibres caused by prolonged intense endurance training and/or prolonged exposure to high altitude conditions. Coyle attributed much of the improvement in efficiency to muscle fibre type adaptations (speculating that the rider's fast twitch fibres took on more of a slow twitch role), even though no muscle biopsies were taken. In contrast, Martin et al. [47] suggested that modifications in diet, training and chemotherapy (with a resultant loss of body and leg mass), were as likely to be responsible.

When training for several hours per day at low force and movement speeds, as in cycling, the possibility of low frequency stimulation-induced transitions from type IIB to type IIA and ultimately type I might be expected [37,59,76]. This has led to the formulation of the 'adaptive range' concept, which describes the adaptive possibilities for each muscle fibre type [27,73,76]. As satellite cells have been shown to be predetermined to end up as a specific muscle fibre type within a certain adaptive range, local genetic factors have been suggested as unimportant [2] in fibre type alteration.

There is a significant difference in tension cost (how much ATP is consumed per unit force generated during an isometric contraction) between human type I and type II fibres [65]. However, during active shortening, thermodynamic efficiency is the same (21–27%), although peak efficiency is reached at different velocities (at around 15% of maximum shortening velocity in both cell types) [29]. Thus, the effect of fibre-type distribution on

whole-body efficiency when cycling is probably not due to differences in contractile efficiency between myosin isoforms. Rather, it may be due to the shortening velocities during cycling being closest to those associated with peak efficiency in type I fibres (as has been suggested elsewhere [16]). Interestingly, marathon training increases the peak shortening velocity of type I fibres [70], and therefore, presumably, the shortening velocity at which peak efficiency is attained. This would improve efficiency during running if the new, most efficient, shortening velocity was better matched with the actual shortening velocity during running. Thus, there is a mechanism by which muscle cells could provide improved contractile efficiency in response to training, without any change in metabolic efficiency.

▼ Factors influencing efficiency in the muscle cell

Oxidative phosphorylation is the main process by which ATP is produced under aerobic conditions. Changes in the efficiency of oxidative phosphorylation will therefore affect cycling efficiency. Adaptations that might affect efficiency which are detectable early in a training programme may be related to the myosin ATP supply. Mitochondrial volume and aerobic capacity increase greatly within the first 4–6 weeks, especially in type II fibres, whilst anaerobic capacity decreases [32,37,59]. It could also be suggested that decreases in submaximal oxygen uptake may be due to changes in the working muscle's oxidative capacity and metabolic processes, represented by an increase in activity of the mitochondrial enzymes [10].

A precursor for the aerobic adaptations seen as a result of training is the production of adenosine monophosphate-activated protein kinase. This enzyme is released as a result of intensive training [24,41], and may cause the up-regulation of PGC-1 α . This in turn is thought to regulate mitochondrial biogenesis in type I, IIA and IIX fibers [58,67,68]. However, these observations are limited to low and moderately trained individuals, who demonstrate marked physiological adaptations and improvements in fitness. Whether the same adaptations are typical of a trained population remains to be determined.

Key questions also remain unanswered regarding the efficiency of energy transfer within the mitochondria and the possible role of the uncoupling of oxidative phosphorylation. Energy liberated from respiratory chain reactions are used to translocate protons across the inner mitochondrial membrane, creating an electrochemical potential [7]. This potential is subsequently used to drive the endergonic re-phosphorylation of ADP to ATP by the ATP-synthase, a reaction which is stoichiometric [8]. However, protons are able to leak back across the inner membrane without driving the ATP-synthase, a phenomenon known as uncoupling. In skeletal muscle, the most important proteins mediating this process appear to be the adenine nucleotide translocase (ANT) and uncoupling protein-3 (UCP3). Thus increases in the content or activity of these proteins might have adverse effects on cycling efficiency.

Uncoupling accounts for around 50% of resting oxygen consumption in rodent muscle [57]. Its contribution to efficiency during exercise is unclear, particularly as proton leak is of diminishing importance as mitochondrial respiratory rate rises [9]. In the only study of UCP3 content and efficiency in cyclists to date, muscle UCP3 content and work efficiency were negatively correlated in a cohort of mixed-ability cyclists [48]. UCP3 content was greater in untrained compared to trained individuals.

Therefore, the process of training might have reduced the expression of UCP3, increased the coupling of oxidative phosphorylation, and improved gross efficiency when cycling. However, this mechanism is yet to be observed following a period of training. The contribution of ANT activity to whole-body efficiency remains unexplored. Longitudinal or interventional studies are needed to illuminate the contribution of mitochondrial uncoupling to exercise efficiency in humans.

Conclusions

Until recently it has been assumed that training had no impact on cycling efficiency. This was largely based on the results of investigations that did not find a difference between trained and untrained cyclists. It now appears that insufficient methods (e.g. small sample sizes and inappropriate exercise intensities) may have resulted in type II errors in many of these studies. To enable the identification of training related changes and obtain consistency in efficiency measurements pre-test preparation, ergometer set-up and methods used for data collection must all be valid and reliable. Recent evidence suggests that training (especially at high intensities) improves gross efficiency in cycling. Potential mechanisms which might be responsible for training related increases in GE include muscle fibre type transformation, aerobic enzyme capacity within the muscle and the expression of PGC1 α , ANT and UCP3. Future studies should seek to fully investigate the mechanisms responsible for determining gross efficiency via the use of intervention methodologies and invasive biopsy techniques. Further research is required to clarify the role of specific training regimes on the development of gross efficiency and the underpinning changes.

References

- Andreyev ATO, Bondareva TO, Dedukhova VI, Mokhova EN, Skulachev VP, Tsofina LM, Volkov NI, Vvgodina TV. The ATP/ADP-antiporter is involved in the uncoupling effect of fatty acids on mitochondria. *Eur J Biochem* 1989; 182: 585–592
- Astrand P, Rodahl K, Dahl H, Stromme S. *Textbook of Work Physiology: Physiological Bases of Exercise* (4th Ed.). Illinois: Human Kinetics; 2003
- Atkinson G, Nevill A. Statistical methods for assessing measurement error (reliability) in variables relevant to sports medicine. *Sports Med* 1998; 26: 217–236
- Barbeau P, Serresse O, Boulay M. Using maximal and submaximal aerobic variables to monitor elite cyclists during a season. *Med Sci Sports Exerc* 1993; 25: 1062–1069
- Batterham AM, Atkinson G. How big does my sample need to be? A primer on the murky world of sample size estimation. *Phys Ther Sport* 2005; 6: 153–163
- Boning D, Gonen Y, Maassen N. Relationship between work load, pedal frequency, and physical fitness. *Int J Sports Med* 1984; 5: 92–97
- Brand MD. The efficiency and plasticity of mitochondrial energy transduction. *Biochem Soc Trans* 2005; 33: 897–904
- Brookes PS, Rolfe DFS, Brand MD. The proton permeability of liposomes made from mitochondrial inner membrane phospholipids: Comparison with isolated mitochondria. *J Membr Biol* 1997; 155: 167–174
- Brown GC. Control of respiration and ATP synthesis in mammalian mitochondria and cells. *Biochem J* 1992; 284: 1–13
- Burgomaster KA, Hughes SC, Heigenhauser GJF, Bradwell SN, Gibala MJ. Six sessions of sprint interval training increases muscle oxidative potential and cycle endurance capacity in humans. *J Appl Physiol* 2005; 98: 1985–1990
- Cavanagh P, Kram R. The efficiency of human movement – a statement of the problem. *Med Sci Sports Exerc* 1985; 17: 304–308
- Chavarren J, Calbet JAL. Cycling efficiency and pedalling frequency in road cyclists. *Eur J Appl Physiol* 1999; 80: 555–563
- Chuang ML, Ting H, Otsuka T, Sun XG, Chiu FY, Beaver WL, Hansen JE, Lewis DA, Wasserman K. Aerobically generated CO₂ stored during early exercise. *J Appl Physiol* 1999; 87: 1048–1058
- Coast J, Cox R, Welch H. Optimal pedalling rate in prolonged bouts of cycle ergometry. *Med Sci Sports Exerc* 1986; 18: 225–230
- Coyle EF, Feltner ME, Kautz SA, Hamilton MT, Mountain SJ, Baylor AM, Abraham LD, Petrek GW. Physiological and biochemical factors associated with elite endurance cycling performance. *Med Sci Sports Exerc* 1991; 23: 93–107
- Coyle EF, Sidossis LS, Horowitz JF, Beltz JA. Cycling efficiency is related to the percentage of type I muscle fibers. *Med Sci Sports Exerc* 1992; 24: 782–788
- Coyle EF. Integration of the physiological factors determining endurance performance ability. *Exerc Sport Sci Rev* 1995; 23: 25–64
- Coyle EF. Physiological determinants of endurance exercise performance. *J Sci Med Sport* 1999; 2: 181–189
- Coyle EF. Improved muscular efficiency displayed as Tour de France champion matures. *J Appl Physiol* 2005; 98: 2191–2196
- Echtay KS, Roussel D. Superoxide activates mitochondrial uncoupling proteins. *Nature* 2002; 415: 96–99
- Echtay KS, Winkler E, Frischmuth K, Klingenberg M. Uncoupling proteins 2 and 3 are highly active H⁺ transporters and highly nucleotide sensitive when activated by coenzyme Q (ubiquinone). *Proc Natl Acad Sci USA* 2001; 98: 1416–1421
- Erdfelder E, Faul F, Buchner A. Gpower: A general power analysis program. *Behav Res Methods* 1996; 1: 1–11
- Gaesser GA, Brooks GA. Muscular efficiency during steady-state exercise: effects of speed and work rate. *J Appl Physiol* 1975; 38: 1132–1139
- Gibala MJ, McGee SL, Garnham A, Hargreaves M. Effect of high-intensity interval exercise on signaling proteins involved in skeletal muscle remodeling in humans. *Appl Physiol Nutr Metab* 2006; 31: S37
- Gore CJ, Ashenden MJ, Sharpe K, Martin DT. Delta Efficiency calculation in Tour de France champion is wrong. *J Appl Physiol* 2008; 98: 1020
- Green S, Dawson BT. The oxygen uptake-power regression in cyclists and untrained men: implications for the accumulated oxygen deficit. *Eur J Appl Physiol* 1995; 70: 351–359
- Gundersen K. Determination of muscle contractile properties: The importance of the nerve. *Acta Physiol Scand* 1998; 31: 536–543
- Hagberg J, Mullin J, Giese M, Spitznagel E. Effect of pedalling rate on submaximal exercise responses of competitive cyclists. *J Appl Physiol* 1981; 51: 447–451
- He ZH, Bottinelli R, Pellegrino MA, Ferenczi MA, Reggiani C. ATP consumption and efficiency of human single muscle fibers with different myosin isoform composition. *Biophys J* 2000; 79: 945–961
- Heil DP, Wilcox AP, Quinn CM. Cardiorespiratory responses to seat-tube angle variation during steady-state cycling. *Med Sci Sports Exerc* 1995; 27: 730–735
- Hettinga FJ, De Koning JJ, de Vrijer A, Wust RCI, Daanen HAM, Foster C. The effect of ambient temperature on gross-efficiency in cycling. *Eur J Appl Physiol* 2007; 101: 465–471
- Holloszy JO. Adaptation of skeletal muscle to endurance exercise. *Med Sci Sports Exerc* 1975; 7: 155–164
- Hopker JG, Coleman DA, Wiles J. Differences in efficiency between trained and recreational cyclists. *Appl Physiol Nutr Metab* 2007; 32: 1036–1042
- Hopker JG, Coleman DA, Passfield L. Changes in cycling efficiency over a competitive season. *Med Sci Sports Exerc* 2009; 41: 912–919
- Hopkins WG, Hawley JA, Burke LM. Design and analysis of research on sport performance enhancement. *Med Sci Sports Exerc* 1999; 31: 472–485
- Horowitz JF, Sidossis LS, Coyle EF. High efficiency of type 1 muscle fibres improves performance. *Int J Sports Med* 1994; 15: 152–157
- Jansson E, Kaijser L. Muscle adaptation to extreme endurance training in man. *Acta Physiol Scand* 1977; 100: 315–324
- Jansson E, Kaijser L. Substrate utilisation and enzymes in skeletal muscle of extremely endurance-trained men. *J Appl Physiol* 1987; 62: 999–1005
- Jeukendrup AE, Craig N, Hawley J. The bioenergetics of world class cycling. *J Sci Med Sport* 2000; 3: 414–433
- Kautz SA, Neptune RR. Biomechanical determinants of pedaling energetics: internal and external work are not independent. *Exerc Sport Sci Rev* 2002; 30: 159–165
- Koulmann N, Bigard AX. Interaction between signalling pathways involved in skeletal muscle responses to endurance exercise. *Pflugers Arch* 2006; 452: 125–139
- Künstlinger U, Ludwig H-G, Stegmann J. Force kinetics and oxygen consumption during bicycling ergometer work in road cyclists and reference group. *Eur J Appl Physiol* 1985; 54: 58–61

- 43 Kyröläinen H, Pullinen T, Candau R, Avela J, Huttunen P, Komi PV. Effects of marathon running on running economy and kinematics. *Eur J Appl Physiol* 2000; 82: 297–304
- 44 Lucia A, Hoyos J, Pardo J, Chicharro JL. Metabolic and neuromuscular adaptations to endurance training in professional cyclists: A longitudinal study. *Jpn J Physiol* 2000; 50: 381–388
- 45 Marsh AP, Martin PE. The association between cycling experience and preferred and most economical cadences. *Med Sci Sports Exerc* 1993; 25: 1269–1274
- 46 Marsh AP, Martin PE, Foley KO. Effect of cadence, cycling experience, and aerobic power on delta efficiency during cycling. *Med Sci Sports Exerc* 2000; 32: 1630–1634
- 47 Martin D, Quod M, Gore C. Has Armstrong's cycle efficiency improved? *J Appl Physiol* 2005; 99: 1628–1629
- 48 Mogensen M, Bagger M, Pedersen PK, Fernstrom M, Sahlin K. Cycling efficiency in humans is related to low UCP3 content and to type I fibres but not to mitochondrial efficiency. *J Physiol* 2006; 571: 669–681
- 49 Moseley L, Jeukendrup A. The reliability of cycling efficiency. *Med Sci Sports Exerc* 2001; 33: 621–627
- 50 Moseley L, Achten J, Martin JC, Jeukendrup AE. No differences in cycling efficiency between world-class and recreational cyclists. *Int J Sports Med* 2004; 25: 374–379
- 51 Murphy MP, Echtay KS, Blaikie FH, Asin-Cayuela J, Cocheme HM, Green K, Buckingham JA, Taylor ER, Hurrell F, Hughes G, Miwa S, Cooper CE, Svistunenko DA, Smith RA, Brand MD. Superoxide activates uncoupling proteins by generating carbon-centered radicals and initiating lipid peroxidation: studies using a mitochondria-targeted spin trap derived from {alpha}-phenyl-N-tert-butyl-nitron. *J Biol Chem* 2003; 278: 48534–48545
- 52 Nickleberry BL, Brooks GA. No effect of cycling experience on leg cycle ergometer efficiency. *Med Sci Sports Exerc* 1996; 28: 1396–1401
- 53 Olds T, Norton K, Craig N, Olive S, Lowe E. The limits of the possible: Models of power supply and the demands in cycling. *Aust J Sci Med Sport* 1995; 27: 29–33
- 54 Passfield L, Doust JH. Changes in cycling efficiency and performance after endurance exercise. *Med Sci Sports Exerc* 2000; 32: 1935–1941
- 55 Poole D, Gaesser M, Hogan D, Knight D, Wagner P. Pulmonary and leg VO₂ during submaximal exercise: implications for muscular efficiency. *J Appl Physiol*. 1992; 72: 805–810
- 56 Price D, Donne B. Effect of variation in seat tube angle at different seat heights on submaximal cycling performance in man. *J Sport Sci* 1997; 15: 395–402
- 57 Rolfe DF, Brand MD. Contribution of mitochondrial proton leak to skeletal muscle respiration and to standard metabolic rate. *Am J Physiol* 1996; 271: C1380–1389
- 58 Russell AP, Feilchenfeldt J, Schreiber S, Praz M, Crettenand A, Gobelet C, Meier CA, Bell DR, Kralli A, Giacobino JP, Deriaz O. Endurance training in humans leads to fiber type-specific increases in levels of peroxisome proliferator-activated receptor-γ coactivator-1 and peroxisome proliferator-activated receptor-α in skeletal muscle. *Diabetes* 2003; 52: 2874–2881
- 59 Saltin B, Gollnick PD. Skeletal muscle adaptability: significance for metabolism and performance. In: Peachey LD, Adrian RH, Geiger SR (eds) *Skeletal Muscle. Handbook of Physiology*, sect. 10. Bethesda: American Physiological Society; 1983; 555–631
- 60 Sassi A, Impellizzeri FM, Morelli A, Menaspà P, Rampinini E. Seasonal changes in aerobic fitness indices in elite cyclists. *Appl Physiol Nut Metab* 2008; 33: 735–742
- 61 Schumacher YO, Vogt S, Roecker K, Schmid A. Scientific considerations for physiological evaluations of elite athletes. *J Appl Physiol* 2005; 99: 1630–1632
- 62 Seabury J. Influence of pedalling rate and power output on energy expenditure during bicycle ergometry. *Ergonomics* 1977; 20: 491–498
- 63 Sidossis LS, Horowitz JF, Coyle EF. Load and velocity of contraction influence gross and delta mechanical efficiency. *Int J Sports Med* 1992; 13: 407–411
- 64 Stainsby W, Gladden L, Barclay J, Wilson B. Exercise efficiency: validity of base-line subtractions. *J Appl Physiol* 1980; 48: 518–522
- 65 Stienen GJ, Kiers JL, Bottinelli R, Reggiani C. Myofibrillar ATPase activity in skinned human skeletal muscle fibres: fibre type and temperature dependence. *J Physiol* 1996; 493: 299–307
- 66 Suzuki Y. Mechanical efficiency of fast and slow twitch muscle fibres in man during cycling. *J Appl Physiol* 1979; 47: 263–267
- 67 Taylor EB, Lamb JD, Hurst RW, Chesser DG, Ellingson WJ, Greenwood LJ, Porter BB, Herway ST, Winder WW. Endurance training increases skeletal muscle LKB1 and PGC-1α protein abundance: effects of time and intensity. *Am J Physiol Endocrinol Metab* 2005; 289: E960–E968
- 68 Terada S, Kawanaka K, Goto M, Shimokawa T, Tabata I. Effects of high-intensity intermittent swimming on PGC-1α protein expression in rat skeletal muscle. *Acta Physiol Scand* 2005; 184: 59–65
- 69 Thomas JR, Nelson JK, Silverman SJ. *Research Methods in Physical Activity* (5th Ed.). Champaign IL: Human Kinetics; 2005
- 70 Trappe S, Harber M, Creer A, Gallagher P, Sliwka D, Minchev K, Whitsett D. Single muscle fiber adaptations with marathon training. *J Appl Physiol* 2006; 101: 721–727
- 71 Twist C, Easton RG. The effect of exercise-induced muscle damage on perceived exertion and cycling endurance performance. *Eur J Appl Physiol* 2009; 105: 559–567
- 72 Wasserman KW, Hansen JE, Sue DY, Stinger WW, Whipp BJ. *Principles of Exercise Testing and Interpretation*. Philadelphia: Lippincott, Williams and Wilkins; 2005
- 73 Westgaard RH, Lomo T. Control of contractile properties within adaptive ranges by patterns of impulse activity in the rat. *J Neurosci* 1988; 8: 4415–4426
- 74 Whipp BJ. The slow component of O₂ uptake kinetics during heavy exercise. *Med Sci Sports Exerc* 1994; 26: 1319–1326
- 75 Whipp BJ, Wasserman K. Oxygen uptake kinetics for various intensities of constant-load work. *J Appl Physiol* 1972; 33: 351–356
- 76 Windisch A, Gundersen K, Szabolcs MJ, Gruber H, Lomo T. Fast to slow transformation of denervated and electrically stimulated rat muscle. *J Physiol* 1998; 510: 623–632
- 77 Zameziati K, Mornieux G, Roufflet D, Belli A. Relationship between the increase of effectiveness indexes and the increase of muscular efficiency with cycling power. *Eur J Appl Physiol* 2006; 96: 274–281
- 78 Zoladz J, Rademaker ACHJ, Sargeant AJ. Non-linear relationship between O₂ uptake and power output at high intensities of exercise in humans. *J Physiol* 1995; 488: 211–217