



# Kent Academic Repository

Kengne, Andre Pascal, Bentham, James, Zhou, Bin, Peer, Nasheeta, Matsha, Tandi E., Bixby, Honor, Cesare, Mariachiara Di, Hajifathalian, Kaveh, Lu, Yuan, Taddei, Cristina and others (2017) *Trends in obesity and diabetes across regions in Africa from 1980 to 2014: an analysis of pooled population-based studies*. *International Journal of Epidemiology*, 46 (5). pp. 1421-1432. ISSN 0300-5771.

## Downloaded from

<https://kar.kent.ac.uk/64446/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1093/ije/dyx078>

## This document version

Author's Accepted Manuscript

## DOI for this version

## Licence for this version

UNSPECIFIED

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

### Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

**Trends in obesity and diabetes across regions in Africa from 1980 to 2014: an analysis of pooled population-based studies**

NCD Risk Factor Collaboration (NCD-RisC) – Africa Working Group\*

\* Members listed at end of article

Correspondence to Prof Andre Pascal Kengne, Non-Communicable Diseases Research Unit, South African Medical Research Council, Francie van Zijl Drive, Parow Valley, 7505, Cape Town, South Africa. [Andre.kengne@mrc.ac.za](mailto:Andre.kengne@mrc.ac.za)

## Abstract

**Background:** The 2016 *Dar Es Salaam Call to Action on Diabetes and other NCDs* advocates national multi-sectoral NCD strategies and action plans based on available data and information from countries of sub-Saharan Africa and beyond. We estimated trends, from 1980 to 2014, in age-standardised mean body mass index (BMI) and diabetes prevalence in these countries in order to assess the co-progression and assist policy formulation.

**Methods:** We pooled data from African and world-wide population-based studies which measured height, weight, and biomarkers to assess diabetes status in adults aged  $\geq 18$  years. A Bayesian hierarchical model was used to estimate trends, by sex, [for 200 countries and territories including 53 countries across five African regions, \(central, eastern, northern, southern and western\)](#) in mean BMI and diabetes prevalence (defined as either fasting plasma glucose of  $\geq 7.0$  mmol/L, history of diabetes diagnosis, or use of insulin or oral glucose control agents).

## Results

African data came from 245 population-based surveys (1.2 million participants) for BMI and 76 surveys (182 000 participants) for diabetes prevalence estimates. Countries with the highest number of data [points-sources](#) for BMI were South Africa (n=17), Nigeria (n=15) and Egypt (n=13); and for diabetes estimates, Tanzania (n=8), Tunisia (n=7), Cameroon, Egypt and South Africa (all n=6).

The age-standardised mean BMI increased from 21.0 kg/m<sup>2</sup> (95% credible interval: 20.3-21.7) to 23.0 kg/m<sup>2</sup> (22.7-23.3) in men, and from 21.9 kg/m<sup>2</sup> (21.3-22.5) to 24.9 kg/m<sup>2</sup> (24.6-25.1) in women. The age-standardised prevalence of diabetes increased from 3.4% (1.5-6.3) to 8.5% (6.5-10.8) in men, and from 4.1% (2.0-7.5) to 8.9% (6.9-11.2) in women.

Estimates in northern and southern regions were mostly higher than the global average; those in central, eastern and western regions were lower than global averages. A positive association (correlation coefficient  $\approx 0.9$ ) was observed between mean BMI and diabetes prevalence in both sexes in 1980 and 2014.

**Conclusion:** These estimates, based on limited data sources, confirm the rapidly increasing burden of diabetes in Africa. This rise is being driven, at least in part, by increasing adiposity, with regional

variations in observed trends. African countries' efforts to prevent and control diabetes and obesity should integrate the setting up of reliable monitoring systems, consistent with the WHO's Global Monitoring System Framework.

**Key words** – diabetes, adiposity, body mass index, Africa, prevalence, trends

*Word count - 360*

## **Introduction**

Despite a high burden of infectious diseases in Africa, non-communicable diseases (NCDs), including type 2 diabetes mellitus (hereafter referred to as ‘diabetes’), are rising steadily, contributing substantially to morbidity and mortality. This is partly driven by a growing proportion of elderly people [1]. Furthermore, economic growth and accompanying rapid urbanisation have led to a burgeoning wealthier middle-class, possibly with shifts in diet and physical activity – contributing to the development of obesity which, together with a genetic predisposition and epigenetic changes [2, 3], increases the risk of diabetes [4].

Healthcare systems in Africa are now facing new disease burdens they are ill-equipped to manage. A number of NCD initiatives, informed by global resolutions, including disease-specific initiatives such as for diabetes in Africa [5-8], have been developed [4, 9]. However, tangible responses from national and international stakeholders have been slow, and have largely overlooked adiposity and the unhealthy food environment. Driven by maternal, new-born and child health (MNCH), the traditional focus of investment in preventive health has been on undernutrition and food insecurity.

To highlight the increasing diabetes prevalence and the key contributory role played by obesity requires quality local data. Furthermore, despite the strong link between adiposity and diabetes, efforts to co-examine the burden of the two conditions have been limited. Here, we estimate trends in age-standardised mean body mass index (BMI) and diabetes prevalence across Africa from 1980 to 2014. We aimed to understand (a) the extent and quality of available data, (b) differences in patterns by regions and sex, and (c) the association of adiposity with diabetes prevalence.

## **Methods**

The full methods of data collection, processing and analysis in the NCD Risk Factor Collaboration are available elsewhere [10, 11] and in the Supplementary Textbox 1. We estimated trends for mean BMI and diabetes prevalence from 1980 to 2014, in  $\geq 18$ -year-old adults across 53 countries organised into five regions (Supplementary Table 1).

We included data sources representative of national, sub-national, or community populations, with measured weight and height, or at least one of the following bio-markers: FPG, oral glucose tolerance test (OGTT) or glycosylated haemoglobin (HbA1c). We converted diabetes measured using different biomarkers and definitions to a common definition of fasting plasma glucose (FPG)  $\geq 7.0$  mmol/L or a history of diabetes with ongoing treatment as described previously [11]. Analyses were done separately for men and women, using a statistical model described and validated previously [10, 11]. The model had a hierarchical structure in which estimates for each country and year were informed by its own data, data from other years in the same country and in other countries in the same region. The model also accounted for non-linear time trends and age associations. The reported credible intervals represent the 2.5th - 97.5th percentiles of the posterior distributions. Estimates were standardised to the World Health Organization (WHO) standard population.

## Results

### *Data available*

The analysis used data from 245 population-based surveys encompassing 1.2 million participants aged  $\geq 18$  years for BMI estimates, and 76 surveys (182,000 participants) for diabetes estimates [10]. These data were collected between 1984 and 2014 in 50 countries (94% of African countries) for BMI, and between 1981 and 2014 in 32 countries (60% of African countries) for diabetes (Figure 1) [11].

The distribution of surveys per region for BMI estimates was as follows: 19 surveys (55,000 participants) for central Africa, 68 (325 000) for eastern, 26 (170,000) for northern, 36 (299 000) for southern and 96 (394 000) for western (Figure 1a). South Africa (n=17), Nigeria (n=15) and Egypt (n=13) had the highest number of data points (regardless of sources), while Angola, Djibouti and Somalia had none (Figure 2).

Equivalent figures for diabetes estimates were one survey (3600 participants) for central Africa, 24 (77 000) for eastern, 20 (43 000) for northern, 10 (14 000) for southern and 21 (45 000) for western (Figure 1b). Tanzania (n=8), Tunisia (n=7), Cameroon, Egypt and South Africa (all n=6) had the highest number of data points. No data were available from 21 countries: Angola, Burundi, Chad,

Congo, Cote d'Ivoire, Djibouti, DR Congo, Equatorial Guinea, Ethiopia, Gabon, Guinea Bissau, Kenya, Madagascar, Mali, Mozambique, Namibia, Niger, Sao Tome and Principe, Senegal, Sierra Leone, Somalia (Figure 2).

Regarding the representativeness of data sources, 173 surveys (1.1 million participants) for BMI and 32 surveys (104 000 participants) for diabetes prevalence estimates had national coverage, while the remainder were sub-national or community based (Figure 1). 76 data sources (203 000 participants) for BMI and 28 data sources (60 000 participants) for diabetes (Figure 1) were collected prior to 2000. These data were augmented with 1,453 sources for BMI and 675 for diabetes from other regions. This helped the interpretation of features, such as age patterns and the effects of co-variables through the hierarchical structure, which also influenced trends.

### ***Obesity trends***

From 1980 to 2014, the age-standardised mean BMI increased from 21.0 kg/m<sup>2</sup> (95% credible interval 20.3-21.7) to 23.0 kg/m<sup>2</sup> (22.7-23.3) in men, and from 21.9 kg/m<sup>2</sup> (21.3-22.5) to 24.9 kg/m<sup>2</sup> (24.6-25.1) in women. Mean BMI increased over time across all regions, paralleling the global average; however, there were suggestions, over recent years, of steeper curves in women compared with world-wide trends (Figure 3a). In both men and women, the lowest mean BMI, over time, was usually recorded in central Africa, while the highest, for men, was recorded in northern Africa, and, for women, in northern and southern Africa.

Mean BMI was higher than the global average in northern and southern Africa, and lower in central, eastern and western Africa. In many cases, credible intervals around mean estimates overlapped across regions and with that for the global mean, due to a combination of limited available data and the similarity of the estimates. A scatterplot of mean BMI in men versus women in 2014 is shown in Figure 4. Across the five regions, mean BMI appeared to be mostly higher in women than in men. Per country, mean BMI in 2014 ranged from 20.1 kg/m<sup>2</sup> (CI: 19.2-21.0) in Ethiopia to 27.3 kg/m<sup>2</sup> (26.5-28.1) in Egypt in men, and from 21.0 kg/m<sup>2</sup> in Eritrea, Ethiopia and Madagascar to 30.6 kg/m<sup>2</sup> (30.0-31.2) in Egypt in women (Figure 5a and Supplementary Table 2).

### ***Diabetes prevalence trends***

Age-standardised diabetes prevalence increased, from 1980 to 2014, from 3.4% (1.5-6.3) to 8.5 % (6.5-10.8) in men, and from 4.1% (2.0-7.5) to 8.9 % (6.9-11.2) in women. Across regions and in both sexes, age-standardised diabetes prevalence increased over time along with the global average (Figure 3b).

Prevalences in northern Africa were higher than the global average, while those in central, eastern and western Africa were lower. In southern Africa, compared with the global average, figures were higher in women, but similar in men. While regional time-trends mirrored the global pattern in most regions, curves were steeper than the global average in recent years in northern African men and women, and possibly in southern African women.

Credible intervals around the trend curves for most regions overlapped, suggesting that regional differences, while possible, were, due to limited data availability and large variability in available data (Figure 3b) not statistically distinguishable. Diabetes prevalence by sex was mostly similar in eastern, western and southern Africa, but might have been higher in men than in women in central Africa, and higher in women than in men in northern Africa (Figure 4). Within countries, age-standardised diabetes prevalence in 2014 ranged from 4.2% (1.8-7.9) in Burundi to 16.0% (10.0-23.6) in Egypt for men, and from 4.1% (1.8-7.7) in Burundi to 19.8% (12.9-28.2) in Egypt for women (Figure 5b and Supplementary Table 2).

### ***Associations between diabetes and BMI***

Across countries, there was a strong positive association between mean BMI level and diabetes prevalence in 2014 for men and women (correlation coefficients of 0.86 and 0.88, respectively) (Figure 6). This is slightly weaker than the associations observed in 1980 (0.88 and 0.91 respectively). A moderately strong association between absolute change in BMI and relative change in diabetes was observed in men ( $r=0.76$ ), but the association was weak in women ( $r=0.34$ ).

## **Discussion**

Over the last three and half decades, age-standardised mean BMI and diabetes prevalence have steadily increased across African regions, at least as steeply as the global average. There might have been an acceleration of trends for women's BMI in all regions, and for diabetes prevalence in northern African men and women. Estimates in northern and southern Africa were mostly higher than global averages, while those in other regions were mostly lower. This probably reflects different stages of the epidemiological transitions and other determinants of high BMI and diabetes. Arab countries not only had the highest diabetes prevalence in Africa, but also have some of the highest worldwide prevalences [11], explained to some extent by differences in diet, physical (in)activity and obesity [10]. Mean BMI was consistently higher in women than in men, while male-female differences in diabetes prevalence varied across regions.

The fact that average BMI increased steadily over the same period, as did evidence of an association between BMI and diabetes in 2014, affirms the role of adiposity as a driver of diabetes across Africa. Estimates were, however, based on limited data sources, particularly in sub-Saharan Africa (SSA), with nearly 20 SSA countries having no data for diabetes.

Other investigators have estimated diabetes prevalence in Africa. Manne-Goehler analysed individual participant data from surveys conducted in mostly 15 to 64-year-olds across twelve SSA countries between 2005 and 2015 (STEPwise approach applied in 11 surveys), and reported a median diabetes prevalence of 5% (range 2% to 14%) [12]. The International Diabetes Federation (IDF) Diabetes Atlas, using substantially fewer data sources than ours, estimated the age-standardised diabetes prevalence in 20 to 79-year-old adults in 2015 to be 3.8% (95% CI: 2.1-6.7%) in SSA, and 10.7% (7.4-14.2%) in the Middle East and North Africa. In a meta-analysis of prevalence studies, published between 2000 and 2015, of diabetes among older SSA persons, the overall prevalence was 13.7% (11.3-16.3%) among  $\geq 55$  year olds [13]. Published studies and ours have used overlapping data sources, but different time-periods, geographic coverage, diabetes definitions, population coverage, modelling/estimation strategies, and the results are, therefore, not directly comparable.

Our findings of increasing diabetes prevalence across Africa between 1980 and 2014 are consistent with successive IDF estimates [14]. However, the IDF does not estimate trends, relying instead on

successive cross-sectional reports, while an earlier trend study focused on FPG [15]. Therefore, our estimates are the first consistent diabetes trend estimates for Africa, including by its regions.

Studies in populations of European origin have consistently reported higher diabetes prevalence in men than in women, with widening gaps as populations undergo the economic transition from less developed to more affluent societies [16, 17]. In the US National Health Interview Surveys, diabetes prevalence in African-Americans was higher in women than in men over the years 1963-1985.

However, during this same period, diabetes prevalence increased more steeply in African-American men than in women [18]. We found variable patterns of sex differences, with diabetes prevalence in 2014 being mostly similar or higher in men at a low prevalence, and higher in women at a high prevalence. Several mechanisms, reviewed elsewhere [19], have been suggested to explain sex differences in diabetes incidence (or the lack thereof) across populations and settings [20].

Five years after the adoption of the United Nations' resolution on NCDs [21], and three years after the introduction of the Global NCD Action Plan [8], good quality nationally representative data on the prevalence of such a major NCD are still lacking in most African countries. Our findings, in consequence, were based on just a few data points per country and on extrapolations for many countries. STEPs surveys are growing data sources for diabetes and obesity studies in Africa. Their advantages include national or regional representativeness for some, and the use of standardised questionnaires and protocols which facilitate cross-country comparisons. STEPs data, however, have limitations discussed extensively elsewhere [12, 13]. Furthermore, measurement of bio-markers has not been implemented across all STEPs surveys in Africa. This is unfortunate because up to two-thirds of cases remain undiagnosed [14] and are uncovered only through biomarker measurements. Where biomarkers have been measured, the WHO recommends testing fasting venous blood glucose for diabetes [22]. There are concerns that fasting glucose alone may not capture all cases of undiagnosed diabetes in Africans, whose early stages of the disease are probably expressed through elevated 2-hour post-OGTT glucose. A study in 'Coloured' (i.e. mixed-race) South Africans indicated that nearly 50% with undiagnosed diabetes would be missed in the absence of OGTT [23].

Notwithstanding the challenges of implementing optimal diabetes biomarker measurements in population-based surveys, the many uncertainties around diabetes's true burden invites cross-country efforts to generate reliable data. This, together with pleas for data from surveys conducted across Africa to be more publicly available, to maximise their use for research and policy formulation [12, [2423](#)], might serve as baseline support for the introduction of successful diabetes prevention and control efforts.

Other limitations in the interpretation of these study findings warrant consideration. Regions, as defined in this study, are general geographical entities comprising countries with substantial heterogeneities which can affect diabetes and BMI estimates. However, each region reflects existing common entities recognised by the African Union, and includes some form of concerted action on health.

The current study, however, has major strengths. This is the first time estimates have been made, from a broad African perspective, of trends in diabetes and in BMI. This provides an opportunity for detailed cross-regional comparisons. Despite the limited number of data sources, estimates in this study are based on a larger amount of population-based data than previous initiatives [12, 14].

A high-level response to the growing diabetes threat includes the 2007 Strategy for Diabetes Prevention and Control in Africa [5], the 2009 Mauritius Call for Action [6], and the 2016 Dar es Salaam Call to Action on Diabetes and other NCDs [7]. More generally, all countries adopted the WHO Global Action Plan for the Prevention and Control of NCDs 2013-2020 [8]. Despite the increasing awareness of the growing diabetes burden, ensuing investment and actions are not yet commensurate with the magnitude of the challenge [[2524](#)]. Going forward, the Institute of Medicine has recognised the importance of locally relevant evidence to successfully contextualise and implement prevention and control lessons from other parts of the world [[2625](#)].

Africa's continuing urbanisation offers an opportunity to develop urban environments conducive to healthy choices which could reduce obesity and diabetes risk. Tailoring the MNCH platforms which fight undernutrition to address healthy eating in general could prevent and control both under- and

over-nutrition. Locally relevant solutions are yet to be devised to promote effective diabetes risk reversal [2726] in Africa, where weight loss-related HIV/AIDS stigma is common. However, the platforms created to address common infectious diseases such as HIV/AIDS and tuberculosis offer excellent opportunities for implementing diabetes risk-screening and reduction strategies [2827]. Integrating prevention and care services for common chronic infectious and non-infectious diseases in Africa [2928] could facilitate, for NCD prevention and control, the transfer of skills, and leverage some of the abundant infectious disease resources.

The current study has refined, for Africa, the previous global efforts of the NCD Risk Factor Collaboration [10, 11]. The prevalence of diabetes is substantial and possibly growing faster than elsewhere, probably triggered by increasing obesity. This study highlights the urgent need for locally relevant knowledge, supportive policies, adequately equipped health services, and optimal social and physical environments to address the dual burden of rising obesity and diabetes.

#### **Key message box**

- This is the first detailed analysis of trend in adiposity and diabetes prevalence in Africa, both overall and by major geographic regions.
- We used validated methods to re-analyse and pooled population-based sources with measured data on at least one diabetes biomarker and/or body weight and height, including over 250 African sources.
- Age-standardised mean body mass index and diabetes prevalence have steadily increased across Africa since 1980, tracking in general with the global averages.
- At any given time, estimates in northern Africa (driven by Egypt) and in southern Africa (driven by South Africa) appeared higher than the global average, while estimates for other regions were mostly lower.
- There was a weak association between change in gross domestic product and change in adiposity and diabetes prevalence, highlighting the coping challenges facing African

countries, and the need for innovative approaches to diabetes and obesity prevention and control.

**Funding** – This study was supported by the Wellcome Trust and the South African Medical Research Council

**Conflict of interest** – ~~M Ezzati received a charitable grant from Youth Health Program of Astra Zeneca, and personal fees from Third bridge and Prudential, outside the submitted work.~~ [N Levitt's unit received funding for a diabetes educator from Lilly Laboratories and Roche Diagnostics, and N Levitt received funding from Sanofi Aventis to attend a regional diabetes meeting. A Schutte has received speaker fees for presentations on hypertension management from Boehringer-Ingelheim, Novartis and Omron Healthcare.](#) ~~M Ezzati received a charitable grant from Youth Health Program of Astra Zeneca, and personal fees from Third bridge and Prudential, outside the submitted work.~~ All other authors declared no conflict of interest.

### **Contributors**

APK, JB and ME conceived the study. All authors contributed to data collection, pooling analysis, and/or preparation of the results. APK and JB wrote the first draft with input from BZ, NP, TEM, PB, ME and JCM. All other authors commented on the paper and contributed to the revision.

**NCD Risk Factor Collaboration (NCD-RisC) – Africa Working Group** (\* joint first authors; † joint senior authors; ‡ equal contribution, listed alphabetically)

Andre Pascal Kengne (South African Medical Research Council, South Africa)\*; James Bentham (Imperial College London, UK)\*; Bin Zhou (Imperial College London, UK); Nasheeta Peer (South African Medical Research Council, South Africa); Tandi E Matsha (Cape Peninsula University of Technology, South Africa); Honor Bixby (Imperial College London, UK); Mariachiara Di Cesare (Middlesex University, UK); Kaveh Hajifathalian (Cleveland Clinic, USA); Yuan Lu (Yale University, USA); Cristina Taddei (Imperial College London, UK); Pascal Bovet (University of Lausanne, Switzerland; Ministry of Health, Seychelles); Catherine Kyobutungi (African Population and Health

Research Center, Kenya); Charles Agyemang (University of Amsterdam, The Netherlands)<sup>‡</sup>; Hajer Aounallah-Skhiri (National Institute of Public Health, Tunisia)<sup>‡</sup>; Felix K Assah (University of Yaoundé 1, Cameroon)<sup>‡</sup>; Amina Barkat (Mohammed V University de Rabat, Morocco)<sup>‡</sup>; Habiba Ben Romdhane (University Tunis El Manar, Tunisia)<sup>‡</sup>; Queenie Chan (Imperial College London, UK)<sup>‡</sup>; [Nishi Chaturvedi \(University College London, UK\)](#)<sup>‡</sup>; Albertino Damasceno (Eduardo Mondlane University, Mozambique)<sup>‡</sup>; Hélène Delisle (University of Montreal, Canada)<sup>‡</sup>; Francis Delpuech (Institut de Recherche pour le Développement, France)<sup>‡</sup>; Kouamelan Doua (Ministère de la Santé et de la Lutte Contre le Sida, Côte d'Ivoire)<sup>‡</sup>; Eruke E Egbagbe (University of Benin, Nigeria)<sup>‡</sup>; Jalila El Ati (National Institute of Nutrition and Food Technology, Tunisia)<sup>‡</sup>; Paul Elliott (Imperial College London, UK)<sup>‡</sup>; Reina Engle-Stone (University of California Davis, USA)<sup>‡</sup>; Rajiv T Erasmus (University of Stellenbosch, South Africa)<sup>‡</sup>; Heba M Fouad (World Health Organization Regional Office for the Eastern Mediterranean, Egypt)<sup>‡</sup>; Dickman Gareta (University of KwaZulu-Natal, South Africa)<sup>‡</sup>; Oye Gureje (University of Ibadan, Nigeria)<sup>‡</sup>; Marleen Elisabeth Hendriks (Academic Medical Center of University of Amsterdam, The Netherlands)<sup>‡</sup>; Leila Houti (University of Oran 1, Algeria)<sup>‡</sup>; Mohsen M Ibrahim (Cairo University, Egypt)<sup>‡</sup>; Han CG Kemper (VU University Medical Center, The Netherlands)<sup>‡</sup>; Japhet Killewo (Muhimbili University of Health and Allied Sciences, Tanzania)<sup>‡</sup>; Sudhir Kowlessur (Ministry of Health and Quality of Life, Mauritius)<sup>‡</sup>; Herculina S Kruger (North-West University, South Africa)<sup>‡</sup>; Fatima Zahra Laamiri (Higher Institute of Nursing Professions and Technical Health Morocco, Morocco)<sup>‡</sup>; Youcef Laid (National Institute of Public Health of Algeria, Algeria)<sup>‡</sup>; Naomi S Levitt (University of Cape Town, South Africa)<sup>‡</sup>; Nuno Lunet (Universidade do Porto, Portugal)<sup>‡</sup>; Dianna J Magliano (Baker IDI Heart and Diabetes Institute, Australia)<sup>‡</sup>; [Bernard Maire \(Institut de Recherche pour le Développement, France\)](#)<sup>‡</sup>; Yves Martin-Prevel (Institut de Recherche pour le Développement, France)<sup>‡</sup>; Sounnia Mediene-Benchekor (University of Oran 1, Algeria)<sup>‡</sup>; Mostafa K Mohamed (Ain Shams University, Egypt, deceased)<sup>‡</sup>; Charles K Mondo (Mulago Hospital, Uganda)<sup>‡</sup>; Kotsedi Daniel K Monyeki (University of Limpopo, South Africa)<sup>‡</sup>; Aya Mostafa (Ain Shams University, Egypt)<sup>‡</sup>; [Martin Nankap \(Helen Keller International, Cameroon\)](#)<sup>‡</sup>; Ellis Owusu-Dabo (Kwame Nkrumah University of Science and Technology, Ghana)<sup>‡</sup>; Tobias F Rinke de Wit (PharmAccess Foundation, The Netherlands)<sup>‡</sup>; Olfa Saidi (University Tunis El Manar, Tunisia)<sup>‡</sup>;

Constance Schultsz (Academic Medical Center of University of Amsterdam, The Netherlands)<sup>‡</sup>; Aletta E Schutte (MRC North-West University, South Africa)<sup>‡</sup>; Idowu O Senbanjo (Lagos State University College of Medicine, Nigeria)<sup>‡</sup>; Jonathan E Shaw (Baker IDI Heart and Diabetes Institute, Australia)<sup>‡</sup>; Liam Smeeth (London School of Hygiene & Tropical Medicine, UK)<sup>‡</sup>; Eugène Sobngwi (University of Yaoundé 1, Cameroon)<sup>‡</sup>; Charles Sossa Jérôme (Institut Régional de Santé Publique, Benin)<sup>‡</sup>; Karien Stronks (University of Amsterdam, The Netherlands)<sup>‡</sup>; Frank Tanser (University of KwaZulu-Natal, South Africa)<sup>‡</sup>; Félicité Tchibindat (UNICEF, Cameroon)<sup>‡</sup>; Pierre Traissac (Institut de Recherche pour le Développement, France)<sup>‡</sup>; Lechaba Tshepo (Sefako Makgatho Health Science University, South Africa)<sup>‡</sup>; Fikru Tullu (Addis Ababa University, Ethiopia)<sup>‡</sup>; Flora AM Ukoli (Meharry Medical College, USA)<sup>‡</sup>; Bharathi Viswanathan (Ministry of Health, Seychelles)<sup>‡</sup>; Alisha N Wade (University of the Witwatersrand, South Africa)<sup>‡</sup>; Goodarz Danaei (Harvard TH Chan School of Public Health, USA); Gretchen A Stevens (World Health Organization, Switzerland); [Leanne M Riley \(World Health Organization, Switzerland\)](#); Majid Ezzati (Imperial College London, UK)<sup>†</sup>; Jean Claude N Mbanya (University of Yaoundé 1, Cameroon)<sup>†</sup>

## References

1. Lekoubou A, Echouffo-Tcheugui JB, Kengne AP. Epidemiology of neurodegenerative diseases in sub-Saharan Africa: a systematic review. *BMC Public Health* 2014; 14:653.
2. Yako YY, Guewo-Fokeng M, Balti EV, et al. Genetic risk of type 2 diabetes in populations of the African continent: A systematic review and meta-analyses. *Diabetes Res Clin Pract* 2016; 114:136-150.
3. Yako YY, Echouffo-Tcheugui JB, Balti EV, et al. Genetic association studies of obesity in Africa: a systematic review. *Obes Rev* 2015; 16:259-272.
4. World Health Organisation. *Global Report on Diabetes*. Geneva: WHO Press, 2016.
5. WHO Regional Office for Africa. *Diabetes prevention and control: A strategy for the WHO African Region*. AFR/RC57/7. Brazzaville: WHO African Region, 2007.
6. International Conference on Diabetes and Associated Diseases. *The Mauritius Call for Action*. Mauritius: World Health Organization, 2009.

7. East African Diabetes Study Group, Tanzania Ministry of Health, WHO African Region, World Diabetes Foundation. The 2016 Dar es Salaam Call to Action on Diabetes and Other Non-Communicable Diseases Dar es Salaam: WHO Regional Office for Africa 2016 [updated 25.12.2016; cited 2016 25.12]. Available from: <http://www.afro.who.int/en/tanzania/press-materials/item/8595-who-joins-partners-congress-in-the-fight-against-diabetes-and-non-communicable-diseases.html>.
8. World Health Organisation. Global action plan for the prevention and control of noncommunicable diseases 2013-2020. Geneva, Switzerland: WHO; 2013.
9. Hospedales CJ, Kengne AP, Legetic B, Marquizo AB. Noncommunicable diseases (NCDs): moving toward health promotion. In: Zimmerman RS, DiClemente RJ, Andrus JK, Hosein EN, editors. Introduction to Global Health Promotion: Wiley; 2016. p. 303-338.
10. NCD Risk Factor Collaboration. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* 2016; 387:1377-1396.
11. NCD Risk Factor Collaboration. Worldwide trends in diabetes since 1980: a pooled analysis of 751 population-based studies with 4.4 million participants. *Lancet* 2016; 387:1513-1530.
12. Manne-Goehler J, Atun R, Stokes A, et al. Diabetes diagnosis and care in sub-Saharan Africa: pooled analysis of individual data from 12 countries. *Lancet Diabetes Endocrinol* 2016; 4:903-912.
13. Werfalli M, Engel ME, Musekiwa A, Kengne AP, Levitt NS. The prevalence of type 2 diabetes among older people in Africa: a systematic review. *Lancet Diabetes Endocrinol* 2016; 4:72-84.
14. International Diabetes Federation. IDF Diabetes Atlas, 7th Edition. Brussels: International Diabetes Federation, 2015.
15. Danaei G, Finucane MM, Lu Y, et al. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet* 2011; 378:31-40.
16. Gale EA, Gillespie KM. Diabetes and gender. *Diabetologia* 2001; 44:3-15.

17. Wandell PE, Carlsson AC. Gender differences and time trends in incidence and prevalence of type 2 diabetes in Sweden--a model explaining the diabetes epidemic worldwide today? *Diabetes Res Clin Pract* 2014; 106:e90-92.
18. Tull E, Roseman J. Diabetes in African Americans. *Diabetes in America*. NIH Publications. 2nd Edition ed. Bethesda 1995. p. 613-630.
19. Goedecke JH, Mtintsilana A, Dlamini SN, Kengne AP. Type 2 diabetes mellitus in African women. *Diabetes Res Clin Pract* 2017; 123:87-96.
- ~~20. Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014; 384:766-781.~~
- ~~21~~20. United Nations. Political declaration of the high-level meeting of the general assembly on the prevention and control of non-communicable diseases. New York: United Nations, 2011.
- ~~22~~21. World Health Organisation. STEPwise approach to surveillance (STEPS) Geneva [cited 2013 15.07]. Available from: <http://www.who.int/chp/steps/en/>.
- ~~23~~22. Kengne AP, Erasmus RT, Levitt NS, Matsha TE. Alternative indices of glucose homeostasis as biochemical diagnostic tests for abnormal glucose tolerance in an African setting. *Prim Care Diabetes* 2017; 11:119-131.
- ~~24~~23. Kengne AP, Echouffo-Tcheugui J, Sobgnwi E, Mbanya JC. New Insights on Diabetes Mellitus and Obesity in Africa. Part 1: Prevalence, Pathogenesis and comorbidities. *Heart* 2013; 99:979-983.
- ~~25~~24. Kengne AP, June-Rose Mchiza Z, Amoah AG, Mbanya JC. Cardiovascular diseases and diabetes as economic and developmental challenges in Africa. *Prog Cardiovasc Dis* 2013; 56:302-313.
- ~~26~~25. Institute of Medicine. *Promoting Cardiovascular Health in the Developing World: A Critical Challenge to Achieve Global Health*. Washington, DC: The National Academies Press, 2010.
- ~~27~~26. Colagiuri R, Girgis S, Gomez M, Walker K, Colagiuri S, O'Dea K. Evidence based guideline for the primary prevention of type 2 diabetes. Canberra: NHMRC and Diabetes Australia, 2009.

[2827](#). Haregu TN, Setswe G, Elliott J, Oldenburg B. National Responses to HIV/AIDS and Non-Communicable Diseases in Developing Countries: Analysis of Strategic Parallels and Differences. *J Public Health Res* 2014; 3:99.

[2928](#). Haregu TN, Setswe G, Elliott J, Oldenburg B. Developing an action model for integration of health system response to HIV/AIDS and noncommunicable diseases (NCDs) in developing countries. *Glob J Health Sci* 2013; 6:9-22.

## **Legend for figures**

**Figure 1.** Number of data sources available for body mass index and diabetes by African region and year

**Figure 2.** Number of data sources available for body mass index and diabetes by country in Africa

**Figure 3.** Trends in age-standardised mean body mass index and age-standardised diabetes prevalence by sex and region in Africa

**Figure 4.** Scatterplot of male versus female age-standardised mean body mass index and age-standardised diabetes prevalence by African region in 2014

**Figure 5.** Age-standardised mean body mass index and age-standardised diabetes prevalence by sex and country in Africa in 2014

**Figure 6.** Age-standardised diabetes prevalence versus age-standardised mean body mass index in African men and women in 2014, by region and per-capita gross domestic product (GDP)