



Kent Academic Repository

Bennett, T., Howls, C.J., Nemes, G. and Olde Daalhuis, A.B. (2018) *Globally Exact Asymptotics for Integrals with Arbitrary Order Saddles*. SIAM Journal on Mathematical Analysis, 50 (2). pp. 2144-2177. ISSN 0036-1410.

Downloaded from

<https://kar.kent.ac.uk/68713/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1137/17M1154217>

This document version

Publisher pdf

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

GLOBALLY EXACT ASYMPTOTICS FOR INTEGRALS WITH ARBITRARY ORDER SADDLES*

T. BENNETT[†], C. J. HOWLS[‡], G. NEMES[§], AND A. B. OLDE DAALHUIS[§]

Abstract. We derive the first exact, rigorous but practical, globally valid remainder terms for asymptotic expansions about saddles and contour endpoints of arbitrary order degeneracy derived from the method of steepest descents. The exact remainder terms lead naturally to sharper novel asymptotic bounds for truncated expansions that are a significant improvement over the previous best existing bounds for quadratic saddles derived two decades ago. We also develop a comprehensive hyperasymptotic theory, whereby the remainder terms are iteratively reexpanded about adjacent saddle points to achieve better-than-exponential accuracy. By necessity of the degeneracy, the form of the hyperasymptotic expansions is more complicated than in the case of quadratic endpoints and saddles and requires generalizations of the hyperterminants derived in those cases. However, we provide efficient methods to evaluate them, and we remove all possible ambiguities in their definition. We illustrate this approach for three different examples, providing all the necessary information for the practical implementation of the method.

Key words. integral asymptotics, asymptotic expansions, hyperasymptotics, error bounds, saddle points

AMS subject classifications. 41A60, 41A80, 58K05

DOI. 10.1137/17M1154217

1. Introduction. From catastrophe theory it is well known that integrals with saddle points may be used to compactly encapsulate the local behavior of linear wavefields near the underlying organizing caustics; see, for example, [44, 4]. The saddle points correspond to rays of the underpinning ODEs or PDEs. Their coalescence corresponds to tangencies of the rays at the caustics, leading to nearby peaks in the wave amplitude. On the caustics, the coalesced saddle points are degenerate. The local analytical behavior on the caustic may be derived from an asymptotic expansion about the degenerate saddle [18, Chap. 36]. An analytical understanding of the asymptotic expansions involving degenerate saddles is thus essential to an examination of the wavefield behavior on caustics. A modern approach to this includes the derivation of globally exact remainders, sharp error bounds, and the exponential improvement of the expansions to take into account the contributions of terms beyond all orders.

Recent work in quantum field and string theories, e.g., [20, 14, 1, 2], has led to a major increase in interest in such resurgent approaches in the context of integral asymptotics. One of the reasons to study the higher orders of expansions in quantum field theory is that they can reveal information as to the location of remote critical points, corresponding to physical quantities that can give rise to nonperturbative

*Received by the editors October 27, 2017; accepted for publication (in revised form) February 7, 2018; published electronically April 17, 2018.

<http://www.siam.org/journals/sima/50-2/M115421.html>

Funding: The first author was sponsored by an EPSRC studentship. The third and fourth authors were supported by a research grant (GRANT11863412/ 70NANB15H221) from the National Institute of Standards and Technology.

[†]SMSAS, University of Kent, Sibson Building, Parkwood Road, Canterbury CT2 7FS, UK (T.B.Bennett@kent.ac.uk).

[‡]Mathematical Sciences, University of Southampton, Highfield, Southampton SO17 1BJ, UK (C.J.Howls@maths.soton.ac.uk).

[§]School of Mathematics, University of Edinburgh, James Clerk Maxwell Building, The King's Buildings, Edinburgh EH9 3FD, UK (Gergo.Nemes@ed.ac.uk, A.OldeDaalhuis@ed.ac.uk).

effects. A notable recent success is that high order calculations can be used to uncover previously unknown functional relationships between perturbative and nonperturbative effects within quantum eigenvalue expansions [19].

The first globally exact remainders for asymptotic expansions of integrals possessing simple saddle points were derived by Berry and Howls [7]. The remainder terms were expressed in terms of self-similar integrals over doubly infinite contours passing through a set of adjacent simple saddles. Boyd [12] provided a rigorous justification of the exact remainder terms, together with significantly improved error bounds.

The remainder terms automatically incorporated and precisely accounted for the Stokes phenomenon [46], whereby exponentially subdominant asymptotic contributions are switched on as asymptotics or other parametric changes cause the contour of integration to deform to pass through the adjacent saddles. The Stokes phenomenon occurs across subsets in parameter space called Stokes lines.

Reexpansion of the exact remainder term about the adjacent saddles, using their own exact remainder terms, led to a hyperasymptotic expansion, which delivered better-than-exponential numerical accuracy.

Subsequent work extended globally exact remainder terms and hyperasymptotic analysis to integrals over contours with finite endpoints [23] and multiple integrals [24], [16]. Parallel approaches to differential equations using Cauchy–Heine and Borel transforms were taken by Olde Daalhuis and Olver [41], [37]. This resulted in efficient methods for computation of the universal hyperterminants [38]. The efficient computation of hyperterminants not only made hyperasymptotic expansions numerically feasible but, more importantly, in the absence of the geometric information present in single dimensional integral calculations, allowed them to be used to calculate the Stokes constants that are required in an exponentially accurate asymptotic calculation involving, for example, the solution satisfying given boundary data.

However, the general case of globally exact remainder terms and hyperasymptotic expansions of a single-dimensional integral possessing a set of arbitrary order degenerate saddle points has not yet been considered. The purpose of this paper is to fill this surprising gap.

Hence, in this paper, we provide the first comprehensive globally exact asymptotic theory for integrals with analytic integrands involving finite numbers of arbitrarily degenerate saddle points. It incorporates the special case of Berry and Howls [7] and Howls [23]. However, the complexity of the situation uncovers several new features that were not present in the simple saddle case.

First, the nature of the steepest paths emerging from degenerate saddles gives multiple choices as to which contours might be integrated over, or which might contribute to the remainder term. It is necessary to adopt a stricter convention regarding the choice of steepest paths to clarify the precise nature of the contributions to the remainder and hyperasymptotic expansions.

Second, the degenerate nature requires us to explore additional Riemann sheets associated to the local mappings about the saddle points. This gives rise to additional complex phases, not obviously present in the simple saddle case, that must be taken into account depending on the relative geometrical disposition of the contours.

Third, we provide sharp, rigorous bounds for the remainder terms in the Poincaré asymptotic expansions of integrals with arbitrary critical points. In particular, we improve the results of Boyd [12], who considered integrals with only simple saddles. Our bounds are sharper and have larger regions of validity.

Fourth, the hyperasymptotic tree structure that underpins the exponential improvements in accuracy is *prima facie* more complicated. At the first reexpansion of

a remainder term, for each adjacent degenerate saddle there are two contributions arising from the choice of contour over which the remainder may be taken. At the second reexpansion, each of these two contributions may give rise to another two, and so on. Hence, while the role of the adjacency of saddles remains the same, the numbers of terms required at each hyperasymptotic level increase twofold for each degenerate saddle at each level. Fortunately these terms may be related, and so the propagation of computational complexity is controllable.

Fifth, the hyperterminants in the expansion are more complicated than those in [7], [33], [37], or [38]. However, we provide efficient methods to evaluate them.

Sixth, the results of this integral analysis reveal new insights into the asymptotic expansions of higher order differential equations.

There have been several near misses at a globally exact remainder term for degenerate saddles arising from single dimensional integrals.

Ideas similar to those employed by Berry and Howls were used earlier by Meijer. In a series of papers [27], [31], [32] he derived exact remainder terms and realistic error bounds for specific special functions, namely, Bessel, Hankel, and Anger–Weber-type functions. Nevertheless, he missed the extra step that would have led him to more general remainder terms of [7].

Dingle [17], whose pioneering view of resurgence underpins most of this work, considered expansions around cubic saddle points and gave formal expressions for the higher order terms. However, he did not provide exact remainder terms or consequent (rigorous) error estimates.

Berry and Howls [8], [9] considered the cases of exponentially improved uniform expansions of single dimensional integrals as saddle points coalesced. The analysis [8] focused on the form of the late terms in the more complicated uniform expansions. They [9] provided an approximation to the exact remainder term between a simple and an adjacent cluster of saddles illustrating the persistence of the error function smoothing of the Stokes phenomenon [6] as the Stokes line was crossed. Neither of these works gave globally exact expressions for remainder terms involving coalesced, degenerate saddles.

Olde Daalhuis [39] considered a Borel plane treatment of uniform expansions but did not extend the work to include arbitrary degenerate saddles.

Breen [13] briefly considered the situation of degenerate saddles. The work restricted attention to cubic saddles and, like all the above work, did not provide rigorous error bounds or develop a hyperasymptotic expansion.

Other notable work dealing with exponential asymptotics includes [10], [11], [22], [25], [26], [45], and [48].

It should be stressed that the purpose of a hyperasymptotic approach is not *per se* to calculate functions to high degrees of numerical accuracy: there are alternative computational methods. Rather, hyperasymptotics is an analytical tool to incorporate exponentially small contributions into asymptotic approximations, so as to widen the domain of validity, to understand better the underpinning singularity structures, and to compute invariants of the system such as Stokes constants whose values are often assumed or left as unknowns by other methods (see, for example, [21]).

The idea for this paper emerged from the recent complementary and independent thesis work of [3], [35], which gave rise to the current collaboration. This collaboration has resulted in the present work which incorporates not only a hyperasymptotic theory for both expansions arising from nondegenerate and degenerate saddle points but also significantly improved rigorous and sharp error bounds for the progenitor asymptotic expansions.

The structure of the paper is as follows.

In section 2, we introduce arbitrary finite integer degenerate saddle points. In section 3, we derive the exact remainder term for an expansion about a semi-infinite steepest descent contour emerging from a degenerate saddle and running to a valley at infinity. The remainder term is expressed as a sum of terms of contributions from other, adjacent saddle points of the integrand. Each of these contributions is formed from the difference of two integrals over certain semi-infinite steepest descent contours emerging from the adjacent saddles.

In section 4, we iterate these exact remainder terms to develop a hyperasymptotic expansion. We introduce novel hyperterminants (which simplify to those of Olde Daalhuis [38] when the saddles are nondegenerate).

In section 5, we provide explicit rigorous error bounds for the zeroth hyperasymptotic level. These novel bounds are sharper than those derived by Boyd [12].

In section 6, we illustrate the degenerate hyperasymptotic method with an application to an integral related to the Pearcey function, evaluated on its cusp caustic. The example involves a simple and doubly degenerate saddle. In section 7, we provide an illustration of the extra complexities of a hyperasymptotic treatment of degeneracies with an application to an integral possessing triply and quintuply degenerate saddle points. In this example, we also illustrate the increased size of the remainder near a Stokes line as predicted in section 5. In section 8, we give an example of how it is possible to make an algebraic (rather than geometric) determination of the saddles that contribute to the exact remainder terms in a swallowtail-type integral through a hyperasymptotic examination of the late terms in the saddle point expansion.

In section 9, we conclude with a discussion on the application of the results of this paper to the (hyper-) asymptotic expansions of higher order differential equations.

2. Definitions and assumptions. Let ω_j be a positive integer, with $j = 1, 2, \dots$ an integer index. Consider a function $f(t)$, analytic in a domain of the complex plane. The point $t^{(j)}$ is called a critical point of order $\omega_j - 1$ of $f(t)$ if

$$f^{(p)}(t^{(j)}) = 0 \quad \text{but} \quad f^{(\omega_j)}(t^{(j)}) \neq 0 \quad \text{for all } p = 1, \dots, \omega_j - 1.$$

When $\omega_j = 1, 2, > 2$, $t^{(j)}$ is, respectively, a linear endpoint, a simple saddle point, and a degenerate saddle point. For analytic $f(t)$, the saddle points are then all isolated. Henceforth we denote the value of $f(t)$ at $t = t^{(j)}$ by f_j .

We shall derive the steepest descent expansion, together with its exact remainder term, of integrals of the type

$$(1) \quad I^{(n)}(z; \alpha_n) = \int_{\mathcal{P}^{(n)}} e^{-zf(t)} g(t) dt, \quad z = |z|e^{i\theta}, \quad |z| \rightarrow \infty,$$

where $\mathcal{P}^{(n)} = \mathcal{P}^{(n)}(\theta; \alpha_n)$ is one of the ω_n paths of steepest descent emanating from the $(\omega_n - 1)$ st-order critical point $t^{(n)}$ of $f(t)$ and passing to infinity in a valley of $\text{Re}[-e^{i\theta}(f(t) - f_n)]$.

Suppose we use the notation of $(\omega_n \rightarrow \omega_{\mathbf{m}})$ to indicate the remainder term that rises from an asymptotic expansion about a endpoint/saddle point n of order ω_n in terms of the adjacent (in a sense to be defined later) set of saddles $\mathbf{m} = \{m_1, m_2, m_3, \dots\}$, of orders corresponding to the values $\omega_{\mathbf{m}} = \{\omega_{m_1}, \omega_{m_2}, \dots\}$. Thus Berry and Howls [7] dealt with $(\omega_n \rightarrow \omega_{\mathbf{m}}) = (2 \rightarrow \mathbf{2})$ for doubly infinite contours. Howls [23] dealt with $(1 \rightarrow \mathbf{2})$ and the $(2 \rightarrow \mathbf{2})$. Our goal here is to derive the exact remainder terms for arbitrary integers $(\omega_n \rightarrow \omega_{\mathbf{m}})$.




FIG. 1. The ω_n paths of steepest descent emanating from the $(\omega_n - 1)$ st-order critical point $t^{(n)}$ of $f(t)$.

On the steepest path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ emerging from $t^{(n)}$, we have

$$(2) \quad \arg [e^{i\theta}(f(t) - f_n)] = 2\pi\alpha_n$$

for a suitable integer α_n (see Figure 1).

The local behavior of $f(t)$ at the critical point $t^{(n)}$ of order $\omega_j - 1$ is given by

$$(3) \quad f(t) - f_n = \frac{f^{(\omega_n)}(t^{(n)})}{\omega_n!} (t - t^{(n)})^{\omega_n} + \mathcal{O}(|t - t^{(n)}|^{\omega_n+1}).$$

From (2) and (3), we hence find that

$$(4) \quad \alpha_n = \frac{\theta + \arg(f^{(\omega_n)}(t^{(n)})) + \omega_n\varphi}{2\pi},$$

where $-\pi < \arg(f^{(\omega_n)}(t^{(n)})) \leq \pi$, and φ ($-\pi < \varphi \leq \pi$) is the angle of the slope of $\mathcal{P}^{(n)}(\theta; \alpha_n)$ at $t^{(n)}$, i.e., $\lim(\arg(t - t^{(n)}))$ as $t \rightarrow t^{(n)}$ along $\mathcal{P}^{(n)}(\theta; \alpha_n)$.

The functions $f(t)$ and $g(t)$ are assumed to be analytic in the closure of a domain $\Delta^{(n)}$. We suppose further that $|f(t)| \rightarrow \infty$ as $t \rightarrow \infty$ in $\Delta^{(n)}$, and $f(t)$ has several other saddle points in the complex t -plane at $t = t^{(j)}$ labeled by $j \in \mathbb{N}$.

The domain $\Delta^{(n)}$ is defined by considering all the steepest descent paths for different values of θ , which emerge from the critical point $t^{(n)}$. In general these paths can end either at infinity or at a singularity of $f(t)$. We assume that all of them end at infinity. Since there are no branch points of $f(t)$ along these paths, any point in the t -plane either cannot be reached by any path of steepest descent issuing from $t^{(n)}$ or else can be reached by only one. A continuity argument shows that the set of all the points which can be reached by a steepest descent path from $t^{(n)}$ forms the closure of a domain in the t -plane. It is this domain which we denote by $\Delta^{(n)}$; see, for example, Figure 2.

Instead of considering the raw integral (1), it will be convenient to consider its slowly varying part, defined by

$$(5) \quad T^{(n)}(z; \alpha_n) := \omega_n z^{1/\omega_n} e^{zf_n} I^{(n)}(z; \alpha_n) = \omega_n z^{1/\omega_n} \int_{\mathcal{P}^{(n)}} e^{-z(f(t)-f_n)} g(t) dt.$$




FIG. 2. Contours used in the derivation of the exact remainder terms. (a) The contour $\Gamma^{(n)}(\theta)$ relative to the integration contour $\mathcal{P}^{(n)}(\theta; \alpha_n)$ as used in (17). (b) A schematic representation of the saddle points $t^{(m_j)}$ that are adjacent to $t^{(n)}$ and the adjacent contours $\mathcal{P}^{(m_j)}$ emanating from them in (20), together with the domain $\Delta^{(n)}$.

The ω_n th root is defined to be positive on the positive real line and is defined by analytic continuation elsewhere. We call $T^{(n)}(z; \alpha_n)$ the slowly varying part because it is $\mathcal{O}(1)$ as $z \rightarrow \infty$ (cf. (9)).

It is convenient to introduce the following notation for the special double integrals and their coefficients in the asymptotic expansions:

$$(6) \quad \begin{aligned} \mathbf{T}^{(m)}(u; \alpha_n) &= T^{(m)}(u; \alpha_n) - T^{(m)}(u; \alpha_n + 1), \\ \mathbf{T}_r^{(m)}(\alpha_n) &= T_r^{(m)}(\alpha_n) - T_r^{(m)}(\alpha_n + 1). \end{aligned}$$

The path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ passes through certain other saddle points $t^{(m)}$ when $\theta = \theta_{nm}^{[1]}, \theta_{nm}^{[2]}, \theta_{nm}^{[3]}, \dots$, with $\theta_{nm}^{[j]} = \theta_{nm}^{[k]} \bmod 2\pi\omega_n$. Such saddle points are defined as being “adjacent” to $t^{(n)}$.

Initially we chose the value of θ so that the steepest descent path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ in (1) does not encounter any of the saddle points of $f(t)$ other than $t^{(n)}$. We define

$$\theta_{nm}^+ := \min \left\{ \theta_{nm}^{[j]} : j \geq 1, \theta < \theta_{nm}^{[j]} \right\} \quad \text{and} \quad \theta_{nm}^- := \max \left\{ \theta_{nm}^{[j]} : j \geq 1, \theta_{nm}^{[j]} < \theta \right\}.$$

Note that $\theta_{nm}^+ = \theta_{nm}^- + 2\pi\omega_n$. Thus, in particular, θ is restricted to an interval

$$(7) \quad \theta_{nm_1}^- < \theta < \theta_{nm_2}^+,$$

where $\theta_{nm_1}^- := \max_m \theta_{nm}^-$ and $\theta_{nm_2}^+ := \min_m \theta_{nm}^+$. We shall suppose that $f(t)$ and $g(t)$ grow sufficiently rapidly at infinity so that the integral (1) converges for all values of θ in the interval (7).

Let $\Gamma^{(n)} = \Gamma^{(n)}(\theta)$ be an infinite contour that encircles the path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ in the positive direction within $\Delta^{(n)}$ (see Figure 2(a)). This contour $\Gamma^{(n)}(\theta)$ is now deformed by expanding it onto the boundary of $\Delta^{(n)}$. We assume that the set of saddle points which are adjacent to $t^{(n)}$ is nonempty and finite. Under this assumption, it is shown in Appendix C that the boundary of $\Delta^{(n)}$ can be written as a union of contours $\bigcup_m \mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+) \cup -\mathcal{P}^{(m)}(\theta_{nm}^-, \alpha_{nm}^-)$, where $\mathcal{P}^{(m)}(\theta_{nm}^\pm, \alpha_{nm}^\pm)$ are steepest descent paths emerging from the adjacent saddle $t^{(m)}$ (see Figure 2(b)). These paths are called the adjacent contours. The integers α_{nm}^\pm are computed analogously to α_n (cf. (4)) as

$$(8) \quad \alpha_{nm}^\pm = \frac{\theta_{nm}^\pm + \arg(f^{(\omega_m)}(t^{(m)})) + \omega_m \varphi^\pm}{2\pi},$$

where $-\pi < \arg(f^{(\omega_m)}(t^{(m)})) \leq \pi$, and φ^\pm ($-\pi < \varphi^\pm \leq \pi$) is the angle of the slope of $\mathcal{P}^{(m)}(\theta_{nm}^\pm, \alpha_{nm}^\pm)$ at the $(\omega_m - 1)$ st-order saddle point $t^{(m)}$ to the positive real axis. We assume initially that each adjacent contour contains only one saddle point.¹ The other steepest descent paths from $t^{(m)}$ are always external to the domain $\Delta^{(n)}$.

Finally, we introduce the so-called singulants \mathcal{F}_{nm}^\pm (originally defined by Dingle [17, pp. 147–149]) via

$$\mathcal{F}_{nm}^\pm := |f_m - f_n| e^{i \arg \mathcal{F}_{nm}^\pm}, \quad \arg \mathcal{F}_{nm}^\pm = -\theta_{nm}^\pm + 2\pi\alpha_n.$$

3. Derivation of exact remainder term. In this section we will show that

$$(9) \quad T^{(n)}(z; \alpha_n) = \sum_{r=0}^{N-1} \frac{T_r^{(n)}(\alpha_n)}{z^{r/\omega_n}} + R_N^{(n)}(z; \alpha_n),$$

where

$$(10) \quad T_r^{(n)}(\alpha_n) = e^{\frac{2\pi i \alpha_n (r+1)}{\omega_n}} \frac{\Gamma\left(\frac{r+1}{\omega_n}\right)}{2\pi i} \oint_{t^{(n)}} \frac{g(t)}{(f(t) - f_n)^{(r+1)/\omega_n}} dt$$

$$= e^{\frac{2\pi i \alpha_n (r+1)}{\omega_n}} \left(\frac{\omega_n!}{f^{(\omega_n)}(t^{(n)})}\right)^{(r+1)/\omega_n} \frac{\Gamma\left(\frac{r+1}{\omega_n}\right)}{\Gamma(r+1)}$$

$$(11) \quad \times \left[\frac{d^r}{dt^r} \left(g(t) \left(\frac{f^{(\omega_n)}(t^{(n)})}{\omega_n!} \frac{(t - t^{(n)})^{\omega_n}}{f(t) - f_n} \right)^{(r+1)/\omega_n} \right) \right]_{t=t^{(n)}},$$

and for the remainder we have

$$(12) \quad R_N^{(n)}(z; \alpha_n) = \sum_{m(n)} \frac{z^{(1-N)/\omega_n}}{2\pi i \omega_m} \int_0^\infty e^{i\theta_{nm}^+} \frac{e^{-\mathcal{F}_{nm}^+ u} u^{\frac{N+1}{\omega_n} - \frac{1}{\omega_m} - 1}}{z^{1/\omega_n} - u^{1/\omega_n}} \mathbf{T}^{(m)}(u; \alpha_{nm}^+) du,$$

in which $m(n)$ means that we sum over all saddles that are adjacent to n . The result (12) for the exact remainder term of the asymptotic expansion around the degenerate saddle $t^{(n)}$, expressed in terms of the adjacent (other degenerate) saddles $t^{(m)}$, is one of the main results of this paper.

If we omit the remainder term $R_N^{(n)}(z; \alpha_n)$ in (9) and formally extend the sum to infinity, the result becomes the asymptotic expansion of an integral with $(\omega_n - 1)$ st-order endpoint (cf. [42, eqn. (1.2.16), p. 12]). A representation equivalent to (10) was given, for example, by Copson [15, p. 69]. The expression (11) is a special case of Perron’s formula (see, e.g., [34]).

In the examples below we use (11) to compute conveniently and analytically the exact coefficients. However, we remark that (10) may be combined with the trapezoidal rule evaluated at periodic points on the loop contour about $t^{(n)}$ (see, for example, [47]) to give an efficient approximation for the coefficients as

$$(13) \quad T_r^{(n)}(\alpha_n) \approx e^{\frac{2\pi i \alpha_n (r+1)}{\omega_n}} \frac{\Gamma\left(\frac{r+1}{\omega_n}\right)}{2M} \sum_{m=0}^{2M-1} \frac{g(t_m)}{w_m^r} \left(\frac{(t_m - t^{(n)})^{\omega_n}}{f(t_m) - f_n} \right)^{(r+1)/\omega_n},$$

¹This condition may be relaxed by extending the definition of integrals of the form (5) to include the limiting case when the steepest descent path connects to other saddle points. Also, a limiting case, such as (28), has to be used for the generalized hyperterminants in the corresponding re-expansions.

in which $t_m = t^{(n)} + w_m$ and $w_m = \rho e^{\pi i m/M}$. Typically this approximation converges exponentially fast with M . Note that in hyperasymptotics n can be large and so we would need to take at least $M > n$.

For the proof of the results above we will obtain new integral representations for $T^{(n)}(z; \alpha_n)$ via several changes of variables. The local behavior (3) of $f(t)$ at the critical point $t^{(n)}$ suggests the parameterization

$$(14) \quad s^{\omega_n} = z(f(t) - f_n)$$

of the integrand in (5) along $\mathcal{P}^{(n)}(\theta; \alpha_n)$. Substitution of (14) in (5) yields

$$(15) \quad \begin{aligned} T^{(n)}(z; \alpha_n) &= \omega_n z^{1/\omega_n} \int_0^{\infty e^{\frac{2\pi i \alpha_n}{\omega_n}}} e^{-s^{\omega_n}} g(t) \frac{dt}{ds} ds \\ &= \omega_n \int_0^{\infty e^{\frac{2\pi i \alpha_n}{\omega_n}}} e^{-s^{\omega_n}} \frac{\omega_n s^{\omega_n - 1}}{z^{1 - 1/\omega_n}} \frac{g(t(s/z^{1/\omega_n}))}{f'(t(s/z^{1/\omega_n}))} ds, \end{aligned}$$

where $t = t(s/z^{1/\omega_n})$ is the unique solution of (14) with $t(s/z^{1/\omega_n}) \in \mathcal{P}^{(n)}(\theta; \alpha_n)$. Since the contour $\mathcal{P}^{(n)}(\theta; \alpha_n)$ does not pass through any of the saddle points of $f(t)$ other than $t^{(n)}$, the quantity

$$(16) \quad \frac{\omega_n s^{\omega_n - 1}}{z^{1 - 1/\omega_n}} \frac{g(t(s/z^{1/\omega_n}))}{f'(t(s/z^{1/\omega_n}))} = \frac{\omega_n (f(t(s/z^{1/\omega_n})) - f(t^{(n)}))^{1 - 1/\omega_n}}{f'(t(s/z^{1/\omega_n}))} g(t(s/z^{1/\omega_n}))$$

is an analytic function of t in a neighborhood of $\mathcal{P}^{(n)}(\theta; \alpha_n)$. (We examine the analyticity of the factor $(f(t) - f_n)^{1/\omega_n}$ in $\Delta^{(n)}$, after (18) below.) Hence, according to the residue theorem, the right-hand side of (16) is²

$$\text{Res}_{t=t(s/z^{1/\omega_n})} \frac{g(t)}{(f(t) - f_n)^{1/\omega_n} - s/z^{1/\omega_n}} = \frac{1}{2\pi i} \oint_{\Gamma^{(n)}(t(s/z^{1/\omega_n}))} \frac{g(t)}{(f(t) - f_n)^{1/\omega_n} - s/z^{1/\omega_n}} dt.$$

Substituting this expression into (15) leads to an alternative representation for the integral $T^{(n)}(z; \alpha_n)$ of the form

$$(17) \quad T^{(n)}(z; \alpha_n) = \int_0^{\infty e^{\frac{2\pi i \alpha_n}{\omega_n}}} e^{-s^{\omega_n}} \frac{\omega_n}{2\pi i} \oint_{\Gamma^{(n)}} \frac{g(t)}{(f(t) - f_n)^{1/\omega_n} - s/z^{1/\omega_n}} dt ds.$$

The infinite contour $\Gamma^{(n)} = \Gamma^{(n)}(\theta)$ encircles the path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ in the positive direction within $\Delta^{(n)}$ (see Figure 2(a)). This integral will exist provided that $g(t)/f^{1/\omega_n}(t)$ decays sufficiently rapidly at infinity in $\Delta^{(n)}$. Otherwise, we can define $\Gamma^{(n)}(\theta)$ as a finite loop contour surrounding $t(s/z^{1/\omega_n})$ and consider the limit

$$(18) \quad \lim_{S \rightarrow \infty} \int_0^{Se^{\frac{2\pi i \alpha_n}{\omega_n}}} e^{-s^{\omega_n}} \frac{\omega_n}{2\pi i} \oint_{\Gamma^{(n)}} \frac{g(t)}{(f(t) - f_n)^{1/\omega_n} - s/z^{1/\omega_n}} dt ds.$$

The factor $(f(t) - f_n)^{1/\omega_n}$ in (17) is carefully defined in the domain $\Delta^{(n)}$ as follows. First, we observe that $f(t) - f_n$ has an ω_n th-order zero at $t = t^{(n)}$ and is nonzero

²If $P(t)$ and $Q(t)$ are analytic in a neighborhood of t_0 with $P(t_0) = 0$ and $P'(t_0) \neq 0$, then $Q(t_0)/P'(t_0) = \text{Res}_{t=t_0} Q(t)/P(t)$.

elsewhere in $\Delta^{(n)}$ (because any point in $\Delta^{(n)}$, different from $t^{(n)}$, can be reached from $t^{(n)}$ by a path of descent). Second, $\mathcal{P}^{(n)}(\theta; \alpha_n)$ is a periodic function of θ with (least) period $2\pi\omega_n$. Hence, we may define the ω_n th root so that $(f(t) - f_n)^{1/\omega_n}$ is a single-valued analytic function of t in $\Delta^{(n)}$. The correct choice of the branch of $(f(t) - f_n)^{1/\omega_n}$ is determined by the requirement that $\arg s = 2\pi\alpha_n/\omega_n$ on $\mathcal{P}^{(n)}(\theta; \alpha_n)$, which can be fulfilled by setting $\arg[(f(t) - f_n)^{1/\omega_n}] = (2\pi\alpha_n - \theta)/\omega_n$ for $t \in \mathcal{P}^{(n)}(\theta; \alpha_n)$. With any other definition of $(f(t) - f_n)^{1/\omega_n}$, the representation (17) would be invalid.

Now, we employ the finite expression for nonnegative integer N

$$\frac{1}{1-x} = \sum_{r=0}^{N-1} x^r + \frac{x^N}{1-x}, \quad x \neq 1,$$

to expand the denominator in (17) in powers of $s/[z(f(t) - f_n)]^{1/\omega_n}$. We thus obtain

$$T^{(n)}(z; \alpha_n) = \sum_{r=0}^{N-1} \frac{1}{z^{r/\omega_n}} \int_0^{\infty e^{\frac{2\pi i \alpha_n}{\omega_n}}} e^{-s^{\omega_n}} s^r \frac{\omega_n}{2\pi i} \oint_{\Gamma^{(n)}} \frac{g(t)}{(f(t) - f_n)^{(r+1)/\omega_n}} dt ds + R_N^{(n)}(z; \alpha_n)$$

with

$$(19) \quad R_N^{(n)}(z; \alpha_n) = \frac{\omega_n}{2\pi i z^{N/\omega_n}} \int_0^{\infty e^{\frac{2\pi i \alpha_n}{\omega_n}}} e^{-s^{\omega_n}} s^N \times \oint_{\Gamma^{(n)}} \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} \frac{dt}{1 - \frac{s}{(z(f(t) - f_n))^{1/\omega_n}}} ds.$$

Again, a limiting process is used in (19) if necessary. Throughout this work, if not stated otherwise, empty sums are taken to be zero.

For each term in the finite sum, the contour $\Gamma^{(n)}(\theta)$ can be shrunk into a small positively oriented circle with center $t^{(n)}$ and radius ρ , and we arrive at (9), where the coefficients are given by (10) and (11).

By expanding $\Gamma^{(n)}(\theta)$ to the boundary of $\Delta^{(n)}$ (see section 2), we obtain

$$(20) \quad R_N^{(n)}(z; \alpha_n) = \frac{\omega_n}{2\pi i z^{N/\omega_n}} \sum_{m(n)} \int_0^{\infty e^{\frac{2\pi i \alpha_n}{\omega_n}}} e^{-s^{\omega_n}} s^N \times \left(\int_{\mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+)} \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} \frac{dt}{1 - \frac{s}{(z(f(t) - f_n))^{1/\omega_n}}} - \int_{\mathcal{P}^{(m)}(\theta_{nm}^-, \alpha_{nm}^-)} \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} \frac{dt}{1 - \frac{s}{(z(f(t) - f_n))^{1/\omega_n}}} \right) ds.$$

The expansion process is justified provided that (i) $f(t)$ and $g(t)$ are analytic in the domain $\Delta^{(n)}$, (ii) the quantity $g(t)/f^{(N+1)/\omega_n}(t)$ decays sufficiently rapidly at infinity in $\Delta^{(n)}$, and (iii) there are no zeros of the denominator $1 - s/[z(f(t) - f_n)]^{1/\omega_n}$ within the region R through which the loop $\Gamma^{(n)}(\theta)$ is deformed.

The first condition is already satisfied by prior assumption. The second condition is met by requiring that $g(t)/f^{(N+1)/\omega_n}(t) = o(1/|t|)$ as $t \rightarrow \infty$ in $\Delta^{(n)}$ which

we shall assume to be the case. The third condition is satisfied according to the following argument. The zeros of the denominator are those points of the t -plane for which $\arg[e^{i\theta}(f(t) - f_n)] = 2\pi\alpha_n$, in particular the points of the path $\mathcal{P}^{(n)}(\theta; \alpha_n)$. Furthermore, no components of the set defined by the equation $\arg[e^{i\theta}(f(t) - f_n)] = 2\pi\alpha_n$ other than $\mathcal{P}^{(n)}(\theta; \alpha_n)$ can lie within $\Delta^{(n)}$, otherwise $f(t)$ would have branch points along those components. By observing that $\mathcal{P}^{(n)}(\theta; \alpha_n)$ is different for different values of $\theta \bmod 2\pi\omega_n$, we see that the locus of the zeros of the denominator $1 - s/[z(f(t) - f_n)]^{1/\omega_n}$ inside $\Delta^{(n)}$ is precisely the contour $\mathcal{P}^{(n)}(\theta; \alpha_n)$, which is wholly contained within $\Gamma^{(n)}(\theta)$ and so these zeros are external to R .

We now consider the convergence of the double integrals in (20) further. To do this, we change variables from t to v by

$$(21) \quad f(t) - f_n = ve^{(-\theta_{nm}^\pm + 2\pi\alpha_n)i},$$

where $v \geq |\mathcal{F}_{nm}^\pm|$. Since $e^{(\theta_{nm}^\pm - 2\pi\alpha_n)i}(f(t) - f_n)$ is a monotonic function of t on the contour $\mathcal{P}^{(m)}(\theta_{nm}^\pm, \alpha_{nm}^\pm)$, corresponding to each value of v , there is a value of t , say, $t_\pm(v)$, that satisfies (21). The assumption (7) implies that the factor $[1 - s/[z(f(t) - f_n)]^{1/\omega_n}]^{-1}$ in (20) is bounded above by a constant. Hence, the convergence of the double integrals in (20) will be ensured provided the real double integrals

$$\int_0^\infty \int_{|\mathcal{F}_{nm}^\pm|}^\infty \frac{e^{-|s|^{\omega_n}} |s|^N}{v^{(N+1)/\omega_n}} \left| \frac{g(t_\pm(v))}{f'(t_\pm(v))} \right| dv ds$$

exist. In turn, these real double integrals will exist if and only if the single integrals

$$(22) \quad \int_{|\mathcal{F}_{nm}^\pm|}^\infty \frac{1}{v^{(N+1)/\omega_n}} \left| \frac{g(t_\pm(v))}{f'(t_\pm(v))} \right| dv$$

exist. Henceforth, we assume that the integrals in (22) exist for each of the adjacent contours.

On each of the contours $\mathcal{P}^{(m)}(\theta_{nm}^\pm, \alpha_{nm}^\pm)$ in (20), we perform the change of variable from s and t to u and t via

$$s^{\omega_n} = u(f(t) - f_n) = \mathcal{F}_{nm}^\pm u + u(f(t) - f_m)$$

to obtain

$$(23) \quad R_N^{(n)}(z; \alpha_n) = \sum_{m(n)} \frac{z^{(1-N)/\omega_n}}{2\pi i} \times \left(\int_0^{\infty e^{i\theta_{nm}^+}} \frac{e^{-\mathcal{F}_{nm}^+ u} u^{\frac{N+1}{\omega_n} - 1}}{z^{1/\omega_n} - u^{1/\omega_n}} \int_{\mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+)} e^{-u(f(t)-f_m)} g(t) dt du \right. \\ \left. - \int_0^{\infty e^{i\theta_{nm}^-}} \frac{e^{-\mathcal{F}_{nm}^- u} u^{\frac{N+1}{\omega_n} - 1}}{z^{1/\omega_n} - u^{1/\omega_n}} \int_{\mathcal{P}^{(m)}(\theta_{nm}^-, \alpha_{nm}^-)} e^{-u(f(t)-f_m)} g(t) dt du \right).$$

This change of variable is permitted because the infinite double integrals in (20) are assumed to be absolutely convergent, which is a consequence of the requirement that the integrals (22) exist. Hence the exact remainder of the expansion (9) about the

critical point $t^{(n)}$ is expressible in terms of similar integrals over infinite contours emanating from the adjacent saddles $t^{(m)}$ as

$$(24) \quad R_N^{(n)}(z; \alpha_n) = \sum_{m(n)} \frac{z^{(1-N)/\omega_n}}{2\pi i \omega_m} \left(\int_0^{\infty e^{i\theta_{nm}^+}} \frac{e^{-\mathcal{F}_{nm}^+ u} u^{\frac{N+1}{\omega_n} - \frac{1}{\omega_m} - 1}}{z^{1/\omega_n} - u^{1/\omega_n}} T^{(m)}(u; \alpha_{nm}^+) du - \int_0^{\infty e^{i\theta_{nm}^-}} \frac{e^{-\mathcal{F}_{nm}^- u} u^{\frac{N+1}{\omega_n} - \frac{1}{\omega_m} - 1}}{z^{1/\omega_n} - u^{1/\omega_n}} T^{(m)}(u; \alpha_{nm}^-) du \right).$$

Since $\theta_{nm}^+ = \theta_{nm}^- + 2\pi\omega_n$, a simple change of integration variable in (23) then yields

$$(25) \quad R_N^{(n)}(z; \alpha_n) = \sum_{m(n)} \frac{z^{(1-N)/\omega_n}}{2\pi i} \times \left(\int_0^{\infty e^{i\theta_{nm}^+}} \frac{e^{-\mathcal{F}_{nm}^+ u} u^{\frac{N+1}{\omega_n} - 1}}{z^{1/\omega_n} - u^{1/\omega_n}} \int_{\mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+)} e^{-u(f(t)-f_m)} g(t) dt du - \int_0^{\infty e^{i\theta_{nm}^-}} \frac{e^{-\mathcal{F}_{nm}^- u} u^{\frac{N+1}{\omega_n} - 1}}{z^{1/\omega_n} - u^{1/\omega_n}} \int_{\mathcal{P}^{(m)}(\theta_{nm}^-, \beta_{nm})} e^{-u(f(t)-f_m)} g(t) dt du \right).$$

The path $\mathcal{P}^{(m)}(\theta_{nm}^+, \beta_{nm})$ is geometrically identical to $\mathcal{P}^{(m)}(\theta_{nm}^-, \alpha_{nm}^-)$, and since the angle of the slope of $\mathcal{P}^{(m)}(\theta_{nm}^-, \alpha_{nm}^-)$ to the positive real axis at $t^{(m)}$ is $2\pi/\omega_m$ higher than the corresponding angle of $\mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+)$, we find (cf. (8))

$$\begin{aligned} \beta_{nm} &= \frac{\theta_{nm}^+ + \arg(f^{(\omega_m)}(t^{(m)})) + \omega_m(\varphi^+ + 2\pi/\omega_m)}{2\pi} \\ &= \frac{\theta_{nm}^+ + \arg(f^{(\omega_m)}(t^{(m)})) + \omega_m\varphi^+}{2\pi} + 1 = \alpha_{nm}^+ + 1. \end{aligned}$$

With the notation in (6), integral representation (25) can be written as (12). The observation that

$$(26) \quad R_N^{(n)}(z; \alpha_n + 1) = \sum_{m(n)} \frac{z^{(1-N)/\omega_n}}{2\pi i \omega_m} \int_0^{\infty e^{i(\theta_{nm}^+ + 2\pi)}} \frac{e^{-\mathcal{F}_{nm}^+ u} u^{\frac{N+1}{\omega_n} - \frac{1}{\omega_m} - 1}}{z^{1/\omega_n} - u^{1/\omega_n}} \mathbf{T}^{(m)}(u; \alpha_{nm}^+ + 1) du$$

will also be useful.

In previous publications [7, 24] there were issues with the exact sign of the terms on the right-hand side of (12). These were referred to as ‘‘orientation anomalies.’’ Here we do not encounter these issues because of the careful definitions of the phases on the contours (4), (8).

4. Hyperasymptotic iteration of the exact remainder. In this section we reexpand the exact remainder terms (12) and (26) to derive a template for hyperasymptotic calculations.

First, we begin by defining a set of universal, but generalized, hyperterminant functions $\mathbf{F}^{(j)}$, that form the basis of the template.

Let us introduce the notation $\int_0^{[n]} = \int_0^{\infty e^{i\eta}}$. Then, for k a nonnegative integer, we define

$$\begin{aligned}
 \mathbf{F}^{(0)}(z) &:= 1, & \mathbf{F}^{(1)}\left(z; \begin{matrix} M_0 \\ \omega_0 \\ \sigma_0 \end{matrix}\right) &:= \int_0^{[\pi - \arg \sigma_0]} \frac{e^{\sigma_0 t_0} t_0^{M_0 - 1}}{z^{1/\omega_0} - t_0^{1/\omega_0}} dt_0, \\
 (27) & & & \\
 \mathbf{F}^{(k+1)}\left(z; \begin{matrix} M_0, & \dots, & M_k \\ \omega_0, & \dots, & \omega_k \\ \sigma_0, & \dots, & \sigma_k \end{matrix}\right) & \\
 &:= \int_0^{[\pi - \arg \sigma_0]} \dots \int_0^{[\pi - \arg \sigma_k]} \frac{e^{\sigma_0 t_0 + \dots + \sigma_k t_k} t_0^{M_0 - 1} \dots t_k^{M_k - 1}}{(z^{1/\omega_0} - t_0^{1/\omega_0})(t_0^{1/\omega_1} - t_1^{1/\omega_1}) \dots (t_{k-1}^{1/\omega_k} - t_k^{1/\omega_k})} dt_k \dots dt_0
 \end{aligned}$$

for arbitrary sets of complex numbers M_0, \dots, M_k and $\sigma_0, \dots, \sigma_k$ such that $\text{Re}(M_j) > 1/\omega_j$ and $\sigma_j \neq 0$ for $j = 0, \dots, k$ and for an arbitrary set of positive integers $\omega_0, \dots, \omega_k$. The multiple integrals converge when $|\arg(\sigma_0 z)| < \pi\omega_0$. The $\mathbf{F}^{(j)}$ is termed a “generalized j th-level hyperterminant.” If $\omega_0 = \dots = \omega_{j-1} = 1$, $\mathbf{F}^{(j)}$ reduces to the much simpler j th-level hyperterminant $F^{(j)}$ discussed in the paper [38].

Note that in the case that two successive σ ’s have the same phase the choice of integration path over the poles in (27) needs to be defined more carefully. In those cases we can define the hyperterminant via a limit. For example,

$$(28) \quad \lim_{\varepsilon \rightarrow 0^+} \mathbf{F}^{(k+1)}\left(z; \begin{matrix} M_0, & M_1, & \dots, & M_{k-1}, & M_k \\ \omega_0, & \omega_1, & \dots, & \omega_{k-1}, & \omega_k \\ \sigma_0 e^{-k\varepsilon i}, & \sigma_1 e^{-(k-1)\varepsilon i}, & \dots, & \sigma_{k-1} e^{-\varepsilon i}, & \sigma_k \end{matrix}\right)$$

is an option. Other limits are also possible.

The efficient computation of these generalized hyperterminant functions is outlined in Appendix A.

4.1. Superasymptotics and optimal number of terms. A necessary step in hyperasymptotic reexpansions is to determine the “optimal” number of terms in the original Poincaré expansion (9), defined as the index of the least term in magnitude.

For this section it reasonable to denote the original number of terms in the truncated asymptotic expansion as $N = N_0^{(n)}$ and we denote the associated remainder as $R_0^{(n)}(z; \alpha_n)$. With this notation the integrands in (12) will have a factor $u^{N_0^{(n)}/\omega_n}$. Therefore, when $N_0^{(n)}$ is large, the main contribution to the integrals in (12) comes from infinity where $\mathbf{T}^{(m)}(u; \alpha_{nm}^+) = \mathcal{O}(1)$. In the case that z and u are collinear, i.e., on a Stokes line, it is well known (see, e.g., [12, sect. 8] or [40, sect. 5]) that the Stokes phenomenon produces an extra factor of $\mathcal{O}(\sqrt{N_0^{(n)}})$ when estimating $R_0^{(n)}(z; \alpha_n)$ (see also the proof of Proposition B.1). Thus, we have

$$R_0^{(n)}(z; \alpha_n) = \sqrt{N_0^{(n)}} \frac{\Gamma\left(\frac{N_0^{(n)} + 1}{\omega_n}\right)}{|z|^{N_0^{(n)}/\omega_n}} \sum_{m^{(n)} \in \mathcal{F}_{nm}^+} \frac{1}{|N_0^{(n)}|^{1/\omega_m}} \mathcal{O}(1)$$

for large $N_0^{(n)}$ and $\theta_{nm_1}^- \leq \theta \leq \theta_{nm_2}^+$. Let $N_0^{(n)} = \eta_0^{(n)} \omega_n |z| + \nu_0^{(n)}$ with $\nu_0^{(n)}$ being bounded. Then, with the help of Stirling’s formula,

$$(29) \quad R_0^{(n)}(z; \alpha_n) = e^{-\eta_0^{(n)}|z|} \sum_{m(n)} |z|^{\frac{1}{\omega_n} - \frac{1}{\omega_m}} \left(\frac{\eta_0^{(n)}}{|\mathcal{F}_{nm}^+|} \right)^{\eta_0^{(n)}|z|} \mathcal{O}(1),$$

as $|z| \rightarrow \infty$ in the sector $\theta_{nm_1}^- \leq \theta \leq \theta_{nm_2}^+$. For a fixed m the magnitude of the right-hand side of (29) is minimal in the case that $\eta_0^{(n)} = |\mathcal{F}_{nm}^+|$. Since we sum over all the adjacent saddles we obtain that for the optimal number of terms we have $\eta_0^{(n)} = r_0^{(n)} := \min_{m(n)} |\mathcal{F}_{nm}^+|$, and with that choice we have

$$(30) \quad R_0^{(n)}(z; \alpha_n) = e^{-r_0^{(n)}|z|} |z|^{\frac{1}{\omega_n} - \frac{1}{\tilde{\omega}}} \mathcal{O}(1),$$

as $|z| \rightarrow \infty$ in the sector $\theta_{nm_1}^- \leq \theta \leq \theta_{nm_2}^+$ with $\tilde{\omega} = \max_j \omega_j$.

In the hyperasymptotic process below, we will reexpand this remainder and each of these reexpansions will be truncated and reexpanded and so on. Correspondingly we have to determine the number of terms to take in the original expansion $N_0^{(n)}$, in the first reexpansions $N_1^{(m)}$, and so on. The criterion for determining the ‘‘optimal’’ $N_0^{(n)}, N_1^{(m)}, \dots$, is that the overall error obtained by summing all the contributing expansions should be minimized. This may be determined from considering estimates such as (29) and (35), (37) below. The procedure for determining these optimal numbers of terms is very similar to that of [37] and may be summarized as follows.

Let $G = (V, E)$ be a graph with for the vertices V all the f_j and for the edges $E = \{(f_m, f_n) : t^{(m)} \text{ is adjacent to } t^{(n)}\}$. We define $r_k^{(n)}$ to be the length of the shortest path of $k + 1$ steps in this graph starting at $t^{(n)}$. For a hyperasymptotic expansion of Level k the optimal number of terms is

$$(31) \quad N_0^{(m_0)} = \eta_0^{(m_0)} \omega_{m_0} |z| + \nu_0^{(m_0)}, \quad \dots, \quad N_k^{(m_k)} = \eta_k^{(m_k)} \omega_{m_k} |z| + \nu_k^{(m_k)},$$

with $m_0 = n$, in which

$$\eta_0^{(m_0)} := r_k^{(m_0)}, \quad \eta_j^{(m_j)} := \max \left(0, \eta_{j-1}^{(m_{j-1})} - |\mathcal{F}_{m_{j-1}m_j}^+| \right), \quad j = 1, \dots, k,$$

and the ν_j are all bounded as $|z| \rightarrow \infty$, with estimate

$$(32) \quad R_k^{(n)}(z; \alpha_n) = e^{-r_k^{(n)}|z|} |z|^{\frac{1}{\omega_n} - \frac{1}{\tilde{\omega}}} \mathcal{O}(1),$$

for the remainder as $|z| \rightarrow \infty$ in the sector $\theta_{nm_1}^- \leq \theta \leq \theta_{nm_2}^+$. The main difference from the results in [37] is that here in (31) we have the extra factors ω_j .

4.2. Level 1 hyperasymptotics. We now derive the Level 1 hyperasymptotic expansion. In the integral representation (12) for this remainder we substitute (9) into the $\mathbf{T}^{(m)}$ function. We obtain the reexpansion

$$(33) \quad R_0^{(n)}(z; \alpha_n) = \sum_{m(n)} \frac{z^{(1-N_0^{(n)})/\omega_n}}{2\pi i \omega_m} \sum_{r=0}^{N_1^{(m)}-1} \mathbf{T}_r^{(m)}(\alpha_{nm}^+) \mathbf{F}^{(1)} \left(z; \begin{matrix} \frac{N_0^{(n)}+1}{\omega_n} - \frac{r+1}{\omega_m} \\ \omega_n \\ |\mathcal{F}_{nm}^+| e^{i(\pi-\theta_{nm}^+)} \end{matrix} \right) + R_1^{(n)}(z; \alpha_n).$$

The remainder $R_1^{(n)}(z; \alpha_n)$ depends on the number of terms $N_0^{(n)}$ and $N_1^{(m)}$ and can be represented as

$$\begin{aligned}
 R_1^{(n)}(z; \alpha_n) &= \sum_{m(n)} \sum_{\ell(m)} \frac{z^{(1-N_0^{(n)})/\omega_n}}{(2\pi i)^2 \omega_m \omega_\ell} \\
 &\times \left(\int_0^\infty e^{i\theta_{nm}^+} \int_0^\infty e^{i\theta_{nm\ell}^+} \frac{e^{-\mathcal{F}_{nm}^+ u - \mathcal{F}_{m\ell}^+ v} u^{\frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}}{\omega_m} - 1} v^{\frac{N_1^{(m)}+1}{\omega_m} - \frac{1}{\omega_\ell} - 1}}{(z^{1/\omega_n} - u^{1/\omega_n})(u^{1/\omega_m} - v^{1/\omega_m})} \right. \\
 (34) \quad &\quad \times \mathbf{T}^{(\ell)}(v; \alpha_{nm\ell}^+) dv du \\
 &\quad \left. - \int_0^\infty e^{i\theta_{nm}^+} \int_0^\infty e^{i\theta_{nm\ell}^+ + 2\pi i} \frac{e^{-\mathcal{F}_{nm}^+ u - \mathcal{F}_{m\ell}^+ v} u^{\frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}}{\omega_m} - 1} v^{\frac{N_1^{(m)}+1}{\omega_m} - \frac{1}{\omega_\ell} - 1}}{(z^{1/\omega_n} - u^{1/\omega_n})(u^{1/\omega_m} - v^{1/\omega_m})} \right. \\
 &\quad \left. \times \mathbf{T}^{(\ell)}(v; \alpha_{nm\ell}^+ + 1) dv du \right),
 \end{aligned}$$

in which $\theta_{nm\ell}^+(\theta_{nm}^+)$ corresponds to the path $\mathcal{P}^{(n)}(\theta_{nm}^+; \alpha_{nm}^+)$ and is defined similarly as $\theta_{nm}^+ = \theta_{nm}^+(\theta)$. The $\alpha_{nm\ell}^+$ is the corresponding α_{nm}^+ , which is defined (8). In this derivation we have used the observation (26).

We can estimate the remainder $R_1^{(n)}(z; \alpha_n)$ in a similar way as we did $R_0^{(n)}(z; \alpha_n)$, and one finds

$$\begin{aligned}
 R_1^{(n)}(z; \alpha_n) &= \frac{1}{|z|^{\frac{N_0^{(n)}}{\omega_n}}} \sum_{m(n)} \sqrt{(N_0^{(n)} - N_1^{(m)}) N_1^{(m)}} \\
 &\times \frac{\Gamma\left(\frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}+1}{\omega_m}\right) \Gamma\left(\frac{N_1^{(m)}+1}{\omega_m}\right)}{|\mathcal{F}_{nm}^+|^{\frac{N_0^{(n)}}{\omega_n} - \frac{N_1^{(m)}}{\omega_m}}} \sum_{\ell(m)} \frac{1}{|\mathcal{F}_{m\ell}^+|^{\frac{N_1^{(m)}}{\omega_m}} (N_1^{(m)})^{\frac{1}{\omega_\ell}}} \mathcal{O}(1).
 \end{aligned}$$

Then

$$\begin{aligned}
 R_1^{(n)}(z; \alpha_n) &= e^{-\eta_0^{(n)}|z|} \sum_{m(n)} \left(\frac{\eta_0^{(n)} - \eta_1^{(m)}}{|\mathcal{F}_{nm}^+|} \right)^{(\eta_0^{(n)} - \eta_1^{(m)})|z|} \\
 (35) \quad &\quad \times \sum_{\ell(m)} |z|^{\frac{1}{\omega_n} - \frac{1}{\omega_\ell}} \left(\frac{\eta_1^{(m)}}{|\mathcal{F}_{m\ell}^+|} \right)^{\eta_1^{(m)}|z|} \mathcal{O}(1),
 \end{aligned}$$

as $|z| \rightarrow \infty$ in the sector $\theta_{nm_1}^- \leq \theta \leq \theta_{nm_2}^+$. For fixed m and ℓ , using an approach similar to subsection 4.1 above, it is easy to show that the optimal number of terms is obtained when $\eta_0^{(n)} - \eta_1^{(m)} = |\mathcal{F}_{nm}^+|$ and $\eta_1^{(m)} = |\mathcal{F}_{m\ell}^+|$.

Rigorous bounds for Level 1 hyperterminants are derived in Appendix B.

4.3. Level 2 hyperasymptotics. The Level 2 hyperasymptotic expansion is now derived by reexpanding the Level 1 expansion. Again we substitute (9) into the $\mathbf{T}^{(\ell)}$ functions on the right-hand side of (34) and obtain the reexpansion

$$\begin{aligned}
 R_1^{(n)}(z; \alpha_n) &= \sum_{m(n)} \sum_{\ell(m)} \frac{z^{(1-N_0^{(n)})/\omega_n}}{(2\pi i)^2 \omega_m \omega_\ell} \sum_{r=0}^{N_2^{(\ell)}-1} \\
 (36) \quad &\left\{ \mathbf{T}_r^{(\ell)}(\alpha_{nm\ell}^+) \mathbf{F}^{(2)} \left(z; \begin{matrix} \frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}}{\omega_m}, & \frac{N_1^{(m)}+1}{\omega_m} - \frac{r+1}{\omega_\ell} \\ \omega_n, & \omega_m, & \omega_\ell \end{matrix} \right) \right. \\
 &\quad \left. - \mathbf{T}_r^{(\ell)}(\alpha_{nm\ell}^+ + 1) \mathbf{F}^{(2)} \left(z; \begin{matrix} \frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}}{\omega_m}, & \frac{N_1^{(m)}+1}{\omega_m} - \frac{r+1}{\omega_\ell} \\ \omega_n, & \omega_m, & \omega_\ell \end{matrix} \right) \right\} \\
 &\quad + R_2^{(n)}(z; \alpha_n).
 \end{aligned}$$

We also obtain an exact integral representation for the remainder, and this can be used to obtain the estimate

$$\begin{aligned}
 (37) \quad R_2^{(n)}(z; \alpha_n) &= e^{-\eta_0^{(n)}|z|} \sum_{m(n)} \left(\frac{\eta_0^{(n)} - \eta_1^{(m)}}{|\mathcal{F}_{nm}^+|} \right)^{(\eta_0^{(n)} - \eta_1^{(m)})|z|} \sum_{\ell(m)} \left(\frac{\eta_1^{(m)} - \eta_2^{(\ell)}}{|\mathcal{F}_{m\ell}^+|} \right)^{(\eta_1^{(m)} - \eta_2^{(\ell)})|z|} \\
 &\quad \times \sum_{k(\ell)} |z|^{\frac{1}{\omega_n} - \frac{1}{\omega_k}} \left(\frac{\eta_2^{(\ell)}}{|\mathcal{F}_{\ell k}^+|} \right)^{\eta_2^{(\ell)}|z|} \mathcal{O}(1),
 \end{aligned}$$

as $|z| \rightarrow \infty$ in the sector $\theta_{nm_1}^- \leq \theta \leq \theta_{nm_2}^+$.

4.4. Level 3 hyperasymptotics. We can continue with this process and will obtain at Level 3 the expansion

$$\begin{aligned}
 (38) \quad R_2^{(n)}(z; \alpha_n) &= \sum_{m(n)} \sum_{\ell(m)} \sum_{k(\ell)} \frac{z^{(1-N_0^{(n)})/\omega_n}}{(2\pi i)^3 \omega_m \omega_\ell \omega_k} \sum_{r=0}^{N_3^{(k)}-1} \\
 &\left(\mathbf{T}_r^{(k)}(\alpha_{nm\ell k}^+) \mathbf{F}^{(3)} \left(z; \begin{matrix} \frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}}{\omega_m}, & \frac{N_1^{(m)}+1}{\omega_m} - \frac{N_2^{(\ell)}}{\omega_\ell}, & \frac{N_2^{(\ell)}+1}{\omega_\ell} - \frac{r+1}{\omega_k} \\ \omega_n, & \omega_m, & \omega_\ell \end{matrix} \right) \right. \\
 &\quad \left. - \mathbf{T}_r^{(k)}(\alpha_{nm\ell k}^+ + 1) \mathbf{F}^{(3)} \left(z; \begin{matrix} \frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}}{\omega_m}, & \frac{N_1^{(m)}+1}{\omega_m} - \frac{N_2^{(\ell)}}{\omega_\ell}, & \frac{N_2^{(\ell)}+1}{\omega_\ell} - \frac{r+1}{\omega_k} \\ \omega_n, & \omega_m, & \omega_\ell \end{matrix} \right) \right. \\
 &\quad \left. - \mathbf{T}_r^{(k)}(\alpha_{nm\ell k}^+ + 1) \mathbf{F}^{(3)} \left(z; \begin{matrix} \frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}}{\omega_m}, & \frac{N_1^{(m)}+1}{\omega_m} - \frac{N_2^{(\ell)}}{\omega_\ell}, & \frac{N_2^{(\ell)}+1}{\omega_\ell} - \frac{r+1}{\omega_k} \\ \omega_n, & \omega_m, & \omega_\ell \end{matrix} \right) \right. \\
 &\quad \left. + \mathbf{T}_r^{(k)}(\alpha_{nm\ell k}^+ + 2) \mathbf{F}^{(3)} \left(z; \begin{matrix} \frac{N_0^{(n)}+1}{\omega_n} - \frac{N_1^{(m)}}{\omega_m}, & \frac{N_1^{(m)}+1}{\omega_m} - \frac{N_2^{(\ell)}}{\omega_\ell}, & \frac{N_2^{(\ell)}+1}{\omega_\ell} - \frac{r+1}{\omega_k} \\ \omega_n, & \omega_m, & \omega_\ell \end{matrix} \right) \right) \\
 &\quad + R_3^{(n)}(z; \alpha_n).
 \end{aligned}$$

An estimate for the remainder $R_3^{(n)}(z; \alpha_n)$, similar to those of (29), (35), and (37), may be obtained, and further iterations to higher hyper-levels derived. We spare the reader these details as the pattern should now be clear.

Initially, this expansion might seem overcomplicated. However, inspection of the terms shows that once we have line 2 of (38) the details of the other lines can be easily deduced. It follows from (6) and (11) that the coefficients follow from the coefficients in line 2 by just multiplying by a simple exponential. The generalized hyperterminants only differ by a change in the phases of two (bottom center and right) arguments.

4.5. Late coefficients and resurgence. The reexpansion (33) is suitable for obtaining an asymptotic expansion for the late (large- N) coefficients $T_N^{(n)}(\alpha_n)$. Indeed, if we combine the identity

$$T_N^{(n)}(\alpha_n) = z^{N/\omega_n} \left(R_N^{(n)}(z; \alpha_n) - R_{N+1}^{(n)}(z; \alpha_n) \right)$$

with (33), we deduce

$$(39) \quad T_N^{(n)}(\alpha_n) = \sum_{m(n)} \frac{1}{2\pi i \omega_m} \sum_{r=0}^{N_1^{(m)}-1} \mathbf{T}_r^{(m)}(\alpha_{nm}^+) \frac{e^{i\theta_{nm}^+ \left(\frac{N+1}{\omega_n} - \frac{r+1}{\omega_m} \right)} \Gamma \left(\frac{N+1}{\omega_n} - \frac{r+1}{\omega_m} \right)}{|\mathcal{F}_{nm}^+|^{\frac{N+1}{\omega_n} - \frac{r+1}{\omega_m}}} + \tilde{R}_1^{(n)}(N; \alpha_n).$$

Note that the coefficients in this expansion are the coefficients of the asymptotic expansions of integrals over doubly infinite contours passing through the adjacent saddles, a manifestation of “resurgence.” The form (39) is of a generalized sum of factorials over powers. Note the careful representation of the phases of the singulants. Various special cases of (39) were derived, using nonrigorous methods, by Dingle (see [17, Chap. VII], including exercises). See also [7], [23].

When we eliminate $|z|$ in the definitions (31) we obtain for the optimal numbers of terms in (39) that

$$N_1^{(m)} = \frac{\eta_1^{(m)} \omega_m}{\eta_0^{(n)} \omega_n} N + \mathcal{O}(1),$$

as $N \rightarrow \infty$.

In the swallowtail example below we shall illustrate how this result can be used to determine the adjacency of the saddles algebraically rather than geometrically.

5. Error bounds. In this section we derive rigorous, novel, and sharp error bounds for the exact remainder $R_N^{(n)}(z; \alpha_n)$ of asymptotic expansions of the form (9) derived from integrals of the class (1).

The remainder term (20) can be written as

$$(40) \quad R_N^{(n)}(z; \alpha_n) = \frac{\omega_n}{2\pi i z^{N/\omega_n}} \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} \int_0^{\infty e^{\frac{2\pi i \alpha_n}{\omega_n}}} \frac{e^{-s \omega_n} s^N}{1 - \frac{s}{(z(f(t) - f_n))^{1/\omega_n}}} ds dt = \frac{e^{2\pi i \frac{N+1}{\omega_n} \alpha_n}}{2\pi i z^{N/\omega_n}} \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} \int_0^{\infty} \frac{e^{-u} u^{\frac{N+1}{\omega_n} - 1}}{1 + \left(\frac{ue^{\pi i(2\alpha_n - \omega_n)}}{z(f(t) - f_n)} \right)^{1/\omega_n}} du dt,$$

where $\mathcal{C}^{(m)}(\theta_{nm}^+) := \mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+) \cup -\mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+ + 1)$. We note that

$$\begin{aligned} \arg\left(\frac{ue^{\pi i(2\alpha_n - \omega_n)}}{z(f(t) - f_n)}\right) &= 2\pi\alpha_n - \pi\omega_n - \theta - (-\theta_{nm}^+ + 2\pi\alpha_n) \\ &= -\pi\omega_n - \theta + \theta_{nm}^+ > -\pi\omega_n \end{aligned}$$

and

$$\begin{aligned} \arg\left(\frac{ue^{\pi i(2\alpha_n - \omega_n)}}{z(f(t) - f_n)}\right) &= 2\pi\alpha_n - \pi\omega_n - \theta - (-\theta_{nm}^+ + 2\pi\alpha_n) = -\pi\omega_n - \theta + \theta_{nm}^+ \\ &= -\pi\omega_n - \theta + \theta_{nm}^- + 2\pi\omega_n = \pi\omega_n - \theta + \theta_{nm}^- < \pi\omega_n \end{aligned}$$

whenever $t \in \mathcal{C}^{(m)}(\theta_{nm}^+)$. Thus,

$$\left| \arg\left(\frac{ue^{\pi i(2\alpha_n - \omega_n)}}{z(f(t) - f_n)}\right) \right| < \pi\omega_n.$$

Consequently, the u -integral may be expressed in terms of the generalized first-level hyperterminant as

$$\begin{aligned} &\int_0^\infty \frac{e^{-u} u^{\frac{N+1}{\omega_n} - 1}}{1 + \left(\frac{ue^{\pi i(2\alpha_n - \omega_n)}}{z(f(t) - f_n)}\right)^{1/\omega_n}} du \\ &= e^{-\pi \frac{N+1}{\omega_n} i} \left(e^{\pi i(\omega_n - 2\alpha_n)} z(f(t) - f_n) \right)^{\frac{1}{\omega_n}} \mathbf{F}^{(1)} \left(e^{\pi i(\omega_n - 2\alpha_n)} z(f(t) - f_n); \begin{matrix} \frac{N+1}{\omega_n} \\ \omega_n \\ 1 \end{matrix} \right). \end{aligned}$$

Inserting this expression into (40), we obtain the following alternative representation of $R_N^{(n)}(z; \alpha_n)$:

$$\begin{aligned} (41) \quad R_N^{(n)}(z; \alpha_n) &= \frac{e^{(2\alpha_n - 1)\pi i \frac{N+1}{\omega_n}}}{2\pi i z^{N/\omega_n}} \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} \\ &\times \left(e^{\pi i(\omega_n - 2\alpha_n)} z(f(t) - f_n) \right)^{\frac{1}{\omega_n}} \mathbf{F}^{(1)} \left(e^{\pi i(\omega_n - 2\alpha_n)} z(f(t) - f_n); \begin{matrix} \frac{N+1}{\omega_n} \\ \omega_n \\ 1 \end{matrix} \right) dt. \end{aligned}$$

This representation is valid when $\theta_{nm_1}^- - \frac{\pi}{2} < \theta < \theta_{nm_2}^+ + \frac{\pi}{2}$ (cf. (42) below). We may then bound the t integral as follows:

$$\begin{aligned} \left| R_N^{(n)}(z; \alpha_n) \right| &\leq \frac{\Gamma\left(\frac{N+1}{\omega_n}\right)}{2\pi |z|^{N/\omega_n}} \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \left| \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} dt \right| \\ &\times \sup_{r \geq 1} \left| \frac{\left(z |\mathcal{F}_{nm}^+| e^{(\pi\omega_n - \theta_{nm}^+)i} r \right)^{\frac{1}{\omega_n}} \mathbf{F}^{(1)} \left(z |\mathcal{F}_{nm}^+| e^{(\pi\omega_n - \theta_{nm}^+)i} r; \begin{matrix} \frac{N+1}{\omega_n} \\ \omega_n \\ 1 \end{matrix} \right)}{\Gamma\left(\frac{N+1}{\omega_n}\right)} \right|. \end{aligned}$$

A further simplification of this bound is possible by employing the estimates for the generalized first-level hyperterminant given in Appendix B. In this way, we obtain

(42)

$$\begin{aligned}
 \left| R_N^{(n)}(z; \alpha_n) \right| &\leq \frac{\Gamma\left(\frac{N+1}{\omega_n}\right)}{2\pi |z|^{N/\omega_n}} \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \left| \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} dt \right| \\
 &\times \begin{cases} 1 & \text{if } |\theta - \theta_{nm}^+ + \pi\omega_n| \leq \frac{\pi}{2}\omega_n, \\ \min\left(\left| \csc\left(\frac{\theta - \theta_{nm}^+}{\omega_n}\right) \right|, \omega_n \sqrt{e\left(\frac{N+1}{\omega_n} + \frac{1}{2}\right)}\right) & \text{if } \frac{\pi}{2}\omega_n < |\theta - \theta_{nm}^+ + \pi\omega_n| \leq \pi\omega_n, \\ \frac{\sqrt{2\pi\omega_n(N+1)}}{|\cos(\theta - \theta_{nm}^+)|^{\frac{N+1}{\omega_n}}} + \omega_n \sqrt{e\left(\frac{N+1}{\omega_n} + \frac{1}{2}\right)} & \text{if } \pi\omega_n < |\theta - \theta_{nm}^+ + \pi\omega_n| < \pi\omega_n + \frac{\pi}{2}. \end{cases}
 \end{aligned}$$

In the case of linear endpoint ($\omega_n = 1$), the quantity $\sqrt{e(N+\frac{3}{2})}$ in (42) can be replaced by (51) with $M = N + 1$.

In (10) we may expand the loop contour of integration around the critical point $t^{(n)}$ across the domain $\Delta^{(n)}$ to obtain a representation of the asymptotic coefficients in terms of integrals over the contours $\mathcal{C}^{(m)}(\theta_{nm}^+)$ as follows:

$$(43) \quad \left| \frac{T_N^{(n)}(\alpha_n)}{z^{N/\omega_n}} \right| = \frac{\Gamma\left(\frac{N+1}{\omega_n}\right)}{2\pi |z|^{N/\omega_n}} \left| \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \frac{g(t)}{(f(t) - f_n)^{(N+1)/\omega_n}} dt \right|.$$

This representation illustrates the close relation between the form of the bound (42) and the absolute value of the first neglected term. The modulus bars are inside the integral in (42), whereas they are at the outside of the integral in (43). However, in (43) we integrate along steepest descent paths $\mathcal{C}^{(m)}(\theta_{nm}^+)$ on which $f(t) - f_n$ is monotonically decreasing. This means that only when $g(t)$ is highly oscillatory will the integral in (42) be considerably larger than the integral in (43). The larger the value of N , the smaller the difference in size of the two integrals.

Figure 4, for our first example below, clearly demonstrates the asymptotic property that sizes of the exact terms and the corresponding remainders are approximately the same. This follows from the factor 1 in the second line of (42). In Figure 6, which is for our second example, the remainders are considerably larger than the terms. That example illustrates the effect of the additional factor $\omega_n \sqrt{e(\frac{N+1}{\omega_n} + \frac{1}{2})}$ in the third line of (42) pertaining to the parameters θ , ω_n , and θ_{nm}^+ of that particular calculation.

5.1. Bounds for simple saddles. If $t^{(n)}$ is a simple saddle, then the integral over the double infinite contour through $t^{(n)}$ can be expanded as

$$\mathbf{T}^{(n)}(z, 0) = \sum_{r=0}^{N-1} \frac{\mathbf{T}_{2r}^{(n)}(0)}{z^r} + \mathbf{R}_N^{(n)}(z, 0)$$

with $\mathbf{R}_N^{(n)}(z, 0) = R_{2N}^{(n)}(z; 0) - R_{2N}^{(n)}(z; 1)$. The estimation of $\mathbf{R}_N^{(n)}(z, 0)$ was considered by Boyd [12] in the case that all the adjacent saddles are simple. Employing (41) and simplifying the result, we obtain

$$\begin{aligned}
 \mathbf{R}_N^{(n)}(z, 0) &= \frac{(-1)^{N+1}}{\pi z^N} \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \frac{g(t)}{(f(t) - f_n)^{N+\frac{1}{2}}} \\
 &\quad \times e^{\pi i z(f(t) - f_n)} F^{(1)}\left(e^{\pi i z(f(t) - f_n)}; \begin{matrix} N + \frac{1}{2} \\ 1 \end{matrix}\right) dt.
 \end{aligned}$$

This representation is valid when $\theta_{nm_1}^- - \frac{\pi}{2} < \theta < \theta_{nm_2}^+ + \frac{\pi}{2}$. We may then bound the t integral as follows:

$$\begin{aligned} \left| \mathbf{R}_N^{(n)}(z, 0) \right| &\leq \frac{\Gamma(N + \frac{1}{2})}{\pi |z|^N} \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \left| \frac{g(t)}{(f(t) - f_n)^{N + \frac{1}{2}}} dt \right| \\ &\quad \times \sup_{r \geq 1} \left| \frac{z |\mathcal{F}_{nm}^+| e^{(\pi - \theta_{nm}^+)i_r}}{\Gamma(N + \frac{1}{2})} F^{(1)} \left(z |\mathcal{F}_{nm}^+| e^{(\pi - \theta_{nm}^+)i_r}; \begin{matrix} N + \frac{1}{2} \\ 1 \end{matrix} \right) \right|. \end{aligned}$$

A further simplification of this bound is possible by applying the estimates for the generalized first-level hyperterminant given in Appendix B. In this way, we deduce

$$\begin{aligned} \left| \mathbf{R}_N^{(n)}(z, 0) \right| &\leq \frac{\Gamma(N + \frac{1}{2})}{\pi |z|^N} \sum_{m(n)} \int_{\mathcal{C}^{(m)}(\theta_{nm}^+)} \left| \frac{g(t)}{(f(t) - f_n)^{N + \frac{1}{2}}} dt \right| \\ (44) \quad &\times \begin{cases} 1 & \text{if } |\theta - \theta_{nm}^+ + \pi| \leq \frac{\pi}{2}, \\ \min(|\csc(\theta - \theta_{nm}^+)|, \sqrt{e(N + 1)}) & \text{if } \frac{\pi}{2} < |\theta - \theta_{nm}^+ + \pi| \leq \pi, \\ \frac{\sqrt{2\pi(N + \frac{1}{2})}}{|\cos(\theta - \theta_{nm}^+)|^{N + \frac{1}{2}}} + \sqrt{e(N + 1)} & \text{if } \pi < |\theta - \theta_{nm}^+ + \pi| < \frac{3\pi}{2}. \end{cases} \end{aligned}$$

The quantity $\sqrt{e(N + 1)}$ in this bound can be replaced by (51) with $M = N + \frac{1}{2}$.

The bound (44) improves Boyd’s [12] results in three ways. First, it is more general in that the adjacent saddles need not be simple. Second, (44) extends the range of validity of the bound to include $\pi < |\theta - \theta_{nm}^+ + \pi| < \frac{3\pi}{2}$. Third, the new result sharpens the bound with a factor $\sqrt{e(N + 1)}$ in place of Boyd’s larger $2\sqrt{N}$ factor, and for this larger factor to hold he even requires the extra assumption $N \geq \cot^2(\frac{1}{2}(\theta_{nm_2}^+ - \theta_{nm_1}^-))$.

6. Example 1: Pearcey on the cusp. A rescaled Pearcey function (compare [18, sect. 36.2]) is defined by the integral

$$(45) \quad \Psi_2(x, y; z) = \int_{-\infty}^{+\infty} e^{-zf(t;x,y)} dt, \quad f(t; x, y) = -i(t^4 + yt^2 + xt).$$

Due to the polynomial nature of the exponent function and the ability to scale t , z , with x and y , without loss of generality the modulus of the large parameter z may be set to 1. The function represents the wavefield in the neighborhood of the canonically stable cusp catastrophe [5] and occurs commonly in two-dimensional linear wave problems.

The integrand possesses three saddle points $t^{(j)}, j = 1, 2, 3$, satisfying

$$f'(t^{(j)}; x, y) = 4(t^{(j)})^3 + 2yt^{(j)} + x = 0.$$

In [7] a hyperasymptotic expansion of the Pearcey function was calculated in the case of three distinct saddle points. Here we have extended that analysis to cover the case where two of the saddles have coalesced.

Two of the three saddle points coalesce on the cusp-shaped caustic given by

$$f'(t; x, y) = f''(t; x, y) = 0 \quad \Rightarrow \quad 27x^2 = -8y^3, \quad (x, y) \neq 0;$$

see Figure 3(a). (At the origin $(x, y) = (0, 0)$, all three saddles coalesce, where the integral reduces to an exact explicit representation [18, eqn. 36.2.15].)




FIG. 3. (a) Location of the parameter point $(x, y) = (2\sqrt{2}, -3)$ at which we evaluate the integral (45) relative to the caustic of the Pearcey function, satisfying $27x^2 = -8y^3$. (b) The steepest descent paths $\mathcal{P}^{(1)}(-\frac{\pi}{4}, 0)$, $\mathcal{P}^{(1)}(-\frac{\pi}{4}, 1)$ in the complex t -plane emerging from the simple saddle $t^{(1)}$ ($\omega_1 = 2$) and traveling to labeled valleys V_j , $j = 2, 3$, at infinity. Also shown is the degenerate saddle $t^{(2)}$ ($\omega_2 = 3$). (c) The steepest descent paths $\mathcal{P}^{(2)}(\frac{\pi}{2}, \alpha_2)$, $\alpha_2 = 0, 1, 2$, emerging from $t^{(2)}$, as a Stokes phenomenon occurs between $t^{(1)}$ and $t^{(2)}$ when $\theta_{12}^+ = \frac{\pi}{2}$. The bold lines are the steepest paths that are used in the Level 1 hyperasymptotic expansion about $t^{(1)}$ (33), (6). (d) The steepest descent paths $\mathcal{P}^{(2)}(\frac{11}{2}\pi, \alpha_2)$, $\alpha_2 = 0, 1, 2$, emerging from $t^{(2)}$, as a Stokes phenomenon occurs between $t^{(2)}$ and $t^{(1)}$ when $\theta_{12}^+ = \frac{11}{2}\pi$. The bold lines are the steepest paths that are used in the Level 2 hyperasymptotic expansion about $t^{(1)}$ (36), (6). (Or Level 1 hyperasymptotic expansion about $t^{(2)}$.)

We shall choose $x = 2\sqrt{2}$, $y = -3$. There is a simple saddle at $t^{(1)} = -\sqrt{2}$ and a double saddle denoted by $t^{(2)} = 1/\sqrt{2}$. The asymptotic expansion about $t^{(1)}$ has $\omega_1 = 2$ and is controlled by the double saddle at $t^{(2)}$ with $\omega_2 = 3$, and vice versa.

We shall calculate a hyperasymptotic expansion about $t^{(1)}$. We take $z = e^{i\theta}$ and chose $\theta = -\frac{\pi}{4}$. The steepest paths are denoted by $\mathcal{P}^{(1)}(-\frac{\pi}{4}, 0)$ and $\mathcal{P}^{(1)}(-\frac{\pi}{4}, 1)$; see Figure 3(b).

In the calculations below we will use (8) many times and observe that in this case $\arg(f^{(\omega_1)}(t^{(1)})) = \arg(f^{(\omega_2)}(t^{(2)})) = -\frac{\pi}{2}$, and in Figures 3(c), (d) for the curve $\mathcal{P}^{(2)}(\frac{\pi}{2}, 1)$ we have $\varphi = \frac{2}{3}\pi$ and for curve $\mathcal{P}^{(1)}(\frac{11}{2}\pi, 1)$ we have $\varphi = -\frac{\pi}{2}$.

The normalized integrals that we consider are

$$T^{(1)}(z; \alpha_1) = 2z^{1/2} \int_{\mathcal{P}^{(1)}(-\frac{\pi}{4}, \alpha_1)} e^{zi(t^4 - 3t^2 + 2\sqrt{2}t + 6)} dt, \quad \alpha_1 = 0, 1,$$

which posses the asymptotic expansions

$$(46) \quad T^{(1)}(z; \alpha_1) = \sum_{r=0}^{N_0^{(1)}-1} \frac{T_r^{(1)}(\alpha_1)}{z^{r/2}} + R_1^{(1)}(z; \alpha_1),$$

with coefficients

$$(47) \quad \begin{aligned} T_r^{(1)}(0) &= e^{\frac{\pi}{4}(r+1)i} \frac{\Gamma(\frac{r+1}{2})}{\Gamma(r+1)} \left[\frac{d^r}{dt^r} \left(\frac{(t + \sqrt{2})^2}{t^4 - 3t^2 + 2\sqrt{2}t + 6} \right)^{(r+1)/2} \right]_{t=-\sqrt{2}} \\ &= e^{\frac{\pi}{4}(r+1)i} \frac{\Gamma(\frac{r+1}{2})}{\Gamma(r+1)} \left[\frac{d^r}{dt^r} \left(\frac{1}{t^2 - 4\sqrt{2}t + 9} \right)^{(r+1)/2} \right]_{t=0} \\ &= \frac{e^{\frac{\pi}{4}(r+1)i}}{3^{2r+1}} \Gamma\left(\frac{r+1}{2}\right) C_r^{(\frac{r+1}{2})} \left(\frac{2\sqrt{2}}{3}\right), \end{aligned}$$

$T_r^{(1)}(\alpha_1) = e^{2\pi i \alpha_1 (r+1)/2} T_r^{(1)}(0)$. In deriving the coefficients, in the penultimate line of (47) we have recognized the presence of the generating function [18, eqn. 18.12.4] for the ultraspherical polynomials $C_r^{(p)}(w)$.

We will also need the coefficients of the asymptotic expansions of the integrals

$$T^{(2)}(z; \alpha_2) = 3z^{1/3} \int_{\mathcal{P}^{(2)}(-\frac{\pi}{4}, \alpha_2)} e^{zi(t^4 - 3t^2 + 2\sqrt{2}t - \frac{3}{4})} dt, \quad \alpha_2 = 0, 1, 2,$$

which possess the asymptotic expansions

$$T^{(2)}(z; \alpha_2) = \sum_{r=0}^{N_0^{(1)}-1} \frac{T_r^{(2)}(\alpha_2)}{z^{r/3}} + R_0^{(2)}(z; \alpha_2),$$

with coefficients

$$\begin{aligned} T_r^{(2)}(0) &= e^{\frac{\pi}{6}(r+1)i} \frac{\Gamma(\frac{r+1}{3})}{\Gamma(r+1)} \left[\frac{d^r}{dt^r} \left(\frac{(t - 1/\sqrt{2})^3}{t^4 - 3t^2 + 2\sqrt{2}t - 3/4} \right)^{(r+1)/3} \right]_{t=1/\sqrt{2}} \\ &= e^{\frac{\pi}{6}(r+1)i} \frac{\Gamma(\frac{r+1}{3})}{\Gamma(r+1)} \left[\frac{d^r}{dt^r} \left(\frac{1}{t + 2\sqrt{2}} \right)^{(r+1)/3} \right]_{t=0} \\ &= \frac{e^{\frac{\pi}{6}(r+1)i}}{2^{2r+1/2}} \Gamma\left(\frac{r+1}{3}\right) \binom{-\frac{r+1}{3}}{r}, \end{aligned}$$

and $T_r^{(2)}(\alpha_2) = e^{2\pi i \alpha_2 (r+1)/3} T_r^{(2)}(0)$.

For the singulant on the caustic we have

$$|\mathcal{F}_{12}^+| = |f(t^{(2)}; 2\sqrt{2}, -3) - f(t^{(1)}; 2\sqrt{2}, -3)| = \frac{27}{4}.$$

The effective asymptotic parameter in the expansion is thus $|z\mathcal{F}_{12}^+| = 6.75$, and hence the optimal number of terms in (46) is $N_0^{(1)} = \lceil |z\mathcal{F}_{12}^+| \omega_1 \rceil = 13$.

Since $\theta = -\frac{\pi}{4}$ it follows that for the integral $T^{(1)}(z; 0)$, the corresponding $\theta_{12}^+ = \frac{\pi}{2}$. The corresponding contour of integration emanating from adjacent saddle $t^{(2)}$ is $\mathcal{P}^{(2)}(\frac{\pi}{2}, 1)$ (see Figure 3(c)), and hence, the Level 1 reexpansion is of the form

$$R_0^{(1)}(z; \alpha_n) = \frac{z^{(1-N_0^{(1)})/2}}{6\pi i} \sum_{r=0}^{N_1^{(2)}-1} \mathbf{T}_r^{(2)}(1)\mathbf{F}^{(1)} \left(z; \begin{matrix} \frac{N_0^{(1)}+1}{2} - \frac{r+1}{3} \\ \frac{27}{4}e^{\frac{\pi}{2}i} \end{matrix} \right) + R_1^{(1)}(z; 0).$$

The optimal numbers of terms at Level 1 are $N_0^{(1)} = \lfloor 2|z\mathcal{F}_{12}^+|\omega_1 \rfloor = 27$ and $N_1^{(2)} = \lfloor |z\mathcal{F}_{12}^+|\omega_2 \rfloor = 20$.

With $\theta_{12}^+ = \frac{\pi}{2}$ and contour $\mathcal{P}^{(2)}(\frac{\pi}{2}, 1)$ it follows that $\theta_{121}^+ = \theta_{12}^+ + 5\pi = \frac{11}{2}\pi$, and the corresponding contour of integration emanating from adjacent saddle $t^{(1)}$ is $\mathcal{P}^{(1)}(\frac{11}{2}\pi, 2)$ (see Figure 3(d)), and hence, the Level 2 reexpansion is of the form

$$\begin{aligned} R_1^{(1)}(z; 0) &= \sum_{r=0}^{N_2^{(1)}-1} \frac{z^{(1-N_0^{(1)})/2}}{(2\pi i)^2 6} \\ &\times \left(\mathbf{T}_r^{(1)}(2)\mathbf{F}^{(2)} \left(z; \begin{matrix} \frac{N_0^{(1)}+1}{2} - \frac{N_1^{(2)}}{3}, & \frac{N_1^{(2)}+1}{3} - \frac{r+1}{2} \\ 2, & 3 \\ \frac{27}{4}e^{\frac{\pi}{2}i}, & \frac{27}{4}e^{-\frac{9}{2}\pi i} \end{matrix} \right) \right. \\ &\quad \left. - \mathbf{T}_r^{(1)}(3)\mathbf{F}^{(2)} \left(z; \begin{matrix} \frac{N_0^{(1)}+1}{2} - \frac{N_1^{(2)}}{3}, & \frac{N_1^{(2)}+1}{3} - \frac{r+1}{2} \\ 2, & 3 \\ \frac{27}{4}e^{\frac{\pi}{2}i}, & \frac{27}{4}e^{-\frac{13}{2}\pi i} \end{matrix} \right) \right) \\ &+ R_2^{(1)}(z; 0). \end{aligned}$$

The optimal numbers of terms at Level 2 are given in Table 1.

Finally, with $\theta_{121}^+ = \frac{11}{2}\pi$ and contour $\mathcal{P}^{(1)}(\frac{11}{2}\pi, 2)$ it follows that $\theta_{1212}^+ = \theta_{121}^+ + 3\pi = \frac{17}{2}\pi$, $\alpha_{1212}^+ = 5$, and the optimal numbers in (38) are again given in Table 1.

When we compute our integral numerically with high precision for these values of x , y , and z we obtain

$$T^{(1)}(z, 0) = 0.37277007370182291370 + 0.47493131741141216950i.$$

The numerics of the hyperasymptotic approximations are given in Table 1, and for the Level 3 expansion we display the terms and errors in Figure 4. We observe in this

TABLE 1

The numbers of terms in each series of the hyperasymptotic expansion that are required to minimize overall the absolute error for the (1 → 2) Pearcey example derived from (31). Note that each row corresponds to a decision to stop the reexpansion at that stage. Hence the table row corresponding to Level 2 corresponds to the truncations required at each level up to two, after deciding to stop after two reexpansions of the remainder. Note that all the truncations change with the decision to stop at a particular level.

Level	$N_0^{(1)}$	$N_1^{(2)}$	$N_2^{(1)}$	$N_3^{(2)}$	Error
0	13				1.9×10^{-4}
1	27	20			9.5×10^{-9}
2	40	40	13		3.8×10^{-14}
3	54	60	27	20	9.0×10^{-17}




FIG. 4. For Example 1: The modulus of the n th term in the Level 3 hyperasymptotic expansion (blue dots), and the modulus of the remainder after taking n terms in the approximation (red crosses).

figure that the remainders in the original Poincaré expansions are of the same size as the first neglected terms, as predicted in section 5. In fact at all levels the remainders are of a similar size as the first neglected terms. Occasionally, the remainders are considerably smaller.

In this section we derived hyperasymptotic approximations for $T^{(1)}(z, 0)$. Note that we can repeat the calculation for the integral $T^{(1)}(z, 1)$. The *only* changes in the reexpansions are that all the θ^+ are increased by 2π and all the α^+ are increased by 1. The optimal numbers of terms will remain the same.

7. Example 2: Higher order saddles. In the second main example we take an integral of the form (1), but now with $g(t) \equiv 1$ and

$$f(t) = \frac{15}{28}t^7 - 5t^6 + 18t^5 - 30t^4 + 20t^3 \quad \implies \quad f'(t) = \frac{15}{4}t^2 (t - 2)^4 .$$

The saddle points are $t^{(1)} = 0$ and $t^{(2)} = 2$, with $\omega_1 = 3$ and $\omega_2 = 5$. Hence this example is an example of the hyperasymptotic method when both saddles are degenerate.

Once again, due to the scaling properties of the polynomial $f(t)$ we may take $z = e^{i\theta}$ and also choose $\theta = -\frac{\pi}{4}$. The steepest descent paths are displayed in Figure 5(a). For the coefficients in the asymptotic expansions we have

$$\begin{aligned} T_r^{(1)}(0) &= \frac{\Gamma(\frac{r+1}{3})}{\Gamma(r+1)} \left[\frac{d^r}{dt^r} \left(\frac{1}{\frac{15}{28}t^4 - 5t^3 + 18t^2 - 30t + 20} \right)^{(r+1)/3} \right]_{t=0} , \\ T_r^{(2)}(0) &= \frac{\Gamma(\frac{r+1}{5})}{\Gamma(r+1)} \left[\frac{d^r}{dt^r} \left(\frac{(t-2)^5}{\frac{15}{28}t^7 - 5t^6 + 18t^5 - 30t^4 + 20t^3 - \frac{32}{7}} \right)^{(r+1)/5} \right]_{t=2} \\ &= \frac{\Gamma(\frac{r+1}{5})}{\Gamma(r+1)} \left[\frac{d^r}{dt^r} \left(\frac{1}{\frac{15}{28}t^2 + \frac{5}{2}t + 3} \right)^{(r+1)/5} \right]_{t=0} \\ &= \frac{(5/28)^{r/2}}{3^{(r+1)/5}} \Gamma\left(\frac{r+1}{5}\right) C_r^{(\frac{r+1}{5})} \left(-\sqrt{\frac{35}{36}}\right) , \end{aligned}$$

and the other coefficients are defined via $T_r^{(m)}(\alpha_m) = e^{2\pi i \alpha_m (r+1) / \omega_m} T_r^{(m)}(0)$.




FIG. 5. (a) Steepest descent paths in the complex t -plane passing through the third order saddle $t^{(1)}$ ($\omega_1 = 3$) and the fifth order saddle $t^{(2)}$ ($\omega_2 = 5$) between labeled valleys V_j , $j = 1, 2, \dots, 6$, at infinity for $\theta = -\frac{\pi}{4}$. The path of integration chosen is $\mathcal{P}^{(1)}(-\frac{\pi}{4}, 0)$ which runs between $t^{(1)}$ and V_3 . (b) The rotated steepest descent path $\mathcal{P}^{(1)}(0, 0)$ emerging from $t^{(1)}$ connects with $t^{(2)}$ at the Stokes phenomenon $\theta_{12}^+ = 0$. The bold lines are the steepest paths that are used in the Level 1 hyperasymptotic expansion about $t^{(1)}$ (33), (6). (c) The steepest descent path $\mathcal{P}^{(2)}(9\pi, \alpha_2)$, emerging from $t^{(2)}$ connects with $t^{(1)}$ at the Stokes phenomenon $\theta_{121}^+ = 9\pi$. The bold lines are the steepest paths that are used in the Level 2 hyperasymptotic expansion about $t^{(1)}$ (36), (6) (or Level 1 hyperasymptotic expansion about $t^{(2)}$).

For the singulant we have

$$|\mathcal{F}_{12}^+| = |f(2) - f(0)| = \frac{32}{7}.$$

The effective asymptotic parameter in the expansion is thus $|z\mathcal{F}_{12}^+| = \frac{32}{7}$, and hence the optimal number of terms in

$$(48) \quad T^{(1)}(z; \alpha_1) = \sum_{r=0}^{N_0^{(1)}-1} \frac{T_r^{(1)}(\alpha_1)}{z^{r/3}} + R_1^{(1)}(z; \alpha_1)$$

is $N_0^{(1)} = \lceil |z\mathcal{F}_{12}^+| \omega_1 \rceil = 13$.

We will focus again on $T^{(1)}(z; 0)$ and give only the main details, which are

$$\theta_{12}^+ = 0, \quad \theta_{121}^+ = 9\pi, \quad \theta_{1212}^+ = 14\pi, \quad \alpha_{12}^+ = 2, \quad \alpha_{121}^+ = 4, \quad \alpha_{1212}^+ = 9.$$

When we compute this integral numerically for this value of z with high precision, we obtain

$$T^{(1)}(z, 0) = 1.244081553113296 + 0.145693991003805i.$$

The numerics of the hyperasymptotic approximations are given in Table 2, and for the Level 2 expansion we display the terms and errors in Figure 6. We observe that this time the remainders in the original Poincaré expansion are considerably larger than the first neglected terms, again, as predicted in section 5. However, in the higher levels the remainders are again of a similar size as the first neglected terms.

8. Example 3: Swallowtail and the adjacency of the saddles. In this example we apply hyperasymptotic techniques to determine the relative adjacency, and hence which saddles would contribute to the exact remainder terms of an expansion, using algebraic, rather than geometric means. We choose to illustrate this using the swallowtail integral [18, sect. 36.2].

TABLE 2

The numbers of terms required to minimize the absolute error at each level of the hyperasymptotic reexpansions for the (3 → 5) degenerate example.

Level	$N_0^{(1)}$	$N_1^{(2)}$	$N_2^{(1)}$	$N_3^{(1)}$	Error
0	13				6.9×10^{-3}
1	27	22			3.7×10^{-7}
2	41	45	13		2.0×10^{-10}
3	54	68	27	22	1.1×10^{-13}




FIG. 6. For Example 2: The modulus of the n th term in the Level 2 hyperasymptotic expansion (blue dots), and the modulus of the remainder after taking n terms in the approximation (red crosses).

For the swallowtail integral the bifurcation set is given in [18, eqn. 36.4.7] and with the notation in this reference we take $t = \frac{1}{2}i - \frac{1}{4}$ and $z = \frac{5}{6}i - \frac{25}{8}$. (The choice of complex parameters is to force one of the saddles to be nonadjacent; see below.)

The resulting semi-infinite contour integral that we will study is again integral (1), but now with $g(t) \equiv 1$ and

$$f(t) = t^5 + \frac{5}{24}(4i - 15)t^3 + \frac{45}{16}(2i - 1)t^2 + \frac{5}{256}(101 + 168i)t.$$

The saddle points are $t^{(1)} = \frac{7}{4} - \frac{1}{2}i$, $t^{(2)} = -\frac{5}{4} - \frac{1}{2}i$, and $t^{(3)} = \frac{1}{2}i - \frac{1}{4}$, with $\omega_1 = \omega_2 = 2$ and $\omega_3 = 3$. Once again, the polynomial form of $f(t)$ means that we may take $z = e^{i\theta}$ with the choice of $\theta = -\frac{\pi}{4}$. To obtain the Level 1 hyperasymptotic approximation we find that

$$|\mathcal{F}_{12}^+| = \frac{9\sqrt{109}}{4}, \quad |\mathcal{F}_{13}^+| = \frac{125\sqrt{5}}{12}, \quad \theta_{12}^+ = 3\pi - \arctan \frac{10}{3}, \quad \theta_{13}^+ = 3\pi - \arctan \frac{278}{29}.$$

It follows that $\alpha_{12}^+ = 1$ and $\alpha_{13}^+ = 0$. We write the Level 1 hyperasymptotic approximation as

(49)

$$T^{(1)}(z; 0) = \sum_{r=0}^{N-1} \frac{T_r^{(n)}(0)}{z^{r/2}} + K_{12} \frac{z^{(1-N)/2}}{4\pi i} \sum_{r=0}^{N_1^{(2)}-1} \mathbf{T}_r^{(2)}(1) \mathbf{F}^{(1)} \left(z; \begin{matrix} \frac{N+1}{2} - \frac{r+1}{2} \\ 2 \\ |\mathcal{F}_{12}^+| e^{i(\pi-\theta_{12}^+)} \end{matrix} \right) \\ + K_{13} \frac{z^{(1-N)/2}}{6\pi i} \sum_{r=0}^{N_1^{(3)}-1} \mathbf{T}_r^{(3)}(0) \mathbf{F}^{(1)} \left(z; \begin{matrix} \frac{N+1}{2} - \frac{r+1}{3} \\ 2 \\ |\mathcal{F}_{13}^+| e^{i(\pi-\theta_{13}^+)} \end{matrix} \right) + R_1^{(1)}(z; 0).$$

Note that we have here introduced unknown constant prefactors K_{nm} into the expression for the Level 1 hyperasymptotic expansion (33). Each constant will be equal to 1 if the saddles $t^{(n)}$ and $t^{(m)}$ are adjacent and zero otherwise. We could determine these constants by examining how the steepest descent contours deform as θ is varied. However, here we illustrate their algebraic calculation. These constants appear in the late term expansion (39) (which also follows from (49)) as follows:

$$T_N^{(1)}(0) = \frac{K_{12}}{4\pi i} \sum_{r=0}^{N_1^{(2)}-1} \mathbf{T}_r^{(2)}(1) \frac{e^{i\theta_{12}^+(\frac{N+1}{2} - \frac{r+1}{2})} \Gamma(\frac{N+1}{2} - \frac{r+1}{2})}{|\mathcal{F}_{12}^+|^{\frac{N+1}{2} - \frac{r+1}{2}}} + \frac{K_{13}}{6\pi i} \sum_{r=0}^{N_1^{(3)}-1} \mathbf{T}_r^{(3)}(0) \frac{e^{i\theta_{13}^+(\frac{N+1}{2} - \frac{r+1}{3})} \Gamma(\frac{N+1}{2} - \frac{r+1}{3})}{|\mathcal{F}_{13}^+|^{\frac{N+1}{2} - \frac{r+1}{3}}} + \tilde{R}_1^{(1)}(N; 0).$$

In this (asymptotic) expression, everything is known except K_{12} and K_{13} . Hence if we take two high orders $N = 50$ and $N = 51$ and set $\tilde{R}_1^{(1)}(N; 0) = 0$ we obtain two linear algebraic equations with two unknowns. The optimal numbers of terms on the right-hand side may be calculated from (31) and are $N_1^{(2)} = 7$ and $N_1^{(3)} = 11$. Hence we can solve this simultaneous set of equations to obtain numerical approximations for K_{12} and K_{13} as

$$K_{12} = -0.00123 + 0.00095i, \quad K_{13} = 1.00076 + 0.00060i.$$

Given that the K_{nm} are quantized as integers, within the limits of the errors at this stage, we may infer that $K_{12} = 0$ and $K_{13} = 1$.

Hence we may assert that $t^{(3)}$ is adjacent to $t^{(1)}$, but $t^{(2)}$ is not. This may be confirmed geometrically by consideration of the steepest paths.

9. Discussion. The main results of this paper are the exact remainder terms (12), (26), the hyperasymptotic reexpansions (33), (36), (38), with novel hyperterminants (27), the asymptotic form for the late coefficients (39), and the improved error bounds for the remainder of an asymptotic expansion involving saddle points (42), degenerate or otherwise. We have illustrated the application of these results to the better-than-exponential asymptotic expansions and calculations of integrals with semi-infinite contours and degenerate saddles.

The results of this paper are more widely applicable, for example, to broadening the class of differential equations for which a hyperasymptotic expansion may be derived using a Borel transform approach. We observe that all the examples in this paper are of the form

$$w(z) = \int_{t^{(1)}}^{\infty} e^{-zf(t)} g(t) dt,$$

in which $f(t)$ and $g(t)$ are polynomials in t . (In fact $g(t) \equiv 1$.) Using computer algebra, it is not difficult to construct the corresponding inhomogeneous linear ordinary differential equations for $w(z)$:

$$(50) \quad \sum_{p=0}^P a_p(z) w^{(p)}(z) = h(z),$$

in which the $a_p(z)$'s and $h(z)$ are polynomials.

For our second example with $(\omega_1, \omega_2) = (3, 5)$ we find $P = 6$, the $a_p(z)$'s are polynomials of order 9, and $h(z)$ is of order 6. Integrals involving combinations pairs of the contours $\mathcal{P}^{(n)}$ are solutions of the homogeneous version of (50).

In that example, for the first saddle point we have $\omega_1 = 3$, and hence, there are 2 independent double infinite integrals through this saddle, and for the second saddle point we have $\omega_2 = 5$, and hence, there are 4 independent double infinite integrals through the second saddle. Thus, $P = 2 + 4$.

The differential equation (50) has an irregular singularity of rank one at infinity, but we are dealing with the exceptional cases. That is, the solutions all have initial terms proportional to $\exp(\lambda_p z) z^{\mu_p}$ but now with coinciding λ_p 's. For example, in our second example we have two distinct solutions with $\lambda_1 = \lambda_2 = 0$ and four other different solutions but each with $\lambda_3 = \lambda_4 = \lambda_5 = \lambda_6 = \frac{32}{7}$.

Note also that $h(z)$ in (50) is a polynomial in z . Hence we should expect a particular integral of (50) to involve only integer powers of z . However, the particular integral $w(z) = z^{-1/3} T^{(1)}(z; 0)$ has, according to (48), an asymptotic expansion in inverse powers of $z^{1/3}$. The resolution of this paradox is that the combination of such solutions

$$w(z) = \frac{z^{-1/3}}{3} \left(T^{(1)}(z; 0) + T^{(1)}(z; 1) + T^{(1)}(z; 2) \right)$$

is itself a particular integral but contains only integer powers. This solution involves a star-shaped contour of integration, typically not studied if the problem is posed in terms of integrals alone.

We also remark that differential equations of the form (50) will give us recurrence relations for the coefficients in the asymptotic expansions, and these are, of course, much more efficient than our formula (11).

Appendix A. Computation of the generalized hyperterminants. In this appendix we relate the generalized hyperterminants (27) to the simpler ones given in [38] and thereby develop an efficient method to calculate them.

First, the following theorem improves on the main theorem in [38].

THEOREM A.1. *For $k \geq 0$, $|\arg z + \arg \sigma_0| < \pi$ and $0 < \arg \sigma_j - \arg \sigma_{j-1} < 2\pi$, $j \geq 1$, $\text{Re}(M_1) > 2$ and $\text{Re}(M_j) > 1$, $j \neq 1$, we have the convergent expansion*

$$F^{(k+1)} \left(z; \begin{matrix} M_0, \dots, M_k \\ \sigma_0, \dots, \sigma_k \end{matrix} \right) = \sum_{n=0}^{\infty} A^{(k+1)} \left(n; \begin{matrix} M_0, \dots, M_k \\ \sigma_0, \dots, \sigma_k \end{matrix} \right) U(n+1, 2-M_0, z\sigma_0),$$

where

$$\begin{aligned} A^{(1)} \left(n; \begin{matrix} M_0 \\ \sigma_0 \end{matrix} \right) &= \delta_{n,0} e^{M_0 \pi i} \sigma_0^{1-M_0} \Gamma(M_0), \\ A^{(2)} \left(n; \begin{matrix} M_0, & M_1 \\ \sigma_0, & \sigma_1 \end{matrix} \right) &= -e^{\pi M_0 i} \sigma_0^{2-M_0-M_1} \left(e^{-\pi i} \frac{\sigma_1}{\sigma_0} \right)^{n-M_1+1} \Gamma(M_0+n) \Gamma(M_1) \\ &\quad \times \frac{n! \Gamma(M_0+M_1-1)}{\Gamma(M_0+M_1+n)} {}_2F_1 \left(\begin{matrix} M_0+n, n+1 \\ M_0+M_1+n \end{matrix}; 1 + \frac{\sigma_1}{\sigma_0} \right), \end{aligned}$$

and when $k \geq 1$,

$$A^{(k+1)} \left(n; \begin{matrix} M_0, \dots, M_k \\ \sigma_0, \dots, \sigma_k \end{matrix} \right) = e^{\pi M_0 i} \sigma_0^{1-M_0} \left(e^{-\pi i} \frac{\sigma_1}{\sigma_0} \right)^n \Gamma(M_0 + n) \Gamma(M_0 + M_1 - 1) \\ \times \sum_{m=0}^{\infty} \frac{(n+m)! A^{(k)} \left(m; \begin{matrix} M_1, \dots, M_k \\ \sigma_1, \dots, \sigma_k \end{matrix} \right)}{m! \Gamma(M_0 + M_1 + n + m)} {}_2F_1 \left(\begin{matrix} M_0 + n, n + m + 1 \\ M_0 + M_1 + n + m \end{matrix}; 1 + \frac{\sigma_1}{\sigma_0} \right).$$

Here ${}_2F_1$ stands for the hypergeometric function [18, sect. 15.2].

The proof of this theorem is very similar to the one for Theorem 2 in [38]. The main difference here is that we must be more careful with the definitions of the phases and use the restrictions $0 < \arg \sigma_j - \arg \sigma_{j-1} < 2\pi$. This removes any phase-related ambiguity in the calculation of the hyperterminants.

Note that the representation of $A^{(k+1)}$ is a convergent infinite series. In a practical implementation it is necessary to truncate the series appropriately. In our numerical examples we took 40 terms of the convergent series and checked, by taking successively more terms, that this truncation gave us sufficient correct digits in the evaluation of the corresponding hyperterminants.

In the theorem above we also require, for example, that $0 < \arg \sigma_1 - \arg \sigma_0 < 2\pi$. In fact, one often encounters the case $\sigma_0 = \sigma_1$ and some care is then needed to evaluate the ${}_2F_1$. Numerical methods to evaluate the (confluent) hypergeometric function may be found in [21].

With these phase clarifications, the generalized hyperterminants (27) can be expressed in terms of the ones above as follows.

First, by rationalization, we have

$$\mathbf{F}^{(k+1)} \left(\begin{matrix} M_0, \dots, M_k \\ z; \omega_0, \dots, \omega_k \\ \sigma_0, \dots, \sigma_k \end{matrix} \right) \\ = \sum_{\ell_0=0}^{\omega_0-1} z^{1-(\ell_0+1)/\omega_0} \int_0^{[\pi-\arg \sigma_0]} \dots \int_0^{[\pi-\arg \sigma_k]} \prod_{j=1}^k \frac{e^{\sigma_0 t_0} t_0^{M_0+\ell_0/\omega_0-1}}{z-t_0} \\ \times \sum_{\ell_j=0}^{\omega_j-1} \frac{e^{\sigma_j t_j} t_j^{1-(\ell_j+1)/\omega_j} t_j^{M_j+\ell_j/\omega_j-1}}{t_{j-1}-t_j} dt_k \dots dt_0 \\ = \sum_{\ell_0=0}^{\omega_0-1} \dots \sum_{\ell_k=0}^{\omega_k-1} z^{1-(\ell_0+1)/\omega_0} \int_0^{[\pi-\arg \sigma_0]} \dots \int_0^{[\pi-\arg \sigma_k]} \frac{e^{\sigma_0 t_0} t_0^{M_0+\ell_0/\omega_0-(\ell_1+1)/\omega_1}}{z-t_0} \\ \times \left(\prod_{j=1}^{k-1} \frac{e^{\sigma_j t_j} t_j^{M_j+\ell_j/\omega_j-(\ell_{j+1}+1)/\omega_{j+1}}}{t_{j-1}-t_j} \right) \frac{e^{\sigma_k t_k} t_k^{M_k+\ell_k/\omega_k-1}}{t_{k-1}-t_k} dt_k \dots dt_0.$$

We make the changes of integration variables from t_0 to s_0 and from t_j to s_j ($1 \leq j \leq k$) via $t_0 = s_0 e^{2\pi\gamma_0 i}$ and $t_j = s_j e^{2\pi(\gamma_{j-1}+\gamma_j) i}$. Here, the integers γ_0 and γ_j are chosen so that $|\arg z + \arg \sigma_0 + 2\pi\gamma_0| < \pi$ and $0 < \arg \sigma_j - \arg \sigma_{j-1} + 2\pi\gamma_j < 2\pi$.

Thus, we can finally relate the $\mathbf{F}^{(k+1)}$ to the $F^{(k+1)}$ with the result:

$$\begin{aligned} & \mathbf{F}^{(k+1)} \left(\begin{matrix} M_0, & \dots, & M_k \\ z; & \omega_0, & \dots, & \omega_k \\ & \sigma_0, & \dots, & \sigma_k \end{matrix} \right) \\ &= \sum_{\ell_0=0}^{\omega_0-1} \dots \sum_{\ell_k=0}^{\omega_k-1} z^{1-(\ell_0+1)/\omega_0} e^{2\pi i \left(\gamma_{k-1} (M_{k-1} + M_k + \frac{\ell_{k-1}}{\omega_{k-1}} - \frac{1}{\omega_k}) + \gamma_k (M_k + \frac{\ell_k}{\omega_k}) \right)} \\ & \quad \times \prod_{j=0}^{k-2} e^{2\pi i \gamma_j (M_j + M_{j+1} + \frac{\ell_j}{\omega_j} - \frac{1}{\omega_{j+1}} - \frac{\ell_{j+2} + 1}{\omega_{j+2}})} \\ & \quad \times \int_0^{[\pi - \arg \sigma_0 - 2\pi \gamma_0]} \dots \int_0^{[\pi - \arg \sigma_k - 2\pi (\gamma_{k-1} + \gamma_k)]} \frac{e^{\sigma_0 s_0 + \dots + \sigma_k s_k} s_0^{M_0 + \ell_0 / \omega_0 - (\ell_1 + 1) / \omega_1}}{z - s_0} \\ & \quad \times \left(\prod_{j=1}^{k-1} \frac{s_j^{M_j + \ell_j / \omega_j - (\ell_{j+1} + 1) / \omega_{j+1}}}{s_{j-1} - s_j} \right) \frac{s_k^{M_k + \ell_k / \omega_k - 1}}{s_{k-1} - s_k} ds_k \dots ds_0 \\ &= \sum_{\ell_0=0}^{\omega_0-1} \dots \sum_{\ell_k=0}^{\omega_k-1} z^{1-(\ell_0+1)/\omega_0} e^{2\pi i \left(\gamma_{k-1} (M_{k-1} + M_k + \frac{\ell_{k-1}}{\omega_{k-1}} - \frac{1}{\omega_k}) + \gamma_k (M_k + \frac{\ell_k}{\omega_k}) \right)} \\ & \quad \times \prod_{j=0}^{k-2} e^{2\pi i \gamma_j (M_j + M_{j+1} + \frac{\ell_j}{\omega_j} - \frac{1}{\omega_{j+1}} - \frac{\ell_{j+2} + 1}{\omega_{j+2}})} \\ & \quad \times F^{(k+1)} \left(z; \begin{matrix} M_0 + \frac{\ell_0}{\omega_0} - \frac{\ell_1 + 1}{\omega_1} + 1, & M_1 + \frac{\ell_1}{\omega_1} - \frac{\ell_2 + 1}{\omega_2} + 1, & \dots, & M_k + \frac{\ell_k}{\omega_k} \\ \sigma_0 e^{2\pi i \gamma_0 i}, & \sigma_1 e^{2\pi i (\gamma_0 + \gamma_1) i}, & \dots, & \sigma_k e^{2\pi i (\gamma_{k-1} + \gamma_k) i} \end{matrix} \right). \end{aligned}$$

Appendix B. Bounds for the generalized first-level hyperterminant.

PROPOSITION B.1. *For any positive real M and positive integer ω , we have*

$$\left| \frac{z^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(z; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) \right| \leq \begin{cases} 1 & \text{if } |\theta| \leq \frac{\pi}{2}\omega, \\ \min \left(\left| \csc \left(\frac{\theta}{\omega} \right) \right|, \omega \sqrt{e \left(M + \frac{1}{2} \right)} \right) & \text{if } \frac{\pi}{2}\omega < |\theta| \leq \pi\omega, \\ \frac{\omega \sqrt{2\pi M}}{|\cos \theta|^M} + \omega \sqrt{e \left(M + \frac{1}{2} \right)} & \text{if } \pi\omega < |\theta| < \pi\omega + \frac{\pi}{2}. \end{cases}$$

If $\omega = 1$, the quantity $\sqrt{e(M + \frac{1}{2})}$ can be replaced by

$$(51) \quad \sqrt{\pi} \frac{\Gamma(\frac{M}{2} + 1)}{\Gamma(\frac{M}{2} + \frac{1}{2})} + 1,$$

which is asymptotic to $\sqrt{\frac{\pi}{2}(M + \frac{1}{2})}$ as $M \rightarrow \infty$ and hence yields a sharper bound for large M .

Proof. The case $\omega = 1$ was proved in a recent paper by Nemes [36, Propositions B.1 and B.3]. For the general case, let M be any positive real number and ω be any positive integer. The integral representation of the first generalized hyperterminant can be rewritten

$$(52) \quad \frac{z^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(z; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) = \frac{e^{\pi M i}}{\Gamma(M)} \int_0^\infty \frac{e^{-t} t^{M-1}}{1 + (t/z)^{1/\omega}} dt,$$

provided that $|\theta| < \pi\omega$. For $t \geq 0$, we have

$$(53) \quad \left|1 + \frac{t}{w}\right| \geq \begin{cases} 1 & \text{if } |\arg w| \leq \frac{\pi}{2}, \\ |\sin(\arg w)| & \text{if } \frac{\pi}{2} < |\arg w| < \pi, \end{cases}$$

and therefore

$$\begin{aligned} \left| \frac{z^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(z; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) \right| &\leq \frac{1}{\Gamma(M)} \int_0^\infty \frac{e^{-t} t^{M-1}}{|1 + (t/z)^{1/\omega}|} dt \\ &\leq \begin{cases} 1 & \text{if } |\theta| \leq \frac{\pi}{2}\omega, \\ |\csc(\frac{\theta}{\omega})| & \text{if } \frac{\pi}{2}\omega < |\theta| < \pi\omega. \end{cases} \end{aligned}$$

We continue by showing that the absolute value of the left-hand side of (52) is bounded by $\omega\sqrt{e^{(M+1/2)}}$ when $\frac{\pi}{2}\omega < \theta \leq \pi\omega$. (The analogous bound for the range $-\pi\omega \leq \theta < -\frac{\pi}{2}\omega$ follows by taking complex conjugates.) For this purpose, we deform the contour of integration in (52) by rotating it through an acute angle φ . Thus, by appealing to Cauchy’s theorem and analytic continuation, we have, for arbitrary $0 < \varphi < \frac{\pi}{2}$, that

$$\frac{z^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(z; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) = \frac{e^{\pi Mi}}{\Gamma(M)} \left(\frac{e^{i\varphi}}{\cos \varphi} \right)^M \int_0^\infty \frac{e^{-\frac{e^{i\varphi} u}{z \cos \varphi}} u^{M-1}}{1 + \left(\frac{e^{i\varphi} u}{z \cos \varphi} \right)^{1/\omega}} du$$

when $\frac{\pi}{2}\omega < \theta \leq \pi\omega$. Employing the inequality (53), we find that

$$\begin{aligned} \left| \frac{z^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(z; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) \right| &\leq \frac{1}{\Gamma(M)} \frac{1}{\cos^M \varphi} \int_0^\infty \frac{e^{-u} u^{M-1}}{\left| 1 + \left(\frac{e^{i\varphi} u}{z \cos \varphi} \right)^{1/\omega} \right|} du \\ &\leq \frac{1}{\cos^M \varphi} \times \begin{cases} 1 & \text{if } \frac{\pi}{2}\omega < \theta \leq \frac{\pi}{2}\omega + \varphi, \\ |\csc(\frac{\theta-\varphi}{\omega})| & \text{if } \frac{\pi}{2}\omega + \varphi < \theta \leq \pi\omega. \end{cases} \end{aligned}$$

We now choose the value of φ which approximately minimizes the right-hand side of this inequality when $\theta = \pi\omega$, namely, $\varphi = \arctan(M^{-1/2})$. We may then claim that

$$\frac{1}{\cos^M(\arctan(M^{-1/2}))} = \left(1 + \frac{1}{M}\right)^{M/2} \leq \omega\sqrt{e\left(M + \frac{1}{2}\right)}$$

when $\frac{\pi}{2}\omega < \theta \leq \frac{\pi}{2}\omega + \arctan(M^{-1/2})$, where the last inequality can be obtained by means of elementary analysis. In the remaining case $\frac{\pi}{2}\omega + \arctan(M^{-1/2}) < \theta \leq \pi\omega$, we have

$$\begin{aligned} \frac{\left| \csc\left(\frac{\theta - \arctan(M^{-1/2})}{\omega}\right) \right|}{\cos^M(\arctan(M^{-1/2}))} &\leq \frac{\left| \csc\left(\pi - \frac{\arctan(M^{-1/2})}{\omega}\right) \right|}{\cos^M(\arctan(M^{-1/2}))} \\ &= \left(1 + \frac{1}{M}\right)^{M/2} \csc\left(\frac{\arctan(M^{-1/2})}{\omega}\right) \leq \left(1 + \frac{1}{M}\right)^{M/2} \omega \csc(\arctan(M^{-1/2})) \\ &= \omega \left(1 + \frac{1}{M}\right)^{(M+1)/2} \sqrt{M} \leq \omega\sqrt{e\left(M + \frac{1}{2}\right)}. \end{aligned}$$

Here we have used the convexity of $\csc(x)$ for $0 < x < \frac{\pi}{2}$ and that the quantity $(1 + \frac{1}{M})^{(M+1)/2} \sqrt{\frac{M}{M+a}}$, as a function of $M > 0$, increases monotonically if and only if $a \geq \frac{1}{2}$, in which case it has limit \sqrt{e} .

We finish by proving the claimed bound for the range $\pi\omega < |\theta| < \pi\omega + \frac{\pi}{2}$. It is sufficient to consider the range $\pi\omega < \theta < \pi\omega + \frac{\pi}{2}$, as the estimates for $-\pi\omega - \frac{\pi}{2} < \theta < -\pi\omega$ can be derived by taking complex conjugates. The proof is based on the functional relation

$$\frac{z^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(z; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) = \frac{2\pi i \omega (ze^{-\pi i(\omega-1)})^M}{\Gamma(M)e^{ze^{-\pi i\omega}}} + \frac{(ze^{-2\pi i\omega})^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(ze^{-2\pi i\omega}; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right)$$

(see [43, eqn. (A.13)]). From this functional relation, we can infer that

$$\begin{aligned} \left| \frac{z^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(z; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) \right| &\leq \frac{2\pi\omega |z|^M}{\Gamma(M)e^{|z|\cos\theta|}} + \left| \frac{(ze^{-2\pi i\omega})^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(ze^{-2\pi i\omega}; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) \right| \\ &\leq \frac{2\pi\omega |z|^M}{\Gamma(M)e^{|z|\cos\theta|}} + \omega \sqrt{e \left(M + \frac{1}{2} \right)}. \end{aligned}$$

Notice that the quantity $r^M e^{-ra}$, as a function of $r > 0$, takes its maximum value at $r = M/a$ when $a > 0$ and $M > 0$. We therefore find that

$$\begin{aligned} \left| \frac{z^{1/\omega}}{\Gamma(M)} \mathbf{F}^{(1)} \left(z; \begin{matrix} M \\ \omega \\ 1 \end{matrix} \right) \right| &\leq \frac{\omega \sqrt{2\pi M} M^{M-1/2} e^{-M} \sqrt{2\pi}}{|\cos\theta|^M \Gamma(M)} + \omega \sqrt{e \left(M + \frac{1}{2} \right)} \\ &\leq \frac{\omega \sqrt{2\pi M}}{|\cos\theta|^M} + \omega \sqrt{e \left(M + \frac{1}{2} \right)}. \end{aligned}$$

The second inequality can be obtained from the inequality $M^{M-1/2} e^{-M} \sqrt{2\pi} \leq \Gamma(M)$ for any $M > 0$ (see, for instance, [18, eqn. 5.6.1]). \square

Appendix C. The boundary of the domain $\Delta^{(n)}$. In this appendix, we prove that the boundary of $\Delta^{(n)}$ can be written as a union $\bigcup_m \mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+) \cup -\mathcal{P}^{(m)}(\theta_{nm}^-, \alpha_{nm}^-)$, where $\mathcal{P}^{(m)}(\theta_{nm}^\pm, \alpha_{nm}^\pm)$ are steepest descent paths emerging from the adjacent saddle $t^{(m)}$ (see Figure 2(b)). For α_{nm}^\pm , see (8).

First, we show that as we change θ , the steepest descent path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ varies smoothly, unless, perhaps, it encounters an adjacent saddle point $t^{(m)}$. To see this, consider the map $s(t)$ between the t -plane and the s -surface, defined by

$$s = f(t) - f_n.$$

The steepest descent path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ is mapped into a half line with phase $2\pi\alpha_n - \theta$ emerging from the origin as an ω_n th-order branch point on the s -surface. As this half line is rotated on the s -surface, the corresponding steepest descent path varies smoothly, unless we encounter a singularity of the inverse map $t(s)$. Since $f(t)$ is holomorphic in the closure of $\Delta^{(n)}$, and $|f(t)| \rightarrow \infty$ as $t \rightarrow \infty$ in $\Delta^{(n)}$, the only singularities of $t(s)$ are branch points located at the images of the saddle points of $f(t)$ under the map $s(t)$. When the half line hits a branch point of $t(s)$ on the s -surface, the corresponding steepest descent path hits a saddle point in the t -plane.

If we rotate θ in the positive direction, the steepest descent path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ runs into a saddle point $t^{(m)}$ when $\theta = \theta_{nm}^+$. Likewise, if we rotate θ in the negative direction, the steepest descent path $\mathcal{P}^{(n)}(\theta; \alpha_n)$ hits a saddle $t^{(m)}$ when $\theta = \theta_{nm}^-$. By definition, the domain $\Delta^{(n)}$ is the union $\bigcup_{\theta \neq \theta_{nm}^\pm} \mathcal{P}^{(n)}(\theta; \alpha_n)$, which is precisely the image of the points on the s -surface that can be seen from the branch point at the origin minus half lines with phases $2\pi\alpha_n - \theta_{nm}^\pm$ issuing from the points $s(t^{(m)})$ under the map $t(s)$. The boundary of the domain $\Delta^{(n)}$ therefore consists of the images of these half lines under the map $t(s)$. It is easy to see that the image of the half line with phase $2\pi\alpha_n - \theta_{nm}^+$ emerging from $s(t^{(m)})$ under the map $t(s)$ is precisely the steepest descent path $\mathcal{P}^{(m)}(\theta_{nm}^+, \alpha_{nm}^+)$ emanating from the adjacent saddle $t^{(m)}$. Similarly, the image of the half line with phase $2\pi\alpha_n - \theta_{nm}^-$ emerging from $s(t^{(m)})$ under the map $t(s)$ is the steepest descent path $\mathcal{P}^{(m)}(\theta_{nm}^-, \alpha_{nm}^-)$ emanating from the adjacent saddle $t^{(m)}$. In order to make the orientation of the domain $\Delta^{(n)}$ positive, the orientation of the steepest path $\mathcal{P}^{(m)}(\theta_{nm}^-, \alpha_{nm}^-)$ has to be reversed.

Acknowledgment. The authors thank the referees for very helpful comments and suggestions for improving the presentation.

REFERENCES

- [1] A. BEHTASH, G. V. DUNNE, T. SCHÄFER, T. SULEJMANPASIC, AND M. ÜNSAL, *Complexified path integrals, exact saddles, and supersymmetry*, Phys. Rev. Lett., 116 (2016), 011601, <https://doi.org/10.1103/PhysRevLett.116.011601>.
- [2] A. BEHTASH, G. V. DUNNE, T. SCHÄFER, T. SULEJMANPASIC, AND M. ÜNSAL, *Toward Picard-Lefschetz theory of path integrals, complex saddles and resurgence*, Ann. Math. Sci. Appl., 2 (2017), pp. 95–212.
- [3] T. BENNETT, *Exponential Asymptotics for Integrals with Degenerate and Non-Isolated Critical Points*, Ph.D. thesis, Mathematical Sciences, University of Southampton, 2015.
- [4] M. BERRY AND C. UPSTILL, IV, *Catastrophe Optics: Morphologies of Caustics and Their Diffraction Patterns*, Progr. Optics 18, Elsevier, New York, 1980, pp. 257–346, [https://doi.org/10.1016/S0079-6638\(08\)70215-4](https://doi.org/10.1016/S0079-6638(08)70215-4).
- [5] M. V. BERRY, *Uniform approximation: A new concept in wave theory*, Science Progress (Oxford), 57 (1969), pp. 43–64, <http://www.jstor.org/stable/43423745>.
- [6] M. V. BERRY, *Uniform asymptotic smoothing of Stokes's discontinuities*, Proc. A, 422 (1989), pp. 7–21.
- [7] M. V. BERRY AND C. J. HOWLS, *Hyperasymptotics for integrals with saddles*, Proc. A, 434 (1991), pp. 657–675, <https://doi.org/10.1098/rspa.1991.0119>.
- [8] M. V. BERRY AND C. J. HOWLS, *Unfolding the high orders of asymptotic expansions with coalescing saddles: Singularity theory, crossover and duality*, Proc. A, 443 (1993), pp. 107–126, <https://doi.org/10.1098/rspa.1993.0134>.
- [9] M. V. BERRY AND C. J. HOWLS, *Overlapping Stokes smoothings: Survival of the error function and canonical catastrophe integrals*, Proc. A, 444 (1994), pp. 201–216, <https://doi.org/10.1098/rspa.1994.0012>.
- [10] J. P. BOYD, *Weakly Nonlocal Solitary Waves and Beyond-All-Orders Asymptotics: Generalized Solitons and Hyperasymptotic Perturbation Theory*, Math. Appl 442, Kluwer, Dordrecht, 1998, <https://doi.org/10.1007/978-1-4615-5825-5>.
- [11] J. P. BOYD, *The devil's invention: Asymptotic, superasymptotic and hyperasymptotic series*, Acta Appl. Math., 56 (1999), pp. 1–98, <https://doi.org/10.1023/A:1006145903624>.
- [12] W. G. C. BOYD, *Error bounds for the method of steepest descents*, Proc. A, 440 (1993), pp. 493–518, <https://doi.org/10.1098/rspa.1993.0030>.
- [13] S. BREEN, *Exponentially-Improved Asymptotics of Single and Multidimensional Integrals*, Ph.D. thesis, School of Mathematical Sciences, Dublin City University, 1999.
- [14] P. V. BUIVIDOVICH, G. V. DUNNE, AND S. N. VALGUSHEV, *Complex path integrals and saddles in two-dimensional gauge theory*, Phys. Rev. Lett., 116 (2016), pp. 132001, <https://doi.org/10.1103/PhysRevLett.116.132001>.
- [15] E. T. COPSON, *Asymptotic Expansions*, Cambridge Tracts in Math. 55, Cambridge University Press, Cambridge, UK, 2004.

- [16] E. DELABAERE AND C. J. HOWLS, *Global asymptotics for multiple integrals with boundaries*, Duke Math. J., 112 (2002), pp. 199–264, <https://doi.org/10.1215/S0012-9074-02-11221-6>.
- [17] R. B. DINGLE, *Asymptotic Expansions: Their Derivation and Interpretation*, Academic Press, New York, 1973.
- [18] F. W. J. OLVER, A. B. OLDE DAALHUIS, D. W. LOZIER, B. I. SCHNEIDER, R. F. BOISVERT, C. W. CLARK, B. R. MILLER, AND B. V. SAUNDERS, eds., *NIST Digital Library of Mathematical Functions*, <http://dlmf.nist.gov>, release 1.0.16 (2017).
- [19] G. V. DUNNE AND M. ÜNSAL, *Generating nonperturbative physics from perturbation theory*, Phys. Rev. D, 89 (2014), 041701, <https://doi.org/10.1103/PhysRevD.89.041701>.
- [20] G. V. DUNNE AND M. ÜNSAL, *Deconstructing zero: Resurgence, supersymmetry and complex saddles*, J. High Energy Phys., 2016 (2016), p. 2, [https://doi.org/10.1007/JHEP12\(2016\)002](https://doi.org/10.1007/JHEP12(2016)002).
- [21] A. GIL, J. SEGURA, AND N. M. TEMME, *Numerical Methods for Special Functions*, SIAM, Philadelphia, 2007, <https://doi.org/10.1137/1.9780898717822>.
- [22] R. GRIMSHAW, *Exponential asymptotics and generalized solitary waves*, in *Asymptotic Methods in Fluid Mechanics: Survey and Recent Advances*, CISM Courses and Lectures, Springer, New York, 2010, pp. 71–120, https://doi.org/10.1007/978-3-7091-0408-8_3.
- [23] C. J. HOWLS, *Hyperasymptotics for integrals with finite endpoints*, Proc. A, 439 (1992), pp. 373–396, <https://doi.org/10.1098/rspa.1992.0156>.
- [24] C. J. HOWLS, *Hyperasymptotics for multidimensional integrals, exact remainder terms and the global connection problem*, Proc. A, 453 (1997), pp. 2271–2294, <https://doi.org/10.1098/rspa.1997.0122>.
- [25] D. S. JONES, *Introduction to Asymptotics: A Treatment Using Nonstandard Analysis*, World Scientific, River Edge, NJ, 1997, <https://doi.org/10.1142/9789812779373>.
- [26] E. LOMBARDI, *Oscillatory Integrals and Phenomena Beyond All Algebraic Orders: With Applications to Homoclinic Orbits in Reversible Systems*, Lecture Notes in Math. 1791, Springer-Verlag, Berlin, 2000, <https://doi.org/10.1007/BFb0104102>.
- [27] C. S. MELJER, *Asymptotische Entwicklungen von Besselschen, Hankelschen und verwandten Funktionen I*, Proc. Kon. Akad. Wet. Amsterdam, 35 (1932), pp. 656–667.
- [28] C. S. MELJER, *Asymptotische Entwicklungen von Besselschen, Hankelschen und verwandten Funktionen II*, Proc. Kon. Akad. Wet. Amsterdam, 35 (1932), pp. 852–866.
- [29] C. S. MELJER, *Asymptotische Entwicklungen von Besselschen, Hankelschen und verwandten Funktionen III*, Proc. Kon. Akad. Wet. Amsterdam, 35 (1932), pp. 948–958.
- [30] C. S. MELJER, *Asymptotische Entwicklungen von Besselschen, Hankelschen und verwandten Funktionen IV*, Proc. Kon. Akad. Wet. Amsterdam, 35 (1932), pp. 1079–1090.
- [31] C. S. MELJER, *Über die asymptotische Entwicklung von $\int_0^{\infty-i(\arg w-\mu)} e^{\nu z-w \sinh z} dz$ ($-\frac{\pi}{2} < \mu < \frac{\pi}{2}$) für grosse Werte von $|w|$ und $|\nu|$* , Proc. Kon. Akad. Wet. Amsterdam, 35 (1932), pp. 1170–1180, 1291–1303.
- [32] C. S. MELJER, *Asymptotische Entwicklungen von Besselschen und Hankelschen Funktionen für große Werte des Arguments und der Ordnung*, Math. Ann., 108 (1933), pp. 321–359, <https://doi.org/10.1007/BF01452841>.
- [33] B. T. M. MURPHY AND A. D. WOOD, *Hyperasymptotic solutions of second-order ordinary differential equations with a singularity of arbitrary integer rank*, Methods Appl. Anal., 4 (1997), pp. 250–260, <https://doi.org/10.4310/MAA.1997.v4.n3.a2>.
- [34] G. NEMES, *An explicit formula for the coefficients in Laplace’s method*, Constr. Approx., 38 (2013), pp. 471–487, <https://doi.org/10.1007/s00365-013-9202-6>.
- [35] G. NEMES, *The Role of Resurgence in the Theory of Asymptotic Expansions*, Ph.D. thesis, Department of Mathematics and Its Applications, Central European University, 2015.
- [36] G. NEMES, *Error bounds for the large-argument asymptotic expansions of the Hankel and Bessel functions*, Acta Appl. Math., 150 (2017), pp. 141–177, <https://doi.org/10.1007/s10440-017-0099-0>.
- [37] A. B. OLDE DAALHUIS, *Hyperasymptotic solutions of higher order linear differential equations with a singularity of rank one*, Proc. A, 454 (1998), pp. 1–29, <https://doi.org/10.1098/rspa.1998.0145>.
- [38] A. B. OLDE DAALHUIS, *Hyperterminants. II*, J. Comput. Appl. Math., 89 (1998), pp. 87–95, [https://doi.org/10.1016/S0377-0427\(97\)00220-3](https://doi.org/10.1016/S0377-0427(97)00220-3).
- [39] A. B. OLDE DAALHUIS, *On the asymptotics for late coefficients in uniform asymptotic expansions of integrals with coalescing saddles*, Methods Appl. Anal., 7 (2000), pp. 727–745.
- [40] A. B. OLDE DAALHUIS AND F. W. J. OLVER, *Exponentially improved asymptotic solutions of ordinary differential equations. II. Irregular singularities of rank one*, Proc. A, 445 (1994), pp. 39–56, <https://doi.org/10.1098/rspa.1994.0047>.
- [41] A. B. OLDE DAALHUIS AND F. W. J. OLVER, *Hyperasymptotic solutions of second-order linear differential equations. I*, Methods Appl. Anal., 2 (1995), pp. 173–197.

- [42] R. B. PARIS, *Hadamard Expansions and Hyperasymptotic Evaluation: An Extension of the Method of Steepest Descents*, Encyclopedia Math. Appl. 141, Cambridge University Press, Cambridge, UK, 2011, <https://doi.org/10.1017/CBO9780511753626>.
- [43] R. B. PARIS, *Exponentially small expansions of the Wright function on the Stokes lines*, Lith. Math. J., 54 (2014), pp. 82–105, <https://doi.org/10.1007/s10986-014-9229-9>.
- [44] T. POSTON AND I. STEWART, *Catastrophe Theory and Its Applications*, Dover, Mineola, NY, 1996.
- [45] H. SEGUR, S. TANVEER, AND H. LEVINE, EDS., *Asymptotics Beyond All Orders*, NATO Adv. Sci. Inst. Ser. B Phy. 284, Plenum Press, New York, 1991, <https://doi.org/10.1007/978-1-4757-0435-8>.
- [46] G. G. STOKES, *On the discontinuity of arbitrary constants which appear in divergent developments*, Trans. Cambridge Philosophical Society, 10 (1857), pp. 105–128.
- [47] L. N. TREFETHEN AND J. A. C. WEIDEMAN, *The exponentially convergent trapezoidal rule*, SIAM Rev., 56 (2014), pp. 385–458, <https://doi.org/10.1137/130932132>.
- [48] M. J. WARD AND M.-C. KROPINSKI, *Asymptotic methods for PDE problems in fluid mechanics and related systems with strong localized perturbations in two-dimensional domains*, in *Asymptotic Methods in Fluid Mechanics: Survey and Recent Advances*, CISM Courses and Lectures, Springer, New York, 2010, pp. 23–70, https://doi.org/10.1007/978-3-7091-0408-8_2.