



Kent Academic Repository

Pearce, Kyle, Borys, Andryj M., Clark, Ewan R. and Shepherd, Helena J. (2018) *Exploring the reactivity of donor-stabilised phosphonium cations: Lewis acid catalysed reduction of chlorophosphanes by silanes*. *Inorganic Chemistry*, 57 (18). pp. 11530-11536. ISSN 0020-1669.

Downloaded from

<https://kar.kent.ac.uk/68938/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1021/acs.inorgchem.8b01578>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Exploring the reactivity of donor-stabilised phosphonium cations: Lewis acid catalysed reduction of chlorophosphanes by silanes

Kyle G. Pearce, Andryj M. Borys, Ewan R. Clark* and Helena J. Shepherd

School of Physical Sciences, Ingram Building, University of Kent, Canterbury, Kent, CT2 7NH, United Kingdom of Great Britain and Northern Ireland

ABSTRACT: Phosphane-stabilised phosphonium cations react with silanes to effect either reduction to primary or secondary phosphanes, or formation of P-P bonded species depending upon counter-anion. This operates for *in situ* generated phosphonium cations, allowing catalytic reduction of P(III)-Cl bonds in the absence of strong reducing agents. Anion and substituent dependence studies have allowed insight into the competing mechanisms involved.

Introduction

Organophosphorus species find use as optoelectronic materials,¹ pharmaceuticals,² ligands,³ and many other applications, and have historically been prepared primarily by reaction of organometallic nucleophiles with chlorophosphane electrophiles.⁴ Complimentary routes have been developed exploiting phosphane reaction with electrophiles,⁵ transition metal catalysed cross-coupling,⁶ or hydrophosphination of unsaturated species.⁷ These processes all rely on the presence of a P-H bond for later functionalisation but P-H species are not generally commercially available for any but the simplest derivatives. P-H species are accessible by reductive cleavage of P-C bonds using alkali metals (Na/NH₃ or Li/THF) followed by aqueous workup.⁸ This approach shows poor functional group tolerance and selectivity in heteroleptic phosphanes, however, so is typically used only with simple, homoleptic phosphane precursors. It has been reported that P-Cl bonds may be reduced under milder conditions using Zn metal.⁹ P-H species are instead typically synthesised by the milder reduction of phosphorus-halogen, phosphorus-oxygen, or phosphorus-nitrogen bonds using stoichiometric reduction by main group metal hydrides, with a single reference in the literature reporting Pd-catalysed reduction of P-Cl bonds under H₂,¹⁰ P=O moieties are resistant to Pd catalysed hydrogenation.¹¹ Aluminium hydride reducing agents are effective at reducing a wide range of P-X bonds (X = halide, OR, O), with reductive coupling to form P-P bonds a common side reaction.¹² The most common such reagent, LiAlH₄, is pyrophoric and its use is made hazardous by the exothermic aqueous work-up which releases dihydrogen as a byproduct. Reductive coupling can be avoided by using the milder reagent DIBAL, but with a significant increase in cost and retention of the hazardous workup.¹³ NaBH₄ has been reported to reduce secondary chlorophosphanes to directly form the protected secondary phosphane-borane adduct,^{14,15} which may then be deprotected if required, but does not reduce other P-X bonds (X = OR, O).¹⁶ Borane itself, BH₃, does not reduce P-Cl bonds, instead forming chlorophosphane-borane adducts which may then be cleanly reduced to P-H species with the protecting group intact,^{17,18} a

mixture of LiAlH₄ and NaBH₄ may also be used to form phosphane-borane adducts, generating the required BH₃ *in situ*.¹⁹

Silanes have been extensively used as mild reducing agents, with and without catalysts,^{20,21,22} for the reduction of P=O bonds to convert phosphane oxides to phosphanes, a reaction driven by the formation of strong Si-O bonds.²³ This has been used not just in phosphane synthesis, but to develop variations of the Wittig,¹¹ Mitsunobu²⁴ and Appel²⁵ reactions which are catalytic in phosphane. Investigation has shown at least two competing mechanisms for this process,^{26,27} both of which rely on the nucleophilicity of the terminal oxygen to drive the reaction. For this reason, more Lewis acidic halosilanes (*e.g.* HSiCl₃, Si₂Cl₆, PhSi(Cl)H₂) are in general more effective reducing agents rather than the more hydridic species as might be expected.²⁸ Unhalogenated silanes therefore require extended reaction times and higher temperatures for less nucleophilic phosphine oxides.²⁵ These silanes are insufficiently reducing, however, to directly reduce P-Cl bonds due to the combined low nucleophilicity of the unactivated Si-H moiety and the reduced thermodynamic driving force of Si-Cl bond formation.


We reasoned that unactivated hydrosilanes should nevertheless react with a sufficiently electrophilic P(III) centre; Vidović²⁹ and Stephan³⁰ recently reported analogous reaction of P(III) dications with silanes. Whilst ligand exchange reactions about donor-stabilised phosphonium cations have been studied in the past,^{31,32} and they have been investigated as ligands in transition metal complexes,^{33,34} comparatively little is known about their other reactivity, in part due to their perceived instability and high Lewis acidity.³⁵ The principle exceptions to this are the N-heterocyclic phosphonium cations where chelation and nitrogen π -donor ligands stabilise the resultant cations to give catalytically useful species.^{36,37,38} For phosphane-stabilised phosphonium cations, the empty p orbital on phosphorus is quenched by donation of a lone pair from a second phosphane. These species may also be regarded as phosphino-phosphonium species³⁹ but as they remain electrophilic at the three-coordinate phosphorus centre due to the low-lying and minimally hindered P-P σ^* orbital, the phosphane-

phosphonium nomenclature is used herein as a better representation of the observed reactivity.

Results and Discussion

To our delight, on reaction of the known adduct of the weak donor ligand **1**, $[\text{Ph}_2(\text{Cl})\text{P}-\text{PPh}_2][\text{GaCl}_4]$,³⁹ with one equivalent of Et_3SiH in PhCl (Scheme 1), the ^{31}P NMR showed immediate P-H bond formation and, after heating at 60°C for 1 hour, clean conversion to $[\text{Ph}_2(\text{H})\text{P}-\text{PPh}_2][\text{GaCl}_4]$, **[3]** GaCl_4 with transformation of Et_3SiH to Et_3SiCl . No immediate reaction was observed on addition of a second equivalent of Et_3SiH , but further heating at 60°C overnight lead to almost complete conversion to Ph_2PH , **4**, with trace formation of $\text{Ph}_2\text{P}-\text{PPh}_2$, **5**; all Et_3SiH was converted to Et_3SiCl with GaCl_3 liberated overall. We subsequently tested the more stable $[\text{Ph}_3\text{P}-\text{PPh}_2][\text{GaCl}_4]$ ³⁹ with Et_3SiH and, while this required 2 hours at 60°C to go to completion, **4** and **5** were formed in 49:1 ratio. When **1** and Et_3SiH were reacted with 25 mol% GaCl_3 (*i.e.* a catalytic loading) in PhCl , immediate formation of **[3]** GaCl_4 was evident, and heating overnight at 100°C gave complete conversion to **4**, implying catalytic behaviour.

Scheme 1: Successive hydride transfer from silane to phosphorus centre.



A range of Lewis acids, silanes, and halophosphanes were screened to probe the scope of this potentially useful catalytic reactivity. The Lewis acids were screened by reaction of **1** and Et_3SiH with an initial 25% loading of Lewis acid and heated for up to 7 days at 100°C , with daily monitoring (Table 1). The exception to this was FeCl_3 , for which a 5% loading was initially tested to avoid issues with paramagnetic broadening in the NMR. Of these results, GaCl_3 was found to be the optimal Lewis acid for P-H bond formation, giving essentially quantitative yields even at 5% catalyst loading. The use of Weakly Coordinating Anions (WCAs) resulted in very different reactivity from that observed for GaCl_3 and AlCl_3 . Me_3SiOTf is an insufficiently strong halide abstraction agent to form **[2]** OTf but, reasoning that a small thermal population may be formed on heating, was nevertheless tested as a potential Lewis acid. Prolonged heating at 100°C lead to clean conversion to a sharp singlet at δ -15.3 ppm, indicating the formation of $\text{Ph}_2\text{P}-\text{PPh}_2$, **5**, and growth of a peak at 4.5 ppm in the ^1H NMR corresponding to the formation of H_2 , for an effective dehydrocoupling reaction (Scheme 2).

Scheme 2: Reductive coupling in the presence of TMSOTf, leading to overall dehydrocoupling



In comparison, at elevated temperatures both NaBAR^{F} and NaBAR^{Cl} (BAR^{F} = tetrakis(3,5-trifluoromethylphenyl)borate, BAR^{Cl} = tetrakis(3,5-dichlorophenyl)borate) give simultaneous dehydrocoupling and P-H bond formation, in direct contrast to the behaviour of OTf , coupled with anion decomposition, either by hydrodehalogenation (BAR^{F}) or protodeboronation (BAR^{Cl}). On heating at 100°C , the BAR^{F} anion undergoes fluo-

ride abstraction, leading to the formation of partially fluorinated phosphane centres and Et_3SiF , confirmed by ^{11}B , ^{19}F and ^{29}Si NMR, but no change in the final ^{11}B NMR spectrum is observed on addition of excess pyridine, indicating an absence of free 3° boron species. In contrast, the BAR^{Cl} system showed almost complete loss of signal intensity in the ^{11}B NMR, indicating protodeboronation and formation of BAR_3 species. Together, these confirm the presence of anion-dependent reaction mechanisms.

Table 1: Screening Lewis acids for catalytic efficacy

		Lewis Acid				
$\text{Ph}_2\text{P}(\text{Cl})\text{PPh}_2$	$+$	Et_3SiH	$\xrightarrow{\text{PhCl, } 100^\circ\text{C}}$	Ph_2PH	$+$	$\text{Ph}_2\text{P}-\text{PPh}_2$
1				4		5
			Conversion ^a			
Lewis Acid	Loading	Time	4	5		
GaCl_3	5%	7 days	> 99%	-		
	10%	5 days	> 99%	-		
	25%	1 day	> 99%	-		
AlCl_3	25%	7 days	61 %	-		
	100%	1 day	92 %	8%		
FeCl_3	5%	7 days	-	7 %		
TMSOTf	25%	5 days	-	82 %		
	100 %	7 days	-	> 99%		
$\text{NaBAR}^{\text{F}^{\text{b}}}$	25%	1 day	68 %	26 %		
$\text{NaBAR}^{\text{Cl}^{\text{b}}}$	25%	1 day	62 %	38 %		

^a. NMR conversion by relative ^{31}P NMR intensity (see ESI for full details). ^b. Anion decomposition observed.

Having identified a suitable Lewis acid and loading, several commercially available silanes were screened as hydride donors (Table 2). At a 5% catalyst loading, Et_3SiH proved the most effective donor, but increasing the catalyst loading could be used to improve yield of **4** with other, cheaper silanes. For Et_3SiH to PHMS, the trend in reactivity follows that predicted by Mayr's nucleophilicity index,^{40,41} but this trend is reversed for Ph_3SiH to PhSiH_3 . This may indicate that the steric hindrance about Si is such that the assumptions about rate of reaction in Mayr's scale are not valid for the very hindered phosphonium electrophiles, as seen for other bulky electrophiles.⁴²

Screening some simple aryl-alkyl and alkyl-alkyl chlorophosphanes revealed significant influence of steric bulk and electron donating substituents on reduction (Table 3). Surprisingly, $\text{Ph}^{\text{tBu}}\text{P}(\text{Cl})\text{PPh}_2$ was reduced more efficiently than the less bulky $\text{Ph}^{\text{nBu}}\text{P}(\text{Cl})\text{PPh}_2$, which even after 7 days at 100°C with an increased 25% catalyst loading showed only a complex, unresolved dynamic mixture in the ^{31}P NMR which is attributed to free and rapid exchange between the many diastereomeric possibilities of $[\text{Ph}^{\text{nBu}}(\text{H})\text{P}-\text{P}^{\text{nBu}}\text{Ph}]^+$ and $[\text{Ph}^{\text{nBu}}\text{P}-\text{P}^{\text{nBu}}\text{Ph}-\text{P}^{\text{nBu}}\text{Ph}]^+$.

Table 2: Screening silanes as hydride donors

Ph ₂ PCl + Silane		Lewis Acid	Ph ₂ PH + Ph ₂ P-PPh ₂ + Chlorosilanes	
1		100 °C up to 7 days	4	5
Silane	Catalyst	Time	Conversion ^a	
			4	5
Et ₃ SiH	GaCl ₃ (5 %)	7 days	> 99%	-
PhMe ₂ SiH	GaCl ₃ (5 %)	7 days	40%	34 %
	AlCl ₃ (100 %)	5 days	87 %	13 %
Me ₂ Si(H)-O-Si(H)Me ₂	GaCl ₃ (5 %)	7 days	31 %	42 %
Me ₂ Si(H)-O-Si(H)Me ₂	GaCl ₃ (25 %)	7 days	97 %	-
PHMS	GaCl ₃ (5 %)	7 days	20 %	32 %
	AlCl ₃ (100 %)	7 days	85 %	12 %
Ph ₃ SiH	GaCl ₃ (5 %)	7 days	19 %	18 %
Ph ₂ SiH ₂	GaCl ₃ (5 %)	7 days	51 %	26 %
PhSiH ₃	GaCl ₃ (5 %)	7 days	82 %	18 %


a. NMR conversion by relative ³¹P NMR intensity.

Addition of excess base at this point caused the spectra to resolve to cleanly give exclusive formation of the *rac*- and *meso*- P-P coupled species, indicating that whilst transfer of the first hydride to form the protio-phosphane-stabilised phosphonium is easily achieved, the transfer of the second is not. Repetition at 50% GaCl₃ loading (*i.e.* preforming the halophosphane-phosphonium) did give a small yield of Ph^{(n)Bu}PH (see ESI) but the dimers remained the dominant product. To probe the influence of anion, the reduction of both Ph^{(n)Bu}PCl and Ph^{(n)Bu}PCl was repeated in the presence of 25 % TMSOTf, and the same phenomena were observed. The reduction of P-N and P-O bonds using a 5% loading of GaCl₃ does proceed but much more slowly and with unwanted side reactions. For the reduction of Ph₂PNⁱPr₂, the consumption of Et₃SiH is coupled with the formation of Et₃SiCl rather than Et₃SiNⁱPr₂, indicating consumption of the GaCl₃ and the formation of less Lewis acidic gallium amido species – this ultimately is not therefore a catalytic reaction, and a different mechanism may be in play, driven by the relative difference in Si-N vs Ga-N bond strengths. On reaction of Ph₂POEt with Et₃SiH in the presence of 5% GaCl₃, slow formation of 4 and 5 is observed, along with Ph₂P(O)Et and Ph₂PEt and a number of unknown by-products, indicating simultaneous competing Arbuzov-type reactivity.

Reduction of primary dichlorophosphanes by Et₃SiH in the presence of catalytic Lewis acid is also achievable, with cyclic species and the *rac*- and *meso*-R(H)P-P(H)R products of incomplete reduction as side products; no R(Cl)P-P(Cl)R were observed. Use of a higher loading of Lewis acid leads to improved yield of primary phosphane and, as before, use of TMSOTf provides only reductive coupling products. PCl₃ reacts rapidly with effervescence and a marked exotherm on

introduction of the GaCl₃ to produce a red precipitate, with PH₃ and P₄ the sole observable species in solution; the red solid was confirmed as a polymeric phosphorus species by chemical testing (see ESI for details). Given the stability of P-N bonds under these reaction conditions, we attempted the reduction of the heteroleptic species Ph(ⁱPr₂N)PCl. Slow cyclisation was observed with a 5% GaCl₃ loading, but increasing the loading to 50% (*i.e.* preforming the phosphane-phosphonium) lead to rapid reaction at room temperature to form [(ⁱPr₂N)Ph(H)-PPh(NⁱPr₂)]⁺ as the dominant species with small quantities of *rac*- and *meso*-(ⁱPr₂N(Ph)P)₂, and cyclic byproducts. On heating to 100 °C, however, the protio-phosphane-stabilised phosphonium decomposed into a complex mixture and no Ph(ⁱPr₂N)PH was observed.

Table 3: Reduction of secondary P-Cl, P-O, and P-N bonds



			Conversion ^a	
Substrate	Catalyst	Time	R ₂ PH	R ₂ P-PR ₂
Ph ₂ PCl	none	3 days	-	trace
Ph ₂ PCl	GaCl ₃ (5 %)	7 days	> 99 %	-
Ph ^{(n)Bu} PCl	GaCl ₃ (25 %)	7 days	-	> 99%
Ph ^{(n)Bu} PCl	GaCl ₃ (50 %)	7 days	26 %	74 %
Ph ^{(n)Bu} PCl	TMSOTf (25 %)	7 days	3 %	95 %
Ph ^{(t)Bu} PCl	GaCl ₃ (5 %)	7 days	84 %	16 %
Ph ^{(t)Bu} PCl	TMSOTf (100 %)	9 days	23 %	34 %
ⁱ Bu ₂ PCl	GaCl ₃ (5 %)	7 days	66 %	-
ⁱ Bu ₂ PCl	TMSOTf (25 %)	7 days	8 %	-
Ph ₂ POEt ^b	GaCl ₃ (5 %)	7 days	3 %	10 %
Ph ₂ PN ⁱ Pr ₂	GaCl ₃ (5 %)	7 days	15 %	5 %

a. NMR conversions by relative ³¹P NMR intensity. b. Other products observed. See ESI for details.

The potential utility of silane/Lewis acid reduction of chlorophosphanes to practical synthesis was explored by the reduction of **1** on a 2 mmol scale using AlCl₃ and PMHS as the reductive system. Following work-up with Me₂S.BH₃, the desired product, Ph₂P(BH₃)H, was isolated in unoptimised 59% yield. This augurs well for the potential future application of this reactivity, given the comparatively mild reaction conditions and cheap, easily handled reagents, but the yield remains low and reaction times long compared to other reductive approaches to this compound (*e.g.* LiAlH₄, NaBH₄). Additional work on optimisation of reaction conditions and Lewis acid are required before this can be considered a generally useful method for primary and secondary phosphane synthesis.





Figure 1 – Variable Temperature NMR Studies. a) VT studies on reaction mixture showing decoalescence on cooling. B) Comparison of ^{31}P and $^{31}\text{P}\{^1\text{H}\}$ NMR at $-60\text{ }^\circ\text{C}$ showing clean formation of $[\mathbf{3}]^+$ and $[\mathbf{6}]^+$



Substrate	Catalyst	Time	Conversion ^a		
			RPH ₂	R(H)P-P(H)R	(RP) _n ^b
PhPCl ₂	GaCl ₃ (5 %)	7 days	65 %	23 %	10 %
PhPCl ₂	TMSOTf (25 %)	7 days	-	-	86 %
PhPCl ₂ ^c	NaBAR ^{Cl} (100 %)	~ 5 minutes	42 %	13 %	45 %
^t BuPCl ₂	GaCl ₃ (5 %)	7 days	18 %	-	44 %
^t BuPCl ₂	GaCl ₃ (25 %)	7 days	30 %	8 %	62 %
PCl ₃ ^c	GaCl ₃ (5 %)	~ 5 minutes	23 %	-	77 %

a. NMR conversions by relative ^{31}P NMR intensity. b. cyclic systems including (RP)₄, and larger rings. See ESI for details c. Proceeds at ambient temperature.

Table 4: Reduction of primary chlorophosphanes and PCl₃

Mechanistic Concerns

Given the dramatic influence on anion and substituents on reaction products, we sought a deeper understanding of the mechanisms involved. During reductions with substoichiometric Lewis acid, the ^{31}P NMR shows a number of broad product signals, indicative of multiple exchanging species. Reductions were performed with a 25% Lewis acid loading (GaCl₃ and TMSOTf) and heated at $60\text{ }^\circ\text{C}$ to allow the reactions to proceed, after which variable temperature NMR studies were used to freeze out the exchange processes and identify the intermediates.

For the GaCl₃ catalysed reaction, on cooling to $-30\text{ }^\circ\text{C}$ the ^{31}P spectra resolve to show $[\mathbf{3}]\text{GaCl}_4$ and the known adduct $[\text{Ph}_2\text{P}-\text{P}(\text{Ph}_2)-\text{PPh}_2]\text{GaCl}_4$, $[\mathbf{6}]\text{GaCl}_4$,⁴³ as the exchanging species (see Figure 1); the TMSOTf reaction mixture does not fully resolve to show $^1\text{J}_{\text{P-P}}$ coupling even down to $-70\text{ }^\circ\text{C}$ but the unresolved peaks do correspond to those seen for GaCl₃, confirming the formation of phosphonium intermediates in this reaction, and that transient M-H bond formation is not required for Si-to-P hydride transfer.


Given the difference in reactivity observed for the $[\text{GaCl}_4]^-$ and TfO^- salts, and the anion degradation observed when catalytic loadings of NaBAR^F and NaBAR^{Cl} were used to initiate reduction, we synthesised $[\mathbf{2}]\text{BAR}^{\text{F}}$ and $[\mathbf{2}]\text{BAR}^{\text{Cl}}$ to preform the phosphonium cation with a WCA. In both cases, on addition of Et₃SiH, formation of $[\mathbf{3}]^+$ occurred rapidly at ambient temperature. Reaction stopped at that stage for the BAR^F salt, and $[\mathbf{3}]\text{BAR}^{\text{F}}$ was isolated in 60% yield as a colourless crystalline solid. Although $[\mathbf{3}][\text{B}(\text{C}_6\text{F}_5)_4]$ is known in the literature,⁴⁴ this is the first crystallographically characterised salt of this cation. The cation is disordered about an inversion centre, and the proton could not be located in the difference map, but the proton position can be assigned by comparison to calculated geometry (See ESI for details). The P-P bond length is short at 2.176 (3) Å, compared to that of $[\mathbf{2}]\text{GaCl}_4$ and $[\text{Ph}_3\text{P}-\text{PPh}_2]\text{OTf}$ (2.205(4) Å³¹ and 2.230 (1) Å³⁹ respectively) as expected with the reduction in steric demand. In contrast, the BAR^{Cl} salt continued to react, with slow formation of $[\mathbf{6}]\text{BAR}^{\text{Cl}}$ seen over 19 days. This was accompanied by loss of intensity in the ^{11}B spectrum. On addition of excess pyridine, a new signal formed at $\delta 0.5$ ppm in the ^{11}B NMR, indicating the formation of a four-coordinate boron species and thus that the protio-phosphane-stabilised phosphonium is sufficiently acidic to cause protodeboronation of the BAR^{Cl} anion even at ambient temperature.

Since ambient temperature reaction of the BAR^{Cl} salts with silanes leads to dehydrocoupling whilst **4** is observed when the reaction mixture is heated, a different mechanism is implicated. When a reaction in which BAR^{Cl} had thermally degraded

was recharged with **1** and Et₃SiH and further heated at 100 °C, preferential reduction to form **4** was observed (see ESI for details). Ingleson has shown that tris(3,5-dichlorophenyl)borane is a competent Lewis acid for activating silanes *via* FLP chemistry,⁴⁵ and it is therefore plausible that a borohydride intermediate is involved in this process. Similar behaviour is implicated in the NaBAR^{Cl}-induced reduction of PhPCl₂. When PhPCl₂ and Et₃SiH are premixed before NaBAR^{Cl} addition, rapid reaction ensues giving PhPH₂ as the major product; addition of pyridine confirmed anion degradation and the formation of a py-Ar₃ species *in situ*.

The source of the hydride was confirmed to be the silane by isotopic labelling. Reaction of **1** with catalytic (25%) GaCl₃ in the presence of Et₃SiD gave clean formation of Ph₂PD and a 1:1:1 triplet in the ³¹P NMR, ruling out solvent activation. An analogous experiment combining **1**, 25% Me₃SiOTf and 1:1 mixture of Et₃SiH and Et₃SiD on heating at 100 °C cleanly formed **5** in addition to H₂, HD and D₂ as seen in the ¹H and ²H NMR, confirming hydrogen formation and overall dehydrocoupling. The necessity for phosphonium formation for reduction was confirmed by heating **1** with Et₃SiH at 100 °C in the absence of Lewis acid – no reduction was observed after 3 days. To rule out the possibility that [GaCl₄]⁻ might be acting as a soluble Cl⁻ source interacting with Et₃SiH to form a 5-coordinate activated silane, the combination of **1** and Et₃SiH were heated at 100 °C with 10 % [BnNEt₃]Cl – after 7 days, 3% of the **1** had reacted to form **5** as the sole product. The observation of **6**[GaCl₄]⁻ in the variable temperature studies indicated the formation of **5** as an intermediate, but this is not seen under equivalent catalyst loadings at higher temperatures, implying that reduction of the P-P bond may also occur under these conditions. A control reaction of **5** with 25% GaCl₃ and Et₃SiH showed that Lewis acid mediated cleavage of the P-P bond to form **4** does occur, but that it is slow (47% conversion after 3 days at 100 °C) relative to the formation of **4** from **1** (near quantitative conversion after 1 day at 100 °C), indicating that it is a minor pathway.

Table 5: Relative donor strengths of phosphanes as evaluated by donor exchange.




Donor	Relative Stability / kCalmol ⁻¹
Ph ₂ P(Cl)	0.00
Ph ₂ P(H)	-5.95
Ph ₂ P-PPh ₂	-10.45
Ph ₃ P	-11.83
Ph(ⁿ Bu)P(Cl)	-0.74
Ph(ⁿ Bu)P(H)	-8.50
Ph(^t Bu)P(Cl)	-1.80
Ph(^t Bu)P(H)	-8.18
Ph(ⁱ Pr ₂ N)P(Cl)	-0.51
Ph(ⁱ Pr ₂ N)P(H)	-10.91

All calculations were performed at the M06-2X/6-311g(d,p) level with PCM(Dichloromethane) solvent model. See ESI for details.

Both radical and Lewis acid hydride abstraction were considered as potential mechanisms of hydride transfer. However, as ^tBu₂P-P^tBu₂ is a known species accessible *via* single electron reduction and is not formed under these reaction conditions, this argues strongly against a radical mechanism for P-P coupling in these species.⁴⁶ For all stoichiometric reactions, formation of protio-phosphane-stabilised phosphonium cations is rapid (minutes to hours at ambient temperature) whereas subsequent hydride transfer requires extended heating. Furthermore, the observation of [2]⁺ and [3]⁺, and [3]⁺ and [6]⁺ simultaneously, in conjunction with free **1**, giving well resolved signals in the ³¹P NMR indicates that there is a significant increase in donor strength at each stage of the process. A quantitative assessment of relative donor strength was obtained by DFT evaluation of ligand exchange about [2]⁺ for a selection of relevant donors involved in the reactions observed, as shown in Table 5, where a more negative value indicates an increasingly stable adduct relative to [2]⁺. The results are in agreement with the experimental observation that Ph₃P displaces Ph₂P(Cl) from [2]⁺,³⁹ and the qualitative observation that the rate of hydride transfer to phosphonium (of the order [2]⁺ > [3]⁺ > [6]⁺ ≈ [Ph₃P-PPh₂]⁺) correlates well with the calculated donor strengths of the phosphanes. Furthermore, it can be seen that in each instance the secondary phosphane is a stronger donor than the corresponding secondary chlorophosphane and that the alkyl-substituted phosphanes are stronger donors than diphenylphosphane derivatives. This may in part explain the increased formation of P-P coupled products in these cases as reduced electrophilicity at phosphorus favours competitive deprotonation instead.

Scheme 3: Proposed Catalytic Cycle for Halophosphane Reduction



From these results, it can be seen that anion participation is not required for the transfer of hydride from silane to phosphorus for the halo-phosphonium but that phosphonium stabilisation (or lack thereof) and thus Lewis acidity is important. We therefore propose a Piers-Oestreich type-transfer mechanism,^{47,48} whereby the Si-H bond coordinates to the Lewis acidic site at phosphorus followed by attack of a donor centre (solvent, anion, or one of the many phosphanes in solution) generating a transient silylium cation intermediate. This silylium intermediate can then abstract a halide from another equivalent of chlorophosphane or tetrahalogallate, closing the catalytic cycle (Scheme 3). After formation of protio-phosphane-phosphonium, subsequent reaction could then either proceed *via* a second equivalent of silane reacting at P,

forming a second P-H bond, or reaction with a base to deprotonate the intermediate, forming a diphosphane. This step would appear to be strongly anion dependent. The proposed mechanism is likewise consistent with the less successful reduction of P-N and P-O bonds in the presence of Lewis acid and silane due to the reduced lability of these bonds relative to P-Cl systems, and with the decomposition of the BAr^{F} anion as ArCF_3 groups are known to react with silylium species.⁴⁹

Conclusions

We have investigated the reactivity of phosphane-stabilised phosphonium cations with hydrosilanes and shown that they undergo facile hydride transfer to form protio-phosphane-stabilised phosphonium species. These can then further react through two, anion-dependent reaction pathways; deprotonation effects reductive concatenation with the formation of P-P bonds, whilst hydride transfer to the less electrophilic, protio-phosphane-stabilised phosphonium centre leads to primary or secondary phosphanes. These transformations can be made catalytic in Lewis acid, and can be extended using cheap Lewis acids and silanes potentially offering a mild, operationally simple reduction protocol without reactive M-H bonds. Further catalytic applications of these donor-stabilised phosphonium cations are currently under investigation.

ASSOCIATED CONTENT

Supporting Information.

Complete synthetic details, multinuclear NMR data, computational results and Cartesian coordinates of all optimised species are available in the supporting information. This material is available free of charge via the Internet at <http://pubs.acs.org>. The crystallographic data have been submitted to the Cambridge Crystallographic Database (Deposit number: 1846580).

AUTHOR INFORMATION

Corresponding Author

* e.r.clark@kent.ac.uk

ORCID

Kyle G. Pearce: 0000-0003-1886-4483
Andryj M. Borys: 0000-0001-8437-2562
Ewan R. Clark: 0000-0001-7287-2631
Helena J. Shepherd: 0000-0003-0832-4475

Author Contributions

All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT


The authors wish to thank the University of Kent for funding.

REFERENCES

1. Baumgartner, T.; Réau, R. Organophosphorus π -Conjugated Materials. *Chem. Rev.* **2006**, *106*, 4681.
2. Bellingham, R.; Borrett, G.; Bret, G.; Choudary, B.; Colclough, D.; Hayes, J.; Hayler, J.; Hodnett, N.; Ironmonger, A.; Ochen, A.; Pascoe, D.; Richardson, J.; Vit, E.; Alexandre, F.; Caillet, C.; Amador, A.; Bot, S.; Bonaric, S.; da Costa, D.; Lioure, M.; Roland, A.; Rosinovsky, E.; Parsy, C.; Dousson, C. B. Development and Scale-Up of a Manufacturing Route for the Non-nucleoside Reverse Transcriptase Inhibitor GSK2248761A (IDX-899): Synthesis of an Advanced Key Chiral Intermediate. *Org. Process Res. Dev.* **2018**, *22*, 200.
3. Kamer, P. C. J.; van Leeuwen, Piet W. N. M., Eds.; In *Phosphorus(III) Ligands in Homogeneous Catalysis: Design and Synthesis*; John Wiley & Sons, Ltd: United Kingdom, 2012; .
4. Wauters, I.; Debrouwer, W.; Stevens, C. V. Preparation of phosphines through C-P bond formation. *Beilstein J. Org. Chem.* **2014**, *20*, 1064-1096.
5. Schrems, M. G.; Pfaltz, A. NeoPHOX-an easily accessible P,N-ligand for iridium-catalyzed asymmetric hydrogenation: preparation, scope and application in the synthesis of demethyl methoxycalamenene. *Chem. Commun.* **2009**, 6210.
6. Montchamp, J.; Dumond, Y. R. Synthesis of Monosubstituted Phosphinic Acids: Palladium-Catalyzed Cross-Coupling Reactions of Anilinium Hypophosphite. *J. Am. Chem. Soc.* **2001**, *123*, 510-511.
7. Bange, C. A.; Waterman, R. Challenges in Catalytic Hydrophosphination. *Chem. Eur. J.* **2016**, *22*, 12598.
8. Budzelaar, P. H.; van Doorn, J. A.; Nico, M. Reductive cleavage of the carbon-phosphorus bond with alkali metals. I. Cleavage of functionalised triphenylphosphines; formation of secondary and primary phosphines. *Recl. Trav. Chim. Pays-Bas* **1991**, *110*, 420-432.
9. Kawashima, M. Japan Patent JP 2000109491, 2000.
10. Kawashima, M. Japan Patent JP 2000007689, 2000.
11. O'Brien, C.; Tellez, J.; Nixon, Z.; Kang, L.; Carter, A.; Kunkel, S.; Przeworski, K.; Chass, G. Recycling the Waste: The Development of a Catalytic Wittig Reaction. *Angew. Chem. Int. Ed.* **2009**, *48*, 6836-6839.
12. Blair, S.; Izod, K.; Taylor, R.; Clegg, W. Isolation and structural characterisation of two sterically crowded diphosphanes. *J. Organomet. Chem.* **2002**, *656*, 43-48.
13. Busacca, C. A.; Bartholomeyzyk, T.; Cheekoori, S.; Raju, R.; Eriksson, M.; Kapadia, S.; Saha, A.; Zeng, X.; Senanayake, C. H. Reduction of Phosphinites, Phosphinates, and Related Species with DIBAL-H. *Synlett* **2009**, *2009*, 287-291.
14. Tatarnikov, A. N.; Trubina, L. J. Russia Patent , 2004.
15. Frew, J. J. R.; Karen, D.; Van Rensburg, H.; Slawin, A. M. Z.; Tooze, R. P.; Clarke, M. Palladium(II) Complexes of New Bulky Bidentate Phosphanes: Active and Highly Regioselective Catalysts for the Hydroxycarbonylation of Styrene. *Chem. Eur. J.* **2009**, *15*, 10504-10513.
16. Imamoto, T.; Kusumoto, T.; Suzuki, N.; Sato, K. Phosphine oxides and lithium aluminum hydride-sodium borohydride-cerium(III) chloride: synthesis and reactions of phosphine-boranes. *J. Am. Chem. Soc.* **1985**, *107*, 5301-5303.
17. Imamoto, T.; Oshiki, T.; Onozawa, T.; Kusumoto, T.; Sato, K. Synthesis and reactions of phosphine-boranes. Synthesis of new bidentate ligands with homochiral phosphine centers via optically pure phosphine-boranes. *J. Am. Chem. Soc.* **1990**, *112*, 5244-5252.
18. Clarke, C.; Foussat, S.; Fox, D. J.; Pedersen, D. S.; Warren, S. Asymmetric synthesis of trans-disubstituted cyclopropanes using phosphine oxides and phosphine boranes. *Org. Biomol. Chem.* **2009**, *7*, 1323-1328.
19. Staubitz, A.; Robertson, A. P. M.; Sloan, M. E.; Manners, I. Amine- and Phosphine-Borane Adducts: New Interest in Old Molecules. *Chem. Rev.* **2010**, *110*, 4023-4078.
20. Xu, L.; Kurukulasuriya, R. Trichlorosilane. *EROS* **2006**.
21. Mehta, M.; de la Arada, Isaac Garcia.; Perez, M.; Porwal, D.; Oestreich, M.; Stephan, D. W. Metal-Free Phosphine Oxide Reductions Catalyzed by $\text{B}(\text{C}_6\text{F}_5)_3$ and Electrophilic Fluorophosphonium Cations. *Organometallics* **2016**, *35*, 1030.
22. Berthod, M.; Favre-Réguillon, A.; Mohamad, J.; Mignani, G.; Docherty, G.; Lemaire, M. A Catalytic Method for the Reduction of Secondary and Tertiary Phosphine Oxides. *Synlett* **2007**, *2007*, 1545-1548.
23. Engel, R., Ed.; In *Handbook of Organophosphorus Chemistry*; CRC Press: United States of America, 1992; .

24. Buonomo, J. A.; Aldrich, C. C. Mitsunobu Reactions Catalytic in Phosphine and a Fully Catalytic System. *Angew. Chem. Int. Ed.* **2015**, *54*, 13041-13044.
25. van Kalker, H. A.; Leenders, S. H.; Rianne, H. C.; Rutjes, F. P.; van Delft, F. L. In Situ Phosphine Oxide Reduction: A Catalytic Appel Reaction. *Chem. Eur. J.* **2011**, *17*, 11290-11295.
26. Krenke, E. H. Theoretical Investigation of the Mechanisms and Stereoselectivities of Reductions of Acyclic Phosphine Oxides and Sulfides by Chlorosilanes. *J. Org. Chem.* **2012**, *77*, 3969.
27. Demchuk, O. M.; Jasiński Radomir, Micha, P. K. New Insights into the Mechanism of Reduction of Tertiary Phosphine Oxides by Means of Phenylsilane. *Heteroatom Chem.* **2015**, *26*, 441-448.
28. Pilard, J.; Baba, G.; Gaumont, A.; Denis, J. Reduction of Phosphine Oxides, A Mild and Practical Synthesis of Secondary P-Ethynylphosphines. *SynnLett* **2000**, *1995*, 1168-1170.
29. Đorđević, N.; Ganguly, R.; Petković, M.; Vidović, D. E-H (E = B, Si, C) Bond Activation by Tuning Structural and Electronic Properties of Phosphenium Cations. *Inorg. Chem.* **2017**, *56*, 14671.
30. Chitnis, S. S.; Felix, K.; Stephan, D. W. Catalytic Hydrodefluorination of C-F Bonds by an Air-Stable PIII Lewis Acid. *Chem. Eur. J.* **2018**, *24*, 6543-6546.
31. Burford, N.; Ragogna, P. J.; McDonald, R.; Ferguson, M. J. Phosphine Coordination Complexes of the Diphenylphosphenium Cation: A Versatile Synthetic Methodology for P-P Bond Formation. *J. Am. Chem. Soc.* **2003**, *125*, 14404.
32. Clark, E. R.; Borys, A. M.; Pearce, K. Donor-substituted phosphanes - surprisingly weak Lewis donors for phosphenium cation stabilisation. *Dalton Trans.* **2016**, *45*, 16125.
33. Canac, Y.; Maaliki, C.; Abdallah, I.; Chauvin, R. Carbeniophosphanes and their carbon →phosphorus→metal ternary complexes. *New J. Chem.* **2012**, *36*, 17.
34. Bezpalko, M. W.; Poitras, A. M.; Foxman, B. M.; Thomas, C. M. Cobalt N-Heterocyclic Phosphenium Complexes Stabilized by a Chelating Framework: Synthesis and Redox Properties. *Inorg. Chem.* **2017**, *56*, 503.
35. Slattery, J. M.; Hussein, S. How Lewis acidic is your cation? Putting phosphenium ions on the fluoride ion affinity scale. *Dalton Trans.* **2012**, *41*, 1808-1815.
36. Burck, S.; Forster, D.; Gudat, D. Phosphorus-carbon bond formation catalysed by electrophilic N-heterocyclic phosphines. - *Chem. Commun.* **2006**, 2810.
37. Adams, M. R.; Tien Chieh-Hung; Huchenski, B. S.; Ferguson, M. J.; Speed, A. W. Diazaphospholene Precatalysts for Imine and Conjugate Reductions. *Angew. Chem. Int. Ed.* **2017**, *56*, 6268-6271.
38. Rao, B.; Chong, C. C.; Kinjo, R. Metal-Free Regio- and Chemoselective Hydroboration of Pyridines Catalyzed by 1,3,2-Diazaphosphenium Triflate. *J. Am. Chem. Soc.* **2018**, *140*, 652.
39. Burford, N.; Cameron, T. S.; Ragogna, P. J.; Ocando-Mavarez, E.; Gee, M.; McDonald, R.; Wasylshen, R. E. Phosphine Ligand Exchange at a Phosphine Lewis Acceptor: The First Structural Characterization of Homoleptic Phosphinophosphenium Salts. *J. Am. Chem. Soc.* **2001**, *123*, 7947-7948.
40. Ammer, J.; Nolte, C.; Mayr, H. Free Energy Relationships for Reactions of Substituted Benzhydrylium Ions: From Enthalpy over Entropy to Diffusion Control. *J. Am. Chem. Soc.* **2012**, *134*, 13902.
41. Markus, H.; Schappele, L. H.; Lang-Wittkowski Gabriele; Herbert, M.; Ofial, A. R. Towards a Comprehensive Hydride Donor Ability Scale. *Chem. Eur. J.* **2013**, *19*, 249-263.
42. Minegishi, S.; Mayr, H. How Constant Are Ritchie's "Constant Selectivity Relationships"? A General Reactivity Scale for n-, π-, and σ-Nucleophiles. *J. Am. Chem. Soc.* **2003**, *125*, 286.
43. Burford, N.; Dyker, A. C.; Decken, A. Facile Synthetic Methods for the Diversification of Catena-Polyphosphorus Cations. *Angew. Chem. Int. Ed.* **2005**, *44*, 2364-2367.
44. Dobrovetsky, R.; Takeuchi, K.; Stephan, D. W. Metal-free Lewis acid mediated dehydrocoupling of phosphines and concurrent hydrogenation. *Chem. Commun.* **2015**, *51*, 2396.
45. Valerio, F.; Ingleson, M. J. Expanding Water/Base Tolerant Frustrated Lewis Pair Chemistry to Alkylamines Enables Broad Scope Reductive Aminations. *Chem. Eur. J.* **2017**, *23*, 2217-2224.
46. Goldwhite, H.; Kaminski, J.; Millhauser, G.; Ortiz, J.; Vargas, M.; Vertal, L.; Lappert, M. F.; Smith, S. J. Phosphorus-phosphorus single or double bond formation from PCl_{3-n}R_n (n = 1 or 2) and a bis-imidazolidine reducing agent. *J. Organomet. Chem.* **1986**, *310*, 21-25.
47. Parks, D. J.; Blackwell, J. M.; Piers, W. E. Studies on the Mechanism of B(C₆F₅)₃-Catalyzed Hydrosilylation of Carbonyl Functions. *J. Org. Chem.* **2000**, *65*, 3090.
48. Rendler, S.; Oestreich, M. Conclusive Evidence for an S_N2-Si Mechanism in the B(C₆F₅)₃-Catalyzed Hydrosilylation of Carbonyl Compounds: Implications for the Related Hydrogenation. *Angew. Chem. Int. Ed.* **2008**, *47*, 5997-6000.
49. Scott, V. J.; Çelenligil-Çetin, R.; Ozerov, O. V. Room-Temperature Catalytic Hydrodefluorination of C(sp³)-F Bonds. *J. Am. Chem. Soc.* **2005**, *127*, 2852-2853.

Insert Table of Contents artwork here



Phosphane-stabilised phosphenium cations react with hydrosilanes in an anion-dependent process leading either to secondary phosphanes, or reductive coupling to diphosphanes and liberation of H₂. Studies into the mechanism of the transformation are reported, and catalytic-in-Lewis-acid variants explored.
