



Kent Academic Repository

Azevedo, Ruben T., Aglioti, Salvatore Maria and Lenggenhager, Bigna (2016) *Participants' above-chance recognition of own-heart sound combined with poor metacognitive awareness suggests implicit knowledge of own heart cardiodynamics*. Scientific Reports, 6 . ISSN 2045-2322.

Downloaded from

<https://kar.kent.ac.uk/73812/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1038/srep26545>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

SCIENTIFIC REPORTS



OPEN

Participants' above-chance recognition of own-heart sound combined with poor metacognitive awareness suggests implicit knowledge of own heart cardiodynamics

Ruben T. Azevedo^{1,2,†}, Salvatore Maria Aglioti^{1,2} & Bigna Lenggenhager^{1,*}

Mounting evidence suggests that interoceptive signals are fundamentally important for the experience of the self. Thus far, studies on interoception have mainly focused on the ability to monitor the timing of ongoing heartbeats and on how these influence emotional and self-related processes. However, cardiac afferent signalling is not confined to heartbeat timing and several other cardiac parameters characterize cardiodynamic functioning. Building on the fact that each heart has its own self-specific cardio-dynamics, which cannot be expressed uniquely by heart rate, we devised a novel task to test whether people could recognize the sound of their own heart even when perceived offline and thus not in synchrony with ongoing heartbeats. In a forced-choice paradigm, participants discriminated between sounds of their own heartbeat (previously recorded with a Doppler device) versus another person's heart. Participants identified the sound of their own heart above chance, whereas their metacognition of performance – as calculated by contrasting performance against ratings of confidence – was considerably poorer. These results suggest an implicit access to fine-grained neural representations of elementary cardio-dynamic parameters beyond heartbeat timing.

It is now held that the ever-continuous representation and regulation of bodily signals provide the foundations for the experience of the self¹. We exist in our bodies, as our bodies exist in our minds². Traditionally, experimental work in this field has focused on the processing of exteroceptive (e.g. tactile and visual) and motor bodily signals. Many experimental paradigms have been developed over the last years to manipulate these signals and to investigate their influence on the bodily self^{3–6}. However, the influence of the private internal body, i.e. interoceptive signals⁷, has been largely overlooked and has only rather recently entered the experimental study of the self^{8–10}. This is potentially surprising, as it has long been argued theoretically that self-awareness emerges from an image of the homeostatic state of the body, i.e. from the constantly present interoceptive signals^{11,12}. The reason for this dearth of experimental research may be that inherently private interoceptive sensations are not easily amenable to experimental measurement and manipulation. Cardiac sensations provide a privileged source of information for the study of interoception. The central role of mapping, re-mapping and interpreting one's own cardiac signals has long been recognized in theories of emotion^{11,12}. Yet people differ greatly in their ability to explicitly and accurately identify discrete interoceptive events such as heartbeats¹³. This ability, often named interoceptive accuracy (IAcc;¹⁴) is typically assessed by asking participants to track their heartbeats for short periods of time¹³, or by asking them to discriminate between auditory tones presented either synchronously or asynchronously with their

¹IRCCS Fondazione Santa Lucia, Via Ardeatina 306, 00100 Rome, Italy. ²Department of Psychology, University of Rome "La Sapienza", Via dei Marsi 78, 00185 Rome, Italy. [†]Present address: Lab of Action & Body, Department of Psychology, Royal Holloway, University of London, Egham, UK. ^{*}Present address: Neuropsychology, Department of Neurology, University Hospital Zurich, Frauenklinikstrasse 26, CH-8091 Zurich, Switzerland. Correspondence and requests for materials should be addressed to B.L. (email: bigna.lenggenhager@gmail.com)

heartbeats¹⁵. While individual differences in IAcc have been shown to predict a wide number of cognitive and emotional processes¹⁶, it is widely accepted that the influence of interoceptive signals in, for example, emotional processing^{17–20}, or in the sense of the self^{21–23}, often occurs without full explicit awareness. Indeed, people demonstrate remarkable different levels of *awareness* of their success when performing interoceptive tasks (i.e., their meta-cognitive awareness differs). In other words, their ability to identify interoceptive signals, i.e. heartbeats, is not always paralleled by accurate judgments of their own performance^{14,24}. This lack of explicit awareness, or meta-awareness, makes the study of interoception additionally challenging but also appealing.

Thanks to new experimental paradigms, recent studies indicate that interoceptive signals are constantly integrated with exteroceptive (i.e. visual or auditory) signals - even if often at an implicit level - in order to build a coherent representation of the own body^{21,22}. For example, Aspell and colleagues²¹, presented participants with a virtual body whose silhouette could be flashing either synchronously or asynchronously with respect to their own heartbeats. Even if participants were not aware of this contingency, they identified themselves more with the avatar in the synchronous condition. In a similar line of research, a recent electrophysiological study has shown sensory suppression of heartbeat-related auditory tones - again without the participants' explicit awareness²³. Both studies suggest that cardiac signals presented through an exteroceptive channel can activate and be compared with representations of interoceptive information (see also²⁵). This implies that the bodily awareness is, at least partially, grounded in a successful integration of extero- and interoceptive signals.

Importantly, however, the vast majority of research into interoceptive awareness as well as into implicit integration of intero- and exteroceptive signals, has focused on cardiac timing (i.e. the timing of the occurrence of each heartbeat). Yet cardiac afferent signalling is not confined to simple heartbeat timing. Several other cardiovascular parameters of autonomic reactivity²⁶ as well as parameters related to cardiac inotropy²⁷ have been shown to mediate interoception. Indeed the various cardiac parameters are interrelated and it is their concerted activity that fully characterizes cardiovascular functioning and regulation. It is known that each heart has its own self-specific cardio-dynamics, which cannot be expressed uniquely by mere heart rate. Such a “cardiac fingerprint” is detectable by computer algorithms that can successfully recognize individual hearts independently of their heart rate at the moment of recording²⁸. Accordingly, in order to comprehend the various parameters comprising one's own cardiac signature, a person's representation of their cardiac information will not be restricted to the online processing of the timing of their heartbeats. It is worth noting here that in other domains, e.g. locomotion, it has been shown that people not only are able to monitor their movements online but can also recognize their “motor signature” when asked to recognize their movement on a point light walker^{29,30} as well as on a virtual character³¹ from a third person perspective via offline visual presentation. Moreover, such recognition seems to rely on an individual's implicit knowledge about the intrinsic temporal dynamics of his/her movements, which has been gained, at least partially, through a lifetime of afferent proprioceptive and sensorimotor experiences³¹.

In this study, we aimed to explore whether people have implicit and/or explicit knowledge of their cardiac signature, i.e. whether people can discriminate their own self-specific cardio-dynamics, even when presented offline. In a novel forced-choice paradigm (see Fig. 1), we asked participants to discriminate between sounds of their own heartbeat previously recorded (with a Doppler device, see below) and those of another person's heart. We have previously shown that presenting these heart sounds online to participants can implicitly modulate their behaviour²⁰, but it remains unclear if participants can actually recognize the sound of their own heart. To control for the possibility that own-heart recognition could be merely due to cardiac timings or to a general knowledge about their own heart rate, the samples of the other-heart sound were matched in heart-rate. If participants are able to identify their own cardiac signature, we expected them to perform significantly above chance. In order to assess whether performance is underpinned by implicit or by explicit processes³², participants were further asked to rate their confidence in their decisions after each trial. Comparing objective performance with subjective perceptions of the own accuracy allowed us to calculate participants' (meta) awareness of their ability to recognize their own-heart sounds¹⁴. A heartbeat counting task¹³ was further used to enable us to relate the participant's ability to recognise their own-heart sound recognition to a standard measure of interoceptive accuracy.

Results

Recognition of the sound of the heart. Participants' performance was analysed using binominal distribution probabilities, which indicate that performance above chance would be shown by correct recognition of 20 trials, out of the total 30 trials ($p = 0.028$).

Accordingly, participants were assigned to three groups (see Fig. 2a): 1) the SELF group contained those participants who had accuracy rates above 0.67, i.e. 20 or more trials correct, and therefore performed significantly better than chance. In other words, these participants correctly attributed their own heart to themselves; 2) the NON-DISCRIMINATOR group comprised those participants who performed at chance level (i.e. accuracy rates between 0.33 and 0.66); 3) the OTHER group was made up of those individuals who performed significantly below-chance (accuracy below 0.33). These participants incorrectly identified the other person's heart as their own heart.

The distribution of participants between the different groups is shown in Fig. 2a. 17 out of 27 fell into the SELF group (i.e., correct self-attribution, mean accuracy 0.77 (SEM 0.02), see Table 1), 7 out of 27 fell into the NON-DISCRIMINATOR group (i.e., they responded at chance-level (mean accuracy 0.52 (SEM 0.03)) and 3 out of 27 fell into the OTHER group (i.e., misattribution of the other's heart as their own heart, mean accuracy 0.21 (SEM 0.06)). A chi-square test suggests significant differences in numbers between the groups (chi square = 11.6, $p = 0.003$, Cohen's $\omega = 0.65$). Post-hoc chi square tests revealed that more participants belonged to the SELF group than to the NON-DISCRIMINATOR (chi square = 4.2, $p = 0.04$, $\omega = 0.42$) and OTHER groups (chi square = 9.8, $p = 0.002$, $\omega = 0.70$). The numbers in the two latter groups did not differ significantly.




Figure 1. Experimental procedure: Before the experiment started, we recorded: (a) the sound of the participant's heart using a Doppler device (main experiment); (b) and “beep” sounds triggered by the participant's pulse waves measured with a pulse transducer (control experiment). After pre-processing, short samples of their own or another person's heart sounds were presented, in random order. After each of the 30 trials, participants had to make a forced-choice decision about whether they had heard the sound of their own heartbeats or not, followed by a rating of their confidence in their decision.

Other measures. *Confidence ratings.* Confidence ratings in the experimental condition was an average of 43.2 (SEM 3.5, ranging from 4.9 to 64.5). The three groups did not significantly differ in their confidence ratings (Kruskal-Wallis, chi square = 0.53, $p = 0.77$, $\omega = 0.14$).

Heartbeat Counting task. The performance in the heartbeat counting task was on average 0.75 (SEM 0.04, ranging from 0.31 to 0.97). The three groups did not significantly differ in their performance (Kruskal-Wallis, chi square = 3.37, $p = 0.18$, $\omega = 0.59$).

Meta awareness. Participants' meta-awareness of performance (meta- d') was calculated as a function of objective performance and associated confidence ratings, according to “type 2” Signal Detection Theory indices of sensitivity³². Meta- d' scores (Mean 0.30; SEM; 0.23) were significantly lower ($t = 2.6$, $p = 0.016$, Cohen's $d = 0.43$) than d' (Mean 0.80, s.d. 0.23) showing that participants had poor awareness of their own accuracy, which suggests intuition-based behavior. Moreover, correlation analyses revealed that meta- d' scores ($r = 0.40$, $p = 0.041$) but not d' scores ($r = 0.03$, $p = 0.88$) were significantly correlated to performance in the heartbeat counting task. The former correlation is larger than the latter: $t = 2.19$, $p = 0.028$, using Lee & Preacher, test for dependent correlations³³; see Fig. 2b,c.

Control task. Analysis of accuracy was performed identically as in the experimental task. 3 out of the 16 participants fell into the SELF group (mean accuracy 0.87, see Table 1), 11 into the NON-DISCRIMINATOR group (mean accuracy 0.49) and 2 into the OTHER group (mean accuracy 0.13). A chi square test suggests a significantly different distribution between the groups (chi square = 9.13, $p = 0.01$, $\omega = 0.58$). Post-hoc chi square tests suggest that more participants belonged to the NON-DISCRIMINATOR group than to the SELF group (chi square = 4.6, $p = 0.03$, $\omega = 0.44$) and OTHER group (chi square = 6.2, $p = 0.01$, $\omega = 0.79$), while the two latter groups did not differ significantly from each other (chi square = 0.2, $p = 0.66$, $\omega = 0.1$).

Debriefing. In the semi-structured debriefing interview all participants reported believing that the sounds belonged to 3 or more different hearts. They further reported that their choices were mostly based on guessing or on intuition. Even the few participants who felt considerably confident in having discriminated their own heart correctly still confessed difficulties in identifying the criteria used. Together this evidence supports the finding that the identification of own-heart sound was performed implicitly.

Discussion

Bodily signals, particularly those arising from the interoceptive system, are thought to set the foundations on which the subjective experience of the self unfolds^{2,9,10}. This notion of embodied self has received convincing support from recent empirical research, which has shown that cardiac signals are integrated with external information in order to build a coherent sense of self^{21,22,34}. Here, we aimed to address a previously neglected dimension of interoception, namely the offline representation of the inner body. Specifically, we explored whether




Figure 2. Results: (a) Percentage of participants grouped according to their performance in the sound recognition task; (b) Correlation between performance in the counting task and meta-awareness; (c) Correlation between performance in the counting task and performance in the sound recognition task.

Group	Self	Non-discriminators	Other
Heartbeat	17/27 (63%)	7/27 (26%)	3/27 (11%)
Beep	3/16 (18.75%)	11/16 (68.75%)	2/16 (12.5%)

Table 1. Numbers and percentages of participants falling into the different groups in the experimental task. Upper row using real heartbeat sound; lower row in the control task using R-wave evoked beep sounds.

healthy participants were able to recognize their own cardiac signature, independent of their current interoceptive state, i.e., whether there was offline recognition of self-specific cardiovascular dynamics. Our method was first to record the participants' heart sound with a Doppler device and then to ask them to discriminate the sound of their own heart from that of someone else's heart. The study had two main findings. Firstly, most participants performed significantly above chance, indicating that participants were indeed able to discriminate between previously recorded, and thus offline, sounds of their own and another person's heart. Importantly, this finding could not be based merely on heartbeat timing because this parameter was matched between the participant's own and the other person's heart sound. Moreover, the effect was absent in the control condition, where acoustic tones were delivered in synchrony with the participant's own pre-recorded heart beats. Our findings extend recent empirical research which has indicated that participants integrate externally presented (online) auditory or visual cues with cardiac signals during body-related processing^{21–23}. We show that a representation of an individual's cardiac signature goes beyond mere heartbeat timings. Furthermore, our results also suggest that people can, at least partially, access and translate neural representations of elementary cardio-dynamic parameters that have been formed throughout time (probably through sensory, acoustic or mechano-pressure mechanisms) and can compare these representations with externally presented auditory stimuli (as the sound of the blood flow recorded with a Doppler device). Even though participants could not recognize the criteria they were using, their performance indicates that they were able to identify the sounds that matched their cardiodynamic signature. So far we can only speculate about the underlying mechanisms. However, we believe that they may be comparable to those that underpin the comparison of proprioceptive and sensorimotor representation of own movements when these are presented offline, such as gait²⁹; skilled actions³⁵; or facial motion³¹. Importantly, this ability is thought to rely not merely on previous first-person visual experiences, but principally to involve the proprioceptive and sensorimotor feedback from previous movements³¹. In keeping with models that have extended (from exteroceptive to interoceptive perception) the notion of comparing different types of bodily representations¹⁰, we show in this experiment that the interoception is likely to be grounded both on online and offline body representations of the inside of our body (interoceptive representations). The continuous mapping and re-mapping of afferent signals form and maintain accurate representations of the body's structural properties, as well as of the fine grained dynamic processes that regulate its functioning. Holding coherent and detailed maps of the body allows the brain to efficiently detect, simulate and regulate changes in its state, in order to maintain homeostasis³⁶.

The second result of our study is that the recognition of one's own heartbeat signature seems to be based mostly on implicit processes. Indeed, not only did participants report intuition-based behaviour but their objective performance on the forced-choice task clearly exceeded their metacognitive awareness. Thus, although their behaviour indicates recognition of their own heart sound, participants were less good at judging their ability to perform the task itself. This indicates that the perceptual decision about self vs other heart sound was, at least partially, guided by implicit processes^{32,37}. Tellingly, only metacognitive awareness was correlated to the classical interoceptive accuracy measure that we tested (i.e., accuracy in heartbeat counting¹³). The results of the correlation analyses suggest that even if, contrary to what could be expected, individuals with high interoceptive accuracy were no better at discriminating the sound of their heart, they were more aware of their ability to perform the task. Given the relatively small sample size, these results must be treated with caution. However, we believe that they: i) suggest that offline and online representations of interoceptive signals may rely on different neurocognitive underpinnings; and ii) add to recent literature which has shown a dissociation between the various dimensions

of interoception, such as interoceptive accuracy and interoceptive awareness (i.e. meta-cognition^{14,24,38,39}). In our study, individuals with high interoceptive accuracy seem to rely in a greater extent on explicit criteria to guide their decision^{32,37}. What this suggests is that good interoceptors, who presumably hold more accurate representations of their online bodily states, are also more aware of their elementary bodily dynamics. Even if they did not have higher ability to discriminate their own-heart sound, they showed better appraisal of their decisions, arguably because they relied on a coherent understanding of their ability to perceive bodily signals.

Several limitations of this study should be considered. *First*, the study was not designed to identify the individual differences in cardio-dynamic parameters (e.g. stroke volume, ejection time), which could have contributed to accurate discrimination of own-heart sound. Future research should aim at identifying such parameters, for example by recording longer samples using equipment that allows the extraction and estimation of cardiovascular parameters (e.g., impedance cardiography). Likewise, the interoceptive pathways (e.g. tactile somatosensory, mechanoreceptors within the heart) and mechanisms underlying cardiovascular perception and the ability to recognize own-heart sound have yet to be determined. *Furthermore*, we did not assess factors that are generally known to influence interoceptive accuracy such as for example gender, mean heart rate^{40,41} or BMI⁴².

Nonetheless, we believe that this study represents a significant conceptual step and likewise a departure from previous rather simplistic heartbeat-timing-dependent approaches to cardioception. We suggest that research into interoception, both in the context of its contribution to the bodily self as well as to various cognitive and emotional processes, would benefit from acknowledging the importance of considering the full complexity of cardiovascular activity²⁷ and the need to adopt new methodologies to study cardiac interoception²⁰.

Methods

Participants. Twenty-eight healthy participants took part in the main study. A participant who was a medical doctor ($n = 1$) was excluded from the analyses, as a theoretical knowledge on cardio dynamics or experience in discriminating heart beat sounds could influence performance. The final sample size thus comprised twenty-seven participants (19 female, mean age 28.6 \pm 3.3 STD), none of whom had previously heard their own-heart sound as translated by a doppler device. All participants gave written informed consent. The study and all experimental procedures were approved by the ethics committee of the IRCSS, Fondazione Santa Lucia and were carried out in accordance with the ethical standards of the 1964 Declaration of Helsinki.

A sub-group of these participants ($n = 16$, 8 female, mean age 30.3 \pm 3.4 STD) completed the additional control task (see below).

Experimental procedure. *Stimuli.* The participant's heart rate was recorded during one minute using a pulse transducer (Adinstruments; TN1012/ST), which was attached to the index finger using a velcro fastener. Simultaneously, a Doppler device (Angel sound, Fetal Doppler, <http://www.jumper-medical.com/>) placed over the participant's heart to record the sound of the heart. The Doppler detects blood flow and generates a sound reflecting its dynamics (e.g. varying intensity (loudness), frequency (pitch), quality and duration). Because the direction and velocity of blood flow is different throughout the cardiac cycle, the audible sound represents both diastolic filling and systolic ejection⁴³. Several individual cardiovascular parameters can be inferred from the Doppler sound by a trained person, or objectively extracted by computational algorithms. For example, the first heart sound (i.e. caused by the closure of the tricuspid and mitral valves) and second heart sounds (i.e. caused by the closure of the aortic and pulmonary valves) can be perceived^{44,45}. Moreover, additional parameters related to cardiac inotropy (e.g. ejection velocity, stroke volume) and diastolic/systolic relations can be extracted⁴³. The recorded sound of the heart was treated with a sound processing software (Ableton Live 8.2.2) to reduce noise using low-pass frequency filters (1.35 kHz) and set the sound to a standard volume. This recording was automatically cut into ten 5s-samples using a computer algorithm. The *own heart samples* consisted on these ten 5s samples, five of which were repeated twice. The *other person's samples* consisted on ten 5s samples of another person (again five of these samples were presented twice), individually matched in terms of heart rate. The mean difference in heart rate between the two samples was 1.47 bpm (SD 1.54).

Experimental task. Participants were seated comfortably with headphones on. They were asked to fixate the screen and concentrate on the sound samples presented over the headphones. They were told that they would be listening to various short audio clips of heartbeat sounds. Eprime (Psychology Software Tools, Pittsburgh, PA) was used to randomly present the pre-recorded samples of their own or another person's heart sounds. A total of 30 5-s samples were presented, which contained the presentation of 15 samples of their own and 15 samples of another person's heartbeat sounds. After each sample, the participants indicated in a forced-choice task whether the heard sound had reproduced their own heart, or not. It was explicitly explained that the presented sounds were offline. After each response they rated on a VAS scale ranging from 0 (not confident at all) to 100 (highly confident) their confidence in their decision. A sub-group of the participants further completed the control task (see below), which was added to the experiment in order to test whether self-recognition of one's own heart could be based on the recognition of heartbeat timing, for example, heart rate variability. In those participants the experimental and control task were carried out in counterbalanced order across participants.

Measure and calculation of interoceptive accuracy. Interoceptive accuracy was measured prior to the experimental task using the "heartbeat counting paradigm"¹³. Participants internally counted their heartbeats during four intervals (25s, 35s, 45s, and 100s) presented in random order. The start and end of the period was signalled by an auditory cue delivered by Eprime via the headphones. The real heart rate was recorded using a pulse transducer (Adinstruments; TN1012/ST), which was attached to the index finger using a Velcro fastener. All participants

reported that they could not feel their pulse on the finger. These two measures were used to calculate an accuracy index as described in¹³.

Measure and calculation of metacognitive ability (Meta- d'). We combined confidence ratings with objective performance in order to estimate participant's metacognitive ability. Classical (or Type I) signal detection theory (SDT) measures quantify the subject's ability to discriminate (d') between the presence (e.g. own-heart sound) and the absence (e.g. other-heart sound) of a stimulus, independently of response biases, i.e. the tendency to favor one type of response over the other⁴⁶. d' can be calculated as $d' = z(\text{Hit rate}) - z(\text{false alarm})$. These measures of objective performance can be extended to evaluate the ability to discriminate between one's own correct and incorrect answers, i.e. metacognitive ability (type 2 SDT)^{46,47}. A close correspondence between confidence and accuracy (e.g. correct trials accompanied by high confidence ratings or wrong trials accompanied by low confidence ratings), indicates that the participants have good knowledge on the correctness of their responses (i.e. a high metacognitive ability). Several SDT-based measures have been proposed to quantify metacognition ability. Here, we used a recently developed measure (meta- d' ; sum-square error approach) to calculate type 2 SDT³² (<http://www.columbia.edu/~bsm2105/type2sdt/>). Importantly, unlike standard type 2 SDT measures⁴⁷, meta- d' is robust to response biases and type I performance^{32,37}. In other words, meta- d' scores are reasonably independent of the tendency to provide one answer type over another (e.g. "my heart" vs "not my heart") as well as the ability to correctly discriminate one's own-heart sound. Moreover, meta- d' and d' are expressed on the same scale and therefore can be directly compared to provide an estimate of how much subjects base their decisions on explicit knowledge^{32,37}. If metacognition were to be optimal, the subjects would be expected to make use of all the information available for the type I task when judging their own performance, with the consequence that meta- $d' = d'$. Conversely, in sub-optimal metacognition conditions meta- $d' < d'$. In that case, because subjects are not able to judge the correctness of own response (i.e. provide high-confidence ratings to incorrect responses and low-confidence ratings to correct responses), it is believed that they were not (fully) aware of the criteria used to guide their performance. Thus, their performance will have been (at least partially) underpinned by implicit processes.

A similar approach has been previously used to estimate metacognitive awareness of interoceptive accuracy in classical measures of interoception (ref. 14). However, in the current study we adopted the standard protocol of the heartbeat counting task, which is comprised by only four trials and is therefore not ideal to estimate a reliable index of metacognitive ability.

Control task. A control task was designed to understand a possible contribution of pure timing effects (i.e. heart rate frequency and variability). The procedure of stimuli creation followed the procedure described above but instead of recording the Doppler sound we used a hardware-based function of Powerlab (www.adinstruments.com) to detect the peak of the finger pulse and at each pulse trigger the presentation (Eprime; Psychology Software tools) of a standard tone (1000 Hz, 100 ms). These tones were recorded using Audacity (<http://audacity.sourceforge.net/>). Then the own and other person's heart sounds were again cut in 5s samples, which were presented as described above.

Debriefing. At the end of the experiment, participants were verbally asked in a semi-structured way, to report how many different hearts they thought the audio samples had been obtained from, as well as how difficult they perceived the different tasks to be. The debriefing was included as a complement to confidence ratings, in particular in order to understand if i) participants felt confident about their performance in this task, i.e. if they felt they could identify the sound of their own heart and if yes, based on which subjective criteria and ii) if they could figure out how many different heartbeats there were.

References

1. Blanke, O. Multisensory brain mechanisms of bodily self-consciousness. *Nat. Rev. Neurosci.* **13**, 556–571 (2012).
2. Damasio, A. & Damasio, H. Minding the Body. *Daedalus* **135**, 15–22 (2006).
3. Botvinick, M. & Cohen, J. Rubber hands 'feel' touch that eyes see. *Nature* **391**, 756 (1998).
4. Ehrsson, H. H. The experimental induction of out-of-body experiences. *Science* **317**, 1048 (2007).
5. Franck, N. *et al.* Defective recognition of one's own actions in patients with schizophrenia. *Am J Psychiatry* **158**, 454–9 (2001).
6. Lenggenhager, B., Tadi, T., Metzinger, T. & Blanke, O. Video Ergo Sum: Manipulating Bodily Self-Consciousness. *Science* **317**, 1096–1099 (2007).
7. Craig, A. D. How do you feel? Interoception: the sense of the physiological condition of the body. *Nat. Rev. Neurosci.* **3**, 655–666 (2002).
8. Apps, M. A. J. & Tsakiris, M. The free-energy self: a predictive coding account of self-recognition. *Neurosci. Biobehav. Rev.* **41**, 85–97 (2014).
9. Craig, A. D. B. How do you feel—now? The anterior insula and human awareness. *Nat. Rev. Neurosci.* **10**, 59–70 (2009).
10. Seth, A. K. Interoceptive inference, emotion, and the embodied self. *Trends Cogn Sci* **17**, 565–73 (2013).
11. Damasio, A. Feelings of emotion and the self. *Ann. N. Y. Acad. Sci.* **1001**, 253–261 (2003).
12. James, W. *The Principles of Psychology*. 2vol, (Macmillan, 1890).
13. Schandry, R. Heart beat perception and emotional experience. *Psychophysiology* **18**, 483–488 (1981).
14. Garfinkel, S. N., Seth, A. K., Barrett, A. B., Suzuki, K., & Critchley, H. D. Knowing your own heart: Distinguishing interoceptive accuracy from interoceptive awareness. *Biol. Psychol.* **104C**, 65–74 (2015).
15. Whitehead, D. W. E., Drescher, V. M., Heiman, P. & Blackwell, B. Relation of heart rate control to heartbeat perception. *Biofeedback Self-Regul.* **2**, 371–392 (1977).
16. Herbert, B. M. & Pollatos, O. The body in the mind: on the relationship between interoception and embodiment. *Top. Cogn. Sci.* **4**, 692–704 (2012).
17. Dunn, B. D., Evans, D., Makarova, D., White, J. & Clark, L. Gut feelings and the reaction to perceived inequity: the interplay between bodily responses, regulation, and perception shapes the rejection of unfair offers on the ultimatum game. *Cogn. Affect. Behav. Neurosci.* **12**, 419–429 (2012).

18. Garfinkel, S. N. *et al.* Fear from the heart: sensitivity to fear stimuli depends on individual heartbeats. *J. Neurosci. Off. J. Soc. Neurosci.* **34**, 6573–6582 (2014).
19. Katkin, E. S., Wiens, S. & Ohman, A. Nonconscious fear conditioning, visceral perception, and the development of gut feelings. *Psychol. Sci.* **12**, 366–370 (2001).
20. Lenggenhager, B., Azevedo, R. T., Mancini, A. & Aglioti, S. M. Listening to your heart and feeling yourself: effects of exposure to interoceptive signals during the ultimatum game. *Exp. Brain Res.* **230**, 233–241 (2013).
21. Aspell, J. E. *et al.* Turning body and self inside out: visualized heartbeats alter bodily self-consciousness and tactile perception. *Psychol. Sci.* **24**, 2445–2453 (2013).
22. Suzuki, K., Garfinkel, S. N., Critchley, H. D. & Seth, A. K. Multisensory integration across exteroceptive and interoceptive domains modulates self-experience in the rubber-hand illusion. *Neuropsychologia* **51**, 2909–17 (2013).
23. van Elk, M., Lenggenhager, B., Heydrich, L. & Blanke, O. Suppression of the auditory N1-component for heartbeat-related sounds reflects interoceptive predictive coding. *Biol. Psychol.* doi: 10.1016/j.biopsycho.2014.03.004 (2014).
24. Khalsa, S. S. *et al.* Interoceptive awareness in experienced meditators. *Psychophysiology* **45**, 671–677 (2008).
25. Kleint, N. I., Wittchen, H.-U. & Lueken, U. Probing the Interoceptive Network by Listening to Heartbeats: An fMRI Study. *Plos One* **10**, e0133164 (2015).
26. Herbert, B. M., Pollatos, O., Flor, H., Enck, P. & Schandry, R. Cardiac awareness and autonomic cardiac reactivity during emotional picture viewing and mental stress. *Psychophysiology* **47**, 342–354 (2010).
27. Schandry, R., Bestler, M. & Montoya, P. On the relation between cardiodynamics and heartbeat perception. *Psychophysiology* **30**, 467–474 (1993).
28. Phua, K., Chen, J., Dat, T. H. & Shue, L. Heart sound as a biometric. *Pattern Recognit.* **41**, 906–919 (2008).
29. Jokisch, D., Daum, I. & Troje, N. F. Self recognition versus recognition of others by biological motion: viewpoint-dependent effects. *Perception* **35**, 911–920 (2006).
30. Sevдалис, V. & Keller, P. E. Cues for self-recognition in point-light displays of actions performed in synchrony with music. *Conscious. Cogn.* **19**, 617–626 (2010).
31. Cook, R., Johnston, A. & Heyes, C. Self-recognition of avatar motion: how do I know it's me? *Proc. Biol. Sci.* **279**, 669–674 (2012).
32. Maniscalco, B. & Lau, H. A signal detection theoretic approach for estimating metacognitive sensitivity from confidence ratings. *Conscious. Cogn.* **21**, 422–430 (2012).
33. Lee, I. A. & Preacher, K. J. Calculation for the test of the difference between two dependent correlations with one variable in common [Computer software]. (2013).
34. Tsakiris, M., Tajadura-Jimenez, A. & Costantini, M. Just a heartbeat away from one's body: interoceptive sensitivity predicts malleability of body-representations. *Proc Biol Sci* **278**, 2470–6 (2011).
35. Wöllner, C. Self-recognition of highly skilled actions: a study of orchestral conductors. *Conscious. Cogn.* **21**, 1311–1321 (2012).
36. Damasio, A. *Looking for Spinoza: Joy, Sorrow, and the Feeling Brain* (Houghton Mifflin Harcourt, 2003).
37. Barrett, A. B., Dienes, Z. & Seth, A. K. Measures of metacognition on signal-detection theoretic models. *Psychol. Methods* **18**, 535–552 (2013).
38. Bartfeld, P. *et al.* Distinct patterns of functional brain connectivity correlate with objective performance and subjective beliefs. *Proc. Natl. Acad. Sci. USA* **110**, 11577–11582 (2013).
39. Canales-Johnson, A. *et al.* Auditory Feedback Differentially Modulates Behavioral and Neural Markers of Objective and Subjective Performance When Tapping to Your Heartbeat. *Cereb. Cortex N. Y. N 1991*, doi: 10.1093/cercor/bhv076 (2015).
40. Fairclough, S. H. & Goodwin, L. The effect of psychological stress and relaxation on interoceptive accuracy: Implications for symptom perception. *J. Psychosom. Res.* **62**, 289–295 (2007).
41. Knapp-Kline, K. & Kline, J. P. Heart rate, heart rate variability, and heartbeat detection with the method of constant stimuli: slow and steady wins the race. *Biol. Psychol.* **69**, 387–396 (2005).
42. Herbert, B. M. & Pollatos, O. Attenuated interoceptive sensitivity in overweight and obese individuals. *Eat. Behav.* **15**, 445–448 (2014).
43. Bom, N. *New Concepts in Echocardiography* (Springer Science & Business Media, 1972).
44. Kostis, J. B., Fleischmann, D. & Bellet, S. Use of the Ultrasonic Doppler Method for Timing of Valvular Movement Application in the Differential Diagnosis of Extra Heart Sounds. *Circulation* **40**, 197–207 (1969).
45. Yoshida, T., Mori, M., Nimura, Y., Takagishi, S. & Nakanishi, K. Studies on the Time of Valvular Movements in Mitral Valvular Disease with Ultrasonic Doppler Method. *Jpn. Heart J.* **1**, 261–274 (1960).
46. Macmillan, N. & Creelman, C. D. *Detection theory: A user's guide* (2nd ed.). (Lawrence Erlbaum, 2005).
47. Kunimoto, C., Miller, J. & Pashler, H. Confidence and accuracy of near-threshold discrimination responses. *Conscious. Cogn.* **10**, 294–340 (2001).

Acknowledgements

We thank Jane Aspell and Vivien Ainley for proofreading and their helpful comments. Financial support: Swiss National Science Foundation (grant PBELP3-133268) to BL. Fundação para a Ciência e a Tecnologia (FCT) and POPH/FSE (doctoral grant SFRH/BD/60517/2009) to RTA. EU Information and Communication Technologies Grant (VERE project, FP7-ICT-2009-5, Prot. Num. 257695), the Italian Ministry of Health (and RF-2010-2312912) to SMA.

Author Contributions

All authors conceived the project; R.T.A. and B.L. designed the experiments, collected data and analysed the data and wrote the first version of the manuscript; S.M.A. revised the manuscript; all authors read and approved the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Azevedo, R. T. *et al.* Participants' above-chance recognition of own-heart sound combined with poor metacognitive awareness suggests implicit knowledge of own heart cardiodynamics. *Sci. Rep.* **6**, 26545; doi: 10.1038/srep26545 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>