



Kent Academic Repository

Cottier, Fabien and Hall, Rebecca A. (2020) *Face/Off: The Interchangeable Side of Candida Albicans*. *Frontiers in Cellular and Infection Microbiology*, 9 .

Downloaded from

<https://kar.kent.ac.uk/80212/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.3389/fcimb.2019.00471>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal* , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).



Face/Off: The Interchangeable Side of *Candida Albicans*

Fabien Cottier¹ and Rebecca A. Hall^{1,2*}

¹ School of Biosciences, Institute of Microbiology and Infection, University of Birmingham, Birmingham, United Kingdom,

² Kent Fungal Group, School of Biosciences, University of Kent, Canterbury, United Kingdom

Due to limited mobility, fungi, like most unicellular organisms, have evolved mechanisms to adapt to sudden chemical and/or physical variation in their environment. *Candida albicans* is recognized as a model organism to study eukaryotic responses to environmental changes, as this human commensal yeast but also opportunistic pathogen responds to numerous environmental cues through switching morphologies from yeast to hyphae growth. This mechanism is largely controlled by two major pathways: cAMP-PKA and MAPK, but each environmental signal is sensed by specific sensors. However, morphological switching is not the only response *C. albicans* exerts in response to environmental cues. Recently, fungal cell wall remodeling in response to host-derived environmental cues has been identified as a way for *C. albicans* to manipulate the innate immune system. The fungal cell wall is composed of a chitin skeleton linked to a network of β -glucan, which anchors proteins and mannans to the fungal cell surface. As localized on the cell surface, these molecules drive interactions with the environment and other cells, particularly with host immune cells. *C. albicans* is recognized by immune cells such as neutrophils and macrophages via pathogen recognition receptors (PRRs) that bind different components of the cell wall. While β -glucan and mannan are proinflammatory molecules, chitin can induce anti-inflammatory responses. Interestingly, *C. albicans* is able to regulate the exposure of these pathogen-associated molecular patterns (PAMPs) according to environmental cues resulting in a modulation of the host immune response. This review describes the mechanisms involved in *C. albicans* response to environmental changes and their effect on immune recognition.

Keywords: *Candida*, cell wall, innate immunity, morphogenesis, cell wall remodeling

OPEN ACCESS

Edited by:

Floyd Layton Wormley,
Texas Christian University,
United States

Reviewed by:

Robert T. Wheeler,
University of Maine, United States
Ana Traven,
Monash University, Australia

*Correspondence:

Rebecca A. Hall
r.a.hall@bham.ac.uk

Specialty section:

This article was submitted to
Fungal Pathogenesis,
a section of the journal
Frontiers in Cellular and Infection
Microbiology

Received: 09 September 2019

Accepted: 20 December 2019

Published: 28 January 2020

Citation:

Cottier F and Hall RA (2020) Face/Off:
The Interchangeable Side of *Candida*
Albicans.
Front. Cell. Infect. Microbiol. 9:471.
doi: 10.3389/fcimb.2019.00471

INTRODUCTION

Candida albicans is an opportunistic fungal pathogen of man and is associated with infections ranging from superficial mucosal infections to life-threatening disseminated disease. In 1853, Charles Philippe Robin drew one of the first representations of *C. albicans*, from a muguet infection isolate. On his illustration, yeasts, filaments, and clusters of blastoconidida were represented (Histoire Naturelle des Végétaux Parasites qui Croissent sur l'Homme et sur les Animaux Vivants.). However, since then, *C. albicans* has been described to exist in eight different morphotypes including four yeast morphologies (white, opaque, gray, and gut), two hyphal morphologies (linear and sinusoidal), pseudo-hyphae, and chlamydospores (Hayes, 1966; Slutsky et al., 1987; Brand et al., 2009; Pande et al., 2013; Tao et al., 2014). In addition to this large diversity in shape, *C. albicans* can also exist in unicellular culture, biofilms, and micro-colonies (McCall et al., 2018). This polymorphic nature of *C. albicans* is a great example of a protean organism.

Most of these various morphotypes have been observed clinically, suggesting that each morphotype plays a specialized role in infection. On one hand, *C. albicans* is a commensal yeast, frequently isolated from organs such as the female reproductive tract and the gastrointestinal tract where it can represent up to 0.1% of the microbiome (Beigi et al., 2004; Cottier et al., 2018). However, *C. albicans* is also responsible for mild skin or mucosal infections such as the muguet (oral candidiasis) or vulvovaginal candidiasis, which affects 70% of women (Jeanmonod and Jeanmonod, 2019). On the other end of the spectrum, *C. albicans* bloodstream infections represent over 50% of all reported candidemias and affects eight out of 100,000 people causing life-threatening infections in immunodeficient patients (Pfaller and Diekema, 2007). The breadth and depth of candidiasis bring pressure on medical services as time and cost of treatment amount to several billions dollars per year (Moran et al., 2010).

Historically, the morphological yeast-to-hyphae switch of *C. albicans* has been an intense focus of research as it is one of the main drivers of fungal virulence, and is a visually drastic phenotype. Cells locked in either of these two forms display a lower virulence in murine infection models (Lo et al., 1997; Saville et al., 2003). However, the discovery of the novel hyphal specific fungal cytolytic toxin, candidalysin, revealed that it is the production of this toxin that is essential for virulence, rather than virulence being truly attributed to hyphal formation alone, as *ece1* deficient hyphae are attenuated in virulence (Moyes et al., 2016). Additional work enhanced this dichotomy between yeast-to-hyphae switching and pathogenicity (Noble et al., 2010). Now, it is appreciated that it is not only the morphology of *C. albicans* that is important for virulence, but also the metabolic state of the fungus. Indeed, while hyphae and proteins such as extracellular hydrolytic enzymes and candidalysin could be considered as the spear of *C. albicans*, cell wall remodeling could be viewed as a shield against the innate immune response. This review explores the environmental factors influencing morphological switches in *C. albicans* but also most recent studies demonstrating the role of the environment in regulating the composition of the fungal cell wall. Finally, how this plasticity of *C. albicans* aids exploitation of the innate immune system will be explored.

C. albicans Morphology Opaque, GUT, and Gray Cells

The main morphotype of *C. albicans* is a yeast shape known as white cells. These cells are common in *in vitro* studies, have a round-to-oval shape, and measure around 5–6 μm (Klis et al., 2014). This form is widely observed in *in vivo* studies and clinical isolates and is efficient for dissemination and adhesion to surfaces. The first morphotype derived from the white cell was the opaque cell (Figure 1), the mating competent form of *C. albicans* (Slutsky et al., 1987). Opaque cells have an ellipsoidal shape of around 7 μm long, with pimples on the surface (Lockhart et al., 2002). These cells are the result of a loss of heterozygosity at the *MTL* locus, leading to the formation of α and α cells (Miller and Johnson, 2002). Sexually competent *MTL* α cells produce a pheromone that triggers “shmooing,” the formation of a conjugation tube, in *MTL* α opaque cells (Lockhart et al., 2003).




FIGURE 1 | *Candida albicans* morphologies. Representation of the reported morphotypes of *C. albicans* and the known cues triggering morphological switches.

This initial mating steps ends with the fusion of the two nuclei and formation of a tetraploid cell. Several environmental factors induce white-to-opaque switching like acidic pH, hypercapnia, and exposure to N-acetylglucosamine (Huang et al., 2009, 2010; Sun et al., 2015). These cues are sensed by two systems, Sac7/Rho1 and Ras1, ending in the regulation of the master white to opaque transcriptional regulator Wor1 (Yang et al., 2018).

Recently, the white-to-opaque switch was reported to be part of a tristable phenotypic switch with the discovery of gray cells (Tao et al., 2014). These cells have also an ellipsoidal shape and are similar in length to the opaque cell but with a smaller diameter and do not have pimples on their surface. While gray cells are thousand-fold more efficient than white cells at mating, gray cells are hundred-fold less efficient than opaque cells at mating (Tao et al., 2014). Similar to opaque cells, gray cells are the result of a genetic alteration. Earlier this year, Liang et al. demonstrated that under certain conditions, such as passage through the mammalian gastrointestinal tract, *C. albicans* can gain mutations in the transcription factor *Efg1*, primarily known for its role in morphogenesis, but also promotes the formation of white cells (Liang et al., 2019). A strain of *C. albicans* hemizygous for *EFG1* has a white cell morphology, but any functional loss of the remaining *EFG1* wild-type allele results in formation of gray cells. Such events occur in the human host, with gray cells containing mutations in *EFG1* being isolated from clinical samples (Liang et al., 2019). Furthermore, gray cells display a fitness advantage in mice gastrointestinal colonization models, which correlates with previous work demonstrating the competitive advantage of

an *efg1* mutant over a wild-type strain (Pierce and Kumamoto, 2012). However, these observations were performed in antibiotic-treated mice; therefore, the impact of *efg1* inactivation and gray cells in a complex gut microbiome remains to be characterized.

In addition to gray cells, another *C. albicans* morphotype, gastrointestinal induced transition (GUT), has been associated with gastrointestinal colonization. GUT cells are similar in size and shape to opaque cells but do not possess pimples on their surface (Pande et al., 2013). GUT cells are very similar to gray cells except GUT cells tend to be wider. GUT cells were identified during gastrointestinal colonization studies in mice using a strain over-expressing the transcription factor *WOR1*. Similarly to gray cells, GUT cells present a fitness advantage in gut colonization compared to control white cells (Pande et al., 2013). They are also significantly less efficient at mating than gray cells. The main difference between gray and GUT cells is a lack of stability, with GUT cells transitioning back to white cells outside the gut environment, indicating the requirement for an external stimulus to maintain this phenotype (Pande et al., 2013). Given the similarity between gray and GUT cells, it would be interesting to identify if GUT cells are the result of *EFG1* suppression, which is relieved when removed from the gut environment.

Yeast-to-Hyphal Transition

The most studied morphological switch of *C. albicans* is the formation of hyphae. During hyphal formation, yeast stop forming buds, and grow tube-shaped cells linked to each other by a septum. This structure allows communication between the cytoplasm of the interconnecting cells and creates a mycelium. Hyphae have a lower diameter compared to yeast, around 2.5–3.5 μm (Klis et al., 2014), but their length can easily be over 100 μm , surpassing the size of immune cells. This morphological transition is triggered by various known environmental cues present in the human host including 37°C, pH, hypercapnia, low nitrogen, N-acetylglucosamine, peptidoglycan, mannitol, or amino acids (Figure 1) (Noble et al., 2017). Most of these signals activate either the cAMP-PKA or MAPK pathways resulting in the activation of the transcription factors Efg1 and Cph1, respectively, as reviewed in Noble et al. (2017). Interestingly, deletion of the transcriptional factor *UME6*, which play a central role in morphogenesis of *C. albicans* (Carlisle and Kadosh, 2010), is deficient in hyphal formation in *in vitro* assays but is still able to produce hyphae in the mouse gastrointestinal tract, revealing the existence of at least one unidentified hyphal inducing signal (Witchley et al., 2019).

Hyphae are a major component of *C. albicans* biofilms, which have a higher resistance to antifungal drugs than planktonic cultures (Mathe and Van Dijck, 2013). However, hyphal differentiation is also a strategy for yeast to escape immune cells after phagocytosis. Initially, this escape mechanism was thought to be due to mechanical damage of the phagocyte. However, it is now understood that escape of *C. albicans* from macrophages is due to an intricate mechanism requiring several key processes. For example, hyphae can trigger phagocyte killing through caspase-1-dependent pyroptosis (Uwamahoro et al., 2014; Wellington et al., 2014), and then once extracellular, these hyphae induce glucose starvation resulting in enhanced macrophage cell death (Tucey et al., 2018). Alternatively, hyphal

differentiation in immune cells induces the production of the cytotoxin candidalysin resulting in phagocyte membrane damage (Kasper et al., 2018).

In addition to these straight tubular hyphae, sinusoidal or helical hyphal growth has been reported in *C. albicans* when cultivated on surfaces (Sherwood-Higham et al., 1994; Brand et al., 2009; McCall et al., 2018), and are associated with antifungal resistance (Sherwood-Higham et al., 1994). However, hyphae are not solely a morphological switch, but they are also associated with important transcriptional regulation of virulence factors including secreted aspartyl protease, adhesins and superoxide dismutase (Mayer et al., 2013), and, more recently, the cytotoxic peptide candidalysin encoded by *ECE1* (Moyes et al., 2016).

Pseudo-hyphae are a morphological form sharing aspects of yeast, as cells have an oval to ellipsoid shape, and like hyphae can form a mycelium. Contrary to hyphae, pseudo-hyphae are not tube shaped but are elongated yeasts with separation between each cell that is easily identified by a change in diameter (Sudbery et al., 2004). Pseudo-hyphae are observed in response to various cues inducing hyphae formation, but none are known to differentiate solely pseudo-hyphae. The relevance of this morphotype in *C. albicans* life cycle remains to be studied.

Chlamydospores

The last identified *C. albicans* morphotype is the chlamydospore. These large round cells have a diameter of 7–8 μm and are characterized by a thick cell wall (Hayes, 1966). However, the latter does not provide significant resistance to heat or starvation (Citiulo et al., 2009). Nevertheless, several environmental cues have been identified to trigger chlamydospore differentiation such as light, starvation, or hypoxia, but the biological function of this morphotype remains to be elucidated (Bottcher et al., 2016).

As reported, *C. albicans* responds to various environmental cues by producing different morphotypes. While cell shape is immediately observable, each of these forms is also associated with major transcriptional responses ranging from virulence factors to metabolic pathway regulation (Tuch et al., 2010; Palige et al., 2013; Guedouari et al., 2014; Tao et al., 2014; Grahl et al., 2015; Koch et al., 2018). Several environmental signals have also been shown to influence the expression of secreted, as well as cell wall-associated, proteins. As a result, cell wall remodeling has recently been described to occur in response to these changes (Ballou et al., 2016; Sherrington et al., 2017; Lopes et al., 2018). For example, chitin, which is a polymer of β -(1,4)-N-acetylglucosamine, is regulated by a balance between production via chitin synthases (*CHS1-3*, *CHS8*) and degradation by chitinases (*CHT1-4*), allowing *C. albicans* to reshape its chitin network when needed (Selvaggini et al., 2004). Cell wall characterization and versatility is another aspect of *C. albicans* ever-changing faces whose understanding is critical as cell wall is the initial point of contact between fungal cells and the host immune system (Hopke et al., 2018).

C. albicans Cell Wall Remodeling

C. albicans asymptotically colonize human skin, vagina, and gastrointestinal tract, niches where temperature, gas composition, or carbon source vary greatly through space as well as time (Grahl et al., 2012). As discussed above, *C. albicans*

responds to changes in these environmental cues by altering its shape. Transcriptional responses of *C. albicans* to these signals have been characterized and point toward customized profiles of gene expression. Interestingly, genes involved in cell wall structure have been shown to be regulated in response to pH, morphogenesis, hypoxia, or carbon source (Bensen et al., 2004; Synnott et al., 2010; Ballou et al., 2016).

The *C. albicans* cell wall is composed of three major components: (i) a chitin skeleton, (ii) a network of β -(1,3)- and β -(1,6)-glucan, and (iii) highly glycosylated proteins sometimes referred to as mannoproteins. In *C. albicans* white cells, chitin represents 1–2% of cell wall dry weight, while β -1,6-glucan and β -1,3-glucan amount for, respectively, 20 and 40% and the remaining 35–40% correspond to the mannoproteins (Navarro-Arias et al., 2019). With the help of transmission electron microscopy, we can now visualize the individual layers of the cell wall. The inner layer (chitin and glucan) forms a relatively electron dense region, while the outer layer formed by the mannoproteins is composed of densely packed fibrils extending away from the fungus (Chaffin et al., 1998). Adaptation to environmental conditions alters the composition of both cell wall layers. For example, the thickness of the outer mannan layer in white cells is dependent on temperature and nutrient availability, with cells grown in rich media at 37°C displaying shorter mannan fibrils than cells grown in minimal media at 30°C (Ene et al., 2012; Sherrington et al., 2017). Temperature also regulates the thickness of the inner cell wall layer, again with higher temperatures reducing the overall thickness of the glucan–chitin layer. The thickness of the hyphal cell wall is comparable to the yeast cell wall, but can be greater at the hyphal tip (Knafler et al., 2019). The thickness of the cell wall is also modulated by pH, with growth in acidic media resulting in a 50% reduction in the outer mannan layer, while the inner layer is not significantly affected by pH (Sherrington et al., 2017). Interestingly, alkaline pH has the opposite effect, increasing the outer mannan layer. A change in carbon source, from glucose to lactate in minimum media, or exposure to hypoxic conditions also reduces the thickness of the inner layer by 50% (Ballou et al., 2016).

Cell Wall Remodeling in Response to Lactate

Despite the fact that the inner layer is covered by the outer layer of mannoproteins, which has previously been described as a molecular shield to protect the highly immunogenic glucan from the innate immune system, adaptation to environmental signals results in exposure of the glucan and chitin network, making it accessible to immune cells (Sherrington et al., 2017). Notably, β -(1,3)-glucans are recognized by the pattern recognition receptor (PRR) Dectin-1 triggering phagocytosis of *C. albicans* by neutrophils and macrophages (Taylor et al., 2007). Cell wall remodeling, resulting in the masking and unmasking of PAMPs like β -glucan and chitin, is an intricate mechanism of *C. albicans* survival and virulence in the human body. Similar to cell shape or cell wall thickness, PAMP exposure is influenced by environmental cues like carbon sources. Indeed, cells incubated for 4 h at 30°C in minimal media supplemented with lactate display masking of the β -glucan (>50% reduction in glucan

exposure) (Ballou et al., 2016). This decline in β -glucan exposure resulted in lower phagocytic recruitment (Ene et al., 2013). Cell wall remodeling in response to lactate involves the plasma-membrane receptor Gpr1 and its associated G α protein Gpa2 as a double mutant exhibited a lack of glucan masking in the presence of lactate. Interestingly, these two proteins are known to activate the cAMP-PKA pathway during the yeast-to-hyphal transition, but mutants in this pathway (*cyr1 Δ* and *pde2 Δ*) did not have any impact on cell wall remodeling in response to lactate. Mutants involved in the calcium signaling pathway (*cna1* and *cnb1*) were also unaffected for lactate-induced cell wall remodeling, while the downstream regulator Crz1 was shown to be essential for β -glucan masking (Ballou et al., 2016). These experiments suggest the existence of a novel Crz1-dependent signaling pathway involved in cell wall remodeling in response to lactate.

Cell Wall Remodeling in Response to Hypoxia

A second environmental factor inducing a 2-fold decrease in glucan exposure is hypoxia, a cue particularly important as *C. albicans* encounters low oxygen environments during colonization of internal organs (Pradhan et al., 2018). However, none of the components previously identified for lactate-induced glucan masking play a role in the response to hypoxia. Instead, hypoxia-induced cell wall remodeling is regulated predominately via the cAMP-PKA pathway (Pradhan et al., 2018). Here, hypoxia triggers the production of reactive oxygen species (ROS) from the mitochondria in a Goa1- and Upc2-dependent mechanism. This ROS is then converted to hydrogen peroxide by Sod1, which then activates the cAMP-PKA signaling cascade, through an as yet to be identified mechanism, which then functions to induce glucan masking (Pradhan et al., 2018). Interestingly, inhibition of mitochondrial respiration by exposure to sodium nitroprusside (SNP) and alicyl-hydroxamic acid (SHAM) for 18 h in rich media at 30 or 37°C induces both β -glucan and chitin exposure (Duvenage et al., 2019). The role of the cAMP-PKA pathway in this condition has not been investigated, but contrary to hypoxic exposure, *UPC2* inactivation did not have any impact on PAMP exposure. However, the transcription factor Sko1, involved in the cell wall damage response, displayed an increase in chitin exposure when incubated with SNP and SHAM (Duvenage et al., 2019), suggesting that multiple mitochondrial inputs regulate cell wall remodeling. Importantly, *C. albicans* growth and viability are significantly reduced when exposed to these molecules, which could have secondary effects on PAMP exposure.

Cell Wall Remodeling in Response to Environmental pH

Environmental pH also induces cell wall remodeling, but in contrast to lactate and hypoxia, acidic environments promote the exposure of glucan and chitin exposure (Sherrington et al., 2017). Crz1 was shown to have no role in chitin exposure in response to acidic environment, but the transcription factors Rim101 and, to a lesser extent, Bcr1 play key roles in regulating chitin exposure. This effect was mediated through the regulation of the chitinase, *CHT2*, suggesting that this enzyme is responsible for the unmasking of chitin in the cell wall. Inactivation of

CHT2 induced a significant increase in chitin exposure at pH 6 compared to a control strain (Sherrington et al., 2017). Furthermore, *CHT2* transcript levels are up-regulated during growth at pH 6 compared to pH 4, but this regulation is absent in both *rim101Δ* and *bcr1Δ* mutants (Sherrington et al., 2017). Noticeably, *CHT2* transcript levels have been reported to be a target of She3, an mRNA transporter to the cell wall, and inactivation of *SHE3* resulted in an increase of chitin exposure at pH 6 similar to a *cht2* mutant (Sherrington et al., 2017). Altogether, acidic environments inactivate the Rim101 and Bcr1 pathways, reducing *CHT2* expression, leading to a lack of chitinase in the cell wall and enhancing exposure of chitin in the cell wall (Sherrington et al., 2017).

Cell Wall Remodeling in Response to Multiple Signals

While these environmental cues are studied separately *in vitro*, in the host, *C. albicans* encounters many of these signals, and more, simultaneously. For example, in murine subdermal abscess models, infiltration of neutrophils to the site of *C. albicans* infection contributes to the formation of a hypoxic environment (Lopes et al., 2018). Furthermore, neutrophil-dependent production of lactate is increased by 40% in anoxia compared to normoxia *in vitro* and supernatants of such cultures reduce glucan exposure. Therefore, neutrophil recruitment to sites of *C. albicans* infection could form a local environment that is low on oxygen and rich in lactate (Lopes et al., 2018). This metabolic rewiring was confirmed in human monocytes, as glucose consumption and lactate production were increased in the supernatants of monocytes exposed to *C. albicans* compared to non-stimulated cells (Dominguez-Andres et al., 2017). *In vivo*, glycolysis was also shown to be up-regulated in patients with fungal sepsis compared to healthy controls during genome-wide transcriptional profiles of blood (Cheng et al., 2016). Such conditions would trigger cell wall remodeling, masking critical PAMPs such as glucan, preventing immune recognition, and allowing *C. albicans* colonization and evasion of the host.

Taken together, it appears as though *C. albicans* has evolved several signaling pathways that, in response to particular stimuli, regulate cell wall remodeling (Figure 2). Currently only a few host-derived environmental signals have been investigated for their ability to induce cell wall remodeling, and it is likely that this cell wall plasticity will extend to many other environmental parameters. Given that the cell wall forms the exterior of the fungus, and is in direct contact with the innate immune system, these cell wall perturbations will have a significant effect on how the invading pathogen is sensed by the innate immune cells.

Immune Responses


Using cell wall remodeling, *C. albicans* is able to partially control the innate immune response mounted against it. Masking and unmasking of PAMPs such as β -glucan and chitin can trigger immune suppression as well as immune tolerance (Romani, 2011). Indeed, *C. albicans* is not alone in shielding its glucan to avoid detection from the innate immune system. Upon encountering the host environment, *Histoplasma capsulatum* completely removes its β -glucan from the cell wall, replacing it with the less proinflammatory α -glucan. Residual β -glucan is

trimmed from the cell surface via the Endo and Exoglucanases Eng1 and Exg8 (Garfoot et al., 2017).

Recognition of Glucan

As mentioned previously, β -glucans are the main component of *C. albicans* cell wall as these β -D-glucose polysaccharides represent around 60% of the yeast dried cell wall mass (Ruiz-Herrera et al., 2006). β -(1,3)-glucans are known to be the ligand of the PRR Dectin-1. Interestingly, while live *C. albicans* yeast were able to activate Dectin-1, a similar amount of hyphal biomass did not (Gantner et al., 2005). This was attributed to the fact that budding yeasts have bud scars on their surface that expose glucan, which are lacking in hyphae. This mechanism demonstrates the relevance of PAMP exposure in the activation of the immune response. Indeed, the level of β -glucan exposure correlates with macrophage phagocytic rates, with environments that promote glucan exposure, also promoting phagocytosis (Ballou et al., 2016; Sherrington et al., 2017; Lopes et al., 2018). Similar observations were made when looking at the recruitment of innate immune cells to the site of infection, with glucan-exposed cells recruiting significantly more phagocytes than cells that had masked their glucan (Ballou et al., 2016; Sherrington et al., 2017). Additionally, human peripheral blood mononuclear cells (PBMCs) exposed to *C. albicans* grown under hypoxic conditions secrete a lower amount of IL-10 and RANTES and to a lesser extent TNF α and CCL3 (Pradhan et al., 2018). While for lactate-grown *C. albicans*, PBMCs are shown to produce a lower amount of IL-17, but human M1-activated monocyte-derived macrophages secreted a lower amount of TNF α and CCL3 compared to the glucose grown control (Ballou et al., 2016). As expected, an increase in β -glucan exposure such as in response to acidic environment induced the opposite secretion of cytokines. Indeed, *C. albicans* cells grown at pH 4 induce a higher level of TNF α , IFN- γ , IL-6, and IL-1 β secretion by PBMCs compared to cells grown at pH 6 (Sherrington et al., 2017). Therefore, by controlling the level of glucan exposure, *C. albicans* can affect the level of proinflammatory cytokine produced and the recruitment of immune cells.

Given that Dectin-1 is the predominate β -glucan recognition receptor on innate immune cells, it is not surprising that the above effects are Dectin-1 dependent. However, although blocking Dectin-1 reduced phagocytosis of *C. albicans* with high glucan exposure, this had minimal effect on the ability of *C. albicans* to attach to the macrophage surface (Sherrington et al., 2017). Hence, Dectin-1 is involved in the enhancement of phagocytosis in response to β -glucans exposure but is not the primary receptor for the attachment of macrophages to *C. albicans*. Dectin-1 knockout mice were shown to be significantly more sensitive to *C. albicans* intravenous infection, as well as displaying a large fungal burden in the mouse gastrointestinal tract compared to the control group (Taylor et al., 2007). Furthermore, in an intraperitoneal infection model, Dectin-1 knockout mice were less able to recruit neutrophils, monocytes, and eosinophils to the site of infection than wild-type mice and this was associated with a decrease in IL-6, CCL2, CCL3, G-CSF, and GM-CSF production (Taylor et al., 2007). Interestingly, *C. albicans* virulence as well as kidney



fungal burden in Dectin-1 knockout mice is dependent on the yeast strain (Marakalala et al., 2013). Indeed, *C. albicans* cells from clade 4 do not display increased virulence compared to wild-type mice. This variable influence of Dectin-1 in *C. albicans* recognition seems to be affected by cell wall chitin content (Marakalala et al., 2013).

While Dectin-1 is central to β -glucan recognition on various immune cells such as macrophage, neutrophils, or dendritic cells, other receptors have been identified to act in this process. Galectin-3, a soluble protein, physically interacts with Dectin-1 to enhance $\text{TNF}\alpha$ production in macrophages exposed to *C. albicans* β -glucan (Esteban et al., 2011). Galectin-3 also modulates complement receptor 3 (CR3) activation. CR3 is able to bind β -(1,6)-glucan, inducing phagocytosis and ROS production. However, galectin-3-deficient neutrophils had a higher level of ROS than the control strain after exposure to *C. albicans*. This effect is mediated via a physical interaction between galectin-3 with the CR3 downstream component Syk (Wu et al., 2017). Interestingly, additional receptors are suspected to recognize β -glucan as PBMCs have been identified to produce IL-1Ra in response to *C. albicans* hyphae in a Dectin-1- and CR3-independent manner (Smeekens et al., 2015).

Recognition of Chitin

As a component of the inner cell wall, chitin is normally not accessible to immune cells (Mora-Montes et al., 2011), but acidic environments increase chitin exposure in *C. albicans*. Purified chitin from *C. albicans* alters immune recognition of this yeast. Pre-treatment of PBMCs with chitin before exposure to live *C. albicans* decreases the secretion of proinflammatory cytokines, particularly IL-1 β , IL-6, and $\text{TNF}\alpha$ (Mora-Montes et al., 2011). This effect is mediated by Dectin-1, as addition of chitin to Dectin-1-deficient mouse macrophages failed to inhibit the production of $\text{TNF}\alpha$. However, this effect does not involve direct binding between chitin and Dectin-1, implying cooperation between multiple PRRs (Mora-Montes et al., 2011). In addition to dampening the secretion of proinflammatory cytokines, chitin alone is sufficient to induce the secretion of anti-inflammatory cytokines like IL-10 (Wagener et al., 2014). In conflicting studies, chitin also induced the secretion of proinflammatory cytokines including IL-6 and $\text{TNF}\alpha$ (Wagener et al., 2014). However, it appears as though the size of the chitin particle plays an interesting role here. While small chitin particles are anti-inflammatory and inhibit the secretion of proinflammatory cytokines, larger chitin particles

(40–70 μm) function to induce proinflammatory cytokine production. These opposite effects are driven through the interaction with different PRRs. Smaller chitin particles appear to mediate their effects through Dectin-1/TLR2 signaling and the mannose receptor (MR) (Da Silva et al., 2009). *NOD2*, *TLR9*, and their downstream effectors are also important for chitin-induced secretion of IL-10 (Wagener et al., 2014), suggesting that a complex network is at play here. The anti-inflammatory response induced by chitin could be the result of co-evolution between the host and *C. albicans* preventing further immune inflammation once the yeast has been killed. However, *C. albicans* could take advantage of this mechanism in a certain environment or condition to lessen the immune response.

Recognition of Mannan

Finally, mannoproteins from *C. albicans* outer cell wall layer are themselves targets of PRRs, but their composition also affects the exposure of the underlying network of glucan. Using super resolution imagery of several mannosylation mutants (*mnn2* Δ , *mnn2* Δ /*mmn6* Δ , *mnn2* Δ /*mnn21* Δ /*mnn22* Δ /*mnn23* Δ /*mnn24* Δ /*mnn26* Δ), Graus et al. were able to demonstrate that a decrease in acid labile mannan abundance and α -mannan backbone length results in a larger region of glucan exposure on the yeast surface (Graus et al., 2018). By controlling mannosylation on surface proteins, *C. albicans* possess a mechanism to adjust the level of β -glucan being exposed. These results are in line with previous observations showing that inactivation of mannosyltransferase induces a higher level of glucan exposure and increases the level of TNF α secretion (Mora-Montes et al., 2010; Ueno et al., 2013; Zhang et al., 2016).

PRRs such as DC-SIGN recognize *N*-linked mannans and trigger *C. albicans* phagocytosis by human dendritic cells as well as the secretion of IL-6 (Cambi et al., 2008). The mannose receptor (MR) is a C-type lectin present on the surfaces of macrophages and is able to recognize mannose conformation. MR can be shed from the macrophage surface and be released in the environment, enhancing *C. albicans* recognition by the immune system (Gazi et al., 2011). TLR4, which recognizes *O*-linked mannan, works alongside MR to release TNF α , IL-6, IL-10, and IFN- γ (Netea et al., 2006). Dectin-2 and -3 form heterodimers recognizing α -mannans and are involved in secretion of TNF α in RAW264.7 macrophages (Zhu et al., 2013). Furthermore, knockout mice of either of these receptors display a higher mortality rate following intravenous

infection as well as an increase in kidney fungal burden (Zhu et al., 2013; Ifrim et al., 2016).

In vitro experiments revealed various interactions between fungal molecules and mammalian receptors, reflecting the co-evolution between these organisms. As observed in mice, *C. albicans* recognition triggers immune cell recruitment, which, in the process of killing the yeast, also produces environmental cues that *C. albicans* senses to mask cell surface PAMPs in order to enhance its survival. Additionally, *in vitro* experiments suggest that acidic environments induce glucan exposure, which could be the source for the highly proinflammatory, non-protective innate immune response observed during vulvovaginal candidiasis (Yano et al., 2018).

CONCLUSION

The dynamism and diversity exhibited by *C. albicans* in response to environmental cues impact not only its shape and structure but also its interaction with immune cells and responses mounted against the yeast. This protean behavior might explain the successful colonization by *C. albicans* of so many different human niches, but also provides few options for this commensal organism to become pathogenic. While several morphogenesis pathways and virulence factors have been identified, the exact processes governing *C. albicans* masking and unmasking of PAMPs in response to environmental signals require deeper characterization. This knowledge could help develop alternative or complementing strategies to cure *C. albicans* infection at a time where antifungal-resistant strains are becoming a concern.

AUTHOR CONTRIBUTIONS

FC wrote the initial draft of the manuscript. RH finalized the manuscript.

FUNDING

This work was funded by an MRC Career Development Award fellowship (MR/L00903X/1) and by the BBSRC (BB/R00966X/1).

ACKNOWLEDGMENTS

The authors would like to acknowledge all contributions to the field of sensing, cell wall biogenesis, and immune response to *Candida albicans* that could not be included in this review due to space limitation.

REFERENCES

- Ballou, E. R., Avelar, G. M., Childers, D. S., Mackie, J., Bain, J. M., Wagener, J., et al. (2016). Lactate signalling regulates fungal beta-glucan masking and immune evasion. *Nat. Microbiol.* 2:16238. doi: 10.1038/nmicrobiol.2016.238
- Beigi, R. H., Meyn, L. A., Moore, D. M., Krohn, M. A., and Hillier, S. L. (2004). Vaginal yeast colonization in nonpregnant women: a longitudinal study. *Obstet. Gynecol.* 104, 926–930. doi: 10.1097/01.AOG.0000140687.51048.73
- Bensen, E. S., Martin, S. J., Li, M., Berman, J., and Davis, D. A. (2004). Transcriptional profiling in *Candida albicans* reveals new adaptive responses to extracellular pH and functions for Rim101p. *Mol. Microbiol.* 54, 1335–1351. doi: 10.1111/j.1365-2958.2004.04350.x
- Botcher, B., Pollath, C., Staib, P., Hube, B., and Brunke, S. (2016). *Candida* species rewired hyphae developmental programs for chlamydospore formation. *Front. Microbiol.* 7:1697. doi: 10.3389/fmicb.2016.01697

- Brand, A., Lee, K., Veses, V., and Gow, N. A. (2009). Calcium homeostasis is required for contact-dependent helical and sinusoidal tip growth in *Candida albicans* hyphae. *Mol. Microbiol.* 71, 1155–1164. doi: 10.1111/j.1365-2958.2008.06592.x
- Cambi, A., Netea, M. G., Mora-Montes, H. M., Gow, N. A., Hato, S. V., Lowman, D. W., et al. (2008). Dendritic cell interaction with *Candida albicans* critically depends on N-linked mannan. *J. Biol. Chem.* 283, 20590–20599. doi: 10.1074/jbc.M709334200
- Carlisle, P. L., and Kadosh, D. (2010). *Candida albicans* Ume6, a filament-specific transcriptional regulator, directs hyphal growth via a pathway involving Hgc1 cyclin-related protein. *Eukaryot. Cell* 9, 1320–1328. doi: 10.1128/EC.00046-10
- Chaffin, W. L., Lopez-Ribot, J. L., Casanova, M., Gozalbo, D., and Martinez, J. P. (1998). Cell wall and secreted proteins of *Candida albicans*: identification, function, and expression. *Microbiol. Mol. Biol. Rev.* 62, 130–180. doi: 10.1128/MMBR.62.1.130-180.1998
- Cheng, S. C., Scicluna, B. P., Arts, R. J., Gresnigt, M. S., Lachmandas, E., Giamarellos-Bourboulis, E. J., et al. (2016). Broad defects in the energy metabolism of leukocytes underlie immunoparalysis in sepsis. *Nat. Immunol.* 17, 406–413. doi: 10.1038/ni.3398
- Citiulo, F., Moran, G. P., Coleman, D. C., and Sullivan, D. J. (2009). Purification and germination of *Candida albicans* and *Candida dubliniensis* chlamydospores cultured in liquid media. *FEMS Yeast Res.* 9, 1051–1060. doi: 10.1111/j.1567-1364.2009.00533.x
- Cottier, F., Srinivasan, K. G., Yurieva, M., Liao, W., Poidinger, M., Zolezzi, F., et al. (2018). Advantages of meta-total RNA sequencing (MeTRS) over shotgun metagenomics and amplicon-based sequencing in the profiling of complex microbial communities. *NPJ Biofilms Microbiomes* 4:2. doi: 10.1038/s41522-017-0046-x
- Da Silva, C. A., Chalouni, C., Williams, A., Hartl, D., Lee, C. G., and Elias, J. A. (2009). Chitin is a size-dependent regulator of macrophage TNF and IL-10 production. *J. Immunol.* 182, 3573–3582. doi: 10.4049/jimmunol.0802113
- Dominguez-Andres, J., R., Arts, J. W., Ter Horst, R., Gresnigt, M. S., Smeekens, S. P., Ratter J. M., et al. (2017). Rewiring monocyte glucose metabolism via C-type lectin signaling protects against disseminated candidiasis. *PLoS Pathog.* 13:e1006632. doi: 10.1371/journal.ppat.1006632
- Duvenage, L., Walker, L. A., Bojarczuk, A., Johnston, S. A., MacCallum, D. M., Munro, C. A., et al. (2019). Inhibition of classical and alternative modes of respiration in *Candida albicans* leads to cell wall remodeling and increased macrophage recognition. *MBio* 10:e02535-18. doi: 10.1128/mBio.02535-18
- Ene, I. V., Adya, A. K., Wehmeier, S., Brand, A. C., MacCallum, D. M., Gow, N. A., et al. (2012). Host carbon sources modulate cell wall architecture, drug resistance and virulence in a fungal pathogen. *Cell Microbiol.* 14, 1319–1335. doi: 10.1111/j.1462-5822.2012.01813.x
- Ene, I. V., Cheng, S. C., Netea, M. G., and Brown, A. J. (2013). Growth of *Candida albicans* cells on the physiologically relevant carbon source lactate affects their recognition and phagocytosis by immune cells. *Infect. Immun.* 81, 238–248. doi: 10.1128/IAI.01092-12
- Esteban, A., Popp, M. W., Vyas, V. K., Strijbis, K., Ploegh, H. L., and Fink, G. R. (2011). Fungal recognition is mediated by the association of dectin-1 and galectin-3 in macrophages. *Proc. Natl. Acad. Sci. U.S.A.* 108, 14270–14275. doi: 10.1073/pnas.1111415108
- Gantner, B. N., Simmons, R. M., and Underhill, D. M. (2005). Dectin-1 mediates macrophage recognition of *Candida albicans* yeast but not filaments. *EMBO J.* 24, 1277–1286. doi: 10.1038/sj.emboj.7600594
- Garfoot, A. L., Dearing, K. L., VanSchoick, A. D., Wysocki, V. H., and Rappleye, C. A. (2017). Eng1 and Exg8 Are the Major beta-Glucanases secreted by the fungal pathogen *histoplasma capsulatum*. *J. Biol. Chem.* 292, 4801–4810. doi: 10.1074/jbc.M116.762104
- Gazi, U., Rosas, M., Singh, S., Heinsbroek, S., Haq, I., Johnson, S., et al. (2011). Fungal recognition enhances mannose receptor shedding through dectin-1 engagement. *J. Biol. Chem.* 286, 7822–7829. doi: 10.1074/jbc.M110.185025
- Grahl, N., Demers, E. G., Lindsay, A. K., Harty, C. E., Willger, S. D., Piispanen, A. E., et al. (2015). Mitochondrial activity and Cyr1 are key regulators of Ras1 activation of *C. albicans* virulence pathways. *PLoS Pathog.* 11:e1005133. doi: 10.1371/journal.ppat.1005133
- Grahl, N., Shepardson, K. M., Chung, D., and Cramer, R. A. (2012). Hypoxia and fungal pathogenesis: to air or not to air? *Eukaryot. Cell* 11, 560–570. doi: 10.1128/EC.00031-12
- Graus, M. S., Wester, M. J., Lowman, D. W., Williams, D. L., Kruppa, M. D., Martinez, C. M., et al. (2018). Mannan molecular substructures control nanoscale glucan exposure in *Candida*. *Cell Rep.* 24, 2432–2442.e5. doi: 10.1016/j.celrep.2018.07.088
- Guedouari, H., Gergondey, R., Bourdais, A., Vanparis, O., Bulteau, A. L., Camadro, J. M., et al. (2014). Changes in glutathione-dependent redox status and mitochondrial energetic strategies are part of the adaptive response during the filamentation process in *Candida albicans*. *Biochim. Biophys. Acta* 1842, 1855–1869. doi: 10.1016/j.bbadis.2014.07.006
- Hayes, A. B. (1966). Chlamydospore production in *Candida albicans*. *Mycopathol. Mycol. Appl.* 29, 87–96. doi: 10.1007/BF02055062
- Hopke, A., A., Brown, J. P., Hall, R. A., and Wheeler, R. T. (2018). Dynamic fungal cell wall architecture in stress adaptation and immune evasion. *Trends Microbiol.* 26, 284–295. doi: 10.1016/j.tim.2018.01.007
- Huang, G., Srikantha, T., Sahni, N., Yi, S., and Soll, D. R. (2009). CO(2) regulates white-to-opaque switching in *Candida albicans*. *Curr. Biol.* 19, 330–334. doi: 10.1016/j.cub.2009.01.018
- Huang, G., Yi, S., Sahni, N., Daniels, K. J., Srikantha, T., and Soll, D. R. (2010). N-acetylglucosamine induces white to opaque switching, a mating prerequisite in *Candida albicans*. *PLoS Pathog.* 6:e1000806. doi: 10.1371/journal.ppat.100806
- Ifrim, D. C., Quintin, J., Courjol, F., Verschuere, I., van Krieken, J. H., Koentgen, F., et al. (2016). The role of dectin-2 for host defense against disseminated candidiasis. *J. Interferon Cytokine Res.* 36, 267–276. doi: 10.1089/jir.2015.0040
- Jeanmonod, R., and Jeanmonod, D. (2019). *Vaginal Candidiasis (Vulvovaginal Candidiasis)*. Treasure Island, FL: StatPearls.
- Kasper, L., Konig, A., Koenig, P. A., Gresnigt, M. S., Westman, J., Drummond, R. A., et al. (2018). The fungal peptide toxin Candidalysin activates the NLRP3 inflammasome and causes cytolysis in mononuclear phagocytes. *Nat. Commun.* 9:4260. doi: 10.1038/s41467-018-06607-1
- Klis, F. M., de Koster, C. G., and Brul, S. (2014). Cell wall-related bionumbers and bioestimates of *Saccharomyces cerevisiae* and *Candida albicans*. *Eukaryot. Cell* 13, 2–9. doi: 10.1128/EC.00250-13
- Knafler, H. C., R., Smaczynska-de II, Walker, L. A., Lee, K. K., N., Gow, A. R., Ayscough K. R. (2019). AP-2-dependent endocytic recycling of the chitin synthase Chs3 regulates polarized growth in *Candida albicans*. *MBio* 10:e02421-18. doi: 10.1128/mBio.02421-18
- Koch, B., Barugahare, A. A., Lo, T. L., Huang, C., Schittenhelm, R. B., Powell, D. R., et al. (2018). A metabolic checkpoint for the yeast-to-hyphae developmental switch regulated by endogenous nitric oxide signaling. *Cell Rep.* 25, 2244–2258.e7. doi: 10.1016/j.celrep.2018.10.080
- Liang, S. H., Anderson, M. Z., Hirakawa, M. P., Wang, J. M., Frazer, C., Alaalm, L. M., et al. (2019). Hemizygoty enables a mutational transition governing fungal virulence and commensalism. *Cell Host Microbe* 25, 418–431.e6. doi: 10.1016/j.chom.2019.01.005
- Lo, H. J., Kohler, J. R., DiDomenico, B., Loebenberg, D., Cacciapuoti, A., and Fink, G. R. (1997). Nonfilamentous *C. albicans* mutants are avirulent. *Cell* 90, 939–949. doi: 10.1016/S0092-8674(00)80358-X
- Lockhart, S. R., Pujol, C., Daniels, K. J., Miller, M. G., Johnson, A. D., Pfaller, M. A., et al. (2002). In *Candida albicans*, white-opaque switchers are homozygous for mating type. *Genetics* 162, 737–745.
- Lockhart, S. R., Zhao, R., Daniels, K. J., and Soll, D. R. (2003). Alpha-pheromone-induced “shmooin” and gene regulation require white-opaque switching during *Candida albicans* mating. *Eukaryot. Cell* 2, 847–855. doi: 10.1128/EC.2.5.847-855.2003
- Lopes, J. P., Stylianou, M., Backman, E., Holmberg, S., Jass, J., Claesson, R., et al. (2018). Evasion of immune surveillance in low oxygen environments enhances *Candida albicans* virulence. *MBio* 9:e02120-18. doi: 10.1128/mBio.02120-18
- Marakalala, M. J., Vautier, S., Potrykus, J., Walker, L. A., Shepardson, K. M., Hopke, A., et al. (2013). Differential adaptation of *Candida albicans* in vivo modulates immune recognition by dectin-1. *PLoS Pathog.* 9:e1003315. doi: 10.1371/journal.ppat.1003315
- Mathe, L., and Van Dijck, P. (2013). Recent insights into *Candida albicans* biofilm resistance mechanisms. *Curr. Genet.* 59, 251–264. doi: 10.1007/s00294-013-0400-3

- Mayer, F. L., Wilson, D., and Hube, B. (2013). *Candida albicans* pathogenicity mechanisms. *Virulence* 4, 119–128. doi: 10.4161/viru.22913
- McCall, A. D., Kumar, R., and Edgerton, M. (2018). *Candida albicans* Sfl1/Sfl2 regulatory network drives the formation of pathogenic microcolonies. *PLoS Pathog.* 14:e1007316. doi: 10.1371/journal.ppat.1007316
- Miller, M. G., and Johnson, A. D. (2002). White-opaque switching in *Candida albicans* is controlled by mating-type locus homeodomain proteins and allows efficient mating. *Cell* 110, 293–302. doi: 10.1016/S0092-8674(02)00837-1
- Mora-Montes, H. M., Bates, S., Netea, M. G., Castillo, L., Brand, A., Buurman, E. T., et al. (2010). A multifunctional mannosyltransferase family in *Candida albicans* determines cell wall mannan structure and host-fungus interactions. *J. Biol. Chem.* 285, 12087–12095. doi: 10.1074/jbc.M109.081513
- Mora-Montes, H. M., Netea, M. G., Ferwerda, G., Lenardon, M. D., Brown, G. D., Mistry, A. R., et al. (2011). Recognition and blocking of innate immunity cells by *Candida albicans* chitin. *Infect. Immun.* 79, 1961–1970. doi: 10.1128/IAI.01282-10
- Moran, C., Grussemeyer, C. A., Spalding, J. R., Benjamin, D. K. Jr., and Reed, S. D. (2010). Comparison of costs, length of stay, and mortality associated with *Candida glabrata* and *Candida albicans* bloodstream infections. *Am. J. Infect. Control* 38, 78–80. doi: 10.1016/j.ajic.2009.06.014
- Moyes, D. L., Wilson, D., Richardson, J. P., Mogavero, S., Tang, S. X., Wernecke, J., et al. (2016). Candidalysin is a fungal peptide toxin critical for mucosal infection. *Nature* 532, 64–68. doi: 10.1038/nature17625
- Navarro-Arias, M. J., Hernandez-Chavez, M. J., Garcia-Carnero, L. C., Amezcua-Hernandez, D. G., Lozoya-Perez, N. E., Estrada-Mata, E., et al. (2019). Differential recognition of *Candida tropicalis*, *Candida guilliermondii*, *Candida krusei*, and *Candida auris* by human innate immune cells. *Infect. Drug Resist.* 12, 783–794. doi: 10.2147/IDRS.197531
- Netea, M. G., Gow, N. A., Munro, C. A., Bates, S., Collins, C., Ferwerda, G., et al. (2006). Immune sensing of *Candida albicans* requires cooperative recognition of mannans and glucans by lectin and Toll-like receptors. *J. Clin. Invest.* 116, 1642–1650. doi: 10.1172/JCI27114
- Noble, S. M., French, S., Kohn, L. A., Chen, V., and Johnson, A. D. (2010). Systematic screens of a *Candida albicans* homozygous deletion library decouple morphogenetic switching and pathogenicity. *Nat. Genet.* 42, 590–598. doi: 10.1038/ng.605
- Noble, S. M., Gianetti, B. A., and Witchley, J. N. (2017). *Candida albicans* cell-type switching and functional plasticity in the mammalian host. *Nat. Rev. Microbiol.* 15, 96–108. doi: 10.1038/nrmicro.2016.157
- Palige, K., Linde, J., Martin, R., Bottcher, B., Citiulo, F., Sullivan, D. J., et al. (2013). Global transcriptome sequencing identifies chlamyospore specific markers in *Candida albicans* and *Candida dubliniensis*. *PLoS ONE* 8:e61940. doi: 10.1371/journal.pone.0061940
- Pande, K., Chen, C., and Noble, S. M. (2013). Passage through the mammalian gut triggers a phenotypic switch that promotes *Candida albicans* commensalism. *Nat. Genet.* 45, 1088–1091. doi: 10.1038/ng.2710
- Pfaller, M. A., and Diekema, D. J. (2007). Epidemiology of invasive candidiasis: a persistent public health problem. *Clin. Microbiol. Rev.* 20, 133–163. doi: 10.1128/CMR.00029-06
- Pierce, J. V., and Kumamoto, C. A. (2012). Variation in *Candida albicans* EFG1 expression enables host-dependent changes in colonizing fungal populations. *MBio* 3:e00117-12. doi: 10.1128/mBio.00117-12
- Pradhan, A., Avelar, G. M., Bain, J. M., Childers, D. S., Larcombe, D. E., Netea, M. G., et al. (2018). Hypoxia promotes immune evasion by triggering beta-glucan masking on the *Candida albicans* cell surface via mitochondrial and cAMP-protein kinase A signaling. *MBio* 9:e01318-18. doi: 10.1128/mBio.01318-18
- Romani, L. (2011). Immunity to fungal infections. *Nat. Rev. Immunol.* 11, 275–288. doi: 10.1038/nri2939
- Ruiz-Herrera, J., Elorza, M. V., Valentin, E., and Sentandreu, R. (2006). Molecular organization of the cell wall of *Candida albicans* and its relation to pathogenicity. *FEMS Yeast Res.* 6, 14–29. doi: 10.1111/j.1567-1364.2005.00017.x
- Saville, S. P., Lazzell, A. L., Monteagudo, C., and Lopez-Ribot, J. L. (2003). Engineered control of cell morphology *in vivo* reveals distinct roles for yeast and filamentous forms of *Candida albicans* during infection. *Eukaryot. Cell* 2, 1053–1060. doi: 10.1128/EC.2.5.1053-1060.2003
- Selvaggini, S., Munro, C. A., Paschoud, S., Sanglard, D., and Gow, N. A. (2004). Independent regulation of chitin synthase and chitinase activity in *Candida albicans* and *Saccharomyces cerevisiae*. *Microbiology* 150, 921–928. doi: 10.1099/mic.0.26661-0
- Sherrington, S. L., Sorsby, E., Mahtey, N., Kumwenda, P., Lenardon, M. D., Brown, I., et al. (2017). Adaptation of *Candida albicans* to environmental pH induces cell wall remodelling and enhances innate immune recognition. *PLoS Pathog.* 13:e1006403. doi: 10.1371/journal.ppat.1006403
- Sherwood-Higham, J., Zhu, W. Y., Devine, C. A., Gooday, G. W., Gow, N. A., and Gregory, D. W. (1994). Helical growth of hyphae of *Candida albicans*. *J. Med. Vet. Mycol.* 32, 437–445. doi: 10.1080/02681219480000591
- Slutsky, B., Staebell, M., Anderson, J., Risen, L., Pfaller, M., and Soll, D. R. (1987). “White-opaque transition”: a second high-frequency switching system in *Candida albicans*. *J. Bacteriol.* 169, 189–197. doi: 10.1128/JB.169.1.189-197.1987
- Smeekens, S. P., Gresnigt, M. S., Becker, K. L., Cheng, S. C., Netea, S. A., Jacobs, L., et al. (2015). An anti-inflammatory property of *Candida albicans* beta-glucan: induction of high levels of interleukin-1 receptor antagonist via a Dectin-1/CR3 independent mechanism. *Cytokine* 71, 215–222. doi: 10.1016/j.cyt.2014.10.013
- Sudbery, P., Gow, N., and Berman, J. (2004). The distinct morphogenic states of *Candida albicans*. *Trends Microbiol.* 12, 317–324. doi: 10.1016/j.tim.2004.05.008
- Sun, Y., Cao, C., Jia, W., Tao, L., Guan, G., and Huang, G. (2015). pH regulates white-opaque switching and sexual mating in *Candida albicans*. *Eukaryot. Cell* 14, 1127–1134. doi: 10.1128/EC.00123-15
- Synnott, J. M., Guida, A., Mulhern-Haughey, S., Higgins, D. G., and Butler, G. (2010). Regulation of the hypoxic response in *Candida albicans*. *Eukaryot. Cell* 9, 1734–1746. doi: 10.1128/EC.00159-10
- Tao, L., Du, H., Guan, G., Dai, Y., Nobile, C. J., Liang, W., et al. (2014). Discovery of a “white-gray-opaque” tristable phenotypic switching system in *Candida albicans*: roles of non-genetic diversity in host adaptation. *PLoS Biol.* 12:e1001830. doi: 10.1371/journal.pbio.1001830
- Taylor, P. R., Tsoni, S. V., Willment, J. A., Dennehy, K. M., Rosas, M., Findon, H., et al. (2007). Dectin-1 is required for beta-glucan recognition and control of fungal infection. *Nat. Immunol.* 8, 31–38. doi: 10.1038/ni1408
- Tucey, T. M., Verma, J., Harrison, P. F., Snelgrove, S. L., Lo, T. L., Scherer, A. K., et al. (2018). Glucose homeostasis is important for immune cell viability during *Candida* challenge and host survival of systemic fungal infection. *Cell Metab.* 27, 988–1006.e7. doi: 10.1016/j.cmet.2018.03.019
- Tuch, B. B., Mitrovich, Q. M., Homann, O. R., Hernday, A. D., Monighetti, C. K., De La Vega F. M., et al. (2010). The transcriptomes of two heritable cell types illuminate the circuit governing their differentiation. *PLoS Genet.* 6:e1001070. doi: 10.1371/journal.pgen.1001070
- Ueno, K., Okawara, A., Yamagoe, S., Naka, T., Umeyama, T., Utena-Abe, Y., et al. (2013). The mannan of *Candida albicans* lacking beta-1,2-linked oligomannosides increases the production of inflammatory cytokines by dendritic cells. *Med. Mycol.* 51, 385–395. doi: 10.3109/13693786.2012.733892
- Uwamahoro, N., Verma-Gaur, J., Shen, H. H., Qu, Y., Lewis, R., Lu, J., et al. (2014). The pathogen *Candida albicans* hijacks pyroptosis for escape from macrophages. *MBio* 5:e00003-e00014. doi: 10.1128/mBio.0003-14
- Wagener, J., R., Malireddi, K. S., Lenardon, M. D., Koberle, M., Vautier, S., MacCallum D. M., et al. (2014). Fungal Chitin Dampens Inflammation through IL-10 induction mediated by NOD2 and TLR9 activation. *PLoS Pathog.* 10:e1004050. doi: 10.1371/journal.ppat.1004050
- Wellington, M., Koselny, K., Sutterwala, F. S., and Krysan, D. J. (2014). *Candida albicans* triggers NLRP3-mediated pyroptosis in macrophages. *Eukaryot. Cell* 13, 329–340. doi: 10.1128/EC.00336-13
- Witchley, J. N., Penumetcha, P., Abon, N. V., Woolford, C. A., Mitchell, A. P., and Noble, S. M. (2019). *Candida albicans* morphogenesis programs control the balance between gut commensalism and invasive infection. *Cell Host Microbe* 25, 432–443.e6. doi: 10.1016/j.chom.2019.02.008
- Wu, S. Y., Huang, J. H., Chen, W. Y., Chan, Y. C., Lin, C. H., Chen, Y. C., et al. (2017). Cell intrinsic galectin-3 attenuates neutrophil ROS-dependent killing of *Candida* by modulating CR3 downstream syk activation. *Front. Immunol.* 8:48. doi: 10.3389/fimmu.2017.00048
- Yang, S. L., Zeng, G., Chan, F. Y., Wang, Y. M., Yang, D., and Wang, Y. (2018). Sac7 and Rho1 regulate the white-to-opaque switching in *Candida albicans*. *Sci. Rep.* 8:875. doi: 10.1038/s41598-018-19246-9

- Yano, J., Peters, B. M., Noverr, M. C., and Fidel, P. L. Jr. (2018). Novel mechanism behind the immunopathogenesis of vulvovaginal candidiasis: “neutrophil anergy”. *Infect. Immun.* 86:e00684-17. doi: 10.1128/IAI.00684-17
- Zhang, S. Q., Zou, Z., Shen, H., Shen, S. S., Miao, Q., Huang, X., et al. (2016). Mnn10 maintains pathogenicity in *Candida albicans* by extending alpha-1,6-Mannose backbone to evade host dectin-1 mediated antifungal immunity. *PLoS Pathog.* 12:e1005617. doi: 10.1371/journal.ppat.1005617
- Zhu, L. L., Zhao, X. Q., Jiang, C., You, Y., Chen, X. P., Jiang, Y. Y., et al. (2013). C-type lectin receptors Dectin-3 and Dectin-2 form a heterodimeric pattern-recognition receptor for host defense against fungal infection. *Immunity* 39, 324–334. doi: 10.1016/j.immuni.2013.05.017

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Cottier and Hall. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.