



Kent Academic Repository

Kieninger, Christoph, Wurst, Klaus, Podewitz, Maren, Stanley, Maria, Deery, Evelyne, Lawrence, Andrew D., Liedl, Klaus R., Warren, Martin J. and Kräutler, Bernhard (2020) *Replacement of the Cobalt Center of Vitamin B 12 by Nickel: Nibalamin and Nibyrlic Acid Prepared from MetalFree B 12 Ligands Hydrogenobalam and Hydrogenobyric Acid*. *Angewandte Chemie International Edition* . ISSN 1433-7851.

Downloaded from

<https://kar.kent.ac.uk/82762/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1002/anie.202008407>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal* , Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Macrocycles

How to cite:

International Edition: doi.org/10.1002/anie.202008407

German Edition: doi.org/10.1002/ange.202008407

Replacement of the Cobalt Center of Vitamin B₁₂ by Nickel: Nibalamin and Nibyric Acid Prepared from Metal-Free B₁₂ Ligands Hydrogenobalamin and Hydrogenobyric Acid

Christoph Kieninger, Klaus Wurst, Maren Podewitz, Maria Stanley, Evelyne Deery, Andrew D. Lawrence, Klaus R. Liedl, Martin J. Warren,* and Bernhard Kräutler*

Dedicated to Professor Albert Eschenmoser on the occasion of his 95th birthday

Abstract: The (formal) replacement of Co in cobalamin (**Cbl**) by Ni^{II} generates nibalamin (**Nibl**), a new transition-metal analogue of vitamin B₁₂. Described here is **Nibl**, synthesized by incorporation of a Ni^{II} ion into the metal-free B₁₂ ligand hydrogenobalamin (**Hbl**), itself prepared from hydrogenobyric acid (**Hby**). The related Ni^{II} corrin nibyric acid (**Niby**) was similarly synthesized from **Hby**, the metal-free cobyric acid ligand. The solution structures of **Hbl**, and **Niby** and **Nibl**, were characterized by spectroscopic studies. **Hbl** features two inner protons bound at N2 and N4 of the corrin ligand, as discovered in **Hby**. X-ray analysis of **Niby** shows the structural adaptation of the corrin ligand to Ni^{II} ions and the coordination behavior of Ni^{II}. The diamagnetic **Niby** and **Nibl**, and corresponding isoelectronic Co^I corrins, were deduced to be isostructural. **Nibl** is a structural mimic of four-coordinate base-off **Cbls**, as verified by its ability to act as a strong inhibitor of bacterial adenosyltransferase.

Introduction

Biologically active vitamin B₁₂ derivatives exclusively utilize cobalt as their specific transition metal center, which is bound and activated exquisitely by a helical corrin macrocycle.^[1] The metal-free corrin ligand of vitamin B₁₂, hydrogenobyric acid (**Hby**), has recently been made available as a consequence of engineered B₁₂ biosynthesis in *E. coli*.^[2,3] The availability of **Hby** has provided an unparalleled opportunity for the effective synthesis of metal-free and transition metal analogues of the natural cobalt-corrinoids a previously intractable challenge in bioinorganic and B₁₂ chemistry.^[4] We have recently used **Hby** for the synthesis of the corresponding zinc-corrin zincobyric acid (**Znby**) and the

Zn analogue of vitamin B₁₂ zincobalamin (**Znbl**), of interest as luminescent structural B₁₂ mimics.^[5]

Herein, we report on the first nickel-complexes of natural corrin ligands, including nibalamin (**Nibl**). We also describe the syntheses of crystalline nibyric acid (**Niby**), the novel Ni^{II} complex of **Hby**,^[3] and hydrogenobalamin (**Hbl**), the metal-free complete B₁₂ ligand (see Scheme 1 and Scheme 2). Koppenhagen and co-workers, back in the 1970's, reported the isolation of **Hbl** from a *Chromatium* strain supplemented with 5,6-dimethylbenz-imidazole (DMB). They were able to characterize **Hbl** by UV/Vis-spectroscopy and demonstrated that it could be converted into vitamin B₁₂ by insertion of cobalt,^[4e,7] and later reported its mass spectrum.^[8]

A Ni^{II}-corrin, the **NiCor** (see Scheme 1), was prepared in the Eschenmoser labs as the first synthetic corrin, making use of the Ni^{II} ion as a "template" for the assembly of the corrin macro-ring.^[6] **NiCor** also became the object of the first X-ray crystallographic investigation of the structure of a non-cobalt corrin.^[9] Four coordinate Ni^{II} complexes prefer to adopt a planar geometry and therefore are more structurally related to the corresponding Co^I complexes.^[10] Indeed, recently, there has been a resurgence in the quest for close Ni analogues of the B₁₂ cofactors.^[4c,f] The planar ligand set of **Nibl** potentially represents a structural B₁₂ mimic that is inert to the organometallic transformations typical of B₁₂-dependent enzymes, as suggested by its expected coordination chemistry and structural properties. Specific interest in **Nibl**, the Ni^{II}-analogue of vitamin B₁₂ and of other cobalamins (**Cbls**) (see Scheme 1), is thus a consequence not only of its chemistry, but also of its possible use as a molecular probe in B₁₂ biology and biomedicine, helpful for the investigation of cobalamin-dependent processes and their physiological effects.^[11]

[*] Dr. C. Kieninger, Prof. Dr. B. Kräutler
Institute of Organic Chemistry, University of Innsbruck
6020 Innsbruck (Austria)
E-mail: bernhard.kraeutler@uibk.ac.at

Dr. C. Kieninger, Dr. M. Podewitz, Prof. Dr. K. R. Liedl,
Prof. Dr. B. Kräutler
Center for Molecular Biosciences (CMBI), University of Innsbruck
6020 Innsbruck (Austria)

Dr. K. Wurst, Dr. M. Podewitz, Prof. Dr. K. R. Liedl
Institute of General, Inorganic and Theoretical Chemistry, University
of Innsbruck, 6020 Innsbruck (Austria)


Dr. M. Stanley, Dr. E. Deery, Dr. A. D. Lawrence, Prof. Dr. M. J. Warren
School of Biosciences, University of Kent
Canterbury, CT2 7NJ (UK)

E-mail: m.j.warren@kent.ac.uk


Prof. Dr. M. J. Warren
Quadram Institute Bioscience, Norwich Science Park
Norwich NR4 7UQ (UK)

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202008407>.

© 2020 The Authors. Published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



Scheme 1. Structural formulae of cobalt, zinc, nickel and metal-free corrinoids. Left: cobalamins with “base-on” structures: vitamin B₁₂ (R = CN, **CNCbl**), coenzyme B₁₂ (R = 5'-deoxyadenosyl, **AdoCbl**), methylcobalamin (R = CH₃, **MeCbl**), cob(II)alamin (R = e⁻, **Cbl^{II}**). Center, top: the Ni^{II}-corrin nibrates (**Niby**), hydrogenobyric acid (**Hby**), Co^{II}-cobyric acid (**Cby^{II}**) and zincobyric acid (**Znby**), where Co^{II} and Zn^{II} carry an unspecified axial ligand (e.g., solvent molecule). Center, bottom: Eschenmoser's synthetic racemic Ni^{II}-corrin **NiCor**.^[6] Right: nivalamin (**Nibl**), four-coordinate “base off” cob(II)alamin (**4c-Cbl^{II}**) and cob(II)alamin (**Cbl^{II}**).



Scheme 2. Preparation of the Ni^{II}-corrins **Niby** and **Nibl** from **Hby**.
 i) 0.5 M Ni(OAc)₂ pH 6, 1 h, 90°C, Ar. ii) 3 equiv B₁₂ nucleotide moiety,^[1a,12] HOBT, EDC·HCl, H₂O, 0°C, 4 d. iii) 0.5 M Ni(OAc)₂ pH 6, 1 h, 90°C, Ar (see the SI for details).

Results and Discussion

Nibrinic acid (**Niby**) was prepared by dissolving 1.40 mg (1.6 μmol) of crystalline hydrogenobyric acid (**Hby**)^[3] in 3.5 mL of deoxygenated 0.5 M aqueous Ni^{II}acetate, pH 6, with stirring at 90°C for 75 min. Separation on a short reverse phase column, evaporation and crystallization from aqueous acetonitrile yielded 0.90 mg (0.97 μmol, 61 %) of **Niby**, which was isolated as yellow crystals (see Scheme 2, Exptl. Part and

Supporting Information (SI). The UV/Vis absorption spectrum of an aqueous solution of **Niby** displayed bands at 464 nm (shoulder), 448 nm and 334 nm (Figure 1), and exhibited similar gross features to those observed in an absorption spectrum of the Ni-corrin **NiCor** (but with a slightly red-shifted maxima).^[6a,c] The solution structure of the diamagnetic Ni^{II}-corrin **Niby** (molecular formula C₄₅H₆₄N₁₀O₈Ni, for HR mass spectra see SI, Figure S3) was analyzed by NMR spectroscopy, providing assignment of all 52 non-exchangeable H atoms and 44 C atoms (see SI, Figure S4 and Table S1). A 500 MHz ¹H NMR spectrum of **Niby** in D₂O displayed five high field singlets for the six methyl groups, a singlet of HC10 at 6.30 ppm, as well as several signals at intermediate field for HC19, HC3, HC8 and HC13 (see Figure 2). The data from homonuclear and heteronuclear correlations confirmed the stereostructure of **Niby** (see SI, Figure S5).




Figure 1. Absorption spectra of aqueous solutions of metal-free B₁₂ ligands **Hby** and **Hbl** and of their Ni^{II}-complexes **Niby** and **Nibl** at 298 K. Top: UV/Vis-absorption spectra of **Hby** (*c* = 31.5 μM, pH 5, black trace) and **Niby** (*c* = 34.5 μM, unbuffered, red trace). Bottom: UV/Vis-absorption of **Hbl** (pH 5, black trace) and of **Nibl** (unbuffered, red trace).




Figure 2. 500 MHz ¹H NMR spectrum of **Niby** in D₂O (*c* = 1.9 mM, 298 K); the water signal after presaturation is marked by an X.

The complete metal-free ligand of the cobalamins, hydrogencobalamin (**Hbl**), was assembled by attaching the B₁₂ nucleotide moiety^[1a,12] to the propionate moiety of **Hby** at 0°C through application of the carbodiimide method (Scheme 2).^[4d,13] In brief, an aqueous solution of 9.12 mg (10.4 μmol) of **Hby** and of 14.71 mg (33.4 μmol) of the B₁₂-nucleotide was treated with 9.4 moleq of HOBt and degassed. To the frozen reaction mixture 4.4 moleq of EDC·HCl were added under Ar. Upon subsequent warm-up of the reaction mixture to 0°C, 16 moleq EDC·HCl were added and stirring was continued for 4 d (see SI). Work-up, using RP18-chromatographic purification, precipitation with MeCN and drying, furnished 11.3 mg (8.89 μmol, 85% yield) of **Hbl** as an orange powder. An aqueous solution of **Hbl** at pH 5 exhibited UV/Vis^[4c] and CD spectral features (SI, Figure S8 and S9) similar to those of **Hby**.^[3] The UV/Vis absorption maximum at 525 nm of the α-band of **Hbl** and the fluorescence emission maximum at 554 nm (SI, Figure S10) located the first excited singlet state of **Hbl** at *E*^S near 221 kJ mol⁻¹, marginally lower than for **Hby**.^[3] The structure of **Hbl** (molecular formula C₆₂H₉₀N₁₃O₁₄P, see HR-MS data in SI, Figure S11) in H₂O was characterized by NMR spectroscopy (600 MHz ¹H NMR spectrum in SI, Figure S12), providing assignment of 89 H atoms and of all 62 C atoms (see SI, Table S2). The two “inner” H atoms gave rise to singlets at δ = 12.32 and δ = 12.57 ppm, which were assigned to H(N4) and to H(N2), respectively, indicating a minor up-field shift of both of them when compared to **Hby**.^[3] The methyl group singlet of H₃C1 at δ = 0.81 ppm occurred at 0.47 ppm to higher field, compared to **Hby**, suggesting a temporary residence of the heteroaromatic DMB unit of **Hbl** near to its corrin moiety, a conclusion that was further supported by weak inter-residual correlations in the ¹H, ¹HROESY spectra (see SI, Figure S13). However, the signals of the DMB moiety (HN2 at δ = 8.35 ppm, HN4 at δ = 7.31 ppm, HCN7 at δ = 7.30 ppm) were found at similar chemical shift values to those of the free B₁₂ nucleotide,^[12] effectively incompatible with a time-averaged positioning of the DMB part close to the corrin chromophore, as found for zincobalamin (**Znbl**)^[5] and for typical “base-on” Co^{III}CbIs.^[14]

The Ni^{II}-corrin nivalamin (**Nibl**) was prepared by heating a deoxygenated aqueous solution of **Hbl** and Ni(OAc)₂ for 1 h at 90°C (Scheme 2), furnishing **Nibl** in 77% yield as a yellow powder. An unbuffered aqueous solution of **Nibl** exhibited a UV/Vis spectrum that is incompatible with coordination by the DMB base and nearly indistinguishable (at > 300 nm) from the spectrum of **Niby**, and similar to the spectrum of the Ni^{II}-corrin **NiCor**^[6a,c] (see Figure 1). However, the absorption maxima of **Nibl** occurred at characteristically longer wavelengths when compared to the spectrum of the recently described vitamin B₁₂-derived 5,6-dihydroxy-5,6-dihydro-nivalamin, which features an interrupted corrin π-system.^[4f] Lower pH values affected only the short wavelength part of the **Nibl** UV/Vis-absorption spectrum, which was altered by DMB-protonation, consistent with a p*K*_a = 4.35 ± 0.06 for protonated **Nibl-H**⁺ (see SI, Figure S20).

The structure of **Nibl** (molecular formula C₆₂H₈₈N₁₃O₁₄PNi, SI, Figure S17) was characterized in aqueous solution by heteronuclear NMR spectroscopy (see 500 MHz ¹H NMR spectrum in Figure 3), providing assign-




Figure 3. 500 MHz ¹H NMR spectrum of **Nibl** in D₂O (*c* = 1.4 mM, 298 K); residual water signal after presaturation marked by an X.

ment of all 73 non-exchangeable H atoms and of all 62 C atoms (SI, Table S3). The positions of the singlets of H₃C1A (δ = 1.10 ppm), of HCN2 (δ = 8.51 ppm), HN4 (δ = 7.36 ppm) and HCN7 (δ = 7.39 ppm) of the DMB moiety all indicate a base-off form with a four-coordinate Ni^{II} center. Hence, the UV/Vis and NMR spectral features of **Nibl** characterize it as an isoelectronic and, roughly, isostructural analogue of the diamagnetic cob(I)alamin (**Cbl**^I), which is considered to feature a “base-off” structure with a four-coordinated Co^I center.^[15,16]

The nickel corrin **Niby** was crystallized from aqueous acetonitrile, furnishing yellow single crystals (*P*₂¹₂¹) suitable for X-ray analysis (see Figure 4). The incorporation of a Ni^{II} ion into the corrin macrocycle of the metal-free **Hby** increased the effective symmetry of the corrin ligand as revealed by a comparison of the crystal structures of **Niby** and




Figure 4. Crystal structure of **Niby**. Left. Crystallographic model of **Niby** in two projections; Right. Graphs representing the corrin core with display of N-Ni bond lengths (top) and the coordination geometry around Ni^{II} center, highlighting the arrangement of the four inner corrin N atoms in a flattened tetrahedron around the Ni^{II} center (middle and bottom).

of **Hby** (see SI for details). Coordination of the Ni^{II}-ion largely equalizes the lengths of the two diagonals, where the N2-N4 diagonal of **Niby** exceeds its N1-N3 counterpart by only $\Delta d = 0.047 \text{ \AA}$, far less than in **Hby** ($\Delta d = 0.297 \text{ \AA}$)^[3] or in **Znby** ($\Delta d = 0.197 \text{ \AA}$)^[5]. The somewhat longer N2-N4 diagonals in the metal corrins **Niby** and **Znby** appear to reflect the preferred mode of the conformational adaptation of the coordination hole of the flexible, unsymmetrical corrin ligand to bound metal ions. The radial size of the coordination hole also shrank upon Ni^{II} coordination as the average length of the N1-N3 and N2-N4 diagonals of **Hby** ($d = 3.82 \text{ \AA}$) was reduced to $d = 3.71 \text{ \AA}$ in the complex **Niby**. Hence, the coordination of the Ni^{II} ion in **Niby** contracts the corrin ligand and makes it more symmetrical. This latter effect is also expressed by the regularly alternating bond lengths of the corrin π -system in **Niby**, observations that are compatible with the model Ni-corrin **NiCor**.^[9]

The four-coordinate Ni^{II} ion sits very close to the plane of the four inner corrin N atoms, comparable to the situation in the Ni^{II}-corrin **NiCor**,^[9] and in typical Co^{III}-corrins,^[16a] but contrasting somewhat with the out of plane distance of 0.048 \AA of the five-coordinate Co^{II} center of heptamethylcob(II)yrinate perchlorate (**Cbin^{II}**)^[17] (SI, Table S5). As expected,^[9,18] the metal-N bonds in **Niby** (average Ni-N bond length = 1.86 \AA) are shorter than those found in the Co^{II} analogue **Cbin^{II}** and in the Co^{III}-corrin coenzyme B₁₂ (**AdoCbl**), where average (Co^{II}-N) and (Co^{III}-N) bond lengths of 1.89 \AA ^[19] and of 1.90 \AA ,^[20] respectively, were observed.

The coordination of the Ni^{II} ion barely affects the conformational properties of the metal-free corrin ligand (Figures 5,6). Only a slight reduction of the helicity, h (Figure 7), of the inner corrin N atoms from $h = 12.9^\circ$ in **Hby** to $h = 10.1^\circ$ is seen in **Niby**. Indeed, the effect of the binding of the Ni^{II} ion on the corrin helicity is comparable to the situation in the enzyme-bound four-coordinate cob(II)alamin (**4c-Cbl^{II}(ACA)**) of the adenosyltransferase ACA,^[21] for which $h = 8^\circ$.^[3] In contrast, in five-coordinate Co^{II}-corrins the corrin helicity is significantly smaller, for example, $h = 6.1^\circ$ in the Co^{II}-corrin **Cbin^{II}**, and in typical Co^{III}-




Figure 5. Structural adaptation of the helical corrin ligand to the coordinated metal ions. Comparison of the corrin helicity h in the structures of **Hby**, **NiCor**, **Niby**, of **4c-Cbl^{II}** in adenosyltransferase (ACA), of five-coordinate **Cbin^{II}** and **Cbl^{II}**, and of six-coordinate Co^{III}-corrins **MeCbl**, **CNCbl** and **AdoCbl** (see text for details).

corrins planarization of the corrin ligand is still more pronounced, leading, for example, to $h = 3.5^\circ$ in **AdoCbl**.^[3]

The observed lower drive of the four-coordinate d^8 Ni^{II} ion to planarize the corrin macrocycle is similarly reflected by its own coordination geometry, which deviates strongly in **Niby** from the coplanar arrangement of the coordinating ligand atoms in typical four-coordinate low-spin Ni^{II} complexes.^[9,10,18] In **Niby** a remarkably large interplanar angle ϕ (Figure 7) at Ni^{II} ($\phi = 11.1^\circ$) results from extensive directional coordinative adaptation of the Ni^{II}-center to the geometric requirements imposed by the helical corrin ligand (see Figure 6 and SI). ϕ is significantly larger in **Niby** than in Co^{III}-corrins, which exhibit ϕ 's around 5° or less,^[3] and is comparable to the situation in five-coordinate Co^{II}-corrins **Cbl^{II}** ($\phi = 12^\circ$) and **Cbin^{II}** ($\phi = 7.6^\circ$).




Figure 6. Adaptation of the coordination geometry around the corrin-bound metal ions to the helical corrin ligand. Comparison of the interplanar angles ϕ in the structures of **NiCor**, **Niby**, 5-coordinate **Cbin^{II}** and **Cbl^{II}**, and the six-coordinate Co^{III} corrins **MeCbl**, **CNCbl** and **AdoCbl** (see text for details).

The corrin helicity h and the inter-planar angle ϕ (Figure 7), were introduced recently as two complementary parameters characterizing inner conformational effects of the mutual structural adaptation of the corrin macrocycle and of the coordination geometry at the bound metal ion.^[3] The so called corrin fold of the helical corrin macrocycle,^[22] a classic parameter characterizing the nonplanar corrin ring in CbIs and in other “complete” cobamides (Figure 7), was not used in this current study. Conceived as a measure of the major conformational adaptation of the corrin ring to the cobalt coordination of the (bulky) DMB moiety in “base-on” CbIs, it runs roughly along the Co-C10 (east-west) axis.^[22] However, in four- and five-coordinate metal-corrins lacking the DMB unit, like **Cbin^{II}**, **Niby** and **Znby**, the calculated corrin-fold is dominated by the effects of the corrin helicity and the intersection between the two relevant planes adheres to a north-south direction (see SI Table S5 and Figure S22).

DFT calculations were carried out to test further the proposed close structural similarity between Ni^{II}-corrins and their analogues with four-coordinate cobalt centers (see Figure 8 and SI for further details). In order to minimize the relevance of peripheral H-bonds between the amide functions in the implicit solvent calculations, the five-coordinate lip-






Figure 7. Geometrical parameters describing conformational effects in corrins. Left: corrin helicity (h = the dihedral angle N1-N2-N3-N4 around a virtual bond between N2 and N3 of the corrin ligand), revealing the effect of bound metal-ions on the corrin helix (Figure 5).^[3] Center: interplanar angle (ϕ = angle between the two planes through the metal ion and corrin N's N2/N4 or N1/N4, respectively) characterizing the deviation of the coordination environment at the metal center due to the helical corrin ligand (Figure 6).^[3] Right: corrin fold (angle between the best two planes through the inner corrin atoms from N1 to C10 and from C10 to N4, respectively) describing the main conformational response of the corrin macrocycle to the binding of the DMB base in Cbls.^[22]




Figure 8. Calculated structures (left) of the four-coordinate metal-corrins **Nibin** (black), **Cob'in** (green) and **4c-Cob'in** (orange) and structural superpositions (right) of the corrin cores of the cobalt complexes and **Nibin**, minimized at the four coordinating corrin N atoms.

ophilic heptamethyl-cob(II)yrinate perchlorate (**Cbin^{II}**)^[17] was used as a starting model. Calculations of the structures of the related unknown corrins heptamethyl-nibyrate (**Nibin**), four-coordinate heptamethyl-cob(II)yrinate (**4c-Cob^{II}in**) and heptamethyl-cob(I)yrinate (**Cob^Iin**) were carried out. They indicate extensive structural similarities, but the equatorial metal–N bonds are shorter (by roughly 0.01–0.03 Å) in the Co^I- and Ni^{II}-corrins **Cob^Iin** and **Nibin** in

comparison to the four-coordinate Co^{II}-corrin **4c-Cob^{II}in** and 5-coordinate **Cbin^{II}**. The N1-N3 diagonal was calculated to be shorter than its N2-N4 by roughly 0.01–0.02 Å, which is also in good qualitative agreement with the crystallographic data for **Nibin** and **Cbin^{II}**.^[17] All 4-coordinate metal centers (Ni^{II}, Co^I and Co^{II}) were calculated to be located virtually in the best plane through the four corrin N-atoms. The latter is arranged in a helix with a calculated value of h slightly decreasing from **Nibin** (7.6°) to **Cob^Iin** to **4c-Cob^{II}in** (6.6°), and induced an interplanar angle ϕ that slightly decreased in the same order (from 8.4° to 7.5°). The calculations for five-coordinate **Cbin^{II}** also reproduced, qualitatively, the still smaller value for h (5.4°), a larger value for ϕ (11.3°) and a significant displacement of the Co^{II} center towards the axial β -ligand (calculated as 0.112 Å).

The deduced utility of **Nibl** as a specific new structural mimic of four-coordinate base-off **Cbls** was initially tested in binding studies of **Nibl** to an adenosyltransferase enzyme (ATP:Co^Irrinoid adenosyltransferase), which catalyzes the biosynthetic construction of **AdoCbl** by Co _{β} -adenosylation of bound **Cbl^I-**. Such bacterial^[23] and human adenosyltransferases,^[24] for example, BtuR,^[23a] CobA,^[23b] ACA,^[21] and CblB,^[24] have been shown to facilitate the adenosylation process by first inducing the corrin substrate to adopt a four-coordinate structure, thus raising the redox potential of the Co^{II}/Co^I couple by around 250 mV,^[23,24] thereby allowing the physiologically difficult reduction. We, thus, investigated the effect of **Nibl** on the adenosylation process by incubating the *Brucella melitensis* BtuR^[23a] and the structural Cbl mimic **Nibl** in the presence of an excess of **Cbl^I** (see SI, Figures S27–S29). As expected, **Nibl** itself was not a substrate for the enzyme and was not adenosylated. However, the presence of **Nibl** within the incubation did effectively inhibit the adenosyltransferase, reducing the activity of the enzyme by 38% at a concentration of 1 μ M, and by 60% at 5 μ M (Figure S29). Thus, the four-coordinate Ni^{II} center of **Nibl** affords it the ability to bind within the active site of the adenosyltransferase and to prevent the productive binding of the natural substrate **Cbl**.

Conclusion

Herein, we have described the first Ni^{II} analogues of natural cobalt corrins vitamin B₁₂ (**CNCbl**) and cobyrinic acid (**Cby**). The new Ni-corrins **Nibl** and **Niby** display the same basic structural features and lack of coordinative activity as synthetic model Ni^{II}-corrins, such as **NiCor**.^[1a,6a,c] The Ni^{II}-corrins are known to be exceptionally stable in regard to the chemical removal of their metal center,^[25] and to exhibit no affinity for axially coordinating ligands.^[1a,6c,7] This latter feature has been rationalized by the extraordinary stabilization of the low-spin d⁸ configuration by the ligand field of the ring-contracted corrin ligand.^[1a,9] As shown here, the natural corrin ligand undergoes only a small contraction of its coordination hole by about 0.03 to 0.04 Å to accommodate the low-spin Ni^{II} ion. Indeed, the 15-membered inner perimeter provided by the natural corrin macrocycle, selected for binding low-spin cobalt ions, also binds Ni^{II} consistently in

its low-spin state. In contrast, in the porphyrinoid B_{12} -related nickel complex coenzyme F_{430} ^[1a,18,26] the 16-membered porphyrinoid macrocycle is a key player in the active, specific adjustment of the spin state and coordinative activity of the nickel center to its function in the enzyme catalyzed methane formation.^[27] Indeed, the discovery of coenzyme F_{430} provoked an entirely new look at the structural effect of the tetrapyrrolic macrocycle on the coordination chemistry of bound first-row transition metals.^[18]

A common feature of the valence shell of the low spin states of the transition metal ions Ni^{II} , Co^{III} , Co^{II} and Co^I is their unoccupied $d_{x^2-y^2}$ orbital, a key factor responsible for their strong sigma bonding interactions with the four inner corrin N atoms, leading to similar radial characteristics of their corrin complexes. A differing number of valence shell electrons in Ni^{II} -, Co^{III} -, Co^{II} - and Co^I -corrins is transduced primarily into characteristically different reactivity of the metal centers in the axial direction, strongly affecting their potential binding sites there. Consequently, Ni^{II} -corrins are to be considered particularly well-suited structural mimics of corresponding isoelectronic Co^I -corrins, the critical intermediates in heterolytic organometallic transitions in B_{12} -dependent enzymes.^[28] Crystallographic insights and DFT-based structure calculations also indicate a structural similarity between Ni^{II} -corrins and the exceptional four-coordinate Co^{II} -corrins. This result contrasts strikingly with the mutually different structures of the typical five-coordinate Co^{II} -corrins and their Zn^{II} -analogues^[5] with similarly sized metal ions^[29] that differ by the number of electrons in the valence shell.

The structural analysis of the Nibyrinates predicts that the constitutively robust **Nibl** would likely be an excellent redox-resistant structural mimic for the elusive cob(I)alamin (**Cbl^I**), a highly reactive redox-active intermediate^[30] that is found in B_{12} -dependent methyl group transferases, such as methionine synthase,^[28a] as well as in the biosynthesis of **AdoCbl** from **Cbl^{II}** (via **Cbl^I**) by Cbl-adenosyltransferases.^[23,24] **Nibl** may, likewise, act as a good structural mimic of the recently described natural four-coordinate Co^{II} -corrins, proposed as key intermediates in the enzymatic transformations catalyzed by the vitamin B_{12} tailoring enzyme CblC,^[31] in corrinoid dehalogenases,^[32] or as substrates for the reduction to Co^I -species in enzymatic cobalt alkylation.^[14,33] Indeed, as verified here, **Nibl** is a very effective inhibitor of the bacterial Cbl-adenosyltransferase BtuR.

We have developed a rational and direct synthetic path from hydrogenobyrinic acid (**Hby**) via hydrogenobalamin (**Hbl**) to nibalamin (**Nibl**), a novel transition-metal analogue of the **Cbls**. Our recent studies with the Rh^{III} analogue **AdoRhbl** of **AdoCbl**,^[4d] with the Zn^{II} analogue **Znbl** of **Cbl^{II}**,^[5] and now the Ni^{II} analogue **Nibl** of **Cbl^I**, have furnished a valuable suite of cobalamin mimics for use in the study of B_{12} -dependent enzymatic processes,^[16a,28,30,34,35] and in B_{12} -dependent biological regulation.^[36] Well-characterized and adequately accessible transition metal analogues (**Metbls**) of the **Cbls** provide a promising small-compound platform that may contribute significantly to the ongoing quest for innovative B_{12} -based biological and biomedical applications.^[11,37] Along these lines, some **Metbls** may find applications as effective antivita-

mins B_{12} .^[4d,11] The availability of selected **Metbls** and of related metal corrins (**MetCors**) will also allow more detailed experimental investigations into the chemical relevance of the coordination of transition metal ions by the uniquely skewed, strongly helical and unsymmetric natural corrin ligands.^[3] Such studies will endow a more informed understanding of the specific evolutionary selection of cobalt rather than any other transition metal^[1] for the task of complex organometallic catalysis achieved by the B_{12} cofactors.

Acknowledgements

This work was supported by grants from the Austria Science Fund (FWF, P-28892, P-33059) to BK and (FWF M-2005) to MP, and from the Biotechnology and Biological Sciences Research Council (BBSRC; BB/S014020/1, BB/K009249/1 and BB/S002197/1) and Royal Society (INF/R2180062) to MJW. Part of the computational results presented was obtained using the HPC infrastructure LEO of the University of Innsbruck and the Vienna Scientific Cluster (VSC).

Conflict of interest

The authors declare no conflict of interest.

Keywords: cobalamins · porphyrinoids · transition metals · crystal structures · vitamins

- [1] a) A. Eschenmoser, *Angew. Chem. Int. Ed. Engl.* **1988**, *27*, 5–39; *Angew. Chem.* **1988**, *100*, 5–40; b) A. Eschenmoser, *Angew. Chem. Int. Ed.* **2011**, *50*, 12412–12472; *Angew. Chem.* **2011**, *123*, 12618–12681; c) J. J. R. F. da Silva, R. J. P. Williams, *The Biological Chemistry of the Elements*, Clarendon Press, Oxford, **1991**; d) J. M. Pratt in *B₁₂, Vol. 1* (Ed.: D. Dolphin), Wiley, New York, **1982**, pp. 325–392.
- [2] a) F. J. Leeper, M. J. Warren, J. M. Kelly, A. D. Lawrence in *Handbook of Porphyrin Science, Vol. 25* (Ed.: K. M. Kadish, K. M. Smith, R. Guilard), World Scientific, Singapore, **2012**, pp. 2–83; b) E. Deery, S. Schroeder, A. D. Lawrence, S. L. Taylor, A. Seyedarabi, J. Waterman, K. S. Wilson, D. Brown, M. A. Geeves, M. J. Howard, R. W. Pickersgill, M. J. Warren, *Nat. Chem. Biol.* **2012**, *8*, 933–940.
- [3] C. Kieninger, E. Deery, A. D. Lawrence, M. Podewitz, K. Wurst, E. Nemoto-Smith, F. J. Widner, J. A. Baker, S. Jockusch, C. R. Kreutz, K. R. Liedl, K. Gruber, M. J. Warren, B. Kräutler, *Angew. Chem. Int. Ed.* **2019**, *58*, 10756–10760; *Angew. Chem.* **2019**, *131*, 10869–10873.
- [4] a) H. U. Blaser, E. L. Winnacker, A. Fischli, B. Hardegger, D. Bormann, N. Hashimoto, J. Schossig, R. Keese, A. Eschenmoser, *Helv. Chim. Acta* **2015**, *98*, 1845–1920; b) J. I. Toohey, *Proc. Natl. Acad. Sci. USA* **1965**, *54*, 934–942; c) G. Holze, H. H. Inhoffen, *Angew. Chem.* **1985**, *97*, 887–888; d) F. J. Widner, A. D. Lawrence, E. Deery, D. Heldt, S. Frank, K. Gruber, K. Wurst, M. J. Warren, B. Kräutler, *Angew. Chem. Int. Ed.* **2016**, *55*, 11281–11286; *Angew. Chem.* **2016**, *128*, 11451–11456; e) V. B. Kopenhagen in *B₁₂, Vol. 2* (Ed.: D. Dolphin), Wiley, New York, **1982**, pp. 105–150; f) C. Brenig, L. Prieto, R. Oetterli, F. Zelder, *Angew. Chem. Int. Ed.* **2018**, *57*, 16308–16312; *Angew. Chem.* **2018**, *130*, 16546–16550.



- [5] C. Kieninger, J. A. Baker, M. Podewitz, K. Wurst, S. Jockusch, A. D. Lawrence, E. Deery, K. Gruber, K. R. Liedl, M. J. Warren, B. Kräutler, *Angew. Chem. Int. Ed.* **2019**, *58*, 14568–14572; *Angew. Chem.* **2019**, *131*, 14710–14714.
- [6] a) E. Bertele, H. Boos, J. D. Dunitz, F. Elsinger, A. Eschenmoser, I. Felner, H. P. Gribo, H. Gschwend, E. F. Meyer, M. Pesaro, R. Scheffold, *Angew. Chem. Int. Ed. Engl.* **1964**, *3*, 490–496; *Angew. Chem.* **1964**, *76*, 393–399; b) A. Eschenmoser, *Pure Appl. Chem.* **1969**, *20*, 1–14; c) E. Bertele, R. Scheffold, H. Gschwend, M. Pesaro, A. Fischli, M. Roth, J. Schossig, A. Eschenmoser, *Helv. Chim. Acta* **2015**, *98*, 1755–1844; d) F.-P. Montforts, M. Osmer, D. Leupold in *Handbook of Porphyrin Science, Vol. 25* (Ed.: K. M. Kadish, K. M. Smith, R. Guilard), World Scientific, Singapore, **2012**, pp. 266–308.
- [7] V. B. Kopenhagen, J. J. Pfiffner, *J. Biol. Chem.* **1971**, *246*, 3075–3077.
- [8] L. Grotjahn, V. B. Kopenhagen, L. Ernst, *Z. Naturforsch. B* **1984**, *39*, 248–251.
- [9] J. D. Dunitz, E. F. Meyer, Jr., *Helv. Chim. Acta* **1971**, *54*, 77–89.
- [10] a) S. J. Lippard, J. M. Berg, *Principles of Bioinorganic Chemistry*, University Science Books, Mill Valley, **1994**; b) H.-B. Kraatz, N. Metzler-Nolte, *Concepts and Models in Bioinorganic Chemistry*, Wiley-VCH, Weinheim, **2006**; c) F. A. Cotton, G. Wilkinson, *Advanced Inorganic Chemistry*, Interscience Publishers, New York, **1972**.
- [11] a) F. Zelder, R. Alberto in *Handbook of Porphyrin Science, Vol. 25* (Ed.: K. M. Kadish, K. M. Smith, R. Guilard), World Scientific, Singapore, **2012**, pp. 84–132; b) F. Zelder, *Chem. Commun.* **2015**, *51*, 14004–14017; c) B. Kräutler, *Chem. Eur. J.* **2015**, *21*, 11280–11287; d) M. Ruetz, C. Gherasim, S. N. Fedosov, K. Gruber, R. Banerjee, B. Kräutler, *Angew. Chem. Int. Ed.* **2013**, *52*, 2606–2610; *Angew. Chem.* **2013**, *125*, 2668–2672; e) E. Mutti, M. Ruetz, H. Birn, B. Kräutler, E. Nexo, *Plos One* **2013**, *8*, e75312; f) M. Ruetz, A. Shanmuganathan, C. Gherasim, A. Karasik, R. Salchner, C. Kieninger, K. Wurst, R. Banerjee, M. Koutmos, B. Kräutler, *Angew. Chem. Int. Ed.* **2017**, *56*, 7387–7392; *Angew. Chem.* **2017**, *129*, 7493–7498.
- [12] F. Kreppelt, ETH Zürich, <https://doi.org/10.3929/ethz-a-000626280> (Zürich), **1991**.
- [13] F. J. Widner, F. Gstrein, B. Kräutler, *Helv. Chim. Acta* **2017**, *100*, e1700170.
- [14] a) M. F. Summers, L. G. Marzilli, A. Bax, *J. Am. Chem. Soc.* **1986**, *108*, 4285–4294; b) A. Bax, L. G. Marzilli, M. F. Summers, *J. Am. Chem. Soc.* **1987**, *109*, 566–574; c) M. Rossi, J. P. Glusker, L. Randaccio, M. F. Summers, P. J. Toscano, L. G. Marzilli, *J. Am. Chem. Soc.* **1985**, *107*, 1729–1738; d) M. Tollinger, R. Konrat, B. Kräutler, *Helv. Chim. Acta* **1999**, *82*, 1596–1609.
- [15] D. Lexa, J. M. Savéant, *Acc. Chem. Res.* **1983**, *16*, 235–243.
- [16] a) B. Kräutler, B. Puffer in *Handbook of Porphyrin Science, Vol. 25* (Ed.: K. M. Kadish, K. M. Smith, R. Guilard), World Scientific, Singapore, **2012**, pp. 133–265; b) M. Kumar, P. M. Kozlowski, *Coord. Chem. Rev.* **2017**, *333*, 71–81.
- [17] B. Kräutler, W. Keller, M. Hughes, C. Caderas, C. Kratky, *J. Chem. Soc. Chem. Commun.* **1987**, 1678–1680.
- [18] C. Kratky, R. Waditschatka, C. Angst, J. E. Johansen, J. C. Plaquevent, J. Schreiber, A. Eschenmoser, *Helv. Chim. Acta* **1985**, *68*, 1312–1337.
- [19] B. Kräutler, W. Keller, C. Kratky, *J. Am. Chem. Soc.* **1989**, *111*, 8936–8938.
- [20] L. Ouyang, P. Rulis, W. Y. Ching, G. Nardin, L. Randaccio, *Inorg. Chem.* **2004**, *43*, 1235–1241.
- [21] M. S. St. Maurice, P. Mera, K. Park, T. C. Brunold, J. C. Escalante-Semerena, I. Rayment, *Biochemistry* **2008**, *47*, 5755–5766.
- [22] a) V. B. Pett, M. N. Liebman, P. Murray-Rust, K. Prasad, J. P. Glusker, *J. Am. Chem. Soc.* **1987**, *109*, 3207–3215; b) C. Kratky, G. Färber, K. Gruber, K. Wilson, Z. Dauter, H. F. Noltling, R. Konrat, B. Kräutler, *J. Am. Chem. Soc.* **1995**, *117*, 4654–4670; c) C. Kratky, B. Kräutler in *Chemistry and Biochemistry of B₁₂* (Ed.: R. Banerjee), Wiley, New York, **1999**, pp. 9–41.
- [23] a) F. G. Costa, J. C. Escalante-Semerena, *Biochemistry* **2018**, *57*, 5076–5087; b) T. A. Stich, N. R. Buan, J. C. Escalante-Semerena, T. C. Brunold, *J. Am. Chem. Soc.* **2005**, *127*, 8710–8719; c) K. Park, P. E. Mera, T. C. Moore, J. C. Escalante-Semerena, *Angew. Chem. Int. Ed.* **2015**, *54*, 7158–7161; *Angew. Chem.* **2015**, *127*, 7264–7267.
- [24] a) T. A. Stich, M. Yamanishi, R. Banerjee, T. C. Brunold, *J. Am. Chem. Soc.* **2005**, *127*, 7660–7661; b) C. Gherasim, M. Lofgren, R. Banerjee, *J. Biol. Chem.* **2013**, *288*, 13186–13193.
- [25] N. J. Lewis, A. Pfaltz, A. Eschenmoser, *Angew. Chem. Int. Ed. Engl.* **1983**, *22*, 735–736; *Angew. Chem.* **1983**, *95*, 743–744.
- [26] a) A. Pfaltz, B. Jaun, A. Fassler, A. Eschenmoser, R. Jaenchen, H. H. Gilles, G. Diekert, R. K. Thauer, *Helv. Chim. Acta* **1982**, *65*, 828–865; b) C. Kratky, A. Fässler, A. Pfaltz, B. Kräutler, B. Jaun, A. Eschenmoser, *J. Chem. Soc. Chem. Commun.* **1984**, 1368–1371.
- [27] B. Jaun, R. K. Thauer, *Metal Ions in Life Sciences, Vol. 6* (Eds.: A. Sigel, H. Sigel, R. K. O. Sigel), deGruyter, Berlin, **2009**, pp. 115–132.
- [28] a) R. G. Matthews, *Acc. Chem. Res.* **2001**, *34*, 681–689; b) K. Gruber, B. Puffer, B. Kräutler, *Chem. Soc. Rev.* **2011**, *40*, 4346–4363; c) T. C. Brunold, K. S. Conrad, M. D. Liptak, K. Park, *Coord. Chem. Rev.* **2009**, *253*, 779–794.
- [29] a) B. Cordero, V. Gomez, A. E. Platero-Prats, M. Reves, J. Echeverria, E. Cremades, F. Barragan, S. Alvarez, *Dalton Trans.* **2008**, 2832–2838; b) R. D. Shannon, *Acta Crystallogr. Sect. A* **1976**, *32*, 751–767.
- [30] B. Kräutler in *Advances in Bioorganometallic Chemistry* (Eds.: T. Hirao, T. Moriuchi), Elsevier, Amsterdam, **2019**, pp. 399–430.
- [31] R. Banerjee, C. Gherasim, D. Padovani, *Curr. Opin. Chem. Biol.* **2009**, *13*, 484–491.
- [32] a) B. Bommer, C. Kunze, J. Fessler, T. Schubert, G. Diekert, H. Dobbek, *Science* **2014**, *346*, 455–458; b) K. A. P. Payne, C. P. Quezada, K. Fisher, M. S. Dunstan, F. A. Collins, H. Sjuts, C. Levy, S. Hay, S. E. J. Rigby, D. Leys, *Nature* **2015**, *517*, 513–516.
- [33] I. G. Pallares, T. C. Moore, J. C. Escalante-Semerena, T. C. Brunold, *Biochemistry* **2014**, *53*, 7969–7982.
- [34] a) R. Banerjee, *Chem. Rev.* **2003**, *103*, 2083–2094; b) T. Toraya, *Chem. Rev.* **2003**, *103*, 2095–2127; c) P. A. Frey, A. D. Hegeman, G. H. Reed, *Chem. Rev.* **2006**, *106*, 3302–3316; d) W. Buckel, B. T. Golding, *Annu. Rev. Microbiol.* **2006**, *60*, 27–49; e) E. N. G. Marsh, C. L. Drennan, *Curr. Opin. Chem. Biol.* **2001**, *5*, 499–505.
- [35] a) J. Bridwell-Rabb, A. Zhong, H. G. Sun, C. L. Drennan, H.-w. Liu, *Nature* **2017**, *544*, 322–326; b) D. G. Fujimori, *Curr. Opin. Chem. Biol.* **2013**, *17*, 597–604; c) Q. Zhang, W. van der Donk, W. Liu, *Acc. Chem. Res.* **2012**, *45*, 555.
- [36] a) A. Nahvi, J. E. Barrick, R. R. Breaker, *Nucleic Acids Res.* **2004**, *32*, 143–150; b) A. Schaffner, X. T. Li, Y. Gomez-Llorente, E. Leandrou, A. Memou, N. Clemente, C. Yao, F. Afsari, L. T. Zhi, N. N. Pan, K. Morohashi, X. L. Hua, M. M. Zhou, C. Wang, H. Zhang, S. G. Chen, C. J. Elliott, H. Rideout, I. Ubarretxena-Belandia, Z. Y. Yue, *Cell Res.* **2019**, *29*, 313–329; c) S. Padmanabhan, R. Perez-Castano, M. Elias-Arnanz, *Curr. Opin. Struct. Biol.* **2019**, *57*, 47–55.
- [37] a) S. M. Clardy, D. G. Allis, T. J. Fairchild, R. P. Doyle, *Expert Opin. Drug Delivery* **2011**, *8*, 127–140; b) M. Hunger, E. Mutti, A. Rieder, B. Enders, E. Nexo, B. Kräutler, *Chem. Eur. J.* **2014**, *20*, 13103–13107; c) T. A. Shell, D. S. Lawrence, *Acc. Chem. Res.* **2015**, *48*, 2866–2874; d) A. D. Lawrence, E. Nemoto-Smith, E. Deery, J. A. Baker, S. Schroeder, D. G. Brown, J. M. A. Tullet, M. J. Howard, I. R. Brown, A. G. Smith, H. I. Boshoff, C. E. Barry, M. J. Warren, *Cell Chem. Biol.* **2018**, *25*, 941–951; e) S. Kainrath, M. Stadler, E. Reichhart, M. Distel, H. Janovjak,

Angew. Chem. Int. Ed. **2017**, *56*, 4608–4611; *Angew. Chem.* **2017**, *129*, 4679–4682; f) M. Giedyk, A. Jackowska, M. Równicki, M. Kolanowska, J. Trylska, D. Gryko, *Chem. Commun.* **2019**, *55*, 763–766; g) E. Braselmann, A. J. Wierzba, J. T. Polaski, M. Chromiński, Z. E. Holmes, S.-T. Hung, D. Batan, J. R. Wheeler,

R. Parker, R. Jimenez, D. Gryko, R. T. Batey, A. E. Palmer, *Nat. Chem. Biol.* **2018**, *14*, 964–971.

Manuscript received: June 14, 2020
Accepted manuscript online: July 20, 2020
Version of record online: ■ ■ ■ ■, ■ ■ ■ ■


Research Articles



Macrocycles

C. Kieninger, K. Wurst, M. Podewitz,
M. Stanley, E. Deery, A. D. Lawrence,
K. R. Liedl, M. J. Warren,*
B. Kräutler* ————— ■■■■-■■■■

Replacement of the Cobalt Center of
Vitamin B₁₂ by Nickel: Nibalamine and
Nibyric Acid Prepared from Metal-Free
B₁₂ Ligands Hydrogenobalamin and
Hydrogenobyric Acid



Nibalamine (**Nibl**), the novel Ni^{II} analogue of vitamin B₁₂, was synthesized from the metal-free B₁₂ ligand hydrogenobalamin, prepared from biosynthetic hydrogenobyric acid (**Hby**). An X-ray crystal analysis of nibyric acid, the Ni^{II} complex of **Hby**, revealed the first structural details of a nickel complex of a natural corrin ligand. **Nibl** is a structural mimic of the corresponding four-coordinate cobalamins, which are of interest in biological investigations.

