



Kent Academic Repository

Gonzales, Rafael, Pech, Clelia, Perrin, Nigel and Samokhin, A (2021) *Geometry of horospherical varieties of Picard rank one*. International Mathematics Research Notices . ISSN 1073-7928.

Downloaded from

<https://kar.kent.ac.uk/83827/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1093/imrn/rnaa331>

This document version

Author's Accepted Manuscript

DOI for this version

Licence for this version

UNSPECIFIED

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

GEOMETRY OF HOROSPHERICAL VARIETIES OF PICARD RANK ONE

R. GONZALES, C. PECH, N. PERRIN, AND A. SAMOKHIN

ABSTRACT. We study the geometry of smooth non-homogeneous horospherical varieties of Picard rank one. These have been classified by Pasquier and include the well-known odd symplectic Grassmannians. We focus our study on quantum cohomology, with a view towards Dubrovin's conjecture. In particular, we describe the cohomology groups of these varieties as well as a Chevalley formula, and prove that many Gromov-Witten invariants are enumerative. This enables us to prove that in many cases the quantum cohomology is semisimple. We give a presentation of the quantum cohomology ring for odd symplectic Grassmannians. The final two sections are devoted to the derived categories of coherent sheaves on horospherical varieties. We first discuss a general construction of exceptional bundles on these varieties. We then study in detail the case of the horospherical variety associated to the exceptional group G_2 , and construct a full rectangular Lefschetz exceptional collection in the derived category.

CONTENTS

Introduction	1
1. Horospherical varieties of Picard rank one - Geometry	5
2. Moduli space of curves	11
3. Quantum cohomology	15
4. Examples	20
5. The odd symplectic Grassmannian	24
6. Recollections on semiorthogonal decompositions	40
7. A construction of exceptional vector bundles on horospherical varieties	42
8. A full rectangular Lefschetz exceptional collection on the G_2 -variety	46
9. Appendix	64
References	65

INTRODUCTION

Schubert calculus and enumerative geometry are very classical. One of their many aspects is the study of intersection theory of projective rational homogeneous spaces G/P with G a reductive group and P a parabolic subgroup. We now have a very good understanding of the cohomology of G/P as well as a good understanding of their quantum cohomology and even their quantum K -theory.

Date: 30.03.2020.

2010 Mathematics Subject Classification. 14M27,14N35,14F05,14M17.

One of the simplest classes of quasi-homogeneous spaces are toric varieties for which many results are known. A common generalisation of rational projective homogeneous spaces and toric varieties are spherical varieties. Recall that a normal variety X is called G -spherical for G a reductive group if X has a dense B -orbit, where $B \subset G$ is a Borel subgroup. Spherical varieties form such a large class of varieties that we do not even have a description of their cohomology ring in general. In this paper, we concentrate on a special class of spherical varieties: *horospherical varieties*, see Definition 1.2. More specifically, we consider smooth projective horospherical varieties of Picard rank one. These varieties have been classified by Pasquier [45], who proved that they are either homogeneous or can be constructed in a uniform way via a triple $(\text{Type}(G), \omega_Y, \omega_Z)$ of representation-theoretic data. Here $\text{Type}(G)$ is the semisimple Lie type of a reductive group G and ω_Y, ω_Z are fundamental weights, see Section 1.3. Pasquier has classified the possible triples in five classes:

- (1) $(B_n, \omega_{n-1}, \omega_n)$ with $n \geq 3$;
- (2) $(B_3, \omega_1, \omega_3)$;
- (3) $(C_n, \omega_m, \omega_{m-1})$ with $n \geq 2$ and $m \in [2, n]$;
- (4) $(F_4, \omega_2, \omega_3)$;
- (5) $(G_2, \omega_1, \omega_2)$.

Following [46], we denote by $X^1(n), X^2, X^3(n, m), X^4$ and X^5 the corresponding varieties, the superscript indicating the class that a variety belongs to. We will refer to this list of varieties as Pasquier's list.

The variety $X^3(n, m)$ from the third class has already attracted some attention since in this case it is isomorphic to the so-called *odd symplectic Grassmannian* $\text{IG}(m, 2n + 1)$ (see Subsection 1.9 and Section 5). They were studied, for instance, in [40, 47, 48] and very recently in [41, 39].

In this paper, we start a systematic study of the geometry of non homogeneous projective smooth horospherical varieties X with $\text{Pic}(X) = \mathbb{Z}$, with a view towards Dubrovin's conjecture. Our first results are concerned with the cohomology ring $H^*(X, \mathbb{Z})$. We describe two *Schubert-like* cohomology basis $(\sigma'_u, \tau'_v)_{u,v}$ and $(\sigma_u, \tau_v)_{u,v}$ and prove that these bases are Poincaré dual to each other, see Proposition 1.10. We also give a Chevalley formula, *i.e.*, a formula for multiplying any of the above classes with the hyperplane class h generating the Picard group, see Proposition 1.14.

We then turn to the study of rational curves, in particular to the study of $M_{d,k}(X)$, the Kontsevich moduli space of degree d , genus zero, stable maps with k marked points. In particular, we prove the following result (Theorem 2.5):

Theorem. Let X be a non-homogeneous projective smooth horospherical variety with $\text{Pic}(X) = \mathbb{Z}$. Then

- (1) $M_{1,k}(X)$ is irreducible of dimension $\dim X + c_1(X) + k - 3$.
- (2) $M_{d,k}(X)$ is irreducible of dimension $\dim X + dc_1(X) + k - 3$ for all d if $X = X^2$ or $X = X^3(n, m)$.
- (3) $M_{2,k}(X)$ is irreducible of dimension $\dim X + 2c_1(X) + k - 3$ for $X = X^1(3)$ and $X = X^4$, and has two irreducible components of dimension $\dim X + 2c_1(X) + k - 3$ if $X = X^1(4)$ or $X = X^5$.

In all cases above, the dimension of the moduli space $M_{d,k}(X)$ is equal to the expected dimension $\dim X + dc_1(X) + k - 3$. This implies that the Gromov-Witten

invariants are enumerative and proves, in particular, a conjecture stated in the second author's PhD thesis (see Corollary 3.4). It is worth noting that the moduli space $M_{d,k}(X)$ is not irreducible and/or has non expected dimension whenever $c_1(Z) > c_1(X)$ and d is large enough, see Corollary 2.3 and the proof of Theorem 2.5.

Corollary. If $X = X^2$ or $X = X^3(n, m)$, then the Gromov–Witten invariants of X are enumerative.

Using these enumerativity results, we are able to compute many Gromov–Witten invariants. In particular, we prove the following result, see Sections 3.3 and 4:

Theorem. If X is one of the varieties $X^1(3)$, X^2 , $X^3(3, 3)$ and X^5 , then

- (1) there is an explicit quantum Chevalley formula;
- (2) the small quantum cohomology ring $\text{QH}(X)$ is semisimple.

Note that for the varieties $X^3(n, m)$, *i.e.* for odd symplectic Grassmannians, such a quantum (and even equivariant quantum) Chevalley formula has been obtained recently by Mihalcea and Shifler [41]. In Section 5, we use our proof of a conjecture by the second author to deduce several results on the specific case of odd symplectic Grassmannians (case (3) of Pasquier's classification). In particular, we prove quantum-to-classical results that enable us to compute Gromov–Witten invariants on $\text{IG}(m, 2n + 1)$ thanks to cohomological (thus classical) invariants on $\text{IG}(m + 1, 2n + 3)$ (see Theorem 5.13). In particular, this result contains the quantum Pieri rule even though we do not have any closed formula at the moment. Using these results we deduce a presentation for the quantum cohomology ring $\text{QH}(\text{IG}(m, 2n + 1))$. Let $(\tau'_p)_{p \in [1, 2n+1-m]}$ the Chern classes of the tautological quotient bundle on $\text{IG}(m, 2n + 1)$ and for $r \geq 1$, define

$$d_r := \det(\tau'_{1+j-i})_{1 \leq i, j \leq r} \quad \text{and} \quad b_r := (\tau'_r)^2 + 2 \sum_{i \geq 1} (-1)^i \tau'_{r+i} \tau'_{r-i},$$

with the convention that $\tau'_0 := 1$ and $\tau'_p := 0$ if $p < 0$ or $p > 2n + 1 - m$.

Theorem. The quantum cohomology ring $\text{QH}(\text{IG}(m, 2n + 1), \mathbb{C})$ is generated by the classes $(\tau'_p)_{1 \leq p \leq 2n+1-m}$ and the quantum parameter q , and the relations are

$$\begin{aligned} d_r &= 0 \text{ for } m + 1 \leq r \leq 2n + 1 - m, \\ d_{2n+2-m} &= \begin{cases} 0 & \text{if } m \text{ is even,} \\ -q & \text{if } m \text{ is odd,} \end{cases} \\ b_s &= (-1)^{2n+1-m-s} q \tau'_{2s-2n-2+m} \text{ for } n + 2 - m \leq s \leq n, \end{aligned}$$

Finally, let $\text{BQH}(X)$ denote the big quantum cohomology ring of X . Recall Dubrovin's conjecture:

Conjecture. Let X be a smooth projective Fano variety. The ring $\text{BQH}(X)$ is generically semisimple if and only if the bounded derived category $\mathcal{D}^b(X)$ has a full exceptional collection.

The above results therefore suggest to look for full exceptional collections in many horospherical varieties of the classification. Some results have already been obtained.

For instance, the second author proved in [48] that Dubrovin’s conjecture holds for $X^3(n, 2)$. Dubrovin’s conjecture for X^2 follows from results by the third author [51] for the semisimplicity of quantum cohomology and results by Kuznetsov [34] for the derived category. Indeed, the variety X^2 is a hyperplane section of the maximal orthogonal Grassmannian $\text{OG}(5, 10)$ (see Subsection 1.9). In the last two sections, we consider the derived category side of Dubrovin’s conjecture. In Section 7, for all cases of Pasquier’s list except the case (2) we construct an exceptional vector bundle on X . In Section 8, we extend this construction and produce a full exceptional collection on the horospherical variety X^5 :

$$(1) \quad \left\langle \begin{array}{cccccc} \widehat{\mathbb{S}}(-3), & \mathbb{U}(-3), & \mathcal{O}_{X^5}(-3), & \widehat{\mathbb{S}}(-2), & \mathbb{U}(-2), & \mathcal{O}_{X^5}(-2), \\ \widehat{\mathbb{S}}(-1), & \mathbb{U}(-1), & \mathcal{O}_{X^5}(-1), & \widehat{\mathbb{S}}, & \mathbb{U}, & \mathcal{O}_{X^5} \end{array} \right\rangle.$$

The object \mathbb{U} is a vector bundle, but $\widehat{\mathbb{S}}$ is a two-term complex.

The exceptional collection (1) has an additional property. Namely, it is of the form $\langle \mathcal{A} \otimes \mathcal{O}_{X^5}(-3), \mathcal{A} \otimes \mathcal{O}_{X^5}(-2), \mathcal{A} \otimes \mathcal{O}_{X^5}(-1), \mathcal{A} \rangle$, where $\mathcal{A} = \{\widehat{\mathbb{S}}, \mathbb{U}, \mathcal{O}_{X^5}\}$. Such collections are called *Lefschetz exceptional collections*, see [34].

Theorem. The collection (1) is a full rectangular Lefschetz exceptional collection on X .

Note that the above theorem, together with the results of Section 4.5, prove for the variety X^5 a refinement of Dubrovin’s conjecture recently proposed by Kuznetsov and Smirnov [37, Conjecture 1.12] (see also Section 6.4 for more details).

Acknowledgements. R.G., N.P., and A.S. would like to thank the Mathematical Institute of Düsseldorf where this work first started. C.P. thanks the University of Kent for support. R.G. and N.P. were supported by a DFG grant Project “Gromov–Witten Theorie, Geometrie und Darstellungen” (PE 2165/1–2). N.P. was supported by a public grant as part of the “Investissement d’Avenir” project, reference ANR–11–LABX–0056–LMH, LabEx LMH. A.S. was supported by the strategic research fund of the University of Düsseldorf (grant SFF F–2015/946–8), by the SFB/TR 45 of the University of Duisburg–Essen, and by Prof. M. Levine’s University Fellowship. A.S. also acknowledges the support of the IHES and of the Klaus Tschira Stiftung at the final stage of the present work. He is indebted to A.Kuznetsov who suggested using Hecke modifications to construct exceptional vector bundles in Sections 7–8. Finally, all the authors are grateful to the anonymous referees for their very careful reading of our paper and for the many improvements suggested.

Notation. In this paper, G denotes a complex connected reductive group. We use notation s from [26]. Choosing a maximal torus T in G , let R denote the set of roots with respect to T . Given a root $\alpha \in R$, denote \mathfrak{g}_α the root subspace so that we have a root decomposition $\mathfrak{g} = \mathfrak{h} \oplus_{\alpha \in R} \mathfrak{g}_\alpha$ where $\mathfrak{h} = \text{Lie}(T)$. We choose a Borel subgroup $B \supset T$ and declare the roots of B to be the negative roots $R^- \subset R$. Denote $R^+ = -R^-$ the positive roots, so that $R = R^+ \cup R^-$. Denote $S \subset R^+$ the subset of simple roots. If needed, the Borel subgroup corresponding to R^+ will be denoted by B^+ . A standard parabolic subgroup is a closed subgroup $P \subset G$ containing B . We denote G/B the full flag variety of G . Denote $X(T)$ the weight lattice and $X^+(T)$ the set of dominant weights relative to positive roots R^+ *i.e.* $\lambda \in X^+(T)$ if $\langle \lambda, \alpha^\vee \rangle \geq 0$ for any simple coroot α^\vee . Given a parabolic subgroup

P and a P -variety F (i.e., a variety equipped with a left action of P), define the crossed product $G \times^P F$ to be the quotient of $G \times F$ modulo the equivalence relation $(g, v) = (gp, p^{-1} \cdot v)$ for $g \in G, p \in P$ and $v \in F$. Given a character $\chi \in X(T)$, set \mathbb{C}_χ to be the corresponding one-dimensional B -module obtained by the restriction $B \rightarrow T$. We denote \mathcal{L}_χ the G -equivariant line bundle on G/B defined as the quotient of $G \times^B \mathbb{C}_\chi$. Given a dominant weight $\lambda \in X^+(T)$, we denote $V(\lambda)$ the induced module $\text{Ind}_B^G(\mathbb{C}_\lambda)$, the highest weight irreducible representation of G of highest weight λ (see [26, Proposition 2.2, Part II]). Let $\chi \in X(T)$ be a character, let P be a standard parabolic subgroup and L its Levi subgroup containing T . Set $V_P(\chi) = \text{Ind}_{B \cap L}^L(\chi)$ seen as a P -representation via the surjective map $P \rightarrow L$. Note that we have $G \times^P V_P(\chi) = \pi_* \mathcal{L}_\chi$ where $\pi : G/B \rightarrow G/P$ is the canonical map.

Remark 0.1. There is another commonly used notation (for example in [45] where the Borel subgroup B is chosen to be associated to positive roots. In that case, the line bundle \mathcal{L}_λ has to be induced from the representation $\mathbb{C}_{-\lambda}$. In order to avoid this sign and because Borel-Weil-Bott's Theorem takes a simpler form, we decided to choose the above convention following [26].

1. HOROSPHERICAL VARIETIES OF PICARD RANK ONE - GEOMETRY

In this section we recall some basic properties of smooth projective horospherical varieties of Picard rank one. Our reference is [45], see also [44, 50].

1.1. Horospherical varieties. Let G be a complex reductive group. A G -variety is a reduced scheme of finite type over the field of complex numbers \mathbb{C} , equipped with an algebraic action of G . Some of the simplest G -varieties are spherical varieties.

Definition 1.1. Let G be a reductive group and $B \subset G$ be a Borel subgroup. A G -variety X is called *spherical* if X has a dense B -orbit.

Spherical varieties admit many equivalent characterisations. For example, a normal G -variety is spherical if and only if it has finitely many B -orbits. We refer to [29, 50, 22, 52] and references therein for more on the geometry of spherical varieties or their classification.

We focus on a special case of spherical varieties: *horospherical varieties*. We give two equivalent definitions of horospherical varieties, one from a representation-theoretic point of view, the other more geometric. We refer to Pasquier's PhD thesis [44] for details.

Definition 1.2. Let X be a G -spherical variety and let H be the stabiliser of a point in the dense G -orbit in X . The variety X is called *horospherical* if H contains a conjugate of U , the maximal unipotent subgroup of G contained in the Borel subgroup B .

Remark 1.3. A more geometric equivalent definition can be given as follows. Let X be a G -variety. Then X is horospherical if there exists a parabolic subgroup $P \subset G$, a torus S which is a quotient of P , and a toric S -variety F such that there exists a diagram of G -equivariant morphisms

$$\begin{array}{ccc} \tilde{X} = G \times^P F & \xrightarrow{p} & G/P \\ \pi \downarrow & & \\ & & X \end{array}$$

where $G \times^P F$ is the contracted product, π is birational, and p is a fibration with fibers isomorphic to F . The *rank* of the horospherical variety X is the dimension of the toric variety F , namely, $\text{rk}(X) := \dim F$.

1.2. Classification. Horospherical varieties can be classified using *colored fans*, see [44]. We focus here on smooth projective horospherical varieties of Picard rank one. Pasquier [45] constructed all these varieties as well as their minimal embedding.

Theorem 1.4 (Pasquier). *Let X be a smooth projective G -horospherical variety of Picard rank one. Then either*

(0) X is homogeneous,

or X can be constructed in a uniform way from a triple $(\text{Type}(G), \omega_Y, \omega_Z)$ belonging to the following list:

- (1) $(B_n, \omega_{n-1}, \omega_n)$ with $n \geq 3$;
- (2) $(B_3, \omega_1, \omega_3)$;
- (3) $(C_n, \omega_m, \omega_{m-1})$ with $n \geq 2$ and $m \in [2, n]$;
- (4) $(F_4, \omega_2, \omega_3)$;
- (5) $(G_2, \omega_1, \omega_2)$,

where $\text{Type}(G)$ is the semisimple Lie type of G and ω_Y, ω_Z are fundamental weights.

As stated in the above theorem some smooth projective horospherical varieties of Picard rank one are homogeneous (for instance projective spaces). We will not consider them since their moduli space of curves, quantum cohomology and quantum K -theory have been studied before (see [55, 28, 49, 6, 31, 32, 7, 8, 10, 9, 12, 11, 14, 4, 5] and references therein). We will therefore focus on the second part of the list, *i.e.*, on the cases from (1) to (5). In what follows, we denote by $X^1(n)$, X^2 , $X^3(n, m)$, X^4 and X^5 the corresponding varieties.

1.3. Construction. From now on, let X be a smooth projective non-homogeneous horospherical variety of Picard rank one with associated triple $(\text{Type}(G), \omega_Y, \omega_Z)$. We introduce the following notation which will be used throughout the paper.

Notation 1.5. Let G be the reductive group for which X is horospherical. Then we denote by V_Y and V_Z the irreducible G -representations with highest weights ω_Y and ω_Z and we let v_Y and v_Z be the corresponding lowest weight vectors.

We write P_Y for the stabiliser of $[v_Y]$ in $\mathbb{P}(V_Y)$ and P_Z for the stabiliser of $[v_Z]$ in $\mathbb{P}(V_Z)$. Both stabilizers are standard parabolic subgroups of G . We denote by Y the G -orbit of $[v_Y]$ in $\mathbb{P}(V_Y)$ and by Z that of $[v_Z]$ in $\mathbb{P}(V_Z)$, so that $Y \simeq G/P_Y$ and $Z \simeq G/P_Z$.

Pasquier [45] proves that X is obtained as follows.

Proposition 1.6 (Pasquier). *Let X be one of the varieties in cases (1) to (5) of Theorem 1.4. Then $X = \overline{G \cdot [v_Y + v_Z]} \subset \mathbb{P}(V_Y \oplus V_Z)$ is the closure of the G -orbit $G \cdot [v_Y + v_Z]$ in $\mathbb{P}(V_Y \oplus V_Z)$.*

1.4. Orbits. Let $\text{Aut}(X)$ be the automorphism group of X . The group G acts on X with three orbits: Y , Z , and the complement \mathcal{U} of $Y \cup Z$ in X , which is open and dense in X .

The group $\text{Aut}(X)$ is a semi-direct product of G with its unipotent radical. It acts on X with two orbits. While until then the roles of Y and Z were interchangeable,

we now choose to use the notation Z for the unique closed $\text{Aut}(X)$ -orbit and we call it the *closed orbit of X* . The other $\text{Aut}(X)$ -orbit is $\mathcal{U}_Y = X \setminus Z = \mathcal{U} \cup Y$. We also define $\mathcal{U}_Z = X \setminus Y = \mathcal{U} \cup Z$.

The G -orbits Y and Z are projective and respectively isomorphic to G/P_Y and G/P_Z , where we choose P_Y and P_Z parabolic subgroups containing the same Borel subgroup B . The open G -orbit \mathcal{U} is a \mathbb{G}_m -bundle over the incidence variety $G/(P_Y \cap P_Z)$. We refer to [45] for proofs.

1.5. Blow-ups and projections. We refer to [45] for the results in this subsection. Let $\pi_Z : \tilde{X}_Z \rightarrow X$ be the blow-up of Z in X . It is obtained via base change from the blow-up of $\mathbb{P}(V_Z)$ in $\mathbb{P}(V_Y \oplus V_Z)$. In particular there is a natural projection $p_Y : \tilde{X}_Z \rightarrow Y$, obtained as the restriction of the projection of the former blow-up to $\mathbb{P}(V_Y)$. The projection $p_Y : \tilde{X}_Z \rightarrow Y$ restricts to a projection $p_Y : \mathcal{U}_Y \rightarrow Y$, which realises \mathcal{U}_Y as a vector bundle over Y . This bundle is G -equivariant and thus it is obtained from a P_Y -representation. We write N_Y for the corresponding locally free sheaf; it is the normal bundle to Y in X . The projection $p_Y : \tilde{X}_Z \rightarrow Y$ realises \tilde{X}_Z as the projective bundle $\mathbb{P}_Y(N_Y \oplus \mathcal{O}_Y)$. Write E_Z for the exceptional divisor.

Exchanging the roles of Y and Z we denote by $\pi_Y : \tilde{X}_Y \rightarrow X$ the blow-up of Y in X . In particular there is a natural projection $p_Z : \tilde{X}_Y \rightarrow Z$, which restricts to a projection $p_Z : \mathcal{U}_Z \rightarrow Z$. This projection realises \mathcal{U}_Z as a G -equivariant vector bundle over Z , which is thus obtained from a P_Z -representation. We write N_Z for the corresponding locally free sheaf; it is the normal bundle to Y in X . The projection $p_Z : \tilde{X}_Y \rightarrow Z$ realises \tilde{X}_Y as the projective bundle $\mathbb{P}_Z(N_Z \oplus \mathcal{O}_Z)$. Write E_Y for the exceptional divisor.

In both cases, the exceptional divisors E_Y and E_Z are isomorphic to the incidence variety $E = G/(P_Y \cap P_Z)$. The maps p_Y and π_Z from E_Z to Y and Z and the maps π_Y and p_Z from E_Y to Y and Z are the projections p and q from E to G/P_Y and G/P_Z .

Summarising, we get the commutative diagrams

$$\begin{array}{ccc}
 & \xrightarrow{p} & \\
 E_Z & \xrightarrow{j_Z} \tilde{X}_Z \xrightarrow{p_Y} & Y \\
 q \downarrow & & \downarrow \pi_Z \\
 Z & \xrightarrow{i_Z} & X
 \end{array}
 \qquad
 \begin{array}{ccc}
 & \xrightarrow{q} & \\
 E_Y & \xrightarrow{j_Y} \tilde{X}_Y \xrightarrow{p_Z} & Z \\
 p \downarrow & & \downarrow \pi_Y \\
 Y & \xrightarrow{i_Y} & X
 \end{array}$$

Let $\pi_{YZ} : \tilde{X}_{YZ} \rightarrow X$ be the blow-up of $Y \cup Z$ in X . There is a morphism $p_{YZ} : \tilde{X}_{YZ} \rightarrow E$ which is a \mathbb{P}^1 -bundle, and both connected components of the exceptional divisor are mapped isomorphically onto E . We will denote this last map by $\xi = p_{YZ} : \tilde{X}_{YZ} \rightarrow E$.

1.6. Numerical invariants. Using the above description, it is easy to compute some numerical invariants for X , Y and Z . For example, their dimensions are given in Table 1.

If W is a Fano variety of Picard rank one and H is the ample generator of $\text{Pic}(W)$, we denote by $c_1(W)$ the unique positive integer such that $-K_W = c_1(W)H$, where K_W is the canonical divisor of W . We describe $c_1(X)$, $c_1(Y)$ and $c_1(Z)$ in Table 2.

Fact 1.7. We have $c_1(X) = \text{codim}_X(Y) + \text{codim}_X(Z)$.

Case	$\dim X$	$\operatorname{codim}_X Y$	$\operatorname{codim}_X Z$
$X^1(n)$	$n(n+3)/2$	2	n
X^2	9	4	3
$X^3(n, m)$	$m(2n+1-m) - m(m-1)/2$	m	$2(n+1-m)$
X^4	23	3	3
X^5	7	2	2

TABLE 1. Dimension of X .

Case	$c_1(X)$	$c_1(Y)$	$c_1(Z)$	$\operatorname{codim}_X Y$	$\operatorname{codim}_X Z$
$X^1(n)$	$n+2$	$n+1$	$2n$	2	n
X^2	7	5	6	4	3
$X^3(n, m)$	$2n+2-m$	$2n+1-m$	$2n+2-m$	m	$2(n+1-m)$
X^4	6	5	7	3	3
X^5	4	3	5	2	2

TABLE 2. Numerical invariants.

1.7. Cohomology classes. The description in Subsection 1.5 implies that X has a cellular decomposition. It is therefore easy to describe geometric bases of the cohomology groups of X .

Since Y and Z are projective and homogeneous under G , we can consider their Schubert varieties $(Y_u)_{u \in W^{P_Y}}$ and $(Z_v)_{v \in W^{P_Z}}$ for a given Borel subgroup $B \subset G$. These varieties define cohomology classes as follows:

$$\bar{\sigma}_u = [Y_u] \in H^{2\ell(u)}(Y, \mathbb{Z}) \text{ and } \bar{\tau}_v = [Z_v] \in H^{2\ell(v)}(Z, \mathbb{Z}).$$

We also introduce $Y'_u = \pi_Z(p_Y^{-1}(Y_u))$ and $Z'_v = \pi_Y(p_Z^{-1}(Z_v))$. Denoting by $i_Y : Y \rightarrow X$ and $i_Z : Z \rightarrow X$ the closed embeddings, we may define cohomology classes on X as follows:

$$\begin{aligned} \sigma'_u &= [Y'_u] \in H^{2\ell(u)}(X, \mathbb{Z}) \text{ and } \sigma_u = i_*[Y_u] \in H^{2\ell(u)+2\operatorname{codim}_X(Y)}(X, \mathbb{Z}) \\ \tau'_v &= [Z'_v] \in H^{2\ell(v)}(X, \mathbb{Z}) \text{ and } \tau_v = j_*[Z_v] \in H^{2\ell(v)+2\operatorname{codim}_X(Z)}(X, \mathbb{Z}). \end{aligned}$$

Using the cellular decompositions by Schubert cells in Y and Z and the cellular decomposition induced on \mathcal{U}_Y and \mathcal{U}_Z we obtain the following result.

Fact 1.8. We have the following bases of $H^*(X, \mathbb{Z})$.

1. The classes $((\sigma'_u)_{u \in W^{P_Y}}, (\tau_v)_{v \in W^{P_Z}})$ form a \mathbb{Z} -basis of $H^*(X, \mathbb{Z})$.
2. The classes $((\tau'_v)_{v \in W^{P_Z}}, (\sigma_u)_{u \in W^{P_Y}})$ form a \mathbb{Z} -basis of $H^*(X, \mathbb{Z})$.

The proof is similar to that of [33, Proposition 11.3.2], and it relies on the following observation: the closure of a cell (in X) is a union of cells. Indeed, this is already true for the cells in Y and Z . For the cells in \mathcal{U}_Y and \mathcal{U}_Z , the cells are B -stable (actually they are the inverse image of B -orbits in the vector bundle $p_Y : \mathcal{U}_Y \rightarrow Y$ or $p_Z : \mathcal{U}_Z \rightarrow Z$) so their closures in X are unions of cells in \mathcal{U}_Y or \mathcal{U}_Z or unions of B -orbits in the other orbit: in Z and in Y , respectively.

Notation 1.9. Recall that p and q are the projections from the incidence variety E to Y and Z respectively.

1. Let $u \in W^{PY}$, then $q(p^{-1}(Y_u))$ is a Schubert variety in Z of the form $Z_{\hat{v}}$. We define $\hat{u} \in W^{PZ}$ such that $q(p^{-1}(Y_u)) = Z_{\hat{u}}$. For $v \in W^{PZ}$ we define \hat{v} in a similar way: $p(q^{-1}(Z_v)) = Y_{\hat{v}}$.
2. Let $u \in W^{PY}$, then $p^{-1}(Y_u)$ is the Schubert variety $E_{\tilde{u}}$ in E associated to an element $\tilde{u} \in W^{PY \cap PZ}$. For $v \in W^{PZ}$ we define \tilde{v} in a similar way.
3. Let h_Y and h_Z the hyperplane classes in Y and Z . We define coefficients $c_Y(u, u')$ and $c_Z(v, v')$ as follows:

$$h_Y \cup \bar{\sigma}_u = \sum_{u' \in W^{PY}} c_Y(u, u') \bar{\sigma}_{u'} \quad \text{and} \quad h_Z \cup \bar{\tau}_v = \sum_{v' \in W^{PZ}} c_Z(v, v') \bar{\tau}_{v'}.$$

These coefficients are well known since they are given by the Chevalley formula for homogeneous spaces, see [15].

4. If M is a complete nonsingular irreducible variety of dimension n , then the Poincaré duality map $H^k(M, \mathbb{Z}) \rightarrow H_{2n-k}(M, \mathbb{Z})$, taking σ to $\sigma \cap [M]$, is an isomorphism, where $[M] \in H_{2n}(M, \mathbb{Z}) \simeq \mathbb{Z}$ is the fundamental class of M . If, moreover, $H_*(M, \mathbb{Z})$ is torsion free, then, by Poincaré duality, there is a perfect pairing $\langle \cdot, \cdot \rangle : H^k(M, \mathbb{Z}) \otimes H^{2n-k}(M, \mathbb{Z}) \rightarrow \mathbb{Z}$, given by $\langle \alpha, \beta \rangle := (\alpha \cup \beta) \cap [M]$. This is called the Poincaré duality pairing. If $\{x_u\}$ is a homogeneous basis for $H^*(M, \mathbb{Z})$, then the Poincaré dual basis is the basis $\{x_u^\vee\}$ dual for this pairing, so $\langle x_u, x_v^\vee \rangle = \delta_{u,v}$.
5. For $u \in W^{PY}$, we define $u^\vee \in W^{PY}$ such that $\bar{\sigma}_{u^\vee}$ is Poincaré dual to $\bar{\sigma}_u$ in $H^*(Y, \mathbb{Z})$. That is, $\bar{\sigma}_{u^\vee} = \bar{\sigma}_u^\vee$, with the notation of item 4. It is well-known that $u^\vee = w_0 u w_Y$, where w_0 is the longest element of W and w_Y is the longest element in W_{PY} . For $v \in W^{PZ}$ we define v^\vee in a similar fashion.

Proposition 1.10. *In $H^*(X, \mathbb{Z})$, the singular cohomology of X , the two bases defined above: $((\sigma'_u)_{u \in W^{PY}}, (\tau_v)_{v \in W^{PZ}})$ and $((\tau'_v)_{v \in W^{PZ}}, (\sigma_u)_{u \in W^{PY}})$ are dual for the Poincaré duality pairing. We have*

$$\sigma_u^\vee = \sigma'_{u^\vee} \quad \text{and} \quad \tau_v^\vee = \tau'_{v^\vee}.$$

Proof. Using the projection formula we get $\pi_{Z*}(p_Y^* \bar{\sigma}_{u_1} \cup \pi_Z^* \sigma_{u_2}) = \sigma'_{u_1} \cup \sigma_{u_2}$. But, again by the projection formula, we have $p_{Y*}(p_Y^* \bar{\sigma}_{u_1} \cup \pi_Z^* \sigma_{u_2}) = \bar{\sigma}_{u_1} \cup \bar{\sigma}_{u_2} = \delta_{u_1, u_2^\vee}$. We thus have $\sigma'_{u_1} \cup \sigma_{u_2} = \delta_{u_1, u_2^\vee}$. The same method works for the τ classes. We are left with proving $\sigma_u \cup \tau_v = 0$ and $\sigma'_u \cup \tau'_v = 0$. To compute these intersections, we use general translates of Y_u, Y'_u, Z_v and Z'_v under the action of G . First note that since Y and Z do not meet, we get $\sigma_u \cup \tau_v = 0$. Next we want to compute the intersection of general G -translates of Y'_u and Z'_v . We look for intersection points in Y, Z and \mathcal{U} . Since $Y'_u \cap Z = Z_{\hat{u}}$ is a Schubert variety of dimension at most $\dim X - \ell(u) - 1$, its intersection with a general translate of $Z'_v \cap Z = Z_v$ is of dimension at most $\dim X - \ell(u) - 1 - \ell(v) = -1$ and therefore empty. The same argument proves that no point of the intersection is contained in Y . Finally, if there was an intersection point in \mathcal{U} , then using the restriction of p_{YZ} to \mathcal{U} would give a point in the intersection of general translates of $p_{YZ}(\pi_{YZ}^{-1}(Y'_u)) = p^{-1}(Y_u) = E_{\tilde{u}}$ and $p_{YZ}(\pi_{YZ}^{-1}(Z'_v)) = p^{-1}(Z_v) = E_{\tilde{v}}$ in E . The sum of their codimensions is $\ell(u) + \ell(v) = \dim X = \dim E + 1$ and therefore these general translates do not meet. \square

We state the following cup product formulas for later reference.

Proposition 1.11. *Let $u_1, u_2 \in W^{PY}$ and $v_1, v_2 \in W^{PZ}$ with $\ell(u_1) + \ell(u_2) = \dim X$ and $\ell(v_1) + \ell(v_2) = \dim X$.*

1. If $\text{codim}_X Z = 2$, then $\sigma'_{u_1} \cup \sigma'_{u_2} = \delta_{\widehat{u}_1, \widehat{u}_2^\vee}$.
2. If $\text{codim}_X Y = 2$, then $\tau'_{v_1} \cup \tau'_{v_2} = \delta_{\widehat{v}_1, \widehat{v}_2^\vee}$.

Remark 1.12. The condition $\text{codim}_X Z = 2$ is satisfied for $X^3(n, n)$ and X^5 . The condition $\text{codim}_X Y = 2$ is satisfied in for $X^1(n)$, $X^3(n, 2)$ and X^5 .

Proof. We prove item 1, the proof of item 2 being similar. We again look at general G -translates of Y'_{u_1} and Y'_{u_2} . If they were meeting in \mathcal{U}_Y , then via p_Y there would be an intersection point for general translates of Y_{u_1} and Y_{u_2} in Y . This is not possible for dimension reasons. These varieties can therefore only meet in Z and we have $Y'_{u_i} \cap Z = Z_{\widehat{u}_i}$ which is of codimension at least $\ell(u_i) + 1 - \text{codim}_X Z$ in Z . We look for the intersection of general translates of $Z_{\widehat{u}_1}$ and $Z_{\widehat{u}_2}$. The sum of their codimension in Z is at least $\ell(u_1) + 1 - \text{codim}_X Z + \ell(u_2) + 1 - \text{codim}_X Z = \dim Z + 2 - \text{codim}_X Z = \dim Z$. This proves the result. \square

Remark 1.13. The same proof gives the following more precise result. Let $u_1, u_2 \in W^{P_Y}$ with $\ell(u_1) + \ell(u_2) = \dim X$. Assume that $\text{codim}_Z Z_{\widehat{u}_1} + \text{codim}_Z Z_{\widehat{u}_2} = \dim Z$, then $\sigma'_{u_1} \cup \sigma'_{u_2} = \delta_{\widehat{u}_1, \widehat{u}_2^\vee}$.

1.8. Hasse diagram. In this subsection we describe the multiplication by the hyperplane class h in $H^*(X, \mathbb{Z})$.

Proposition 1.14. *We have*

$$h \cup \sigma'_u = \tau_{\widehat{u}} + \sum_{u' \in W^{P_Y}} c_Y(u, u') \sigma'_{u'} \quad \text{and} \quad h \cup \tau_v = \sum_{v' \in W^{P_Z}} c_Z(v, v') \tau_{v'}.$$

Proof. The first formula follows from the equalities $\pi_Z^* h = [E] + p_Y^* h_Y$ and $\pi_Z(E \cap p_Y^{-1}(Y_u)) = Z_{\widehat{u}}$. The second formula can be directly computed in Z . \square

Remark 1.15. Similar formulas hold for the products $h \cup \sigma_u$ and $h \cup \tau'_v$.

1.9. Special geometry. In two cases of Pasquier's classification, there is a special geometric feature that we want to use: the variety X is the zero locus of a section of a vector bundle. Let us fix first some notation. Denote by $\text{OG}(5, 10)$ a connected component of the Grassmannian of maximal isotropic subspaces in \mathbb{C}^{10} endowed with a non-degenerate quadratic form. Denote by $\text{Gr}(m, n)$ the Grassmannian of m -dimensional subspaces in \mathbb{C}^n and let \mathcal{U}_m be the tautological subbundle. Finally, denote by $\text{IG}(m, 2n + 1)$ the Grassmannian of isotropic subspaces of dimension m in \mathbb{C}^{2n+1} endowed with an antisymmetric form of rank $2n$.

Proposition 1.16. *If $X = X^2$ or $X = X^3(n, m)$, then X is obtained as the zero locus of a vector bundle on a bigger homogeneous space \mathfrak{X} . More precisely*

1. The variety X^2 is a general hyperplane section of $\mathfrak{X} = \text{OG}(5, 10)$.
2. The variety $X^3(n, m)$ is the zero locus of a section of $\Lambda^2 \mathcal{U}_m^\vee$ in $\mathfrak{X} = \text{Gr}(m, 2n + 1)$.

Proof. 1. We use Mukai's classification of Fano varieties of large index (see [25]).

Since X is a Fano variety of dimension 9 and index 7, it has to be a hyperplane section of $\text{OG}(5, 10)$.

2. The variety X is isomorphic to $\text{IG}(m, 2n + 1)$, which is by definition the zero locus of a section of $\Lambda^2 \mathcal{U}_m^\vee$ in $\text{Gr}(m, 2n + 1)$.

\square

2. MODULI SPACE OF CURVES

Let X be as above a projective horospherical variety with $\text{Pic}(X) = \mathbb{Z}$. Let $d \in H_2(X, \mathbb{Z})$ and denote by $M_{d,k}(X)$ the moduli space of stable maps of genus 0, degree d and k marked points. Here the degree of a map $f : C \rightarrow X$ with C a nodal curve is by definition the push-forward $f_*[C]$. Identifying $H_2(X, \mathbb{Z})$ with \mathbb{Z} this moduli space has the expected dimension:

$$\dim X + \langle c_1(T_X), d \rangle + k - 3 = \dim X + dc_1(X) + k - 3.$$

Any irreducible component of $M_{d,k}(X)$ has dimension at least equal to the expected dimension. We refer to [30, 19, 38] for further properties of $M_{d,k}(X)$.

2.1. Curves with irreducible source. We first consider the open subset $M_{d,k}^\circ(X)$ of $M_{d,k}(X)$ consisting of stable maps with irreducible source. In this subsection, we describe the irreducible components of $M_{d,k}^\circ(X)$. We start with an useful lemma on maps to projective bundles. This result was proved for projective bundles of rank one in [49, Proposition 4] and the proof of the general result is very similar. We include this proof for the reader's convenience.

Lemma 2.1. *Let $p : M \rightarrow N$ be a projective bundle, i.e., $M = \mathbb{P}_N(E)$, where E a globally generated vector bundle over a projective variety. Let $\alpha \in H_2(M, \mathbb{Z})$ be such that $a = -\alpha \cap c_1(\mathcal{O}_p(-1)) \geq 0$ (here $\mathcal{O}_p(-1)$ is the tautological rank one subbundle associated to the projective bundle p).*

Then the map $M_{\alpha,k}^\circ(M) \rightarrow M_{p_\alpha,k}^\circ(N)$ induced by composition with p realises $M_{\alpha,k}^\circ(M)$ as a non empty open subset of a projective bundle of rank $p_*\alpha \cap c_1(E) + \text{rk}(E)(a+1) - 1$ over $M_{p_*\alpha,k}^\circ(N)$.*

Proof. The fiber of the map $M_{\alpha,k}^\circ(M) \rightarrow M_{p_*\alpha,k}^\circ(N)$ over $f : C \rightarrow N$ is given by sections of $\mathbb{P}_C(f^*E)$. Since C is rational and irreducible, we may assume that $C = \mathbb{P}^1$ and that these sections are injective maps of vector bundles $f^*\mathcal{O}_p(-1) = \mathcal{O}_C(-a) \rightarrow f^*E$ modulo scalars, where $a = -\alpha \cap c_1(\mathcal{O}_p(-1)) \geq 0$. The fiber is thus the subset of $\mathbb{P}\text{Hom}(\mathcal{O}_C(-a), f^*E)$ of injective maps of vector bundles. Since E is globally generated, so is f^*E . This implies that our sections form a non-empty open subset of this projective space.

This construction can be realised in families, which gives us the assertion. Indeed, for $C = \mathbb{P}^1$ let $f : C \times M_{p_*\alpha,k}^\circ(N) \rightarrow N$ be the universal map. We will need the projection $\text{pr} : C \times M_{p_*\alpha,k}^\circ(N) \rightarrow M_{p_*\alpha,k}^\circ(N)$. Consider the sheaf $\text{pr}_*(f^*E \otimes \mathcal{O}_C(a))$. Since E is globally generated and $a \geq 0$, we have $R^1\text{pr}_*(f^*E \otimes \mathcal{O}_C(a)) = 0$, hence $\text{pr}_*(f^*E \otimes \mathcal{O}_C(a))$ is a vector bundle of rank $\deg f^*E + \text{rk}(E)(a+1) = p_*\alpha \cap c_1(E) + \text{rk}(E)(a+1)$ and $M_{\alpha,k}^\circ(M)$ is an open subset of $\mathbb{P}_{M_{p_*\alpha,k}^\circ(N)}(\text{pr}_*(f^*E \otimes \mathcal{O}_C(a)))$. \square

We use this lemma to describe the irreducible components of $M_{d,k}^\circ(X)$.

Proposition 2.2. *Let X be as above. We have*

- (a) $M_{d,k}^\circ(X) = \overline{M_{d,k}^\circ(\mathcal{U}_Y)} \cup M_{d,k}^\circ(Z)$.
- (b) $\overline{M_{d,k}^\circ(\mathcal{U}_Y)}$ is irreducible of dimension $\dim X + dc_1(X) + k - 3$.
- (c) $M_{d,k}^\circ(Z)$ is irreducible of dimension $\dim Z + dc_1(Z) + k - 3$.

Proof. Recall from [55, 28, 49] that for a homogeneous space, the moduli space of stable maps is irreducible of the expected dimension, therefore $M_{d,k}^\circ(Z)$ is irreducible of dimension $\dim Z + dc_1(Z) + k - 3$ and $M_{d_Y,k}^\circ(Y)$ is irreducible of dimension $\dim Y +$

$d_Y c_1(Y) + k - 3$ for any non negative integer d_Y . This in turn imply that $M_{d,k}^\circ(\mathcal{U}_Y)$ and $\overline{M_{d,k}^\circ(\mathcal{U}_Y)}$ are irreducible of expected dimension. Indeed, in [49, Proposition 3] it is proved that given a globally generated vector bundle $\varphi : W \rightarrow W'$ such that $M_{d,k}^\circ(W')$ is irreducible of expected dimension, then so is $M_{d,k}^\circ(W)$. Since $p_Y : \mathcal{U}_Y \rightarrow Y$ is a globally generated vector bundle, the result follows.

Let $f : C \rightarrow X$ be a stable map, where C is irreducible and nodal. Up to normalising, we may assume that C is smooth, hence $C = \mathbb{P}^1$. If f factors through Z then we are in $M_{d,k}^\circ(Z)$. So we may assume that $f(C)$ meets \mathcal{U}_Y and need to prove that f is in the closure of $M_{d,k}^\circ(\mathcal{U}_Y)$. For this it is enough to prove that stable maps with irreducible source having a non-trivial intersection with \mathcal{U}_Y and Z form a family of dimension smaller than the expected dimension $\dim X + d c_1(X) + k - 3$.

For this we consider stable maps from C to \tilde{X}_Z , the blow-up of X in Z . The map f lifts to a map $\tilde{f} : C \rightarrow \tilde{X}_Z$. We compute its degree with respect to the basis $(\pi_Z^* \mathcal{O}_X(1), p_Y^* \mathcal{O}_Y(1))$ of $\text{Pic}(\tilde{X}_Z)$. We have $d = \deg f^* \mathcal{O}_X(1) = \deg \tilde{f}^* \pi_Z^* \mathcal{O}_X(1)$. Define $d_Y = \deg \tilde{f}^* p_Y^* \mathcal{O}_Y(1)$. Since $p_Y^* \mathcal{O}_Y(1)$ is semiample, we have $d_Y \geq 0$. Since $\mathcal{O}_{\tilde{X}_Z}(E) = \pi_Z^* \mathcal{O}_X(1) - p_Y^* \mathcal{O}_Y(1)$ and $\deg \tilde{f}^* \mathcal{O}_{\tilde{X}_Z}(E)$ is the intersection multiplicity of the curve \tilde{f} with Z , we have $d - d_Y \geq 0$.

Now we consider the open subset $M_{(d,d_Y),k}^\circ(\tilde{X}_Z)$ of stable maps with irreducible source, target \tilde{X}_Z and degree (d, d_Y) with respect to the basis $(\pi_Z^* \mathcal{O}_X(1), p_Y^* \mathcal{O}_Y(1))$ of $\text{Pic}(\tilde{X}_Z)$. We only need to assume that both d and d_Y are non-negative (this is always true for non-empty moduli spaces) and $d - d_Y \geq 0$. We have a natural map $M_{(d,d_Y),k}^\circ(\tilde{X}_Z) \rightarrow M_{d_Y,k}^\circ(Y)$ given by the composition with $p_Y : \tilde{X}_Z \rightarrow Y$. Note that p_Y is the projective bundle $\mathbb{P}_Y(\mathcal{O}_Y \oplus N_{Y/X})$ and that $N_{Y/X}$ is globally generated. We want to apply the previous lemma so we need to compute $a = -\alpha \cap c_1(\mathcal{O}_{p_Y}(-1))$ with $\alpha = (d, d_Y)$. But a map $\tilde{f} : C \rightarrow \tilde{X}_Z$ induces an injective map of vector bundles $\mathcal{O}_C(-a) = \tilde{f}^* \mathcal{O}_{p_Y}(-1) \rightarrow \mathcal{O}_C \oplus \tilde{f}^* N_{Y/X}$, and the intersection of $\tilde{f}(C)$ with E is the vanishing locus of the first factor. In particular $a = d - d_Y \geq 0$. Note that $c_1(N_{Y/X}) = (c_1(X) - c_1(Y))H_Y$, where H_Y is the ample generator of $\text{Pic}(Y)$. Therefore, applying the previous lemma we get that $M_{(d,d_Y),k}^\circ(\tilde{X}_Z)$ is a non-empty open subset of a projective bundle of rank $d_Y(c_1(X) - c_1(Y)) + \text{codim}_X Y(d - d_Y + 1)$ over $M_{d_Y,k}^\circ(Y)$. Therefore $M_{(d,d_Y),k}^\circ(\tilde{X}_Z)$ is irreducible of dimension

$$\dim M_{(d,d_Y),k}^\circ(\tilde{X}_Z) = \text{exp-dim } M_{d,k}(X) - (d - d_Y)(c_1(X) - \text{codim}_X Y).$$

In particular, since $c_1(X) - \text{codim}_X Y > 0$, this dimension is smaller than the expected dimension $\text{exp-dim } M_{d,k}(X)$ of $M_{d,k}(X)$, except for $d_Y = d$, thus for curves not meeting Z . This proves the result. \square

Corollary 2.3. *Let X be as above.*

If $\text{codim}_X Z > d(c_1(Z) - c_1(X))$, then $M_{d,k}^\circ(X)$ is irreducible of dimension $\dim X + d c_1(X) + k - 3$.

Otherwise, $M_{d,k}^\circ(X)$ has two irreducible components $\overline{M_{d,k}^\circ(\mathcal{U}_Y)}$ and $M_{d,k}^\circ(Z)$ of respective dimension $\dim X + d c_1(X) + k - 3$ and $\dim Z + d c_1(Z) + k - 3$.

Proof. This follows from the fact that any irreducible component of $M_{d,k}^\circ(X)$ is of dimension at least $\dim X + d c_1(X) + k - 3$. In the second case, $\overline{M_{d,k}^\circ(\mathcal{U}_Y)}$ will never be in the closure of $M_{d,k}^\circ(Z)$. \square

- Corollary 2.4.** (a) In all the cases of Pasquier's list, $M_{1,k}^\circ(X)$ is irreducible of dimension $\dim X + c_1(X) + k - 3$.
- (b) If $X = X^2$ or $X = X^3(n, m)$, then $M_{d,k}^\circ(X)$ is irreducible of dimension $\dim X + dc_1(X) + k - 3$ for all d .

2.2. Irreducible components. The former results enable us to give a description of the irreducible components of $M_{d,k}(X)$. Depending on the situation it is possible to have many of them. Here we only consider the simplest cases where there are few irreducible components.

- Theorem 2.5.** (a) If $X = X^2$ or $X = X^3(n, m)$, then $M_{d,k}(X)$ is irreducible of dimension $\dim X + dc_1(X) + k - 3$ for any d .
- (b) $M_{1,k}(X)$ is irreducible of dimension $\dim X + c_1(X) + k - 3$ for all X of Pasquier's list. With the exception of X^5 , we have the vanishing $H^1(C, f^*T_X) = 0$ for any stable map $f : C \rightarrow X$.
- (c) $M_{2,k}(X)$ is irreducible of dimension $\dim X + 2c_1(X) + k - 3$ if X is one of the varieties $X^1(3), X^2, X^3(n, m), X^4$.
- (d) $M_{2,k}(X)$ has two irreducible components of dimension $\dim X + 2c_1(X) + k - 3$ if $X = X^1(4)$ or $X = X^5$.
- (e) $M_{3,k}(X)$ has two irreducible components of dimension $\dim X + 3c_1(X) + k - 3$ if $X = X^1(3)$ or $X = X^4$.

Proof. Note that to prove the above results (except for the vanishing in (b)) we only need to prove the equality $M_{d,k}(X) = \overline{M_{d,k}^\circ(X)}$.

For $r \geq 2$, (d_1, \dots, d_r) with $d = \sum_i d_i$ a partition of d and $\prod_{i=1}^r A_i = [1, k]$ a partition, there is a natural map $M_{d_1, |A_1|+1}(X) \times_X \dots \times_X M_{d_r, |A_r|+1}(X) \rightarrow M_{d,k}(X)$. Furthermore the complement $M_{d,k}(X) \setminus M_{d,k}^\circ(X)$ is covered by the union of the images of these maps.

To prove the above results we only need to check that the boundary varieties

$$M_{d_1, |A_1|+1}(X) \times_X \dots \times_X M_{d_r, |A_r|+1}(X)$$

have dimension smaller than the expected dimension $\dim X + dc_1(X) + k - 3$. In all above cases, we have $\dim M_{d,k}^\circ(X) = \dim X + dc_1(X) + k - 3$. So an easy induction argument implies the result.

To prove the vanishing result in (b), we only need to prove that for a line L contained in Z , we have $H^1(L, N_{Z/X}|_L) = 0$. Since $N_{Z/X}$ is the vector bundle $G \times^{P_Z} V(\omega_Y - \omega_Z)$ and since we can choose the line to be the one dimensional Schubert variety in Z , it is enough to check that for any weight λ of $V(\omega_Y - \omega_Z)$ we have $\langle \alpha_Z^\vee, \lambda \rangle \geq -1$, where α_Z is the simple root dual to ω_Z . This is an easy check and is true in all cases except $X = X^5$. \square

- Remark 2.6.** 1. The non-obstruction result (b) above was first proved for the varieties $X^3(n, m)$ in [48]. This in particular implies that the corresponding stack is smooth and thus $M_{1,k}(X)$ has only quotient (and thus rational) singularities, see below for more details.
2. It is worth noting the following result from [38, Proposition 2]: Let X be a smooth projective complex algebraic variety such that $M_{d,k}(X)$ has the expected dimension, then the virtual class is the fundamental class. In particular, this is true in all cases of the previous theorem.

2.3. Singularities. In this section we prove regularity results for $M_{d,k}(X)$.

Theorem 2.7. (i) *With the exception of X^5 , for all X of Pasquier's list the moduli space $M_{1,k}(X)$ only has quotient singularities.*

Let X be either X^2 or $X^3(n, m)$.

- (ii) *If $f : C \rightarrow X$ is a stable map of genus 0 such that no component of C is mapped entirely into Z , then $M_{d,k}(X)$ has quotient singularity at $[f]$.*
- (iii) *The singular locus $\text{Sing}(M_{d,k}(X))$ has codimension at least 2.*
- (iv) *The moduli space $M_{d,k}(X)$ is normal and Cohen-Macaulay.*

Proof. (i) By Theorem 2.5 part (b), the corresponding stack is smooth and the result follows.

- (ii) It follows from [3, Lemma 2.1] that if X has a G -action with a dense orbit and $f : \mathbb{P}^1 \rightarrow X$ meets the dense orbit, then $H^1(\mathbb{P}^1, f^*T_X) = 0$. The result follows as in (i).
- (iii) It follows from (ii) that the singular locus of $M_{d,k}(X)$ lies inside the locus of curves which have a component contained in Z . But for any degree d the moduli space $M_{d,k}(Z)$ is irreducible of dimension $\dim(Z) + dc_1(Z) + k - 3$, which is of codimension at least 2 in $M_{d,k}(X)$. This proves the result.
- (iv) In view of (iii), to prove normality it suffices to show that $M_{d,k}(X)$ is Cohen-Macaulay. For this, recall that in the cases under consideration, X is the zero locus of a vector bundle on a projective homogeneous space $\mathfrak{X} \subset \mathbb{P}^N$ (see Proposition 1.16). That is, there exists a vector bundle E on \mathfrak{X} and a global section $s \in H^0(\mathfrak{X}, E)$ such that $X = V(s)$. Furthermore, note that E is globally generated. Let \mathfrak{Y} be the total space of the vector bundle E . Proceeding as in the proof of [49, Proposition 3] one checks that $M_{d,k}(\mathfrak{Y})$ is a vector bundle over $M_{d,k}(\mathfrak{X})$. Indeed, we have a morphism $\phi : M_{d,k}(\mathfrak{Y}) \rightarrow M_{d,k}(\mathfrak{X})$. Consider the following diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\text{ev}} & \mathfrak{X} \\ \downarrow \pi & & \\ M_{d,k}(\mathfrak{X}) & & \end{array}$$

where $\mathcal{C} = M_{d,k+1}(\mathfrak{X})$ is the universal curve, π is the map that forgets the $(k+1)$ -th marked point, and ev is the evaluation map. Since E is globally generated it follows that $R^1\pi_*\text{ev}^*E = 0$ and that $\pi_*\text{ev}^*E$ is locally free. The fiber of ϕ over $[f : C \rightarrow \mathfrak{X}] \in M_{d,k}(\mathfrak{X})$ is $H^0(C, f^*E)$ and thus $M_{d,k}(\mathfrak{X})$ is the bundle associated to $\pi_*\text{ev}^*E$ over $M_{d,k}(\mathfrak{X})$.

Furthermore, let f^*s denote the pullback section of f^*E . It is clear that $f : C \rightarrow \mathfrak{X}$ factors through $X = V(s)$ if and only if $s \circ f = 0$ or, equivalently, $f^*s \in H^0(C, f^*E)$ and $f^*s = 0$. The section s induces a section $\mathcal{O}_{M_{d,k}(\mathfrak{X})} \rightarrow \pi_*\mathcal{O}_{\mathcal{C}} \rightarrow \pi_*(f^*E)$ defining $M_{d,k}(X)$ as its vanishing locus: $M_{d,k}(X) = M_{d,k}(\mathfrak{X}) \cap V(\pi_*f^*s)$. Since $M_{d,k}(\mathfrak{X})$ has quotient singularities, it has rational singularities and hence it is Cohen-Macaulay. In view of [20, Lemma page 108], $M_{d,k}(X)$ is also Cohen-Macaulay. This fact together with (iii) finally implies that $M_{d,k}(X)$ is normal. \square

Question 2.8. Does $M_{d,k}(X)$ have rational singularities if $X = X^2$ or $X = X^3(n, m)$?

3. QUANTUM COHOMOLOGY

3.1. Reminders. For X a smooth projective complex variety with Picard rank one, we have k evaluation maps $\text{ev}_i : M_{d,k}(X) \rightarrow X$ on $M_{d,k}(X)$, sending a stable map to its value at the i -th marked point for $i \in [1, k]$. The Gromov-Witten invariants are defined as follows:

$$\langle \alpha_1, \dots, \alpha_k \rangle_{X,d,k} := \int_{[M_{d,k}(X)]^{\text{vir}}} \prod_{i=1}^k \text{ev}_i^* \alpha_i$$

for $\sum_i \deg \alpha_i = \dim X + dc_1(X) + k - 3$, and it is equal to zero otherwise. Here $\alpha_i \in H^*(X, \mathbb{R})$ are cohomology classes and $[M_{d,k}(X)]^{\text{vir}}$ is a homology class, the virtual fundamental class (see [2] for a definition). An important feature of that class is the equality $[M_{d,k}(X)]^{\text{vir}} = [M_{d,k}(X)]$ if $M_{d,k}(X)$ has expected dimension $\dim X + dc_1(X) + k - 3$ (see [38, Proposition 2]). The small quantum cohomology is defined as follows. Let $(e_i)_i$ be a basis of $H^*(X, \mathbb{R})$ and let $(e_i^\vee)_i$ be the dual basis for the Poincaré pairing. Then

$$e_i * e_j = \sum_{d \geq 0} \sum_{\ell} \langle e_i, e_j, e_\ell^\vee \rangle_{X,d,3} e_\ell q^d.$$

It is a non trivial result that this product defines an associative ring (see for example [19]). When the moduli space has expected dimension, the Gromov-Witten invariants are enumerative. We make this precise in the next subsection.

Remark 3.1. The varieties Y and Z are homogeneous for the group G . The quantum cohomology algebra of homogeneous spaces has been extensively studied and is well understood in many cases. Of special interest for our explicit computations in Section 4 are minuscule, cominuscule, adjoint and coadjoint varieties. The quantum cohomology of these varieties were studied in [10, 9, 12, 11, 13]. Many of these varieties are isotropic Grassmannians and their quantum cohomology is well understood thanks to the work of Buch, Kresch and Tamvakis [6, 31, 32, 7, 8]. We refer to these papers for explicit computations.

3.2. Enumerability of Gromov-Witten invariants. In this section we use a generalization of the Kleiman-Bertini Transversality Theorem to quasi-homogeneous spaces due to Graber [23, Lemma 2.5].

Lemma 3.2. *Let X be a variety endowed with an action of a connected algebraic group G with only finitely many orbits and Z be an irreducible scheme with a morphism $f : Z \rightarrow X$. Let Y be a subvariety of X that intersects the orbit stratification properly (i.e., for any G -orbit \mathcal{O} of X , we have $\text{codim}_{\mathcal{O}}(Y \cap \mathcal{O}) = \text{codim}_X Y$). Then there exists a dense open subset U of G such that for every $g \in U$, $f^{-1}(gY)$ is either empty or has pure dimension $\dim Y + \dim Z - \dim X$. Moreover, if X , Y and Z are smooth and we denote by Y_{reg} the subset of Y along which the intersection with the stratification is transverse, then the (possibly empty) open subset $f^{-1}(gY_{\text{reg}})$ is smooth.*

We apply the previous lemma to the following situation.

Theorem 3.3. *Let X be a G -variety with finitely many G -orbits and dense G -orbit \mathcal{U} . Let $d \in H_2(X, \mathbb{Z})$ be such that $M_{d,k}(X)$ is irreducible and equals the closure of the*

locus of curves contained in \mathcal{U} . Take Y_1, \dots, Y_k to be subvarieties of X that intersect the orbit stratification properly and that represent cohomology classes $\Gamma_1, \dots, \Gamma_k$ satisfying

$$\sum_i \operatorname{codim}_X Y_i = \dim M_{d,k}(X).$$

Then there is a dense open subset $U \subset G^k$ such that for all $(g_1, \dots, g_k) \in U$, the Gromov-Witten invariant $\langle \Gamma_1, \dots, \Gamma_k \rangle_{X,d,k}$ is equal to the number of stable curves of degree d in X incident to the translates $g_1 Y_1, \dots, g_k Y_k$.

Proof. First, we shall prove by repeatedly applying Lemma 3.2, that for generic (g_1, \dots, g_k) in G^k , the scheme-theoretic intersection

$$\operatorname{ev}^{-1}(g_1 Y_1 \times \dots \times g_k Y_k) = \bigcap_{i=1}^k \operatorname{ev}_i^{-1}(g_i Y_i)$$

consists of a finite number of reduced points supported on any preassigned nonempty open subset M^* of $M = M_{d,k}(X)$ and, in particular, in the smooth locus of M . Here $\operatorname{ev} = \operatorname{ev}_1 \times \dots \times \operatorname{ev}_k$. Bearing this in mind, consider the following diagram

$$\begin{array}{ccc} & M \setminus M^* & \\ & \downarrow \operatorname{ev} & \\ \underline{Y} = Y_1 \times \dots \times Y_k \subset & \longrightarrow & X^k. \end{array}$$

Since M is irreducible, $M \setminus M^*$ has codimension at least one, so Lemma 3.2 implies that there exists an open subset V_1 of G^k such that the inverse image of any of the translates $g \cdot \underline{Y}$ with $g \in V_1$, is empty. Thus in general the intersection is completely supported in M^* .

Next let \mathcal{U} be the dense G -orbit of X . By the above, we can restrict to the locus where ev is in \mathcal{U}^k . Now let M^{sm} be the smooth locus of $M_{d,k}(X)$ and $\underline{Y}^s := \operatorname{Sing} \underline{Y}$ and consider the diagram

$$\begin{array}{ccc} & M^{sm} \cap \operatorname{ev}^{-1}(\mathcal{U}^k) & \\ & \downarrow \operatorname{ev} & \\ \underline{Y}^s \cap \mathcal{U}^k \subset & \longrightarrow & \mathcal{U}^k. \end{array}$$

Applying Kleiman's transversality theorem to the homogeneous space \mathcal{U} , we get an open subset $V_2 \subset G^k$ such that $\operatorname{ev}^{-1}(g \cdot \underline{Y}^s \cap \mathcal{U}^k) = \emptyset$, for any $g \in V_2$. Now let $\underline{Y}^* = (\underline{Y} \setminus \underline{Y}^s) \cap \mathcal{U}^k$. By yet another application of Kleiman's transversality theorem to the diagram

$$\begin{array}{ccc} & M^{sm} \cap \operatorname{ev}^{-1}(\mathcal{U}^k) & \\ & \downarrow \operatorname{ev} & \\ \underline{Y}^* \subset & \longrightarrow & \mathcal{U}^k. \end{array}$$

we get an open subset $V_3 \subset G^k$ such that for all $g \in V_3$, the inverse image in $M^{sm} \cap \operatorname{ev}^{-1}(\mathcal{U}^k)$ of the translate $g \cdot \underline{Y}^*$ is either empty or smooth of the expected dimension (since all varieties involved are smooth). Hence it consists of a finite number of reduced points (possibly zero). We conclude that for any $g \in V_1 \cap V_2 \cap V_3$,

the inverse image of $g \cdot \underline{Y}$ is of the expected dimension, is reduced, and is supported in the given open set. \square

Next we specify the previous result to cases (2) and (3) of Pasquier's list.

Corollary 3.4. *Let X be a horospherical variety of Picard number one. If X belongs to the classes (2) or (3) of Pasquier's list, then Theorem 3.3 holds, i.e., the Gromov-Witten invariants of X are enumerative.*

Remark 3.5. The previous corollary gives a positive answer to a conjecture from [47, Conjecture 5.1] regarding the enumerativity of the degree d Gromov-Witten invariants of the odd symplectic Grassmannians. The case when $d = 1$ was already proven by the second author in [47].

3.3. Quantum hyperplane multiplication. In this section we consider the multiplication with the hyperplane generator $h \in H^2(X, \mathbb{Z})$. Our aim is in particular to deal with the varieties $X^1(3)$, X^2 , $X^3(3, 3)$ and X^5 of Pasquier's list, but we keep the computations as general as possible. Let us start with an easy dimension count.

Lemma 3.6. *Let $\sigma \in H^*(X, \mathbb{Z})$. The highest power of q appearing in a product $h * \sigma$ in $\text{QH}(X)$ satisfies $d \leq \frac{\dim X + 1}{c_1(X)}$. Furthermore, for $X^1(3)$ and X^5 , there is a non-trivial term of degree 2 in the product $h * \sigma$ only if $\sigma = [\text{pt}]$, where $[\text{pt}]$ is the class of a point. Moreover, this term is a multiple of q^2 .*

Proof. The degree of $h * \sigma$ is at most $\dim X + 1$. If there is a q^d appearing in the product, then its degree is at least $dc_1(X)$ leading to the inequality $dc_1(X) \leq \dim X + 1$. The result follows. \square

Remark 3.7. We shall prove in Section 5 that for $X^3(n, m)$, the product $h * \sigma$ has terms of degree at most 1 in q .

3.4. Degree one invariants. In this subsection, we compute the degree one invariants involved in the quantum multiplication with h . We therefore compute Gromov-Witten invariants of the form $\langle h, \sigma, \tau \rangle_{X,1,3}$ with $\sigma, \tau \in H^*(X, \mathbb{Z})$ such that $\deg \sigma + \deg \tau + 1 = \dim M_{1,3}(X) = \dim X + c_1(X)$.

Lemma 3.8. *Let $u_1, u_2 \in W^{Py}$. We have $\langle h, \sigma'_{u_1}, \sigma_{u_2} \rangle_{X,1,3} = 0$.*

Proof. We may assume $\deg \sigma'_{u_1} + \deg \sigma_{u_2} + 1 = \dim M_{1,3}(X) = \dim X + c_1(X)$, which leads to $\ell(u_1) + \ell(u_2) = \dim Y + c_1(X) - 1$. We have $\langle h, \sigma'_{u_1}, \sigma_{u_2} \rangle_{X,1,3} = \pi_{M*}(\text{ev}_1^* h \cup \text{ev}_2^* \sigma'_{u_1} \cup \text{ev}_3^* \sigma_{u_2})$, where π_{M*} denotes the proper pushforward to a point, or integration against the virtual class of the moduli space of stable maps. Choosing general translates H , $g_1 \cdot Y'_{u_1}$ and $g_2 \cdot Y_{u_2}$ with $g_1, g_2 \in G$, we have $\text{ev}_1^* h \cup \text{ev}_2^* \sigma'_{u_1} \cup \text{ev}_3^* \sigma_{u_2} = \text{ev}_1^{-1}(H) \cap \text{ev}_2^{-1}(g_1 \cdot Y'_{u_1}) \cap \text{ev}_3^{-1}(g_2 \cdot Y_{u_2})$. We prove that this set is empty. It is enough to prove that there is no line meeting H , $g_1 \cdot Y'_{u_1}$ and $g_2 \cdot Y_{u_2}$.

We discuss several cases. Let L be such a line. If L were in Z then L would not meet $g_2 \cdot Y_{u_2}$, a contradiction. If L meets Z (necessarily in one point since $Z = X \cap \mathbb{P}(V_Z)$), then $p_Y(\pi_Z^{-1}(L))$ is a point and lies on $p_Y(\pi_Z^{-1}(g_1 \cdot Y'_{u_1})) \cap p_Y(\pi_Z^{-1}(g_2 \cdot Y_{u_2}))$. But $p_Y(\pi_Z^{-1}(g_1 \cdot Y'_{u_1}))$ is a translate of Y_{u_1} and $p_Y(\pi_Z^{-1}(g_2 \cdot Y_{u_2}))$ is a general translate of Y_{u_2} . An easy dimension count gives a contradiction, for $\ell(u_1) + \ell(u_2) > \dim Y$. Finally if L does not meet Z , then $p_Y(\pi_Z^{-1}(L))$ is a line meeting $p_Y(\pi_Z^{-1}(g_1 \cdot Y'_{u_1}))$ and $p_Y(\pi_Z^{-1}(g_2 \cdot Y_{u_2}))$. This is possible if $\ell(u_1) + \ell(u_2) \leq \dim M_{1,2}(Y) = \dim Y + c_1(Y) - 1$. This is a contradiction since $c_1(X) > c_1(Y)$. \square

Lemma 3.9. *Let $u \in W^{Py}$ and $v \in W^{Pz}$. We have $\langle h, \sigma_u, \tau_v \rangle_{X,1,3} = \delta_{\tilde{u}^\vee, \tilde{v}}$.*

Proof. As in the previous lemma, we may assume $\ell(u) + \ell(v) = \dim Y + \dim Z + c_1(X) - \dim X - 1$ and we are looking for lines L meeting general translates $H, g_1.Y_u$ and $g_2.Z_v$. In particular, the line L is not contained in Z (otherwise $L \cap g_1.Y_u = \emptyset$) and not in Y (otherwise $L \cap g_2.Z_v = \emptyset$) but meets Y in $g_1.Y_u$ and Z in $g_2.Z_v$. The line L is thus mapped to a point via p_{YZ} i.e $p_{YZ}(\pi_{YZ}^{-1}(L))$ is a point. Thus $p_{YZ}(\pi_{YZ}^{-1}(g_1.Y_u))$ a general translate of $E_{\tilde{u}}$ and $p_{YZ}(\pi_{YZ}^{-1}(g_2.Z_v))$ a general translate of $E_{\tilde{v}}$ have to meet. The sum of their codimensions is $\ell(u) + \ell(v) = \dim E + c_1(X) - \text{codim}_X Y - \text{codim}_X Z$ and by Fact 1.7 we get $\ell(u) + \ell(v) = \dim E$. In particular this intersection is empty unless $\tilde{v} = \tilde{u}^\vee$. If this intersection is non-empty and therefore a point e , the line is given by $\pi_{YZ}(\pi_{YZ}^{-1}(e))$. \square

Lemma 3.10. *Let $u \in W^{Py}$ and $v \in W^{Pz}$. We have*

$$\langle h, \sigma'_u, \tau'_v \rangle_{X,1,3} = \begin{cases} 0 & \text{if } c_1(X) \geq c_1(Z) \\ \langle h, \bar{\tau}_{\hat{u}}, \bar{\tau}_v \rangle_{Z,1,3} & \text{if } c_1(X) = c_1(Z) - 1 \end{cases}$$

Remark 3.11. Note that $c_1(X) \geq c_1(Z)$ for X^2 and $X^3(n, m)$, while $c_1(X) = c_1(Z) - 1$ for $X^1(3), X^4$ and X^5 .

Proof. As above, we may assume $\ell(u) + \ell(v) = \dim X + c_1(X) - 1$ and we are looking for lines L meeting general translates $H, g_1.Y'_u$ and $g_2.Z'_v$. If $L \subset Y$, then L meets general translates of $H \cap Y, Y_u$ and $Y_{\hat{v}}$. The sum of their codimension is $\ell(u) + \ell(v) + 1 - \text{codim}_X Y + 1 = \dim Y + c_1(X) + 1 > \dim M_{1,3}(Y)$, so there is no such L . If L is contained in \mathcal{U} , then $p_{YZ}(\pi_{YZ}^{-1}(L))$ is a line meeting general translates of $E_{\tilde{u}}$ and $E_{\tilde{v}}$. The sum of their codimensions is $\ell(u) + \ell(v) = \dim X - 1 + c_1(X) = \dim E + c_1(X)$. But the curve $p_{YZ}(\pi_{YZ}^{-1}(L))$ has intersection $c_1(X)$ with $-K_E$, so this sum is one more than the dimension of the moduli space, which implies that there is no such curve. Assume that L meets \mathcal{U} . If the line L meets Z then $p_Y(\pi_Z^{-1}(L))$ is a point meeting general translates of Y_u and $Y_{\hat{v}}$. An easy dimension count proves that this is not possible. The same arguments prove that there is no line L meeting \mathcal{U} and Y .

We thus have the inclusion $L \subset Z$ and since $Y'_u \cap Z = Z_{\hat{u}}$, the number of lines L as above is given by the Gromov-Witten invariant $\langle h, \bar{\tau}_{\hat{u}}, \bar{\tau}_v \rangle_{Z,1,3}$. Again by a dimension count the result follows. \square

Lemma 3.12. *Let $v_1, v_2 \in W^{Pz}$. We have*

$$\langle h, \tau_{v_1}, \tau'_{v_2} \rangle_{X,1,3} = \begin{cases} 0 & \text{if } c_1(X) > c_1(Z) \\ \langle h, \bar{\tau}_{v_1}, \bar{\tau}_{v_2} \rangle_{Z,1,3} & \text{if } c_1(X) = c_1(Z) \end{cases}$$

Remark 3.13. Note that $c_1(X) > c_1(Z)$ for X^2 , while $c_1(X) = c_1(Z)$ for $X^3(n, m)$.

Proof. As above, we may assume $\ell(u) + \ell(v) = \dim Z + c_1(X) - 1$ and we are looking for lines L meeting general translates $H, g_1.Z_{v_1}$ and $g_2.Z'_{v_2}$. An easy dimension count using the composition $p_Z \circ \pi_Y$ proves that L does not meet Y . Projecting to Z we have that $p_Z(\pi_Y^{-1}(L))$ is a line L' which meets general translates of Z_{v_1} and Z_{v_2} . Such a line L' does not exist for dimension reasons if $c_1(X) > c_1(Z)$, and there are $\langle h, \bar{\tau}_{v_1}, \bar{\tau}_{v_2} \rangle_{Z,1,3}$ such lines if $c_1(X) = c_1(Z)$. Since g_1 and g_2 are general in G we may assume that these lines L' are general. We may thus assume that $c_1(X) = c_1(Z)$. We are thus in case (3) of Pasquier's classification and any lift of L' in \mathcal{U}_Z meeting the zero section is a solution for L . But \mathcal{U}_Z is a vector bundle over Z whose restriction

to a general line is trivial. In particular the only lift meeting the zero section is the zero section itself, proving the result. \square

Lemma 3.14. *Let $u \in W^{P_Y}$ and $v \in W^{P_Z}$. For $c_1(X) - c_1(Y) + 1 - \text{codim}_X Y = 0$, we have $\langle h, \sigma_u, \tau'_v \rangle_{X,1,3} = \langle h, \bar{\sigma}_u, \bar{\sigma}_{\hat{v}} \rangle_{Y,1,3}$.*

Remark 3.15. Note that we have $c_1(X) - c_1(Y) + 1 - \text{codim}_X Y \leq 0$ in all cases, with equality for $X = X^1(n), X^3(n, 2)$ and X^5 .

Proof. As above, we may assume $\ell(u) + \text{codim}_X Y + \ell(v) = \dim X + c_1(X) - 1$ and we are looking for lines L meeting general translates $H, g_1.Y_u$ and $g_2.Z'_v$. If L is not contained in Y , then $p_Z(\pi_Y^{-1}(L))$ is a point meeting general translates of $Z_{\hat{u}}$ and Z_v . An easy dimension count proves that this is not possible. If $L \subset Y$, then we are counting curves in Y meeting a hyperplane and general translates of Y_u and $Y_{\hat{v}}$. The condition implies that the classes have the right codimension, which implies the result. \square

Lemma 3.16. *Let $v_1, v_2 \in W^{P_Z}$, we have*

$$\langle h, \tau'_{v_1}, \tau'_{v_2} \rangle_{X,1,3} = \begin{cases} 0 & \text{if } c_1(X) - c_1(Y) > \text{codim}_X Y - 2 \\ \langle h, \bar{\sigma}_{\hat{v}_1}, \bar{\sigma}_{\hat{v}_2} \rangle_{Y,1,3} & \text{if } c_1(X) - c_1(Y) = \text{codim}_X Y - 2. \end{cases}$$

Remark 3.17. Note that we have $c_1(X) - c_1(Y) + 2 - \text{codim}_X Y > 0$ for $X = X^1(n), X^2, X^3(n, 2)$ and X^5 , while $c_1(X) - c_1(Y) + 2 - \text{codim}_X Y = 0$ for $X = X^3(n, 3)$ and $X = X^4$.

Proof. As above, we may assume $\ell(u) + \ell(v) + 1 = \dim X + c_1(X)$ and we are looking for lines L meeting general translates $H, g_1.Z'_{v_1}$ and $g_2.Z'_{v_2}$. If L is contained in Z , then we are computing $\langle h_Z, \bar{\tau}_{v_1}, \bar{\tau}_{v_2} \rangle_{Z,1,3}$. But $\ell(v_1) + \ell(v_2) + 1 = \dim X + c_1(X) > \dim Z + c_1(Z)$ so there is no such L . If L meets Z , then $p_Y(\pi_Z^{-1}(L))$ is a point contained in the intersection of general translates of $p_Y(\pi_Z^{-1}(Z'_{v_1})) = Y_{\hat{v}_1}$ and $p_Y(\pi_Z^{-1}(Z'_{v_2})) = Y_{\hat{v}_2}$. But the sum of their codimensions is $\ell(v_1) + 1 - \text{codim}_X Y + \ell(v_2) + 1 - \text{codim}_X Y = \dim Y + c_1(X) + 1 - \text{codim}_X Y = \dim Y + 1 + \text{codim}_X Z > \dim Y$ so there is no such L . If L meets Y but is not contained in Y , then $p_Z(\pi_Y^{-1}(L))$ is a point meeting general translates of Z_{v_1} and Z_{v_2} . An easy dimension count proves that this is not possible. If L is contained in \mathcal{U} , we consider $L' = p_{YZ}\pi_Y^{-1}(L)$ meeting general translates of $p_{YZ}\pi_Y^{-1}(Z'_{v_1}) = E_{\hat{v}_1}$ and $p_{YZ}\pi_Y^{-1}(Z'_{v_2}) = E_{\hat{v}_2}$. The sum of their codimension in E is $\ell(v_1) + \ell(v_2) = \dim X - 1 + c_1(X) = \dim E + c_1(X)$. The curve L' has intersection $c_1(X)$ with $-K_E$, thus there is no such line. We are left with L contained in Y . So we are computing the number of lines corresponding to the invariant $\langle h_Y, \bar{\sigma}_{\hat{v}_1}, \bar{\sigma}_{\hat{v}_2} \rangle_{Y,1,3}$. The sum of the corresponding codimensions is $\ell(v_1) + 1 - \text{codim}_X Y + \ell(v_2) + 1 - \text{codim}_X Y + 1 = \dim Y + c_1(X) + 2 - \text{codim}_X Y = \dim Y + c_1(Y) + c_1(X) - c_1(Y) + 2 - \text{codim}_X Y$. The result follows. \square

3.5. Degree 2 invariants. In this subsection we compute some degree 2 Gromov-Witten invariants. We concentrate on the case $\dim X = 2c_1(X) - 1$, where there is a unique non-vanishing degree 2 invariant, $\langle h, [\text{pt}], [\text{pt}] \rangle_{X,2,3}$.

Lemma 3.18. *If $\dim X = 2c_1(X) - 1$, then $\langle h, [\text{pt}], [\text{pt}] \rangle_{X,2,3} = 2$.*

Remark 3.19. Note that we have $\dim X = 2c_1(X) - 1$ for $X = X^1(3)$ and $X = X^5$.

Proof. Recall that $\langle h, [\text{pt}], [\text{pt}] \rangle_{X,2,3} = 2\langle [\text{pt}], [\text{pt}] \rangle_{X,2,2}$ so we need to prove that there is a unique conic through two general points in X . Consider two general points p_1

and p_2 in \mathcal{U}_Y . Any stable maps of degree 2 passing through p_1, p_2 give by projection p_Y to Y a stable map of degree at most 2 passing through two general points in Y . With our assumptions we have $\dim Y = 2c_1(Y) - 1$ and there exists exactly one (irreducible) stable map $f : \mathbb{P}^1 \rightarrow Y$ of degree 2 through 2 general points in Y (in our case, Y is an adjoint variety, so this follows from [13, Proposition 10]). We need to prove that f lifts to a unique stable map in \mathcal{U}_Y passing through p_1 and p_2 . This is equivalent to proving that the vector bundle N_Y satisfies $f^*N_Y = \mathcal{O}_{\mathbb{P}^1}(1)^{\text{codim}_X Y}$. The unique irreducible curve passing through the B -stable and the B^+ -stable points in Y is the curve obtained using the cocharacter θ^\vee with θ the highest root of G . Using the fact that $N_Y = p_*\mathcal{L}_{\omega_Y - \omega_Z} = G \times^{P_Y} V_{P_Y}(\omega_Y - \omega_Z)$ it is easy to check that the condition $f^*N_Y = \mathcal{O}_{\mathbb{P}^1}(1)^{\text{codim}_X Y}$ is satisfied. \square

4. EXAMPLES

In this section we compute the quantum hyperplane multiplication for the following examples: $X^1(3), X^2, X^3(3, 3)$ and X^5 . Our aim is to extend the Hasse diagram to a quantum Hasse diagram and deduce semisimplicity results for the quantum cohomology.

4.1. Semisimplicity. Using our quantum Chevalley formulas, we obtain

Theorem 4.1. *If X is one of the varieties $X^1(3), X^2, X^3(3, 3), X^5$, then $\text{QH}(X)$ is semisimple.*

Proof. The proof in all cases goes as follows. We work in the specialisation $q = 1$ of the quantum cohomology. If the algebra obtained for $q = 1$ is semisimple, so is $\text{QH}(X)$ (this is actually an equivalence).

Using the previous section we give in Propositions 4.3, 4.4, 4.5 and 4.6 Chevalley formulas for the quantum multiplication by the hyperplane class h in the above cases.

Using these Chevalley formulas, we can compute the quantum multiplication by $h^{c_1(X)}$ which is an endomorphism of the space

$$\text{QH}^0(X) := \bigoplus_{k \geq 0} \text{H}^{2kc_1(X)}(X, \mathbb{Z}).$$

Writing down the matrix of this endomorphism, which is a matrix of size $qh^0(X) = \dim \text{QH}^0(X)$ we can check in all cases (see below) that it has $qh^0(X)$ different non-vanishing eigenvalues. Since $1 \in \text{QH}^0(X)$, if $\mu(X)$ is the minimal polynomial $h^{c_1(X)}$ on $\text{QH}^0(X)$, then $\mu(X^{c_1(X)})$ is a vanishing polynomial for h on $\text{QH}(X)$ with different (non-vanishing) eigenvalues. Since in all cases, an easy check gives the equality $\dim \text{QH}(X) = c_1(X) \cdot \dim \text{QH}^0(X)$, we see that $\mu(X^{c_1(X)})$ is the minimal and the characteristic polynomial of h on $\text{QH}(X)$. This implies that the endomorphism of $\text{QH}(X)$ defined by multiplication with h is semisimple with simple (non-vanishing) eigenvalues, therefore it generates the algebra which has to be semisimple.

We now explicitly compute the matrix for X^2 and check that it has $qh^0(X)$ different non-vanishing eigenvalues. In the other cases, we only give the matrices obtained from Chevalley formulas. Recall that for $X = X^2$, we have $\dim X = 9$ and $c_1(X) = 7$. Set $\text{QH}^{2i}(X) := \bigoplus_{k \geq 0} \text{H}^{2kc_1(X)+2i}(X, \mathbb{Z})$. We have the following basis:

$$\begin{aligned} \text{QH}^0(X) &= \langle 1, \tau_{v_4} \rangle, & \text{QH}^2(X) &= \langle h, \tau_{v_5} \rangle, & \text{QH}^4(X) &= \langle \sigma'_{u_2}, \tau_{v_6} \rangle, \\ \text{QH}^6(X) &= \langle \sigma'_{u_3}, \tau_{v_0} \rangle, & \text{QH}^8(X) &= \langle \sigma'_{u_4}, \tau_{v_1} \rangle, & \text{QH}^{10}(X) &= \langle \sigma'_{u_5}, \tau_{v_2} \rangle, \\ \text{QH}^{12}(X) &= \langle \tau_{v_3}, \tau_{v'_3} \rangle. \end{aligned}$$

In particular we have $qh^0(X) = 2$ and we see that $\dim \text{QH}(X) = c_1(X) \cdot qh^0(X) = 7 \cdot 2 = 14$. Let A_i be the matrix, in the above basis, of the endomorphism $\text{QH}^{2i}(X) \rightarrow \text{QH}^{2i+2}(X)$ induced by the quantum multiplication by h . By Proposition 4.4, we have

$$A_0 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, A_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, A_2 = \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}, A_3 = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix},$$

$$A_4 = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, A_5 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, A_6 = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

The matrix of $h^{c_1(X)}$ on $\text{QH}^0(X)$ is the product $A_6 A_5 A_4 A_3 A_2 A_1 A_0$. We obtain:

$$\begin{pmatrix} 5 & 12 \\ 12 & 29 \end{pmatrix}$$

which has two distinct non vanishing eigenvalues and minimal polynomial $\mu(X) = x^2 - 34x + 1$.

If X is one of the varieties $X^1(3), X^2, X^3(3, 3), X^5$, the matrices of $h^{c_1(X)}$, in an appropriate basis, are the following ones:

$$\begin{pmatrix} 18 & 20 & 15 & 38 \\ 20 & 35 & 23 & 48 \\ 15 & 33 & 20 & 38 \\ 0 & 10 & 6 & 4 \end{pmatrix}, \begin{pmatrix} 16 & 20 & 63 & 34 \\ 20 & 22 & 69 & 38 \\ 5 & 9 & 26 & 18 \\ 7 & 6 & 19 & 9 \end{pmatrix}, \begin{pmatrix} 18 & 14 & 38 \\ 10 & 10 & 22 \\ 0 & 4 & 4 \end{pmatrix}.$$

Again one can check that they have distinct non vanishing eigenvalues. \square

Remark 4.2. (1) The semisimplicity for $X = X^2$ can also be obtained as follows. It follows from Proposition 1.16 that X is a general linear section of $\text{OG}(5, 10)$ but it was proved in [51], that the quantum cohomology of such hyperplane sections is semisimple.

(2) In the next section, we study more closely the quantum cohomology for $X = X^3(n, m)$ and prove that there is no degree 2 term in the quantum multiplication by h . We also give a presentation of $\text{QH}(X)$ in some cases. For similar varieties with non semisimple small quantum cohomology but semisimple big quantum cohomology, we refer to [51], [21] and [16].

(3) The semisimplicity for $X = X^5$ can also be obtained as follows. In [46], the authors proved that X has a deformation isomorphic to the isotropic Grassmannian $\text{OG}(2, 7)$ and in [13] it was proved that $\text{QH}(\text{OG}(2, 7))$ is semisimple. Since the quantum cohomology is a deformation invariant, this proves the result.

4.2. Case $X = X^1(3)$. We have $\dim X = 9$ and $c_1(X) = 5$. Z is a 6-dimensional quadric. There is only one cohomology class for each degree in Z except in degree 3 so we denote by $(v_i)_{i \in [0, 6]}$ with $\ell(v_i) = i$ and v'_3 with $\ell(v'_3) = 3$ the elements of W^{PZ} . We choose to set $v_3 = s_1 s_2 s_3$ and $v'_3 = s_3 s_2 s_3$. We have $Y = \text{OG}(2, 7)$ and there is one cohomology class in degree 0, 1, 6 and 7 and two cohomology classes otherwise. We denote by $(u_i)_{i \in [0, 7]}$ with $\ell(u_i) = i$ and u'_2, u'_3, u'_4, u'_5 with $\ell(u'_i) = i$ the elements in W^{PY} . We choose to set $u'_2 = s_1 s_2$, $u'_3 = s_3 s_1 s_2$, $u'_4 = s_2 s_3 s_1 s_2$, $u'_5 = s_3 s_2 s_3 s_1 s_2$. Note that $\sigma'_{u_0} = \tau'_{v_0} = 1$, $\sigma'_{u_1} = \tau'_{v_1} = h$, $\sigma_{u_7} = \tau_{v_6} = [\text{pt}]$ and $\sigma_{u_6} = \tau_{v_5} = [\text{line}]$.

Proposition 4.3. *We have the formulas*

(1) $h * 1 = h$

- (2) $h * h = 2\sigma'_{u_2} + \sigma'_{u'_2}$
- (3) $h * \sigma'_{u_2} = \sigma'_{u_3} + \sigma'_{u'_3}$ and $h * \sigma'_{u'_2} = 2\sigma'_{u'_3} + \tau_{v_0}$
- (4) $h * \sigma'_{u_3} = 2\sigma'_{u_4} + \sigma'_{u'_4}$, $h * \sigma'_{u'_3} = \sigma'_{u_4} + 2\sigma'_{u'_4} + \tau_{v_1}$ and $h * \tau_{v_0} = \tau_{v_1}$
- (5) $h * \sigma'_{u_4} = \sigma'_{u_5}$, $h * \sigma'_{u'_4} = \sigma'_{u_5} + 2\sigma'_{u'_5} + \tau_{v_2}$ and $h * \tau_{v_1} = \tau_{v_2} + q$
- (6) $h * \sigma'_{u_5} = 2\sigma'_{u_6} + \tau_{v_3}$, $h * \sigma'_{u'_5} = \sigma'_{u_6} + \tau_{v'_3}$ and $h * \tau_{v_2} = \tau_{v_3} + \tau_{v'_3} + qh$
- (7) $h * \sigma'_{u_6} = \sigma'_{u_7} + \tau_{v_4}$, $h * \tau_{v_3} = \tau_{v_4} + q\sigma'_{u'_2}$ and $h * \tau_{v'_3} = \tau_{v_4} + q\sigma'_{u_2}$
- (8) $h * \sigma'_{u_7} = \tau_{v_5} + q\tau_{v_0}$ and $h * \tau_{v_4} = \tau_{v_5} + q\sigma'_{u'_3}$
- (9) $h * \tau_{v_5} = \tau_{v_6} + q\sigma'_{u'_4} + q\tau_{v_1}$
- (10) $h * \tau_{v_6} = q\sigma'_{u'_5} + q\tau_{v_2} + 2q^2$.

Proof. We will only discuss the quantum corrections, since the classical part follows from the classical Hasse diagram. By Lemma 3.6, there is no power of q higher than 1 except maybe in $h * \tau_{v_6}$. For the first four lines, there is no quantum correction since the degree is smaller than $5 = c_1(X)$. For the fifth line, we only have to compute $\langle h, \sigma'_{u_4}, \sigma_{u_7} \rangle$, $\langle h, \sigma'_{u'_4}, \sigma_{u_7} \rangle$ and $\langle h, \tau_{v_1}, \sigma_{u_7} \rangle$. The result follows from Lemma 3.8 and Lemma 3.9. For line (6), we only have to compute $\langle h, \sigma'_{u_5}, \sigma_{u_6} \rangle$, $\langle h, \sigma'_{u'_5}, \sigma_{u_6} \rangle$ and $\langle h, \tau_{v_2}, \sigma_{u_6} \rangle$. The result follows from Lemma 3.8 and Lemma 3.9. For line (7), we only have to compute $\langle h, \sigma'_{u_6}, \sigma_{u_5} \rangle$, $\langle h, \tau_{v_3}, \sigma_{u_5} \rangle$ and $\langle h, \tau_{v'_3}, \sigma_{u_5} \rangle$. The result follows from Lemma 3.8 and Lemma 3.9.

Line (8) requires more work. We need to compute the Gromov-Witten invariants $\langle h, \sigma'_{u_7}, \sigma_{u_4} \rangle$, $\langle h, \tau_{v_4}, \sigma_{u_4} \rangle$, $\langle h, \sigma'_{u_7}, \tau'_{v_6} \rangle$ and $\langle h, \tau_{v_4}, \tau'_{v_6} \rangle$. The first three are obtained using Lemma 3.8, Lemma 3.9 and Lemma 3.10. For the last one, we prove the equality $\tau_{v_4} = \sigma_{u_5} - \sigma'_{u_7}$. The result then follows from Lemmas 3.10 and 3.14. To prove the equality $\tau_{v_4} = \sigma_{u_5} - \sigma'_{u_7}$, we first write $\sigma'_{u_2} = a\sigma_{u_0} + b\tau'_{v_2}$. Multiplying with σ'_{u_7} and using Remark 1.13, we have $1 = a$. Multiplying with $h^{\cup 7}$, we obtain $56 = 56a + b\tau'_{v_2} \cup h^{\cup 7}$. Finally we get $\sigma'_{u_2} = \sigma_{u_0}$. Now write $\sigma_{u_5} = \lambda\sigma'_{u_7} + \mu\tau_{v_4}$. Multiplying with $\sigma'_{u_2} = \sigma_{u_0}$ we get $1 = \lambda$. Multiplying with $h \cup h$ we get $2 = \lambda + \mu$. It follows that $\lambda = 1$ and $\mu = 1$.

For line (9), we need to compute $\langle h, \tau_{v_5}, \sigma_{u_3} \rangle$, $\langle h, \tau_{v_5}, \sigma_{u'_3} \rangle$ and $\langle h, \tau_{v_5}, \tau'_{v_5} \rangle$. The first two are obtained using Lemma 3.9. For the last one we remark that $\tau_{v_5} = [\text{line}] = \sigma_{u_6}$ and use Lemma 3.14.

For line (10), we need to compute $\langle h, \tau_{v_6}, \sigma_{u_2} \rangle$, $\langle h, \tau_{v_6}, \sigma_{u'_2} \rangle$ and $\langle h, \tau_{v_6}, \tau'_{v_4} \rangle$. The first two are obtained using Lemma 3.9. For the last one we remark that $\tau_{v_6} = [\text{pt}] = \sigma_{u_7}$ and use Lemma 3.14. Finally the q^2 term follows from Lemma 3.18. \square

4.3. Case $X = X^2$. We have $\dim X = 9$ and $c_1(X) = 7$. Both Y and Z are quadrics with $\dim Y = 5$ and $\dim Z = 6$. There is only one cohomology class for each degree in Y so we denote by $(u_i)_{i \in [0,5]}$ with $\ell(u_i) = i$ the elements of W^{P_Y} . In Z , for all degree except 3 there is only one cohomology class for each degree so we denote by $(v_i)_{i \in [0,5]}$ with $\ell(v_i) = i$ and by v'_3 the elements in W^{P_Z} . We choose to set $v_3 = s_1 s_2 s_3$ and $v'_3 = s_3 s_2 s_3$. Note that $\sigma'_{u_0} = \tau'_{v_0} = 1$, $\sigma'_{u_1} = \tau'_{v_1} = h$, $\sigma_{u_5} = \tau_{v_6} = [\text{pt}]$ and $\sigma_{u_4} = \tau_{v_5} = [\text{line}]$.

Proposition 4.4. *We have the formulas*

- (1) $h * 1 = h$
- (2) $h * h = \sigma'_{u_2}$
- (3) $h * \sigma'_{u_2} = 2\sigma'_{u_3} + \tau_{v_0}$

- (4) $h * \sigma'_{u_3} = \sigma'_{u_4} + \tau_{v_1}$ and $h * \tau_{v_0} = \tau_{v_1}$
- (5) $h * \sigma'_{u_4} = \sigma'_{u_5} + \tau_{v_2}$ and $h * \tau_{v_1} = \tau_{v_2}$
- (6) $h * \sigma'_{u_5} = \tau_{v_3}$ and $h * \tau_{v_2} = \tau_{v_3} + \tau_{v'_3}$
- (7) $h * \tau_{v_3} = \tau_{v_4}$ and $h * \tau_{v'_3} = \tau_{v_4} + q$
- (8) $h * \tau_{v_4} = \tau_{v_5} + qh$
- (9) $h * \tau_{v_5} = \tau_{v_6} + q\sigma'_{u_2}$
- (10) $h * \tau_{v_6} = q\sigma'_{u_3}$.

Proof. The first 6 lines follow from the classical Hasse diagram, and the quantum parts follow from Lemmas 3.8, 3.10, 3.9 and 3.12. \square

4.4. **Case $X = X^3(3, 3)$.** We have $\dim X = 9$ and $c_1(X) = 5$. Moreover Y is a 6-dimensional quadric. There is only one cohomology class for each degree in Y except in degree 3, so we denote by $(u_i)_{i \in [0,6]}$ with $\ell(u_i) = i$ and u'_3 with $\ell(u'_3) = 3$ the elements in W^{PY} . We choose to set $u_3 = s_1s_2s_3$ and $u'_3 = s_3s_2s_3$. We have $Z = \text{IG}(2, 6)$ and there is one cohomology class in degree 0, 1, 6 and 7 and two cohomology classes otherwise. We denote by $(v_i)_{i \in [0,7]}$ with $\ell(v_i) = i$ and v'_2, v'_3, v'_4, v'_5 with $\ell(v'_i) = i$ the elements in W^{PZ} . We choose to set $v_2 = s_1s_2$, $v_3 = s_3s_1s_2$, $v_4 = s_2s_3s_1s_2$, and $v_5 = s_3s_2s_3s_1s_2$. Note that $\sigma'_{u_0} = \tau'_{v_0} = 1$, $\sigma'_{u_1} = \tau'_{v_1} = h$, $\sigma_{u_6} = \tau_{v_7} = [\text{pt}]$ and $\sigma_{u_5} = \tau_{v_6} = [\text{line}]$.

Proposition 4.5. *We have the formulas*

- (1) $h * 1 = h$
- (2) $h * h = 2\sigma'_{u_2} + \tau_{v_0}$
- (3) $h * \sigma'_{u_2} = 2\sigma'_{u_3} + \sigma'_{u'_3} + \tau_{v_1}$ and $h * \tau_{v_0} = \tau_{v_1}$
- (4) $h * \sigma'_{u_3} = \sigma'_{u_4} + \tau_{v_2}$, $h * \sigma'_{u'_3} = 2\sigma'_{u_4} + \tau_{v'_2}$ and $h * \tau_{v_1} = \tau_{v_2} + \tau_{v'_2}$
- (5) $h * \sigma'_{u_4} = 2\sigma'_{u_5} + \tau_{v_3}$, $h * \tau_{v_2} = \tau_{v_3} + q$ and $h * \tau_{v'_2} = \tau_{v_3} + \tau_{v'_3}$
- (6) $h * \sigma'_{u_5} = \sigma'_{u_6} + \tau_{v_4}$, $h * \tau_{v_3} = 2\tau_{v_3} + \tau_{v'_4} + qh$ and $h * \tau_{v'_3} = \tau_{v_4} + 2\tau_{v'_4}$
- (7) $h * \sigma'_{u_6} = \tau_{v_5}$, $h * \tau_{v_4} = \tau_{v_5} + \tau_{v'_5} + q\sigma'_{u_2}$ and $h * \tau_{v'_4} = \tau_{v'_5} + q\tau_{v_0}$
- (8) $h * \tau_{v_5} = \tau_{v_6} + q\sigma'_{u_3}$ and $h * \tau_{v'_5} = \tau_{v_6} + q\sigma'_{u'_3} + q\tau_{v_1}$
- (9) $h * \tau_{v_6} = \tau_{v_7} + q\sigma'_{u_4} + q\tau_{v'_2}$
- (10) $h * \tau_{v_7} = q\sigma'_{u_5} + q\tau_{v'_3}$.

Proof. The first 6 lines follows from the classical Hasse diagram. The quantum parts follow from Lemmas 3.8, 3.10, 3.9, 3.12 and the fact (see Corollary 5.11) that there is no q^2 term in the quantum multiplication with h . \square

4.5. **Case $X = X^5$.** We have $\dim X = 7$ and $c_1(X) = 4$. There is only one cohomology class for each degree in Y and Z so we denote by $(u_i)_{i \in [0,5]}$ with $\ell(u_i) = i$ and $(v_i)_{i \in [0,5]}$ with $\ell(v_i) = i$ the elements in W^{PY} and W^{PZ} . Note that $\sigma'_{u_0} = \tau'_{v_0} = 1$, $\sigma'_{u_1} = \tau'_{v_1} = h$, $\sigma_{u_5} = \tau_{v_5} = [\text{pt}]$ and $\sigma_{u_4} = \tau_{v_4} = [\text{line}]$.

Proposition 4.6. *We have the formulas*

- (1) $h * 1 = h$
- (2) $h * h = 3\sigma'_{u_2} + \tau_{v_0}$
- (3) $h * \sigma'_{u_2} = 2\sigma'_{u_3} + \tau_{v_1}$ and $h * \tau_{v_0} = \tau_{v_1}$
- (4) $h * \sigma'_{u_3} = 3\sigma'_{u_4} + \tau_{v_2}$ and $h * \tau_{v_1} = \tau_{v_2} + q$
- (5) $h * \sigma'_{u_4} = \sigma'_{u_5} + \tau_{v_3}$ and $h * \tau_{v_2} = 2\tau_{v_3} + qh$
- (6) $h * \sigma'_{u_5} = \tau_{v_4} + q\tau_{v_0}$ and $h * \tau_{v_3} = \tau_{v_4} + q\sigma'_{u_2}$
- (7) $h * \tau_{v_4} = \tau_{v_5} + q\sigma'_{u_3} + q\tau_{v_1}$

$$(8) \quad h * \tau_{v_5} = q\sigma'_{u_4} + q\tau_{v_2} + 2q^2.$$

Proof. We will only discuss the quantum corrections, since the classical part follows from the classical Hasse diagram. By Lemma 3.6 there is no power of q higher than 1 except maybe in $h * \tau_{v_5}$. For the first three lines there is no quantum correction since the degree is smaller than $4 = c_1(X)$. For the fourth line, we only have to compute $\langle h, \sigma'_{u_3}, \sigma_{u_5} \rangle$ and $\langle h, \tau_{v_1}, \sigma_{u_5} \rangle$. The result follows from Lemma 3.8 and Lemma 3.9. For line (5), we only have to compute $\langle h, \sigma'_{u_4}, \sigma_{u_4} \rangle$ and $\langle h, \tau_{v_2}, \sigma_{u_4} \rangle$. The result follows from Lemma 3.8 and Lemma 3.9.

Line (6) requires more work. We need to compute the Gromov-Witten invariants $\langle h, \sigma'_{u_5}, \sigma_{u_3} \rangle$, $\langle h, \tau_{v_3}, \sigma_{u_3} \rangle$, $\langle h, \sigma'_{u_5}, \tau'_{v_5} \rangle$ and $\langle h, \tau_{v_3}, \tau'_{v_5} \rangle$. The first three are obtained using Lemma 3.8, Lemma 3.9 and Lemma 3.10. For the last one, we prove the equality $\tau_{v_3} = \tau'_{v_5}$. The result then follows from Lemma 3.16. Let us now prove $\tau_{v_3} = \tau'_{v_5}$. Write $\tau'_{v_5} = \lambda\sigma'_{u_5} + \mu\tau_{v_2}$. Multiplying with τ'_{v_2} we get $1 = \mu$. Multiplying with $h \cup h$ we get $1 = \lambda + \mu$. It follows that $\lambda = 0$ and $\mu = 1$.

For line (7) we need to compute $\langle h, \tau_{v_4}, \sigma_{u_2} \rangle$ and $\langle h, \tau_{v_4}, \tau'_{v_4} \rangle$. The first invariant is obtained using Lemma 3.9. For the last one we remark that $\tau_{v_4} = [\text{line}] = \sigma_{u_4}$ and use Lemma 3.14.

For line (8) we need to compute $\langle h, \tau_{v_5}, \sigma_{u_1} \rangle$ and $\langle h, \tau_{v_5}, \tau'_{v_3} \rangle$. The first invariant is obtained using Lemma 3.9. For the last one we remark that $\tau_{v_5} = [\text{line}] = \sigma_{u_5}$ and use Lemma 3.14. Finally the q^2 term follows from Lemma 3.18. \square

5. THE ODD SYMPLECTIC GRASSMANNIAN

In this section we apply our enumerativity result, Corollary 3.4, to deduce some results concerning the quantum multiplication in odd symplectic Grassmannians (class (3) of Pasquier's list). More precisely, in Section 5.4 we use Buch's Ker/Span techniques to relate Gromov-Witten invariants of odd symplectic Grassmannians to intersection numbers in an auxiliary variety. This leads in Section 5.5 to Theorem 5.13, which compares the quantum and the classical multiplication. Finally, in Section 5.6 we obtain a presentation of the quantum cohomology of any odd symplectic Grassmannian, see Theorem 5.17.

5.1. The odd symplectic Grassmannian as a horospherical variety. As in Section 1.9 we let $n \geq 2$ be an integer and ω be an antisymmetric form of maximal rank on a complex vector space V of dimension $2n + 1$. If $2 \leq m \leq n$, we denote by $X = \text{IG}(m, V) = \text{IG}(m, 2n + 1)$ the Grassmannian of vector subspaces of V which are isotropic for ω :

$$\text{IG}(m, V) = \{\Sigma \subset V \mid \dim \Sigma = m, \omega|_{\Sigma} = 0\}.$$

To view X as a horospherical variety we let $K \subset V$ be the kernel of the form ω and W be a complement to K in V , so that $\omega|_W$ is symplectic. Setting $G := \text{Sp}(W) \cong \text{Sp}_{2n}$ we see that X is G -horospherical and corresponds to the triple $(C_n, \omega_m, \omega_{m-1})$, i.e., to the case (3) of Pasquier's classification.

Indeed, X is contained in $\mathbb{P}(\Lambda^m V)$ and as a G -representation we have $V = W \oplus K$, therefore $\Lambda^m V = \Lambda^m W \oplus \Lambda^{m-1} W \otimes K$, which are the irreducible representations of highest weights ω_m and ω_{m-1} , respectively. One easily checks that X is the closure of the G -orbit of the class $[v_Y + v_Z]$ of the sum of the lowest weight vectors v_Y, v_Z of these two representations.

The G -orbits in X are

$$Y = \{\Sigma \in X \mid \Sigma \subset W\} = \text{IG}(m, W) \text{ and } Z = \{\Sigma \in X \mid K \subset \Sigma\} \cong \text{IG}(m-1, W).$$

Moreover, the blow-ups \tilde{X}_Y , \tilde{X}_Z and \tilde{X}_{YZ} are explicitly given by:

$$\begin{aligned} \tilde{X}_Y &= \{(\Sigma, \Sigma') \in X \times Z \mid \Sigma' \cap W \subset \Sigma \cap W\}, \\ \tilde{X}_Z &= \{(\Sigma, \Sigma'') \in X \times Y \mid K + \Sigma \subset K + \Sigma''\}, \\ \tilde{X}_{YZ} &= \{(\Sigma, \Sigma', \Sigma'') \in X \times Z \times Y \mid (\Sigma, \Sigma') \in \tilde{X}_Y \text{ and } (\Sigma, \Sigma'') \in \tilde{X}_Z\}. \end{aligned}$$

The maps $\pi_Y, \pi_Z, p_Y, p_Z, \pi_{YZ}, p_{YZ}$ are given by the obvious projections (for the last one, note that $E \subset Y \times Z$ is the incidence variety).

The automorphism group $\text{Aut}(X)$ is the *odd symplectic group*

$$\text{Sp}_{2n+1} = \text{Sp}(V) := \{M \in \text{GL}(V) \mid M^t \omega M = \omega\}.$$

The odd symplectic group is not reductive, however its properties are closely related to those of the usual symplectic group, see [40].

5.2. Schubert varieties in symplectic Grassmannians. As seen in Section 1.7 we have two Poincaré dual bases of the cohomology of $X := \text{IG}(m, V)$, obtained from the Schubert classes of the closed Sp_{2n} -orbits Y and Z , which are symplectic Grassmannians. There are many indexations for Schubert classes of symplectic Grassmannians; in this section we present those which will play a part later.

Indexation by Weyl group elements. Schubert varieties of the symplectic Grassmannian $\text{IG}(m, 2n)$ are indexed by elements of W^P , i.e., by minimal length coset representatives of W/W_P . Such elements are signed permutations which may be written as

$$w = (w(1), w(2), \dots, w(n)) = (y_1, y_2, \dots, y_{m-\ell}, \bar{z}_\ell, \dots, \bar{z}_2, \bar{z}_1, v_1, v_2, \dots, v_{n-m}),$$

where $0 \leq \ell \leq m$, $0 < y_1 < y_2 < \dots < y_{m-\ell}$, $0 < z_1 < z_2 < \dots < z_\ell$, $0 < v_1 < v_2 < \dots < v_{n-m}$, and $\bar{z}_i := -z_i$.

Indexation by pairs of partitions. This indexation was introduced in [53]. We recall it here since it will be needed in the proof of Theorem 5.17. A *pair of partitions* for $\text{IG}(m, 2n)$ is a pair $\alpha = (\alpha^t, \alpha^b)$, where α^t is a strict partition contained in a $(n-m) \times n$ rectangle, α^b a strict partition contained in a $m \times n$ rectangle, and $\alpha_{n-m}^t \geq \ell(\alpha^b) + 1$. Weyl group elements w and pairs of partitions α are related by the bijective map $w \mapsto \alpha$ given by

- $\alpha_r^t = n + 1 - v_r + \#\{1 \leq j \leq \ell \mid z_j < v_r\}$ for $1 \leq r \leq n - m$;
- $\alpha_j^b = n + 1 - z_j$ for $1 \leq j \leq \ell$.

Indexation by k -strict partitions. This is the indexation from [8, Definition 1.1]. For $\text{IG}(m, 2n)$ we let $k = n - m$, and we say that a partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_m \geq 0)$ is *k -strict* if no part greater than k is repeated, i.e. $\lambda_j > k \Rightarrow \lambda_j > \lambda_{j+1}$. Thereafter we will denote by 1^p the partition with p parts all equal to 1.

We have that k -strict partitions λ are in bijective correspondence with pairs of partitions $\alpha = (\alpha^t, \alpha^b)$, via the map $\lambda \mapsto \alpha$ given by

- $\alpha_r^t = n + 1 - m - r + \#\{1 \leq i \leq m \mid \lambda_i \geq r\}$ for $1 \leq r \leq n - m$;
- $\alpha_j^b = \lambda_j + m - n$ for $1 \leq j \leq \max\{1 \leq i \leq m \mid \lambda_i > n - m\}$.

Indexation by index sets. We say a sequence $\mathbf{p} = (p_1 < \dots < p_m)$ with $p_1 \geq 1$ and $p_m \leq 2n$ is an *index set* for $\mathrm{IG}(m, 2n)$ if $p_i + p_j \neq 2n + 1$ for any i, j . There is a bijection between index sets \mathbf{p} and $(n - m)$ -strict partitions λ with $\lambda_1 \leq 2n - m$. This correspondence is given by

- $\lambda_j = 2n + 1 - m - p_j + \#\{i < j \mid p_i + p_j > 2n + 1\}$;
- $p_j = 2n + 1 - m - \lambda_j + \#\{i < j \mid \lambda_i + \lambda_j \leq 2(n - m) + j - i\}$.

Index sets are most useful for understanding Schubert varieties geometrically. Indeed, let $F_\bullet := (\{0\} = F_0 \subset F_1 \subset F_2 \subset \dots \subset F_{2n} = \mathbb{C}^{2n})$ be an Sp_{2n} -flag, i.e., a complete flag of \mathbb{C}^{2n} such that $F_j^\perp = F_{2n-j}$ for any $0 \leq j \leq 2n$. Then we define the Schubert variety of $\mathrm{IG}(m, 2n)$ associated with the index set $\mathbf{p} := (p_1, \dots, p_m)$ by

$$X_{\mathbf{p}}(F_\bullet) := \{\Sigma \in \mathrm{IG}(m, 2n) \mid \dim(\Sigma \cap F_{p_j}) \geq j, \forall 1 \leq j \leq m\}.$$

5.3. Schubert varieties in odd symplectic Grassmannians. We now come back to the odd symplectic Grassmannian $X := \mathrm{IG}(m, V) = \mathrm{IG}(m, 2n + 1)$ from Section 5.1. In addition to the two Poincaré dual bases introduced in Section 1.7, in 2008 Mihai [40] constructed Schubert-type varieties for X associated with Borel subgroups of the automorphism group $\mathrm{Aut}(X) = \mathrm{Sp}_{2n+1}$. Let us explain how these families of cohomology classes compare.

The Borel subgroups of Sp_{2n+1} are in one-to-one correspondence with complete flags of V of the form

$$F_\bullet := (\{0\} = F_0 \subset K = F_1 \subset F_2 \subset \dots \subset F_{2n} \subset F_{2n+1} = V),$$

where $K = \ker \omega$, and for any $1 \leq j \leq 2n + 1$, $F_j^\perp = F_{2n+2-j}$. We call these flags Sp_{2n+1} -flags. Schubert varieties of X have been defined by Mihai as follows.

Definition 5.1. Let F_\bullet be an Sp_{2n+1} -flag and $1 \leq p_1 < p_2 < \dots < p_m \leq 2n + 1$ be integers such that $p_i + p_j \neq 2n + 3$ for any i, j . We call such a sequence $\mathbf{p} = (p_1, \dots, p_m)$ an *odd index set* for $\mathrm{IG}(m, 2n + 1)$. We define the Schubert variety of X associated with the odd index set \mathbf{p} by

$$X_{\mathbf{p}}(F_\bullet) := \{\Sigma \in X \mid \dim(\Sigma \cap F_{p_j}) \geq j, \forall 1 \leq j \leq m\}.$$

Mihai showed that these varieties form a CW complex; thus the corresponding cohomology classes $v_{\mathbf{p}} := [X_{\mathbf{p}}(F_\bullet)]$ form a \mathbb{Z} -basis for the cohomology ring $H^*(X, \mathbb{Z})$ of X . In [47, Section 2.2] the second author introduced another indexation for these classes analogous to what happens in the symplectic case, see Section 5.2. Namely, an $(n - m)$ -strict partition for $\mathrm{IG}(m, 2n + 1)$ is a sequence of integers $\mu = (\mu_1 \geq \dots \geq \mu_m \geq -1)$ such that $\mu_1 \leq 2n + 1 - m$ and $\mu_1 = 2n + 1 - m$ if $\mu_m = -1$. The next proposition is straightforward.

Proposition 5.2. *There is a bijection between odd index sets \mathbf{p} and $(n - m)$ -strict partitions μ for $\mathrm{IG}(m, 2n + 1)$, given by*

- $\mu_j = 2n + 2 - m - p_j + \#\{i < j \mid p_i + p_j > 2n + 3\}$;
- $p_j = 2n + 2 - m - \mu_j + \#\{i < j \mid \mu_i + \mu_j \leq 2(n - m) + j - i\}$.

It follows from this proposition that we may index the Schubert varieties of $X = \mathrm{IG}(m, 2n + 1)$ by suitable $(n - m)$ -strict partitions. In particular if \mathbf{p} is an odd index set for $\mathrm{IG}(m, 2n + 1)$ and λ is the corresponding $(n - m)$ -strict partition as in Proposition 5.2, we set $v_\lambda := v_{\mathbf{p}}$. In the rest of the section we will sometimes denote by $\mathbf{p}(\lambda)$ the index set corresponding to the strict partition λ .

We now consider the two cohomology bases of X constructed in Section 1.7. We start by checking that the basis (1) from Fact 1.8, namely $\{\sigma'_u, \tau_v\}$, coincides with Mihai's Schubert-type basis $\{(v_{\mathbf{p}})\} = \{(v_\lambda)\}$.

Proposition 5.3. *Let λ be an $(n - m)$ -strict partition of $Y = \text{IG}(m, 2n)$ and μ be an $(n + 1 - m)$ -strict partition of $Z = \text{IG}(m - 1, 2n)$. Then in $H^*(X, \mathbb{Z})$,*

- (1) $\sigma'_\lambda = v_\lambda$;
- (2) $\tau_\mu = v_{2n+1-m, \mu_1-1, \dots, \mu_{m-1}-1}$.

Proof. Recall the decomposition $V = K \oplus W$ and the description of the closed Sp_{2n} -orbits $Y = \{\Sigma \in X \mid \Sigma \subset W\}$ and $Z = \{\Sigma \in X \mid K \subset \Sigma\}$ from Section 5.1. Consider an Sp_{2n} -flag:

$$G_\bullet := (\{0\} \subset G_1 \subset G_2 \subset \dots \subset G_{2n-1} \subset W),$$

where for any $1 \leq j \leq n$ we have that $G_j^\perp \cap W = G_{2n-j}$. Here the orthogonal is taken relatively to the antisymmetric form of maximal rank ω on $V = \mathbb{C}^{2n+1}$. The Schubert varieties of Y and Z from Section 1.7 can be defined with respect to such a flag. To compare them to Mihai's varieties, we extend G_\bullet as follows:

$$K \oplus G_\bullet := (\{0\} \subset K \subset K \oplus G_1 \subset \dots \subset K \oplus G_{2n-1} \subset K \oplus W = V).$$

For $1 \leq j \leq 2n$, we have $(K \oplus G_j)^\perp = G_j^\perp = K \oplus (G_j^\perp \cap W) = K \oplus G_{2n-j}$, hence $K \oplus G_\bullet$ is an Sp_{2n+1} -flag. Now consider the subvariety of Y associated with λ ,

$$Y_{\mathbf{p}(\lambda)}(G_\bullet) = \{\Lambda \in Y \mid \dim(\Lambda \cap G_{p_j(\lambda)}) \geq j, \forall 1 \leq j \leq m\},$$

and similarly for Z ,

$$Z_{\mathbf{p}(\mu)}(G_\bullet) = \{\Lambda \in Z \mid \dim(\Lambda \cap G_{p_j(\mu)}) \geq j, \forall 1 \leq j \leq m - 1\}.$$

Clearly,

$$Y'_{\mathbf{p}(\lambda)}(G_\bullet) := \pi_Z(p_Y^{-1}(Y_{\mathbf{p}(\lambda)}(G_\bullet))) = X_{\mathbf{p}(\lambda)+\mathbf{1}}(K \oplus G_\bullet),$$

where the maps π_Z and p_Y are defined in Section 1.5, and

$$\mathbf{p}(\lambda) + \mathbf{1} := (p_1(\lambda) + 1, \dots, p_m(\lambda) + 1).$$

Since $[X_{\mathbf{p}(\lambda)+\mathbf{1}}(K \oplus G_\bullet)] = v_\lambda$, the first statement follows. Similarly,

$$\tau_\mu = [i_Z(Z_{\mathbf{p}(\mu)}(G_\bullet))] = [X_{\mathbf{1}, \mathbf{p}(\mu)+\mathbf{1}}(K \oplus G_\bullet)] = v_{2n+1-m, \mu_1-1, \dots, \mu_{m-1}-1},$$

which concludes the proof. \square

It is also interesting to see what the basis (2) from Fact 1.8, namely $\{\sigma_u, \tau'_v\}$ corresponds to. We will see that these basis elements no longer correspond to Schubert-type classes of X ; however, they coincide with pullbacks of Schubert classes σ_v^+ from a larger symplectic Grassmannian $X^+ = \text{IG}(m, V^+)$. Here $V^+ \supset V$ is a $(2n + 2)$ -dimensional vector space endowed with a symplectic form ω^+ such that $\omega^+|_V = \omega$, so that X^+ is the Grassmannian of isotropic m -spaces in V^+ .

Consider the embedding $j : X \hookrightarrow X^+$ and the induced surjective pullback map in cohomology $j^* : H^*(X^+, \mathbb{Z}) \rightarrow H^*(X, \mathbb{Z})$. It was already noticed in [40] that X can be identified with the Schubert variety of X^+ associated to the $(n + 1 - m)$ -strict partition 1^m . This immediately implies that the pullback map can be computed as follows, see also [47, Equation (2.3)].

Proposition 5.4. *Let ν be an $(n + 1 - m)$ -strict partition for X^+ , and write*

$$\sigma_\nu^+ \cup \sigma_{1^m}^+ = \sum_{\beta} c_\nu^\beta \sigma_\beta^+,$$

where the coefficients c_ν^β are computed from the Pieri rule of [53]. Then

$$j^* \sigma_\nu^+ = \sum_{\beta} c_\nu^\beta v_{\beta-1^m} \in H^*(X, \mathbb{Z}).$$

Note that the classical Pieri rule of [53] uses the indexation of Schubert classes of X^+ in terms of pairs of partitions introduced in Section 5.2. Therefore in later sections we will use that indexation each time we wish to compute the coefficients c_ν^β explicitly.

We may now identify the basis elements (2) from Fact 1.8 with pullbacks by j .

Proposition 5.5. *Let λ be an $(n - m)$ -strict partition of $Y = \text{IG}(m, 2n)$ and μ be an $(n + 1 - m)$ -strict partition of $Z = \text{IG}(m - 1, 2n)$. Then the classes $\sigma_\lambda, \tau'_\mu \in H^*(X, \mathbb{Z})$ satisfy*

$$\sigma_\lambda = j^* \sigma_{\lambda+1^m}^+ \text{ and } \tau'_\mu = j^* \sigma_\mu^+,$$

where σ_ν^+ denotes the Schubert class of $X^+ = \text{IG}(m, 2n + 2)$ associated with the $(n + 1 - m)$ -strict partition ν .

Proof. From Proposition 1.10 we know that σ_λ is the Poincaré dual of the class $\sigma'_{\lambda^\vee} \in H^*(X, \mathbb{Z})$, where λ^\vee denotes the dual partition with respect to Poincaré duality in $Y = \text{IG}(m, 2n)$. Using Proposition 2.11 from [47] we know that this Poincaré dual class coincides with a pulled-back class, i.e.

$$\sigma_\lambda = j^*(\sigma_{\lambda^\vee+1^m}^+)^\vee = j^* \sigma_{\lambda+1^m}^+,$$

where last equality is obtained by looking at Poincaré duality in terms of index sets. The statement concerning τ'_μ follows by a similar argument. \square

5.4. Ker/Span techniques. In this section we relate degree d Gromov-Witten invariants of the odd symplectic Grassmannian $X = \text{IG}(m, 2n + 1)$ to intersection numbers in an auxiliary variety I_d^+ . To obtain this comparison, following an idea introduced by Buch in [6] and Buch, Kresch and Tamvakis in [7] we replace degree d rational curves on X or X^+ by their *kernel* and *span*.

Definition 5.6. Let $C \subset X$ (resp. $C \subset X^+$) be a degree d rational curve in X (resp. in X^+).

- The *kernel* $\text{Ker } C$ of C is the subspace of V (resp. of V^+) defined by

$$\text{Ker } C := \bigcap_{\Sigma \in C} \Sigma.$$

- The *span* $\text{Span } C$ of C is the subspace of V (resp. of V^+) defined by

$$\text{Span } C := \text{Span}\{\Sigma \mid \Sigma \in C\}.$$

We define a variety $I_{k,s,e}$ of kernel/span pairs in V as a subvariety of the product of Grassmannians $G \times G' = \text{Gr}(m - k, V) \times \text{Gr}(m + s, V)$ as follows:

$$I_{k,s,e} := \left\{ (A, B) \in G \times G' \mid \begin{array}{l} A \subset B \subset A^\perp \subset V, \\ \dim B \cap B^\perp = m - k + e \end{array} \right\}.$$

Here the orthogonal is defined with respect to the antisymmetric form of maximal rank ω on $V = \mathbb{C}^{2n+1}$. The variety $I_{k,s,e}^+$ of kernel/span pairs in $V^+ = \mathbb{C}^{2n+2}$ is defined similarly, replacing V with V^+ and orthogonality in V with orthogonality in V^+ (using the symplectic form ω^+ extending ω). For simplicity we denote $\bigcup_e I_{d,d,e}$ (resp. $\bigcup_e I_{d,d,e}^+$) by I_d (resp. I_d^+).

Lemma 5.7. *For any degree d rational curve $C \subset X$ we have*

$$(\text{Ker } C, \text{Span } C) \in \bigcup_{k,s=0}^d \bigcup_{e=0}^{k+s} I_{k,s,e}.$$

The same result holds for $C \subset X^+$, replacing $I_{k,s,e}$ with $I_{k,s,e}^+$.

Proof. We have the inequalities $\dim \text{Ker } C \geq m - d$ and $\dim \text{Span } C \leq m + d$, see [6]. Moreover, since the curve C is contained in an isotropic Grassmannian we have that $\text{Ker } C \subset \text{Span } C \subset (\text{Ker } C)^\perp$. The inequalities

$$0 \leq \dim(\text{Span } C \cap (\text{Span } C)^\perp) - \dim \text{Ker } C \leq \dim \text{Span } C - \dim \text{Ker } C.$$

follow. □

Requiring that a given degree d curve C should intersect a given Schubert variety X_λ of X (resp. X_λ^+ of X^+) defines a subvariety of $I_{k,s,e}$ (resp. $I_{k,s,e}^+$) as follows. For the sake of readability, we omit here from our notations the flags with respect to which Schubert varieties are defined:

$$\begin{aligned} I_{k,s,e}(\lambda) &= \{(A, B) \in I_{k,s,e} \mid \exists \Sigma \in X_\lambda, A \subset \Sigma \subset B\}, \\ I_{k,s,e}^+(\lambda) &= \{(A, B) \in I_{k,s,e} \mid \exists \Sigma^+ \in X_\lambda^+, A \subset \Sigma^+ \subset B\}. \end{aligned}$$

We also write $I_d(\lambda)$ for $\bigcup_e I_{d,d,e}(\lambda)$ (resp. $I_d^+(\lambda)$ for $\bigcup_e I_{d,d,e}^+(\lambda)$). We may now state our comparison result.

Proposition 5.8. *Let $d \geq 0$ be an integer and α, β, γ be elements of the cohomology basis $\{\sigma_\lambda, \tau'_\mu\}$ of $X = \text{IG}(m, 2n + 1)$ (that is, basis (2) from Fact 1.8, indexed using strict partitions). Assume that*

$$\text{codim } \alpha + \text{codim } \beta + \text{codim } \gamma = \dim M_{d,3}(X) = \dim X + d(2n + 2 - m).$$

Following Proposition 5.5 we denote by $\alpha^+, \beta^+, \gamma^+$ the Schubert classes of $X^+ = \text{IG}(m, 2n + 2)$ such that

$$j^* \alpha^+ = \alpha; j^* \beta^+ = \beta; j^* \gamma^+ = \gamma.$$

Let $X_{\alpha^+}^+, X_{\beta^+}^+$ and $X_{\gamma^+}^+$ be Schubert varieties of X^+ in general position associated with $\alpha^+, \beta^+,$ and γ^+ , respectively, and denote by $I_d^+(\alpha^+), I_d^+(\beta^+), I_d^+(\gamma^+)$ the corresponding subvarieties of I_d^+ . Then

$$\langle \alpha, \beta, \gamma \rangle_{X,d,3} = \int_{I_d^+} [I_d^+(\alpha^+)] \cup [I_d^+(\beta^+)] \cup [I_d^+(\gamma^+)] \cup [I_d].$$

Moreover, the Gromov-Witten invariant $\langle \alpha, \beta, \gamma \rangle_{X,d,3}$ vanishes unless each of the classes $\alpha, \beta, \gamma \in H^(X, \mathbb{Z})$ is either of the form σ_λ with $\lambda \supset \rho_{d-1}$, or of the form τ'_μ with $\mu \supset \rho_d$. Here we denoted by ρ_ℓ the ℓ -staircase partition $(\ell, \ell - 1, \dots, 1)$.*

A proof of Proposition 5.8, provided Corollary 3.4 holds, was already given in [47, Section 5.1] with different notation. For the non-Francophone reader's convenience we repeat here the main steps of the proof.

Sketch of proof of Prop. 5.8. First of all we notice via a dimension count that the intersection $I_{k,s,e}^+(\alpha^+) \cap I_{k,s,e}^+(\beta^+) \cap I_{k,s,e}^+(\gamma^+) \cap I_{k,s,e}$ is empty unless $k = s = d$ and $e = 0$, in which case it consists in a finite number of points. Then we use Corollary 3.4 to prove our first statement, namely that

$$\langle \alpha, \beta, \gamma \rangle_{X,d,3} = \int_{I_d^+} [I_d^+(\alpha^+)] \cup [I_d^+(\beta^+)] \cup [I_d^+(\gamma^+)] \cup [I_d].$$

Let $f : \mathbb{P}^1 \rightarrow X \subset X^+$ be a degree d morphism such that

$$f(0) \in X_{\alpha^+}^+ \cap X, f(1) \in X_{\beta^+}^+ \cap X, f(\infty) \in X_{\gamma^+}^+ \cap X.$$

Then the Ker/Span pair $(\text{Ker } f, \text{Span } f)$ is an element of some intersection

$$I_{k,s,e}^+(\alpha^+) \cap I_{k,s,e}^+(\beta^+) \cap I_{k,s,e}^+(\gamma^+) \cap I_{k,s,e} \subset I_{k,s,e}^+.$$

This means we must have $k = s = d$ and $e = 0$, so that $(\text{Ker } f, \text{Span } f)$ is in fact an element of $I_d^+(\alpha^+) \cap I_d^+(\beta^+) \cap I_d^+(\gamma^+) \cap I_d \subset I_d^+$ as required.

Conversely, consider an element $(A, B) \in I_d^+(\alpha^+) \cap I_d^+(\beta^+) \cap I_d^+(\gamma^+) \cap I_d$. We need to show that there exists a unique degree d morphism $f : \mathbb{P}^1 \rightarrow X$ such that $f(0) \in X_{\alpha^+}^+ \cap X$, $f(1) \in X_{\beta^+}^+ \cap X$, and $f(\infty) \in X_{\gamma^+}^+ \cap X$. By definition of (A, B) there exist elements P, Q, R of X such that $A \subset P, Q, R \subset B$ and

$$P \in X_{\alpha^+}^+, Q \in X_{\beta^+}^+, R \in X_{\gamma^+}^+.$$

By a dimension count, we show that $P \cap Q = P \cap R = Q \cap R = A$. Moreover the set $\{\Sigma^+ \in X^+ \mid A \subset \Sigma^+ \subset B\} \cap X_{\alpha^+}^+$ must be zero-dimensional, otherwise we could choose a space P in it whose intersection with Q is larger than A . Similarly, the intersections with $X_{\beta^+}^+$ and $X_{\gamma^+}^+$ are also zero-dimensional. We then use the following lemma from [7].

Lemma 5.9 ([7, Lemma 1.3]). *Consider*

$$T_d^+ := \{(A, \Sigma^+, B) \mid (A, B) \in I_d^+, \Sigma^+ \in X^+, A \subset \Sigma^+ \subset B\},$$

and for any Schubert variety $X_{\alpha^+}^+$ of X^+ ,

$$T_d^+(\alpha^+) := \{(A, \Sigma^+, B) \in T_d^+ \mid \Sigma^+ \in X_{\alpha^+}^+\}.$$

The projection map $T_d^+(\alpha^+) \rightarrow I_d^+(\alpha^+)$ is generically 1:1 when $\nu \supset \rho_d$, where $\sigma_\nu^+ = \alpha^+$, and has positive-dimensional fibres otherwise.

Applying the lemma to our setting we deduce that $\alpha^+ = \sigma_\nu^+$ with $\nu \supset \rho_d$. Thus the invariant $\langle \alpha, \beta, \gamma \rangle_{X,d,3}$ vanishes when $\nu \not\supset \rho_d$, which is equivalent to the vanishing statement we need to prove. Moreover

$$\{\Sigma^+ \in X^+ \mid A \subset \Sigma^+ \subset B\} \cap X_{\alpha^+}^+ = \{P\} = \{\Sigma \in X \mid A \subset \Sigma \subset B\},$$

and similarly for Q and R . We conclude the proof using the following result.

Lemma 5.10 ([7, Proposition 1]). *Let P, Q, R be three elements of X^+ with pairwise intersections all equal to an $(m-d)$ -dimensional subspace A . Then there exists a unique morphism $f : \mathbb{P}^1 \rightarrow X^+$ of degree d such that*

$$f(0) = P, f(1) = Q, f(\infty) = R.$$

Note that $(A, B) \in I_d$ implies $B \subset V \subset V^+$. Since B is the span of the curve constructed by the former lemma, this curve factors through X . \square

As a consequence of Proposition 5.8 we now obtain that the quantum Pieri rule for odd symplectic Grassmannians only involves classical terms and degree one Gromov-Witten invariants.

Corollary 5.11. *Let $1 \leq p \leq 2n + 1 - m$ be an integer. Then no term of q -degree 2 or higher appears in a quantum product $\tau'_p * \alpha$ with $\alpha = \sigma_\lambda$ or $\alpha = \tau'_\mu$.*

Proof. The terms in q^d in the product $\tau'_p * \alpha$ are of the form $\langle \tau'_p, \alpha, \beta \rangle_{X,d,3} \beta^\vee$ with $\beta = \sigma_\nu$ or $\beta = \tau'_\xi$. By Proposition 5.8, the corresponding Gromov-Witten invariant vanishes for any degree d such that the staircase partition $\rho_d \not\subset (p)$, therefore, as soon as $d \geq 2$. \square

5.5. A quantum-to-classical principle. Proposition 5.8 computes some three-pointed degree d Gromov-Witten invariants of $X = \text{IG}(m, 2n + 1)$ as numbers of points in the intersection of three given subvarieties of the variety I_d^+ of Ker/Span pairs of dimension $(m - d, m + d)$ in $V^+ \cong \mathbb{C}^{2n+2}$. We now focus on degree one invariants and prove a ‘quantum-to-classical principle’ for them, that is, we express such invariants as numbers of intersection points of subvarieties of a larger odd symplectic Grassmannian $\tilde{X} = \text{IG}(m + 1, 2n + 3)$.

Our first step is to relate our invariants to points in $\tilde{X}^+ = \text{IG}(m + 1, 2n + 4)$. We start by introducing some notation for \tilde{X} and \tilde{X}^+ . Denote by S a symplectic vector space of dimension 2, and write

$$\tilde{V} := V \oplus S, \quad \tilde{V}^+ := V^+ \oplus S.$$

Clearly the antisymmetric form of maximal rank ω , respectively ω^+ , extends to an antisymmetric form of maximal rank on \tilde{V} , respectively \tilde{V}^+ , and both extend the symplectic form chosen on S . Thus we may define

$$\tilde{X} := \text{IG}(m + 1, \tilde{V}) \quad \text{and} \quad \tilde{X}^+ := \text{IG}(m + 1, \tilde{V}^+).$$

We also introduce some particular subvarieties of \tilde{X} and \tilde{X}^+ . Let f be a non-zero element of S , F_\bullet^+ be an Sp_{2n+2} -flag in V^+ , and λ be an $(n + 1 - m)$ -strict partition of X^+ . Consider the *augmented* Sp_{2n+4} -flag \tilde{F}_\bullet^+ :

$$\{0\} \subset \tilde{F}_1^+ = \mathbb{C}f \subset \tilde{F}_2^+ = \mathbb{C}f \oplus F_1^+ \subset \cdots \subset \tilde{F}_{2n+3}^+ = \mathbb{C}f \oplus F_{2n+2}^+ \subset \tilde{V}^+$$

and the associated Schubert variety

$$\tilde{X}_\lambda^+ := \left\{ \tilde{\Sigma}^+ \in \tilde{X}^+ \mid \dim \left(\tilde{\Sigma}^+ \cap \tilde{F}_{p_j(\lambda)+1}^+ \right) \geq j, \forall 1 \leq j \leq m \right\}.$$

For simplicity we have omitted the flag \tilde{F}_\bullet^+ from the notation on the left-hand side.

Proposition 5.12. *Let α, β, γ be elements of the cohomology basis $\{\sigma_\lambda, \tau'_\mu\}$ of $X = \text{IG}(m, 2n + 1)$ such that*

$$\text{codim } \alpha + \text{codim } \beta + \text{codim } \gamma = \dim M_{1,3}(X) = \dim X + 2n + 2 - m.$$

Following Proposition 5.5 we denote by $\sigma_\lambda^+, \sigma_\mu^+, \sigma_\nu^+$ the Schubert classes of the symplectic Grassmannian $X^+ = \text{IG}(m, 2n + 2)$ such that

$$j^* \sigma_\lambda^+ = \alpha; j^* \sigma_\mu^+ = \beta; j^* \sigma_\nu^+ = \gamma.$$

If $\ell(\lambda) + \ell(\mu) + \ell(\nu) \leq 2m + 1$, then

$$\langle \alpha, \beta, \gamma \rangle_{X,1,3} = \frac{1}{2} \# \left(\tilde{X}_\lambda^+ \cap \tilde{X}_\mu^+ \cap \tilde{X}_\nu^+ \cap \tilde{X} \right),$$

where \tilde{X}_λ^+ , \tilde{X}_μ^+ and \tilde{X}_ν^+ are associated to generic flags.

Again, providing Corollary 3.4 holds, a proof of this proposition was already given in [47, Section 5.2], and we limit ourselves to giving the main steps of the proof.

Sketch of proof of Proposition 5.12. We first show that for a generic choice of flags we have a well-defined map

$$\begin{aligned} \phi : \tilde{X}_\lambda^+ \cap \tilde{X}_\mu^+ \cap \tilde{X}_\nu^+ \cap \tilde{X} &\rightarrow I_1^+(\sigma_\lambda^+) \cap I_1^+(\sigma_\mu^+) \cap I_1^+(\sigma_\nu^+) \cap I_1 \\ \tilde{\Sigma}^+ &\mapsto \left(\tilde{\Sigma}^+ \cap V^+, (\tilde{\Sigma}^+ \oplus S) \cap V^+ \right). \end{aligned}$$

The group $G := \mathrm{Sp}(V^+) \oplus \mathrm{Sp}(S)$ acts on \tilde{X}^+ with the following orbits:

- (1) $\mathcal{O}_1 \cong \mathrm{IG}(m+1, V^+)$,
- (2) $\mathcal{O}_2 \cong \mathrm{IG}(m, V^+) \times \mathbb{P}(S)$,
- (3) $\mathcal{O}_3 = \left\{ \tilde{\Sigma}^+ \in \tilde{X}^+ \mid \dim(\tilde{\Sigma}^+ \cap V^+) = m \text{ and } \tilde{\Sigma}^+ \cap S = \{0\} \right\}$,
- (4) $\mathcal{O}_4 = \left\{ \tilde{\Sigma}^+ \in \tilde{X}^+ \mid \dim(\tilde{\Sigma}^+ \cap V^+) = m-1 \right\}$.

Using the assumptions on λ, μ, ν , a dimension count shows that the intersection $\tilde{X}_\lambda^+ \cap \tilde{X}_\mu^+ \cap \tilde{X}_\nu^+ \cap \tilde{X}$ must be contained in the open orbit \mathcal{O}_4 , which proves that the map ϕ is well-defined.

It remains to show that for (A, B) in the intersection

$$I_1^+(\sigma_\lambda^+) \cap I_1^+(\sigma_\mu^+) \cap I_1^+(\sigma_\nu^+) \cap I_1,$$

the inverse image $\phi^{-1}(A, B)$ consists in exactly two points. In the proof of Proposition 5.8 we saw that we must have $A = B \cap B^\perp$, so that $U := B/A \oplus S$ is a four-dimensional symplectic vector space. Moreover the corresponding Lagrangian Grassmannian $\mathrm{IG}(2, U)$ identifies with a subvariety of \tilde{X}^+ , namely

$$\mathrm{IG}(2, U) \cong \left\{ \tilde{\Sigma}^+ \mid A \subset \tilde{\Sigma}^+ \subset B \oplus S \right\}.$$

In the proof of Proposition 5.8 we also saw that there exists a unique triple of pairwise distinct subspaces $\Sigma_1^+ \in X_\lambda^+ \cap X$, $\Sigma_2^+ \in X_\mu^+ \cap X$ and $\Sigma_3^+ \in X_\nu^+ \cap X$ such that $A \subset \Sigma_j^+ \subset B$ for $j = 1, 2, 3$. Moreover by a Schubert calculus argument it is easy to see that there exist exactly two elements of $\mathrm{IG}(2, U)$ which are incident to all three of $\Sigma_1/A \oplus \mathbb{C}f$, $\Sigma_2/A \oplus \mathbb{C}g$, and $\Sigma_3/A \oplus \mathbb{C}h$, where f, g, h are generic elements of S . Thus there also exist two elements $\tilde{\Sigma}_{j=1,2}^+ \in \tilde{X}^+$ such that $A \subset \tilde{\Sigma}_j^+ \subset B \oplus S$ and the dimensions

$$\dim \left(\tilde{\Sigma}_j^+ \cap \Sigma_1 \oplus \mathbb{C}f \right), \dim \left(\tilde{\Sigma}_j^+ \cap \Sigma_2 \oplus \mathbb{C}g \right), \dim \left(\tilde{\Sigma}_j^+ \cap \Sigma_3 \oplus \mathbb{C}h \right)$$

are all at least equal to m . Clearly the $\tilde{\Sigma}_j^+$ belong to the quadruple intersection $\tilde{X}_\lambda^+ \cap \tilde{X}_\mu^+ \cap \tilde{X}_\nu^+ \cap \tilde{X}$.

Conversely if $\tilde{\Sigma}^+ \in \tilde{X}_\lambda^+ \cap \tilde{X}_\mu^+ \cap \tilde{X}_\nu^+ \cap \tilde{X}$ is such that $A \subset \tilde{\Sigma}_j^+ \subset B \oplus S$, then we must have $A = \tilde{\Sigma}^+ \cap V^+$ and $B = (\tilde{\Sigma}^+ \oplus S) \cap V^+$, and one can also show that $\Sigma_1 = (\tilde{\Sigma}^+ \oplus \mathbb{C}f) \cap V^+$, and similarly for Σ_2 and Σ_3 . This implies that the dimensions

$$\dim \left(\tilde{\Sigma}_j^+ \cap \Sigma_1 \oplus \mathbb{C}f \right), \dim \left(\tilde{\Sigma}_j^+ \cap \Sigma_2 \oplus \mathbb{C}g \right), \dim \left(\tilde{\Sigma}_j^+ \cap \Sigma_3 \oplus \mathbb{C}h \right)$$

are all at least equal to m , thus $\phi^{-1}(A, B)$ consists in exactly two points. \square

We may now state our quantum-to-classical principle, which is an immediate corollary of Proposition 5.12. In the following theorem we denote by $\tilde{\sigma}_u$ the Schubert class of $\tilde{X} = \text{IG}(m+1, 2n+3)$ associated with an $(n-m)$ -strict partition u , and by $\tilde{\tau}'_v$ the Schubert class of \tilde{X} associated with an $(n+1-m)$ -strict partition v .

Theorem 5.13. *Let α, β, γ be elements of the cohomology basis $\{\sigma_u, \tau'_v\}$ of $X = \text{IG}(m, 2n+1)$ such that*

$$\text{codim } \alpha + \text{codim } \beta + \text{codim } \gamma = \dim X + 2n + 2 - m.$$

Denote by $\sigma_\lambda^+, \sigma_\mu^+, \sigma_\nu^+$ the Schubert classes of $X^+ = \text{IG}(m, 2n+2)$ such that

$$j^* \sigma_\lambda^+ = \alpha; j^* \sigma_\mu^+ = \beta; j^* \sigma_\nu^+ = \gamma.$$

If $\ell(\lambda) + \ell(\mu) + \ell(\nu) \leq 2m + 1$, then

$$\langle \alpha, \beta, \gamma \rangle_{X,1,3} = \frac{1}{2} \langle \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma} \rangle_{\tilde{X},0,3},$$

where

$$\tilde{\alpha} = \begin{cases} \tilde{\sigma}_u & \text{if } \alpha = \sigma_u, \\ \tilde{\tau}'_v & \text{if } \alpha = \tau'_v, \end{cases}$$

and similarly for $\tilde{\beta}$ and $\tilde{\gamma}$.

Remark 5.14. The condition $\ell(\lambda) + \ell(\mu) + \ell(\nu) \leq 2m + 1$ will always be satisfied in the case where one of the partitions, say ν , has length one, i.e. $\nu = (p, 0, \dots, 0)$ for some $1 \leq p \leq 2n + 1 - m$. Keeping in mind the vanishing of invariants of degree larger than one, see Corollary 5.11, it follows that one may use Theorem 5.13 to deduce a *quantum Pieri formula* for the product of a class σ_u or τ'_v by the classes $\tau'_1, \dots, \tau'_{2n+1-m}$.

5.6. Quantum presentation. In this section we give a presentation for the quantum cohomology ring of $\text{IG}(m, 2n+1)$ in terms of the classes $\tau'_1, \tau'_2, \dots, \tau'_{2n+1-m}$ and the quantum parameter q . This presentation is a quantum version of the classical presentation obtained in [47], which we recall here.

Definition 5.15. For $r \geq 1$, define

$$d_r := \det(\tau'_{1+j-i})_{1 \leq i, j \leq r} \quad \text{and} \quad b_r := (\tau'_r)^2 + 2 \sum_{i \geq 1} (-1)^i \tau'_{r+i} \tau'_{r-i},$$

with the convention that $\tau'_0 := 1$ and $\tau'_p := 0$ if $p < 0$ or $p > 2n + 1 - m$.

The following result is proved in the second author's PhD thesis, see [47, Prop. 2.12]. However it is in French, therefore we repeat the proof here.

Proposition 5.16. *The cohomology ring $H^*(\text{IG}(m, 2n+1), \mathbb{Z})$ is generated by the classes $(\tau'_p)_{1 \leq p \leq 2n+1-m}$ and the relations are*

$$\begin{aligned} d_r &= 0 \text{ for } m+1 \leq r \leq 2n+2-m, \\ b_r &= 0 \text{ for } n+2-m \leq r \leq n. \end{aligned}$$

Proof. Let \mathcal{S}_X denote the tautological bundle of $X = \text{IG}(m, 2n+1)$, and \mathcal{S}_{X^+} that of $X^+ = \text{IG}(m, 2n+2)$. Similarly we denote by \mathcal{Q}_X and \mathcal{Q}_{X^+} the corresponding quotient bundles. We know that the Chern classes of \mathcal{Q}_{X^+} generate the cohomology of X^+ , see [8, Section 1.2], and it follows from Proposition 5.5 that the restriction

map $H^*(X^+, \mathbb{Z}) \rightarrow H^*(X, \mathbb{Z})$ is surjective, therefore, the Chern classes $c_p(\mathcal{Q}_X)$ for $1 \leq p \leq 2n+2-m$ generate the cohomology ring of X . Moreover, since \mathcal{Q}_X has rank $2n+1-m$, the Chern class $c_{2n+2-m}(\mathcal{Q}_X)$ vanishes, hence $H^*(X, \mathbb{Z})$ is generated by the classes $c_p(\mathcal{Q}_X) = \tau'_p$ for $1 \leq p \leq 2n+1-m$.

To find the relations we use the method introduced in [8] to obtain presentations of the cohomology of isotropic Grassmannians. Namely, let $R := \mathbb{Z}[a_1, \dots, a_{2n+1-m}]$ be a graded ring with $\deg a_i = i$. Set $a_0 = 0$ and $a_i = 0$ if $i > 2n+1-m$. We let $\delta_0 = 1$ and $\delta_r = \det(a_{1+j-i})_{1 \leq i, j \leq r}$ for $r > 0$. We also set $\beta_r = a_r^2 + 2 \sum_{i \geq 1} (-1)^i a_{r+i} a_{r-i}$ for $r \geq 0$. Now we define a (surjective) graded ring homomorphism $\phi: R \rightarrow H^*(X, \mathbb{Z})$ by setting $\phi(a_i) = \tau'_i$ for $1 \leq i \leq 2n+1-m$. We start by checking that the relations hold, i.e., that $\phi(\delta_r) = 0$ for $r > m$ and $\phi(\beta_r) = 0$ for $n+2-m \leq r \leq n$.

Expanding each determinant δ_r with respect to the first row yields the identity of formal series:

$$\left(\sum_{i=0}^{2n-1} a_i t^i \right) \left(\sum_{i \geq 0} (-1)^i \delta_i t^i \right) = 1.$$

Noticing that the image by ϕ of the leftmost factor is the total Chern class $c(\mathcal{Q}_X)$, and that $c(\mathcal{Q}_X)c(\mathcal{S}_X) = 1$, we deduce that

$$\phi(\mathcal{S}_X) = \left(\sum_{i \geq 0} (-1)^i \phi(\delta_i) t^i \right).$$

Since \mathcal{S}_X has rank m we obtain that $\phi(\delta_r) = d_r = 0$ for $r > m$.

The quadratic relations b_r come from similar relations on $X^+ = \text{IG}(m, 2n+2)$. Indeed it follows from the presentation of $H^*(X^+, \mathbb{Z})$ proved in [8] that the relations

$$\sigma_r^+ + 2 \sum_{i \geq 1} (-1)^i \sigma_{r+i}^+ \sigma_{r-i}^+ = 0$$

hold for $n+2-m \leq r \leq n$, and their restriction to X gives $b_r = 0$.

It now remains to prove that both sets of relations generate the kernel of ϕ . This is done using [8, Lemma 1.1], which states that to do so we only need to check that

- $H^*(X, \mathbb{Z})$ is a free \mathbb{Z} -module of rank $\deg \left(\frac{\prod_r \delta_r \prod_r \beta_r}{\prod_i a_i} \right)$;
- for any field K the K -vector space $(R/I) \otimes_{\mathbb{Z}} K$ has finite dimension, where I is the ideal generated by the relations δ_r and b_r .

The rank of $H^*(X, \mathbb{Z})$ is easily seen to be equal to $2^{m-1} \binom{n}{m} \frac{2n+2-m}{n+1-m}$ as required, see [47, Section 2.2.3]. To check the second condition we prove that R/I is a quotient of $R/(\delta_{m+1}, \dots, \delta_{2n+1})$, which is a free \mathbb{Z} -module of finite rank, see [8, Lemma 1.2]. This means that we need to prove that the relations δ_r for $2n+3-m \leq r \leq 2n+1$ also belong to the ideal I . We use the identities of formal series

$$\left(\sum_{i=0}^{2n+1-m} a_i t^i \right) \left(\sum_{i=0}^{2n+1-m} (-1)^i a_i t^i \right) = \left(\sum_{i=0}^{2n+1-m} (-1)^i \beta_i t^{2i} \right)$$

and

$$\left(\sum_{i=0}^{2n+1-m} (-1)^i a_i t^i \right) \left(\sum_{i \geq 0} \delta_i t^i \right) = 1,$$

so that

$$\left(\sum_{i=0}^{2n+1-m} a_i t^i \right) = \left(\sum_{i=0}^{2n+1-m} (-1)^i \beta_i t^{2i} \right) \left(\sum_{i \geq 0} \delta_i t^i \right).$$

Modding out by the relations in I gives

$$\left(\sum_{i=0}^{2n+1-m} a_i t^i \right) \equiv \left(\sum_{i=0}^{n+1-m} (-1)^i \beta_i t^{2i} + \sum_{i=n+1}^{2n+1-m} (-1)^i \beta_i t^{2i} \right) \left(\sum_{i=0}^m \delta_i t^i + \sum_{i \geq 2n+3-m} \delta_i t^i \right).$$

Looking at the terms in degrees $2n+3-m$ to $2n+1$ we obtain $0 \equiv \delta_{2n+3-m} \equiv \dots \equiv \delta_{2n+1}$ as claimed, which concludes the proof. \square

Theorem 5.17. *The quantum cohomology ring $\mathrm{QH}(\mathrm{IG}(m, 2n+1), \mathbb{C})$ is generated by the classes $(\tau'_p)_{1 \leq p \leq 2n+1-m}$ and the quantum parameter q , and the relations are*

$$\begin{aligned} d_r &= 0 \text{ for } m+1 \leq r \leq 2n+1-m, \\ d_{2n+2-m} &= \begin{cases} 0 & \text{if } m \text{ is even,} \\ -q & \text{if } m \text{ is odd,} \end{cases} \\ b_s &= (-1)^{2n+1-m-s} q \tau'_{2s-2n-2+m} \text{ for } n+2-m \leq s \leq n, \end{aligned}$$

with the convention that $\tau'_p = 0$ if $p < 0$.

Proof. To obtain this quantum presentation from the classical presentation of Proposition 5.16, we use the well-known method due to Siebert and Tian, see [54, Prop. 2.2]. This means that we replace the cup product with the quantum product in the classical relations d_r and b_s . The quantum presentation is then obtained by considering these new relations instead of the original ones.

Let us start by deforming the relations b_s for $n+2-m \leq s \leq n$. We have

$$\begin{aligned} b_s^{\mathrm{quant}} &= \tau'_s * \tau'_s + 2 \sum_{i=1}^{2n+1-m-s} (-1)^i \tau'_{s+i} * \tau'_{s-i} \\ &= q \sum_{\lambda} \left(\langle j^* \sigma_s^+, j^* \sigma_s^+, j^* \sigma_{\lambda}^+ \rangle_{X,1,3} \right. \\ &\quad \left. + 2 \sum_{i=1}^{2n+1-m-s} (-1)^i \langle j^* \sigma_{s+i}^+, j^* \sigma_{s-i}^+, j^* \sigma_{\lambda}^+ \rangle_{X,1,3} \right) (j^* \sigma_{\lambda}^+)^{\vee}, \end{aligned}$$

where $j : X \hookrightarrow X^+ = \mathrm{IG}(m, 2n+2)$ is the natural inclusion. Indeed, the classical part of b_s^{quant} is the classical relation b_s , which vanishes, and by Proposition 5.5 we have that $\tau'_p = j^* \sigma_p^+$. We now use the quantum-to-classical principle from Theorem 5.13.

Write $\tilde{X} := \mathrm{IG}(m+1, 2n+3)$, $\tilde{X}^+ := \mathrm{IG}(m+1, 2n+4)$, and denote by $\tilde{j} : \tilde{X} \hookrightarrow \tilde{X}^+$ the natural inclusion. We also denote by $\tilde{\sigma}_{\lambda}^+$ the Schubert class of \tilde{X}^+ associated with an $(n+1-m)$ -strict partition λ with $\lambda_1 \leq 2n+3-m$. Finally, we introduce the subvarieties \tilde{Y} and \tilde{Z} of \tilde{X} and the cohomology classes $\{\tilde{\sigma}_{\mu}, \tilde{\tau}'_{\nu}\}$ and $\{\tilde{\sigma}'_{\mu}, \tilde{\tau}_{\nu}\}$ of \tilde{X} as in Section 5.3, simply replacing X with \tilde{X} and adding \sim everywhere. It follows

from Theorem 5.13 that

$$\begin{aligned} b_s^{\text{quant}} &= \frac{1}{2}q \sum_{\lambda} \left(\langle \tilde{j}^* \tilde{\sigma}_s^+, \tilde{j}^* \tilde{\sigma}_s^+, \tilde{j}^* \tilde{\sigma}_{\lambda}^+ \rangle_{\tilde{X},0,3} \right. \\ &\quad \left. + 2 \sum_{i=1}^{2n+1-m-s} (-1)^i \langle \tilde{j}^* \tilde{\sigma}_{s+i}^+, \tilde{j}^* \tilde{\sigma}_{s-i}^+, \tilde{j}^* \tilde{\sigma}_{\lambda}^+ \rangle_{\tilde{X},0,3} \right) (j^* \sigma_{\lambda}^+)^{\vee} \\ &= \frac{1}{2}q \sum_{\lambda} \left(\int_{\tilde{X}} \left(\tilde{\tau}'_s \tilde{\tau}'_s + 2 \sum_{i=1}^{2n+1-m-s} (-1)^i \tilde{\tau}'_{s+i} \tilde{\tau}'_{s-i} \right) \cup \tilde{j}^* \tilde{\sigma}_{\lambda}^+ \right) (j^* \sigma_{\lambda}^+)^{\vee}. \end{aligned}$$

In $H^*(\tilde{X}, \mathbb{Z})$, by Proposition 5.16, the classical relation

$$\tilde{\tau}'_s \tilde{\tau}'_s + 2 \sum_{i=1}^{2n+2-m-s} (-1)^i \tilde{\tau}'_{s+i} \tilde{\tau}'_{s-i} = 0$$

is satisfied, thus

$$\tilde{\tau}'_s \tilde{\tau}'_s + 2 \sum_{i=1}^{2n+1-m-s} (-1)^i \tilde{\tau}'_{s+i} \tilde{\tau}'_{s-i} = 2(-1)^{2n+1-m-s} \tilde{\tau}'_{2n+2-m} \tilde{\tau}'_{2s-2n-2+m}.$$

Replacing in the expression of b_s^{quant} , we obtain

$$\begin{aligned} b_s^{\text{quant}} &= (-1)^{2n+1-m-s} q \sum_{\lambda} \left(\int_{\tilde{X}} \tilde{\tau}'_{2n+2-m} \cup \tilde{\tau}'_{2s-2n-2+m} \cup \tilde{j}^* \tilde{\sigma}_{\lambda}^+ \right) (j^* \sigma_{\lambda}^+)^{\vee} \\ &= (-1)^{2n+1-m-s} q \sum_{\lambda} \left(\int_{\tilde{X}} \tilde{j}^* (\tilde{\sigma}_{2n+2-m}^+ \cup \tilde{\sigma}_{2s-2n-2+m}^+) \cup \tilde{j}^* \tilde{\sigma}_{\lambda}^+ \right) (j^* \sigma_{\lambda}^+)^{\vee}. \end{aligned}$$

To simplify notation we introduce the integer $p := 2s - 2n - 2 + m$. We use the classical Pieri rule in \tilde{X}^+ to compute the product $\tilde{\sigma}_{2n+2-m}^+ \cup \tilde{\sigma}_p^+$, see [8, Theorem 1.1]:

$$\tilde{\sigma}_{2n+2-m}^+ \cup \tilde{\sigma}_p^+ = \tilde{\sigma}_{2n+2-m,p}^+ + 2\tilde{\sigma}_{2n+3-m,p-1}^+.$$

We now prove that the second summand will not contribute to b_s^{quant} . Indeed, let us prove that $\tilde{j}^* \tilde{\sigma}_{\lambda}^+ = 0$ when $\lambda_1 = 2n + 3 - m$. From Proposition 5.4 applied to \tilde{X} we have that $\tilde{j}^* \tilde{\sigma}_{\lambda}^+ = 0$ when $\tilde{\sigma}_{\lambda}^+ \cup \tilde{\sigma}_{1^{m+1}}^+ = 0$. The latter cup product can be computed using the Pieri rule of [53]. Namely, let $\alpha = (\alpha^t, \alpha^b)$ be the pair of partitions associated with the $(n + 1 - m)$ -strict partition λ following the correspondence introduced in Section 5.2. Clearly, $\lambda_1 = 2n + 3 - m$ implies that $\alpha_1^b = n + 2$, i.e., the first part of α^b is maximal. Therefore, $\tilde{\sigma}_{\alpha}^+ \cup \tilde{\sigma}_{1^{m+1}}^+ = 0$ by the Pieri rule. For more details we refer to [47, Section 2.3]. Thus we indeed have $\tilde{j}^* \tilde{\sigma}_{\lambda}^+ = 0$ when $\lambda_1 = 2n + 3 - m$. It follows that

$$b_s^{\text{quant}} = (-1)^{2n+1-m-s} q \sum_{\lambda} \left(\int_{\tilde{X}} \tilde{j}^* \tilde{\sigma}_{2n+2-m,p}^+ \cup \tilde{j}^* \tilde{\sigma}_{\lambda}^+ \right) (j^* \sigma_{\lambda}^+)^{\vee}.$$

Let us now express the restricted class $\tilde{j}^* \tilde{\sigma}_{2n+2-m,p}^+$ in terms of the cohomology basis $\{\tilde{\sigma}_{\mu}, \tilde{\tau}'_{\nu}\}$. As before, from Proposition 5.4 it follows that we first need to compute the cup product $\tilde{\sigma}_{2n+2-m,p}^+ \cup \tilde{\sigma}_{1^{m+1}}^+$ in $H^*(\tilde{X}^+, \mathbb{Z})$.

We use again the classical Pieri rule for multiplication by the classes $\tilde{\sigma}_r^+$ from [53], noting that the strict partition $(2n+2-m, p)$ corresponds to the pair of partitions

$$\alpha^b = (n+1, p+m-1-n), \alpha^t = (n+3-m, \dots, 4, 3)$$

if $p > n+1-m$, and to

$$\alpha^b = (n+1), \alpha^t = (n+3-m, n+2-m, \dots, n+4-m-p, n+2-m-p, \dots, 3, 2)$$

otherwise. We deduce that

$$\tilde{\sigma}_{2n+2-m,p}^+ \cup \tilde{\sigma}_{1^{m+1}}^+ = \tilde{\sigma}_{2n+3-m,p+1,1^{m-1}}^+$$

if $p \leq 2n+1-2m$, and otherwise

$$\tilde{\sigma}_{2n+2-m,p}^+ \cup \tilde{\sigma}_{1^{m+1}}^+ = \tilde{\sigma}_{2n+3-m,p+1,1^{m-1}}^+ + \tilde{\sigma}_{2n+3-m,2n+2-m,1^{p-2n-2+2m}}^+.$$

Hence in the first case,

$$\tilde{j}^* \tilde{\sigma}_{2n+2-m,p}^+ = [\tilde{X}_{2n+2-m,p}] = \tilde{\tau}_{p+1,1^{m-1}},$$

while in the second case,

$$\tilde{j}^* \tilde{\sigma}_{2n+2-m,p}^+ = \tilde{\tau}_{p+1,1^{m-1}} + \tilde{\tau}_{2n+2-m,1^{p-2n-2+2m}}.$$

By Poincaré duality, see Proposition 1.10, we know that $\int_{\tilde{X}} \tilde{\tau}_{p+1,1^{m-1}} \cup \tilde{j}^* \tilde{\sigma}_\lambda^+$ is equal to 1 if $\lambda_{m+1} = 0$ and λ is the Poincaré dual partition to $(p+1, 1^{m-1})$ in $\tilde{Z} = \text{IG}(m, 2n+2)$. Otherwise it is equal to zero, and similarly for $\int_{\tilde{X}} \tilde{\tau}_{2n+2-m,1^{p-2n-2+2m}} \cup \tilde{j}^* \tilde{\sigma}_\lambda^+$.

Using [8, Section 4.4], we see that the Poincaré dual partition to $(p+1, 1^{m-1})$ is

$$(2n+1-m, 2n-m, \dots, 2n+3-2m, 2n+2-2m-p)$$

if $p \leq 2n+1-2m$, and

$$(2n+1-m, 2n-m, \dots, p+2, p, p-1, \dots, 2n+2-2m, 1)$$

otherwise. Similarly, the Poincaré dual partition to $(2n+2-m, 1^{p-2n-2+2m})$ is

$$(2n+1-m, 2n-m, \dots, p+1, p-1, p-2, \dots, 2n+2-2m).$$

So if $p \leq 2n+1-2m$ then

$$\begin{aligned} b_s^{\text{quant}} &= (-1)^{2n+1-m-s} q(j^* \sigma_{2n+1-m, 2n-m, \dots, 2n+3-2m, 2n+2-2m-p}^+)^{\vee} \\ &= (-1)^{2n+1-m-s} q \sigma_{2n-m, 2n-1-m, \dots, 2n+2-2m, 2n+1-2m-p}^{\vee}, \end{aligned}$$

and otherwise,

$$\begin{aligned} b_s^{\text{quant}} &= (-1)^{2n+1-m-s} q(j^* \sigma_{2n+1-m, \dots, p+2, p, \dots, 2n+2-2m, 1}^+)^{\vee} \\ &\quad + (-1)^{2n+1-m-s} q(j^* \sigma_{2n+1-m, \dots, p+1, p-1, \dots, 2n+2-2m}^+)^{\vee} \\ &= (-1)^{2n+1-m-s} q \sigma_{2n-m, \dots, p+1, p-1, \dots, 2n+1-2m}^{\vee} \\ &\quad + (-1)^{2n+1-m-s} q(\tau'_{2n+1-m, \dots, p+1, p-1, \dots, 2n+2-2m})^{\vee}. \end{aligned}$$

Moreover, if $p \leq 2n+1-2m$ then the Poincaré dual partition to the partition $(2n-m, 2n-1-m, \dots, 2n+2-2m, 2n+1-2m-p)$ in Y is (p) . On the other hand if $p \geq 2n+2-2m$ the Poincaré dual partition to $(2n-m, \dots, p+1, p-1, \dots, 2n+1-2m)$ in Y is (p) , while the Poincaré dual partition to $(2n+1-m, \dots, p+1, p-1, \dots, 2n+2-2m)$ in Z is $(1^{p-2n-2+2m})$. Hence

$$b_s^{\text{quant}} = \begin{cases} (-1)^{2n+1-m-s} q \sigma'_p & \text{if } p \leq 2n+1-2m, \\ (-1)^{2n+1-m-s} q(\sigma'_p + \tau_{1^{p-2n-2+2m}}) & \text{otherwise.} \end{cases}$$

Let us compare the right-hand side of the above equation with the pullback $j^*\sigma_p^+$ using Proposition 5.4 and the classical Pieri rule from [53]. Note that the strict partition (p) corresponds to the pair of partitions given by

$$\begin{cases} \alpha^b = (p + m - 1 - n), \alpha^t = (n + 2 - m, n + 1 - m, \dots, 2) & \text{if } p > n + 1 - m, \\ \alpha^b = \emptyset, \alpha^t = (n + 1 - m, n - m, \dots, 1) & \text{otherwise.} \end{cases}$$

We obtain

$$\sigma_p^+ \cup \sigma_{1^m}^+ = \begin{cases} \sigma_{p+1, 1^{m-1}}^+ & \text{if } p \leq 2n + 1 - 2m, \\ \sigma_{p+1, 1^{m-1}}^+ + \sigma_{2n+2-m, 1^{p-2n-2+2m}}^+ & \text{otherwise.} \end{cases}$$

It follows from Proposition 5.4 that

$$j^*\sigma_p^+ = \begin{cases} \sigma_p' & \text{if } p \leq 2n + 1 - 2m, \\ \sigma_p' + \tau_{1^{p-2n-2+2m}} & \text{otherwise.} \end{cases}$$

Since by definition $j^*\sigma_p^+ = \tau_p'$ we finally get that

$$b_s^{\text{quant}} = (-1)^{2n+1-m-s} q \tau_{2s-2n-2+m}'$$

as claimed.

In the second part of this proof, let us now deform the relations d_r for $m + 1 \leq r \leq 2n + 2 - m$. Since $\deg q = 2n + 2 - m$, only the relation d_{2n+2-m} is potentially modified. We consider the determinantal identity

$$d_{2n+2-m}^{\text{quant}} = \sum_{p=1}^{2n+1-m} (-1)^{p+1} \tau_p' * d_{2n+2-p-m}.$$

The classical part of the right-hand side of the above equality being trivial, it follows that

$$d_{2n+2-m}^{\text{quant}} = q \sum_{p=1}^{2n+1-m} (-1)^{p+1} \langle \tau_p', d_{2n+2-m-p}, [pt] \rangle_{X,1,3}.$$

Moreover we have that for any $1 \leq s \leq m$, $d_s = j^*\sigma_{1^s}^+$, and $d_s = 0$ if $s > m$, see [47, Section 2.5]. This implies that

$$\begin{aligned} d_{2n+2-m}^{\text{quant}} &= q \sum_{p=2n+2-2m}^{2n+1-m} (-1)^{p+1} \langle \tau_p', d_{2n+2-m-p}, [pt] \rangle_{X,1,3} \\ &= q \sum_{s=1}^m (-1)^{2n+3-m-s} \langle \tau_{2n+2-m-s}', j^*\sigma_{1^s}^+, [pt] \rangle_{X,1,3}. \end{aligned}$$

We have $\tau_{2n+2-m-s}' = j^*\sigma_{2n+2-m-s}^+$, and the class of a point in X is

$$[pt] = \sigma_{2n-m, 2n-1-m, \dots, 2n+1-2m} = j^*\sigma_{2n+1-m, 2n-m, \dots, 2n+2-2m}^+$$

so using the quantum to classical principle we get

$$\begin{aligned}
 d_{2n+2-m}^{\text{quant}} &= \frac{1}{2}q \sum_{s=1}^m (-1)^{2n+3-m-s} \\
 &\quad \langle \tilde{j}^* \tilde{\sigma}_{2n+2-m-s}^+, \tilde{j}^* \tilde{\sigma}_{1^s}^+, \tilde{j}^* \tilde{\sigma}_{2n+1-m, 2n-m, \dots, 2n+2-2m}^+ \rangle_{\tilde{X}, 0, 3} \\
 &= \frac{1}{2}q \int_{\tilde{X}} \left(\sum_{s=1}^m (-1)^{2n+3-m-s} \tilde{j}^* \tilde{\sigma}_{2n+2-m-s}^+ \cup \tilde{j}^* \tilde{\sigma}_{1^s}^+ \right) \\
 &\quad \cup \tilde{j}^* \tilde{\sigma}_{2n+1-m, 2n-m, \dots, 2n+2-2m}^+,
 \end{aligned}$$

where the \sim indicates that we are considering classes in $\tilde{X}^+ = \text{IG}(m+1, 2n+4)$. Let us now look at the *classical* relation \tilde{d}_{2n+2-m} in \tilde{X} . Using a similar determinantal identity as before, we get

$$\begin{aligned}
 \tilde{d}_{2n+2-m} &= \sum_{p=1}^{2n+2-m} (-1)^{p+1} \tilde{\tau}'_p \cup \tilde{d}_{2n+2-p-m} \\
 &= \sum_{p=2n+1-2m}^{2n+2-m} (-1)^{p+1} \tilde{\tau}'_p \cup \tilde{d}_{2n+2-p-m} \\
 &= \sum_{s=0}^{m+1} (-1)^{2n+3-m-s} \tilde{\tau}'_{2n+2-m-s} \cup \tilde{d}_s \\
 &= \sum_{s=0}^{m+1} (-1)^{2n+3-m-s} \tilde{j}^* \tilde{\sigma}_{2n+2-m-s}^+ \cup \tilde{j}^* \tilde{\sigma}_{1^s}^+.
 \end{aligned}$$

Thus replacing in the expression for $d_{2n+2-m}^{\text{quant}}$, we get

$$\begin{aligned}
 d_{2n+2-m}^{\text{quant}} &= \frac{q}{2} \int_{\tilde{X}} \left(\tilde{d}_{2n+2-m} - (-1)^{2n+3-m} \tilde{j}^* \tilde{\sigma}_{2n+2-m}^+ \right. \\
 &\quad \left. - (-1)^{2n+2-2m} \tilde{j}^* \tilde{\sigma}_{2n+1-2m}^+ \cup \tilde{j}^* \tilde{\sigma}_{1^{m+1}}^+ \right) \cup \tilde{j}^* \tilde{\sigma}_{2n+1-m, 2n-m, \dots, 2n+2-2m}^+ \\
 &= (-1)^{2n+2-m} \frac{q}{2} \int_{\tilde{X}} \tilde{j}^* (\tilde{\sigma}_{2n+2-m}^+ \cup \tilde{\sigma}_{2n+1-m, 2n-m, \dots, 2n+2-2m}^+) \\
 &\quad + (-1)^{2n+1-2m} \frac{q}{2} \int_{\tilde{X}} \tilde{j}^* \tilde{\sigma}_{2n+1-2m}^+ \cup \tilde{j}^* (\tilde{\sigma}_{1^{m+1}}^+ \cup \tilde{\sigma}_{2n+1-m, \dots, 2n+2-2m}^+).
 \end{aligned}$$

Using the classical Pieri rule in \tilde{X}^+ , see [8, Theorem 1.1], we have

$$\begin{aligned}
 \tilde{\sigma}_{2n+2-m}^+ \cup \tilde{\sigma}_{2n+1-m, \dots, 2n+2-2m}^+ &= \tilde{\sigma}_{2n+2-m, 2n+1-m, \dots, 2n+2-2m}^+ \\
 &\quad + 2\tilde{\sigma}_{2n+3-m, 2n+1-m, \dots, 2n+3-2m, 2n+1-2m}^+.
 \end{aligned}$$

Moreover, using the classical Pieri rule from [53],

$$\begin{aligned}
 \tilde{\sigma}_{1^{m+1}}^+ \cup \tilde{\sigma}_{2n+1-m, 2n-m, \dots, 2n+2-2m}^+ &= \tilde{\sigma}_{2n+2-m, \dots, 2n+3-2m, 1}^+ \\
 &\quad + \tilde{\sigma}_{2n+3-m, 2n+1-m, \dots, 2n+3-2m}^+.
 \end{aligned}$$

Since we saw that $\tilde{j}^* \sigma_\lambda = 0$ if $\lambda_1 = 2n + 3 - m$, we obtain

$$\begin{aligned} d_{2n+2-m}^{\text{quant}} &= (-1)^{2n+2-m} \frac{q}{2} \int_{\tilde{X}} \tilde{j}^* \tilde{\sigma}_{2n+2-m, 2n+1-m, \dots, 2n+2-2m}^+ \\ &\quad + (-1)^{2n+1-2m} \frac{q}{2} \int_{\tilde{X}} \tilde{j}^* \tilde{\sigma}_{2n+1-2m}^+ \cup \tilde{j}^* \tilde{\sigma}_{2n+2-m, \dots, 2n+3-2m, 1}^+. \end{aligned}$$

The first integral is equal to 1 since $j^* \tilde{\sigma}_{2n+2-m, 2n+1-m, \dots, 2n+2-2m}^+$ is the class of a point in \tilde{X} . Thus it only remains to compute the following cup product using the classical Pieri rule [8, Theorem 1.1],

$$\begin{aligned} \tilde{\sigma}_{2n+1-2m}^+ \cup \tilde{\sigma}_{2n+2-m, \dots, 2n+3-2m, 1}^+ &= \tilde{\sigma}_{2n+2-m, \dots, 2n+2-2m}^+ \\ &\quad + \tilde{\sigma}_{2n+3-m, 2n+1-m, \dots, 2n+3-2m, 2n+1-2m}^+. \end{aligned}$$

Since $\tilde{j}^* \tilde{\sigma}_{2n+3-m, 2n+1-m, \dots, 2n+3-2m, 2n+1-2m}^+ = 0$ and $j^* \tilde{\sigma}_{2n+2-m, \dots, 2n+2-2m}^+$ is the class of a point in \tilde{X} , we get

$$\begin{aligned} d_{2n+2-m}^{\text{quant}} &= ((-1)^{2n+2-m} + (-1)^{2n+1-2m}) \frac{q}{2} \\ &= \begin{cases} 0 & \text{if } m \text{ is even,} \\ -q & \text{if } m \text{ is odd,} \end{cases} \end{aligned}$$

which concludes the proof of the theorem. \square

6. RECOLLECTIONS ON SEMIORTHOGONAL DECOMPOSITIONS

6.1. Semiorthogonal decompositions. We recall here basic definitions on semi-orthogonal decompositions in triangulated categories. For a convenient reference, see, for instance [24, Sections 1.2 and 1.4].

Let k be a field. Assume given a k -linear triangulated category \mathcal{D} , equipped with a shift functor $[1] : \mathcal{D} \rightarrow \mathcal{D}$. For two objects $A, B \in \mathcal{D}$ let $\text{Hom}_{\mathcal{D}}^\bullet(A, B)$ denote the graded k -vector space $\bigoplus_{i \in \mathbb{Z}} \text{Hom}_{\mathcal{D}}(A, B[i])$. Let $\mathcal{A} \subset \mathcal{D}$ be a *full triangulated subcategory*, that is, a full subcategory of \mathcal{D} which is closed under shifts and triangles.

Definition 6.1. A full triangulated subcategory \mathcal{A} of \mathcal{D} is called *right admissible* if the inclusion functor $\mathcal{A} \hookrightarrow \mathcal{D}$ has a right adjoint. Similarly, \mathcal{A} is called *left admissible* if the inclusion functor has a left adjoint. Finally, \mathcal{A} is *admissible* if it is both right and left admissible.

Definition 6.2. The *right orthogonal* of an admissible subcategory $\mathcal{A} \subset \mathcal{D}$ is the full subcategory $\mathcal{A}^\perp \subset \mathcal{D}$ of objects $B \in \mathcal{D}$ such that $\text{Hom}_{\mathcal{D}}(A, B) = 0$ for all $A \in \mathcal{A}$. The *left orthogonal* ${}^\perp \mathcal{A}$ is defined similarly.

If a full triangulated category $\mathcal{A} \subset \mathcal{D}$ is right admissible then every object $X \in \mathcal{D}$ fits into a distinguished triangle

$$\dots \longrightarrow Y \longrightarrow X \longrightarrow Z \longrightarrow Y[1] \longrightarrow \dots$$

with $Y \in \mathcal{A}$ and $Z \in \mathcal{A}^\perp$. One then says that there is a *semiorthogonal decomposition* of \mathcal{D} into the subcategories $(\mathcal{A}^\perp, \mathcal{A})$. More generally, assume given a sequence of full triangulated subcategories $\mathcal{A}_1, \dots, \mathcal{A}_n \subset \mathcal{D}$. Denote by $\langle \mathcal{A}_1, \dots, \mathcal{A}_n \rangle$ the triangulated subcategory of \mathcal{D} generated by $\mathcal{A}_1, \dots, \mathcal{A}_n$.

Definition 6.3. A sequence $(\mathcal{A}_1, \dots, \mathcal{A}_n)$ of full admissible subcategories of \mathcal{D} is called *semiorthogonal* if $\mathcal{A}_i \subset \mathcal{A}_j^\perp$ for $1 \leq i < j \leq n$. The sequence $(\mathcal{A}_1, \dots, \mathcal{A}_n)$ is called a *semiorthogonal decomposition* of \mathcal{D} if via inclusion \mathcal{D} is equivalent to the smallest full triangulated subcategory of \mathcal{D} containing all of them.

Lemma 6.4. [24, Lemma 1.61] *Any semiorthogonal sequence of full admissible triangulated subcategories $(\mathcal{A}_1, \dots, \mathcal{A}_n) \subset \mathcal{D}$ defines a semiorthogonal decomposition of \mathcal{D} , if and only if any object $A \in \mathcal{D}$ with $A \in \mathcal{A}_i^\perp$ for all $i = 1, \dots, n$ is trivial.*

6.2. Mutations. Let \mathcal{D} be a triangulated category and assume \mathcal{D} admits a semiorthogonal decomposition $\mathcal{D} = \langle \mathcal{A}, \mathcal{B} \rangle$.

Definition 6.5. The *left mutation* of \mathcal{B} through \mathcal{A} is defined to be $\mathbf{L}_{\mathcal{A}}(\mathcal{B}) := \mathcal{A}^\perp$. The *right mutation* of \mathcal{A} through \mathcal{B} is defined to be $\mathbf{R}_{\mathcal{B}}(\mathcal{A}) := {}^\perp\mathcal{B}$.

In practice, computing left (resp., right) mutation of an admissible subcategory \mathcal{A} of \mathcal{D} through its right (resp., left) orthogonal \mathcal{A}^\perp (resp., ${}^\perp\mathcal{A}$) amounts to the following. Denote $i : \mathcal{A} \rightarrow \mathcal{D}$ the embedding functor. Since the subcategory \mathcal{A} is admissible there are left and right adjoint functors $\mathcal{D} \rightarrow \mathcal{A}$ which we denote by i^* and $i^!$, respectively. Given an object $F \in \mathcal{D}$, define the *left mutation* $\mathbf{L}_{\mathcal{A}}(F)$ and the *right mutation* $\mathbf{R}_{\mathcal{A}}(F)$ of F through \mathcal{A} by

$$\mathbf{L}_{\mathcal{A}}(F) := \text{Cone}(i_* i^!(F) \rightarrow F), \quad \mathbf{R}_{\mathcal{A}}(F) := \text{Cone}(F \rightarrow i^! i^*(F))[-1].$$

These explicit formulae for mutations can be used to prove the following statement:

Lemma 6.6. [35, Corollary 2.9] *Given a semiorthogonal decomposition $\mathcal{D} = \langle \mathcal{A}, \mathcal{B} \rangle$, there are semiorthogonal decompositions $\mathcal{D} = \langle \mathbf{L}_{\mathcal{A}}(\mathcal{B}), \mathcal{A} \rangle$ and $\mathcal{D} = \langle \mathcal{B}, \mathbf{R}_{\mathcal{B}}(\mathcal{A}) \rangle$.*

Remark 6.7. In case the triangulated subcategory $\mathcal{A} \subset \mathcal{D}$ is generated by an exceptional object E , there are explicit formulas for adjoint functors $i^!, i_*$ of the embedding functor $i : \mathcal{A} \rightarrow \mathcal{D}$. Specifically, there are distinguished triangles:

$$\text{RHom}(E, F) \otimes E \rightarrow F \rightarrow \mathbf{L}_E(F), \quad \mathbf{R}_E(F) \rightarrow F \rightarrow \text{RHom}(F, E)^* \otimes E.$$

6.3. Exceptional collections in geometric categories. We first recall some notation. From now on, our ground field will be the field of complex numbers \mathbb{C} . Let S be a smooth complex projective variety, and $\mathcal{D}^b(S)$ be the bounded derived category of coherent sheaves on S . Given a morphism $f : S \rightarrow T$ between two smooth varieties, we usually write f_*, f^* for the corresponding derived functors of push-forwards and pull-backs between $\mathcal{D}^b(S)$ and $\mathcal{D}^b(T)$, and $R^i f_*$ (resp., $L^i f^*$) for the i -th cohomology of the functor f_* (resp., f^*). We denote $\mathcal{R}\mathcal{H}om_S$ the derived functor of the bi-functor of local $\mathcal{H}om$ -groups on S , and write $\mathcal{E}xt_S^i$ for its cohomology. We denote RHom_S the derived functor of the bi-functor of global $\mathcal{H}om$ -groups on S , and write Ext_S^i for its cohomology. Given an object $A \in \mathcal{D}^b(S)$, we denote $A^\vee := \mathcal{R}\mathcal{H}om_S(A, \mathcal{O}_S)$ its dual. Given two objects $A, B \in \mathcal{D}^b(S)$, we also abbreviate the graded vector space $\text{Hom}_{\mathcal{D}^b(S)}^\bullet(A, B) := \bigoplus_{i \in \mathbb{Z}} \text{Hom}_{\mathcal{D}^b(S)}(A, B[i])$ as $\text{Hom}_S^\bullet(A, B)$.

We denote $\tau_{<i}$ and $\tau_{\geq i}$ the truncation functors with respect to the standard t -structure on $\mathcal{D}^b(S)$, and $\mathcal{H}^i := \tau_{\leq i} \circ \tau_{\geq i} \circ [i]$ the i -th cohomology functor.

Convenient references for the definitions below are [24, Section 1.4] and [34].

Definition 6.8. An object $E \in \mathcal{D}^b(S)$ is said to be *exceptional* if there is an isomorphism of graded k -algebras

$$\text{Hom}_S^\bullet(E, E) = k.$$

Definition 6.9. A collection of exceptional objects (E_0, \dots, E_n) in $\mathcal{D}^b(S)$ is called *exceptional* if for $0 \leq i < j \leq n$ one has

$$\mathrm{Hom}_S^\bullet(E_j, E_i) = 0.$$

An exceptional collection (E_0, \dots, E_n) in $\mathcal{D}^b(S)$ is called *full* if the smallest full triangulated subcategory containing all E_i is $\mathcal{D}^b(S)$.

Let \mathcal{L} be an ample line bundle on S .

Definition 6.10. A *Lefschetz collection with respect to \mathcal{L}* is an exceptional collection which has a block structure

$$E_0, E_1, \dots, E_{\sigma_0}; E_0 \otimes \mathcal{L}, E_1 \otimes \mathcal{L}, \dots, E_{\sigma_1} \otimes \mathcal{L}; E_0 \otimes \mathcal{L}^{\otimes(m-1)}, \dots, E_{\sigma_{m-1}} \otimes \mathcal{L}^{\otimes(m-1)},$$

where $\sigma = (\sigma_0 \geq \sigma_1 \geq \dots \geq \sigma_{m-1} \geq 0)$ is a non-increasing sequence of non-negative integers. The block $(E_0, E_1, \dots, E_{\sigma_0})$ is called the *starting block*. If $\sigma_0 = \sigma_1 = \dots = \sigma_{m-1}$, then the corresponding Lefschetz collection is called *rectangular*.

Given a smooth variety S , let ω_S denote the canonical line bundle on S .

Lemma 6.11. [35, Lemma 2.11] *For S a smooth variety, assume given a semiorthogonal decomposition $\mathcal{D}^b(S) = \langle \mathcal{A}, \mathcal{B} \rangle$. Then*

$$\mathbf{L}_{\mathcal{A}}(\mathcal{B}) = \mathcal{B} \otimes \omega_S \quad \text{and} \quad \mathbf{R}_{\mathcal{A}}(\mathcal{B}) = \mathcal{A} \otimes \omega_S^{-1}.$$

6.4. Exceptional collections on horospherical varieties of Picard rank one.

Some of the varieties in Pasquier's classification have been known to have full Lefschetz exceptional collections. Specifically, the variety X^2 that is isomorphic to a smooth hyperplane section of the orthogonal Grassmannian $\mathrm{OG}(5, 10)$ has been treated in [34, Section 6.2]. The variety $X^3(n, 2)$ that is isomorphic to the odd symplectic Grassmannian $\mathrm{IG}(2, 2n + 1)$ has been treated in [48]. More recently, a full rectangular Lefschetz exceptional collection on $\mathrm{IG}(3, 7)$ (i.e. on $X^3(3, 3)$) has been constructed in [18].

Below, in Theorem 8.20 we construct a full rectangular Lefschetz exceptional collection in the case (5) of the classification, adding another variety to the above list.

Recently, the paper [37] introduced the notion of residual category using the framework of Lefschetz semiorthogonal decompositions. In particular, if a variety X has a full rectangular Lefschetz exceptional collection, then the residual category of X is trivial. Conjecture 1.12 of *loc. cit.* states that if the small quantum cohomology ring of a Fano variety of Picard rank one is generically semisimple, then its residual category should be of some prescribed form, and in the simplest possible case the residual category is trivial. Section 4.5 establishes semisimplicity of the small quantum cohomology for X^5 and together with Theorem 8.20 this confirms the conjecture of [37] for that case.

7. A CONSTRUCTION OF EXCEPTIONAL VECTOR BUNDLES ON HOROSPHERICAL VARIETIES

We start with two general statements concerning the descent of objects along a blow-up morphism.

Proposition 7.1. *Let $j : S \subset T$ be a closed embedding of a smooth subvariety S into a smooth variety T , and $\pi : \tilde{T} \rightarrow T$ be the blow-up of S at T . Denote by $i : E \hookrightarrow \tilde{T}$ the embedding of the exceptional divisor, and let $p : E \rightarrow S$ be the projection. Assume given an object $\mathcal{E} \in \mathcal{D}^b(\tilde{T})$. Then \mathcal{E} is pulled back from T , i.e., $\mathcal{E} = \pi^*\mathbb{E}$ for some $\mathbb{E} \in \mathcal{D}^b(T)$, if and only if the restriction of \mathcal{E} to E is trivial along the fibers of p , i.e., $i^*\mathcal{E} \in p^*\mathcal{D}^b(S)$.*

Proof. Let d be the codimension of S in T . By [42], there is a semiorthogonal decomposition

$$\mathcal{D}^b(\tilde{T}) = \langle i_*(p^*\mathcal{D}^b(S) \otimes \mathcal{O}_p(-d+1)), \dots, i_*(p^*\mathcal{D}^b(S) \otimes \mathcal{O}_p(-1)), \pi^*\mathcal{D}^b(T) \rangle.$$

In other words, the objects of $\pi^*\mathcal{D}^b(T)$ are characterized by the fact that they are left orthogonal to the subcategory $\langle i_*(p^*\mathcal{D}^b(S) \otimes \mathcal{O}_p(-d+1)), \dots, i_*(p^*\mathcal{D}^b(S) \otimes \mathcal{O}_p(-1)) \rangle$. Thus, $\mathcal{E} = \pi^*\mathbb{E}$ for some $\mathbb{E} \in \mathcal{D}^b(T)$ if and only if

$$(2) \quad \text{Hom}_{\tilde{T}}^\bullet(\mathcal{E}, i_*(p^*\mathcal{D}^b(S) \otimes \mathcal{O}_p(-k))) = \text{Hom}_E^\bullet(i^*\mathcal{E}, p^*\mathcal{D}^b(S) \otimes \mathcal{O}_p(-k)) = 0,$$

for $k = 1, \dots, d-1$. By *loc.cit.*, $\mathcal{D}^b(E)$ has a semiorthogonal decomposition

$$\mathcal{D}^b(E) = \langle p^*\mathcal{D}^b(S) \otimes \mathcal{O}_p(-d+1), \dots, p^*\mathcal{D}^b(S) \otimes \mathcal{O}_p(-1), p^*\mathcal{D}^b(S) \rangle.$$

Thus, the equality (2) holds if and only if $i^*\mathcal{E} \in p^*\mathcal{D}^b(S)$, proving the statement. \square

We keep the notation and assumptions of Proposition 7.1.

Proposition 7.2. *Let \mathcal{E} be an object of $\mathcal{D}^b(\tilde{T})$, such that $\mathcal{E} = \pi^*\mathbb{E}$ where $\mathbb{E} \in \mathcal{D}^b(T)$. If \mathcal{E} is a pure object, i.e. is a coherent sheaf on \tilde{T} , then so is the object \mathbb{E} on T . If moreover \mathcal{E} is locally free, then so is the coherent sheaf \mathbb{E} .*

Proof. Since π is the blow-up, there is an isomorphism $\pi_*\mathcal{O}_{\tilde{T}} = \mathcal{O}_T$, and the functor π^* is full and faithful. In particular, π^* sends non-zero objects to non-zero objects. Applying the functor π_* to $\pi^*\mathbb{E}$ and using projection formula, one obtains isomorphisms $\pi_*\pi^*\mathbb{E} = \mathbb{E} \otimes \pi_*\mathcal{O}_{\tilde{T}} = \mathbb{E} = \pi_*\mathcal{E}$. One sees that $\mathbb{E} \in \mathcal{D}^{\geq 0}(T)$, since in particular $\mathcal{E} \in \mathcal{D}^{\geq 0}(\tilde{T})$ and the functor π_* is left t -exact. Assuming that \mathbb{E} had a non-trivial sheaf cohomology outside the degree zero, consider the distinguished triangle $\dots \rightarrow \tau_{< m}\mathbb{E} \rightarrow \mathbb{E} \rightarrow \mathcal{H}^m\mathbb{E}[-m] \rightarrow \dots$ associated with the canonical truncation, the positive integer m being the maximal non-trivial cohomology degree of \mathbb{E} . Applying then the functor π^* to that triangle, one sees that $\pi^*\mathbb{E}$ would also have a non-trivial sheaf cohomology on \tilde{T} in the same degree, the functor π^* being right t -exact. In other words, there is an isomorphism $\mathcal{H}^m(\pi^*\mathbb{E}) = L^0\pi^*\mathcal{H}^m\mathbb{E}$, and the latter coherent sheaf is non-trivial. But the object $\pi^*\mathbb{E} = \mathcal{E}$ being pure by assumption, it does not have non-trivial sheaf cohomology outside the degree zero. Therefore, $m = 0$ and \mathbb{E} is a coherent sheaf.

Assuming that \mathcal{E} is a locally free sheaf, the property of being locally free for \mathbb{E} can be checked against the derived local $\mathcal{H}om$ -groups: a coherent sheaf \mathbb{E} on T is locally free if and only if $\mathcal{R}Hom_T(\mathbb{E}, \mathcal{O}_T)$ is a pure object. Now $\pi^*\mathbb{E} = \mathcal{E}$, and $\pi^*\mathcal{R}Hom_T(\mathbb{E}, \mathcal{O}_T) = \mathcal{R}Hom_{\tilde{T}}(\pi^*\mathbb{E}, \pi^*\mathcal{O}_T) = \mathcal{R}Hom_{\tilde{T}}(\mathcal{E}, \mathcal{O}_{\tilde{T}})$, while the latter object is pure, the sheaf \mathcal{E} being locally free. We can repeat the same argument as above showing that $\mathcal{R}Hom_T(\mathbb{E}, \mathcal{O}_T)$ is then a coherent sheaf, that is a pure object. This forces \mathbb{E} to be locally free. \square

7.1. The bundles \tilde{F}_Y and \tilde{F}_Z . We refer to Section 1.5 for the notation used in this section. Recall that both maps $p : E \rightarrow Y$ and $q : E \rightarrow Z$ are projective bundles associated to vector bundles $N_Y = N_{Y/X}$ and $N_Z = N_{Z/X}$ on Y and Z , respectively. We set $F_Y = N_{Y/X}^\vee = p_* \mathcal{L}_{\omega_Z - \omega_Y}$ and $F_Z = N_{Z/X}^\vee = q_* \mathcal{L}_{\omega_Y - \omega_Z}$, where ω_Y and ω_Z are the fundamental weights associated to Y and Z , respectively. Let $\mathcal{O}_p(1)$ (resp., $\mathcal{O}_q(1)$) denote the line bundle of relative degree one along projection p (resp., along projection q). Note that we also have $\mathcal{O}_p(1) = \mathcal{L}_{\omega_Z - \omega_Y}$ and $\mathcal{O}_q(1) = \mathcal{L}_{\omega_Y - \omega_Z}$. In particular, we have the relative Euler sequences on E :

$$0 \rightarrow \Omega_{E/Y}^1 \otimes \mathcal{O}_p(1) \rightarrow p^* F_Y \rightarrow \mathcal{O}_p(1) \rightarrow 0$$

and

$$0 \rightarrow \Omega_{E/Z}^1 \otimes \mathcal{O}_q(1) \rightarrow q^* F_Z \rightarrow \mathcal{O}_q(1) \rightarrow 0.$$

Recall that the composed morphisms $\xi \circ j_Z : E_Z \rightarrow E$ and $\xi \circ j_Y : E_Y \rightarrow E$ are both identity morphisms $\text{id}_E : E \rightarrow E$. Thus, $j_Z^* \xi^* = j_Y^* \xi^* = \text{id}_E^*$, and the above map $p^* F_Y \rightarrow \mathcal{O}_p(1)$ (resp., $q^* F_Z \rightarrow \mathcal{O}_q(1)$) defines by adjunction a map of coherent sheaves $\xi^* p^* F_Y \rightarrow j_{Z*} \mathcal{O}_p(1)$ (resp., $\xi^* q^* F_Z \rightarrow j_{Y*} \mathcal{O}_q(1)$). We define coherent sheaves \tilde{F}_Y and \tilde{F}_Z via the exact sequences:

$$0 \rightarrow \tilde{F}_Y \rightarrow \xi^* p^* F_Y \rightarrow j_{Z*} \mathcal{O}_p(1) \rightarrow 0$$

and

$$0 \rightarrow \tilde{F}_Z \rightarrow \xi^* q^* F_Z \rightarrow j_{Y*} \mathcal{O}_q(1) \rightarrow 0.$$

Proposition 7.3. *Both \tilde{F}_Y and \tilde{F}_Z are locally free sheaves.*

Proof. A coherent sheaf \mathcal{E} on a smooth variety S is locally free if and only if $\text{Ext}_S^i(\mathcal{E}, \mathcal{O}_S) = 0$ for $i > 0$. The statement follows from applying $\mathcal{H}om(-, \mathcal{O}_{\tilde{X}_{YZ}})$ to the above sequences and taking into account that j_Z and j_Y are divisorial embeddings. \square

Proposition 7.4. *For all the varieties of Pasquier's list except the variety X^2 , the bundle \tilde{F}_Y is the pull-back of a vector bundle on X .*

Proof. Proposition 7.2 ensures that if the vector bundle \tilde{F}_Y is the pull-back of an object of $\mathcal{D}^b(X)$, then that object is also a vector bundle on X . By Proposition 7.1, in order to prove that \tilde{F}_Y is pulled back from X , it is sufficient to check that $j_Y^* \tilde{F}_Y$ and $j_Z^* \tilde{F}_Y$ are trivial on the fibers of p and q , respectively. The first statement is clear: since Y and Z are disjoint we have $j_Y^* \tilde{F}_Y = j_Y^* \xi^* p^* F_Y = p^* F_Y$, proving the result. We now compute $j_Z^* \tilde{F}_Y$. For this we need to compute $\text{Tor}_1^{\tilde{X}}(j_{Z*} \mathcal{O}_p(1), \mathcal{O}_{E_Z}) = \mathcal{O}_{E_Z}(-E_Z) \otimes \mathcal{O}_p(1) = \mathcal{L}_{\omega_Y - \omega_Z} \otimes \mathcal{O}_p(1) = \mathcal{O}_{E_Z}$. We get the exact sequence

$$0 \rightarrow \mathcal{O}_E \rightarrow j_Z^* \tilde{F}_Y \rightarrow p^* F_Y \rightarrow \mathcal{O}_p(1) \rightarrow 0.$$

We are therefore left with proving that $\Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)$ is the pullback of a vector bundle on Z . There is an obvious necessary condition given by the degree, namely, $\Lambda^{\text{top}}(\Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)) = \Lambda^{\text{top}} p^* F_Y \otimes \mathcal{O}_p(1)^{\otimes (-1)}$ should come from Z . Since $F_Y = N_{Y/X}^\vee$ we can compute

$$\Lambda^{\text{top}} p^* F_Y = p^* \mathcal{O}_Y(1)^{\otimes (c_1(Y) - c_1(X))} = \mathcal{L}_{(c_1(Y) - c_1(X))\omega_Y}$$

and therefore

$$\Lambda^{\text{top}}(\Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)) = \mathcal{L}_{(c_1(Y) + 1 - c_1(X))\omega_Y - \omega_Z}.$$

Now for $X \neq X^2$ we have $c_1(X) = c_1(Y) + 1$, so this necessary condition is satisfied (this also proves that \tilde{F}_Y does not descend if $X = X^2$). If furthermore $\Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)$ is of rank one, the above condition is a sufficient condition (since $\Lambda^{\text{top}}(\Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)) = \Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)$). This occurs if and only if $\text{rk}(\Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)) = \text{codim}_X Y - 1 = 1$, therefore for $\text{codim}_X Y = 2$. This is true for $X^1(n)$ and X^5 . We are left with $X^3(n, m)$ and X^4 . For $X = X^3(n, m)$, using the description of X as an odd symplectic Grassmannian we will easily check that p^*F_Y and $\Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)$ are pullbacks of the tautological subbundles in Y and Z , respectively, giving the result. We prove this using representations. Indeed, both p^*F_Y and $\mathcal{L}_{\omega_Z - \omega_Y}$ are homogeneous bundles on E coming from representations of $P_Y \cap P_Z$. The bundle p^*F_Y comes from the P_Y -representation $V_{P_Y}(\omega_Z - \omega_Y)$ while $\mathcal{L}_{\omega_Z - \omega_Y}$ comes from the one-dimensional $P_Y \cap P_Z$ -representation $\mathbb{C}_{\omega_Z - \omega_Y}$ of weight $\omega_Z - \omega_Y$. The exact sequence comes from an exact sequence of representations

$$0 \rightarrow M \rightarrow V_{P_Y}(\omega_Z - \omega_Y) \rightarrow \mathbb{C}_{\omega_Z - \omega_Y} \rightarrow 0.$$

We only need to check that M is a P_Z -representation (and not only a $P_Y \cap P_Z$ -representation). For this we only need to define the action of \mathfrak{g}_{α_Y} , where α_Y is the simple root associated to Y . The only possible action is the trivial action, which is compatible with the $P_Y \cap P_Z$ -action if and only if the cocharacter α_Y^\vee acts trivially. This is an easy check if $X = X^3(n, m)$ or $X = X^4$. \square

Remark 7.5. The above necessary condition proves that \tilde{F}_Z never descends, with maybe the exception of the case $X = X^3(n, m)$ (where $c_1(X) = c_1(Z) + 1$). An easy computation with weights shows that even in that case \tilde{F}_Z does not descend.

Definition 7.6. If $X \neq X^2$, we denote by \mathcal{F}_Y the vector bundle on X such that $\pi_{Y/Z}^* \mathcal{F}_Y = \tilde{F}_Y$.

7.2. The bundle \mathcal{F}_Y is exceptional. Assume that $X \neq X^2$.

Proposition 7.7. *The bundles \tilde{F}_Y and \mathcal{F}_Y are exceptional.*

Proof. Since X is smooth we have $\pi_{Z*} \mathcal{O}_{\tilde{X}_Z} = \mathcal{O}_X$ and $R^i \pi_{Z*} \mathcal{O}_{\tilde{X}_Z} = 0$ for $i > 0$. In particular it is enough to prove that \tilde{F}_Y is exceptional. We apply $\text{Hom}(-, \tilde{F}_Y)$ to the defining exact sequence. We are thus left to consider $\text{Ext}^i(\xi^* p^* F_Y, \tilde{F}_Y)$ and $\text{Ext}^i(j_{Z*} \mathcal{O}_p(1), \tilde{F}_Y)$.

We have $\text{Ext}^i(\xi^* p^* F_Y, \tilde{F}_Y) = \text{Ext}^i(p^* F_Y, \xi_* \tilde{F}_Y)$. Applying ξ_* to the exact sequence defining \tilde{F}_Y , we get $\xi_* \tilde{F}_Y = \Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)$. To compute $\text{Ext}^i(p^* F_Y, \Omega_{E/Y}^1 \otimes \mathcal{O}_p(1))$, we therefore need to compute $\text{Ext}^i(p^* F_Y, p^* F_Y)$ and $\text{Ext}^i(p^* F_Y, \mathcal{O}_p(1))$. But $F_Y = p_* \mathcal{O}_p(1)$ and p is cohomologically trivial, therefore the first one is isomorphic to $\text{Ext}^i(F_Y, F_Y)$ and, by adjunction of p^* and p_* , the second one is also isomorphic to $\text{Ext}^i(F_Y, F_Y)$. This proves that $\text{Ext}^i(p^* F_Y, \tilde{F}_Y) = 0$ for any i .

We have $\text{Ext}^i(j_{Z*} \mathcal{O}_p(1), \tilde{F}_Y) = \text{Ext}^i(\mathcal{O}_p(1), j_Z^! \tilde{F}_Y)$. For the smooth divisor $E_Z \subset X$, we have $j_Z^!(-) = j_Z^*(-) \otimes N_{E/X}[-1] = j_Z^*(-) \otimes \mathcal{L}_{\omega_Z - \omega_Y}[-1]$, thus we get the equalities $\text{Ext}^i(j_{Z*} \mathcal{O}_p(1), \tilde{F}_Y) = \text{Ext}^{i-1}(\mathcal{L}_{\omega_Z - \omega_Y}, j_Z^* \tilde{F}_Y \otimes \mathcal{L}_{\omega_Z - \omega_Y}) = \text{Ext}^{i-1}(\mathcal{O}, j_Z^* \tilde{F}_Y)$. We are therefore left to compute $\text{Ext}^{i-1}(\mathcal{O}, \mathcal{O}) = \delta_{1,i}$ and $\text{Ext}^{i-1}(\mathcal{O}, \Omega_{E/Y}^1 \otimes \mathcal{O}_p(1))$. To compute the last Ext group we need to compute the extensions $\text{Ext}^{i-1}(\mathcal{O}, p^* F_Y)$ and $\text{Ext}^{i-1}(\mathcal{O}, \mathcal{O}_p(1))$. Since $F_Y = p_* \mathcal{O}_p(1)$, both are equal and since $\mathcal{O}_p(1) = \mathcal{L}_{\omega_Z - \omega_Y}$ these groups are trivial for all i . \square

7.3. Exceptional sequence. We construct a 2-term exceptional sequence.

Proposition 7.8. *The sequence $\langle \mathcal{F}_Y, \mathcal{O}_X \rangle$ is exceptional on X .*

Proof. We only need to check the vanishing of the groups $\text{Ext}^i(\mathcal{O}_X, \mathcal{F}_Y)$. This is equivalent to the vanishing of $\text{Ext}^i(\mathcal{O}_{\tilde{X}_Z}, \tilde{F}_Y)$. Via the defining exact sequence we get the group $\text{Ext}^i(\mathcal{O}_{\tilde{X}_Z}, \xi^* p^* F_Y)$ and $\text{Ext}^i(\mathcal{O}_{\tilde{X}_Z}, j_{Z*} \mathcal{O}_p(1))$.

We have $\text{Ext}^i(\mathcal{O}_{\tilde{X}_Z}, j_{Z*} \mathcal{O}_p(1)) = \text{Ext}^i(j_Z^* \mathcal{O}_{\tilde{X}_Z}, \mathcal{O}_p(1)) = \text{Ext}^i(\mathcal{O}, \mathcal{O}_p(1))$ and we have seen that these groups vanish. We have also seen that $\text{Ext}^i(\mathcal{O}_{\tilde{X}_Z}, \xi^* p^* F_Y) = \text{Ext}^i(\mathcal{O}, F_Y)$ vanishes. \square

8. A FULL RECTANGULAR LEFSCHETZ EXCEPTIONAL COLLECTION ON THE G_2 -VARIETY

In this section, we construct a full rectangular Lefschetz exceptional collection on the variety X^5 .

Let G_2 be the exceptional group of rank two. We denote by α and β denote the two simple roots in R_+ , the root β being the long root.

8.1. The flag variety of G_2 . The group G_2 has two standard parabolic subgroups P_α and P_β which correspond to the simple roots α and β . Let $\omega_\alpha, \omega_\beta \in X(T)$ be the two fundamental weights. Recall that given a dominant weight $\lambda \in X(T)$, we denote $V(\lambda)$ the irreducible representation with highest weight λ . The homogeneous spaces G/P_α and G/P_β are isomorphic respectively to the 5-dimensional quadric $\mathbf{Q}_5 \subset \mathbb{P}(V(\omega_\alpha))$ (the projectivization of G_2 -orbit of the highest vector in the irreducible G_2 -representation $V(\omega_\alpha)$), and to the 5-dimensional variety $G_2^{\text{ad}} \subset \mathbb{P}(V(\omega_\beta))$ (the projectivization of G_2 -orbit of the highest vector in $V(\omega_\beta)$, the adjoint representation of G_2). The Levi subgroups of P_α and P_β have a component isomorphic to SL_2 ; its tautological representation in each case gives rise to a homogeneous rank 2 vector bundle on G/P_α (resp., on G/P_β). Denote by π_α and π_β the two projections of G_2/B onto $\mathbf{Q}_5 \subset \mathbb{P}(V(\omega_\alpha))$ and $G_2^{\text{ad}} \subset \mathbb{P}(V(\omega_\beta))$, respectively. By [34, Lemma 8.3], the projection π_α is the projective bundle associated to the stable indecomposable rank two bundle \mathcal{K} on \mathbf{Q}_5 with $\det \mathcal{K} = \mathcal{L}_{-3\omega_\alpha}$, and the projection π_β is the projective bundle over G_2^{ad} associated to the tautological rank two vector bundle \mathcal{U}_2 over $G_2^{\text{ad}} \subset \text{Gr}(2, V(\omega_\alpha))$. The latter embedding is related to an isomorphism of G_2 -modules $\Lambda^2 V(\omega_\alpha) = V(\omega_\alpha)^* \oplus V(\omega_\beta)$: the grassmannian $\text{Gr}(2, V(\omega_\alpha))$ is embedded into $\mathbb{P}(\Lambda^2 V(\omega_\alpha))$ via the Plücker embedding, and the embedding $G_2^{\text{ad}} \subset \mathbb{P}(V(\omega_\beta))$ is obtained via restriction along the natural embedding $\mathbb{P}(V(\omega_\beta)) \subset \mathbb{P}(\Lambda^2 V(\omega_\alpha))$ induced by the above direct sum decomposition.

By [43], there is a short exact sequence

$$0 \rightarrow \mathcal{S} \rightarrow W \otimes \mathcal{O}_{\mathbf{Q}_5} \rightarrow \mathcal{S}^\vee \rightarrow 0,$$

where \mathcal{S} is the spinor bundle on \mathbf{Q}_5 , and W is the spinor representation regarded as a representation of G_2 via the restriction $G_2 \subset \text{Spin}_7$. Next, there is a short exact sequence

$$(3) \quad 0 \rightarrow \mathcal{L}_{-\omega_\beta} \rightarrow \pi_\alpha^* \mathcal{K} \rightarrow \mathcal{L}_{\beta-\omega_\beta} \rightarrow 0.$$

We have $\mathcal{S} \otimes \mathcal{L}_{\omega_\alpha} = \mathcal{S}^\vee$. Denote by $\Psi_1^{\omega_\alpha}$ the vector bundle on \mathbf{Q}_5 which is the pullback of $\Omega_{\mathbb{P}(V(\omega_\alpha))}^1(1)$ along the embedding $\mathbf{Q}_5 \subset \mathbb{P}(V(\omega_\alpha))$. The bundles \mathcal{S} and

\mathcal{K} are related via the short exact sequence

$$(4) \quad 0 \rightarrow \mathcal{K} \rightarrow (\Psi_1^{\omega_\alpha})^\vee \otimes \mathcal{L}_{-\omega_\alpha} \rightarrow \mathcal{S} \rightarrow 0,$$

see [34, Appendix B].

The weights of $V(\omega_\alpha)$ are $\omega_\alpha, -\omega_\alpha + \omega_\beta, 2\omega_\alpha - \omega_\beta, 0, -2\omega_\alpha + \omega_\beta, \omega_\alpha - \omega_\beta, -\omega_\alpha$, while we deduce from (3) that the weights of \mathcal{K} are $-\omega_\beta, \omega_\beta - 3\omega_\alpha$. Thus the weights of \mathcal{S} are $0, -2\omega_\alpha + \omega_\beta, \omega_\alpha - \omega_\beta, -\omega_\alpha$. We have the tautological short exact sequence on G_2^{ad} :

$$(5) \quad 0 \rightarrow \mathcal{U}_2 \rightarrow V(\omega_\alpha) \otimes \mathcal{O}_{G_2^{\text{ad}}} \rightarrow V(\omega_\alpha)/\mathcal{U}_2 \rightarrow 0.$$

Define $\mathcal{U}_2^\perp := (V(\omega_\alpha)/\mathcal{U}_2)^\vee$. There is a short exact sequence for the pullback of \mathcal{U}_2 along the projection $\pi_\beta : G_2/B \rightarrow G_2^{\text{ad}}$:

$$(6) \quad 0 \rightarrow \mathcal{L}_{-\omega_\alpha} \rightarrow \pi_\beta^* \mathcal{U}_2 \rightarrow \mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow 0.$$

8.2. Bott's theorem. Our main cohomological tool is Bott's theorem. Consider a weight $\lambda \in X(T)$ and let \mathcal{L}_λ be the corresponding line bundle on G/B . The weight λ is called *singular* if it lies on a wall of some Weyl chamber defined by $\langle -, \alpha^\vee \rangle = 0$ for some coroot $\alpha^\vee \in R^\vee$. Weights which are not singular are called *regular*. The Weyl group $W = N_G(T)/T$ acts on $X(T)$ via the dot action: if $w \in W$ and $\lambda \in X(T)$, then $w \cdot \lambda = w(\lambda + \rho) - \rho$, where ρ is the sum of fundamental weights.

Theorem 8.1. [17, Theorem 2]

- (1) If $\lambda + \rho$ is singular, then $H^i(G/B, \mathcal{L}_\lambda) = 0$ for all i .
- (2) If $\lambda + \rho$ is regular and dominant, then $H^i(G/B, \mathcal{L}_\lambda) = 0$ for $i > 0$.
- (3) If $\lambda + \rho$ is regular, then $H^i(G/B, \mathcal{L}_\lambda) \neq 0$ for a unique degree i which coincides with $\ell(w)$, the length of a Weyl group element w that takes λ to the dominant chamber, i.e. $w \cdot \lambda \in X_+(T)$. The cohomology group $H^{\ell(w)}(G/B, \mathcal{L}_\lambda)$ is the irreducible G -module of highest weight $w \cdot \lambda$.

8.3. Horospherical G_2 -variety. In what follows, X denotes the horospherical variety X^5 . Recall that X is embedded into $\mathbb{P}(V(\omega_\alpha) \oplus V(\omega_\beta))$ and recall the blow-up construction from Subsection 1.5. In this section we will use the following notation which is better suited to case (5) of the classification:

$$\begin{array}{ccccc} E_1 & \xhookrightarrow{i_1} & \tilde{X} & \xleftarrow{i_2} & E_2 \\ \downarrow \pi_\alpha & & \downarrow \pi & & \downarrow \pi_\beta \\ \mathbb{Q}_5 & \xhookrightarrow{j_1} & X & \xleftarrow{j_2} & G_2^{\text{rad}} \end{array}$$

In the above diagram, we have $Y = G_2^{\text{ad}}$, $Z = \mathbb{Q}_5$. The middle map π is the blow-up of X at $\mathbb{Q}_5 \cup G_2^{\text{ad}}$. The exceptional divisors over \mathbb{Q}_5 and G_2^{ad} are denoted E_1 and E_2 , respectively. The maps π_α and π_β are, respectively, the projections q and p from the incidence variety $E = E_1 = E_2$ to Z and Y . In our case, the incidence variety E is isomorphic to the flag variety G_2/B , and the blow-up \tilde{X} has a \mathbb{P}^1 -bundle structure over G_2/B , namely $\tilde{X} = \mathbb{P}_{G_2/B}(\mathcal{L}_{-\omega_\alpha} \oplus \mathcal{L}_{-\omega_\beta})$. The exceptional divisors E_1 and E_2 are embedded linearly into \tilde{X} with respect to this \mathbb{P}^1 -bundle structure, and we can identify them as $E_1 = \mathbb{P}_{G_2/B}(\mathcal{L}_{-\omega_\alpha}) = G_2/B$ and $E_2 = \mathbb{P}_{G_2/B}(\mathcal{L}_{-\omega_\beta}) = G_2/B$. Denote by ξ the projection of \tilde{X} onto G_2/B .

8.4. Line bundles on \tilde{X} . There are relations among line bundles on \tilde{X} . Firstly, the index of X being equal to 4, the canonical bundle satisfies $\omega_X = \mathcal{O}_X(-4)$. Thus, $\omega_{\tilde{X}} = \pi^* \mathcal{O}_X(-4) \otimes \mathcal{O}_{\tilde{X}}(E_1) \otimes \mathcal{O}_{\tilde{X}}(E_2)$. The adjunction formula gives

$$\omega_{E_1} = (\omega_{\tilde{X}} \otimes \mathcal{O}_{\tilde{X}}(E_1)) \otimes \mathcal{O}_{E_1} = \mathcal{L}_{-4\omega_\alpha} \otimes \mathcal{O}_{E_1}(2E_1).$$

Indeed, $\mathcal{O}_{E_1}(E_2) = \mathcal{O}_{E_1}$ (resp., $\mathcal{O}_{E_2}(E_1) = \mathcal{O}_{E_2}$) since intersection of E_1 and E_2 is empty. On the other hand, $\omega_{E_1} = \mathcal{L}_{-2\rho}$, where $\rho := 5\alpha + 3\beta$ is the half-sum of all positive roots, implying that $\mathcal{O}_{E_1}(E_1) = N_{E_1/\tilde{X}} = \mathcal{L}_{\omega_\alpha - \omega_\beta}$.

Similarly, the normal bundle $\mathcal{O}_{E_2}(E_2) = N_{E_2/\tilde{X}}$ is isomorphic to $\mathcal{L}_{\omega_\beta - \omega_\alpha}$, via the isomorphisms $\omega_{E_2} = (\omega_{\tilde{X}} \otimes \mathcal{O}_{\tilde{X}}(E_2)) \otimes \mathcal{O}_{E_2} = \mathcal{L}_{-4\omega_\beta} \otimes \mathcal{O}_{E_2}(2E_2)$ and $\omega_{E_2} = \mathcal{L}_{-2\rho}$, implying that $\mathcal{O}_{E_2}(E_2) = N_{E_2/\tilde{X}} = \mathcal{L}_{\omega_\beta - \omega_\alpha}$. Finally, recall that $\xi : \tilde{X} \rightarrow G_2/B$ is the \mathbb{P}^1 -bundle associated to $\mathcal{L}_{-\omega_\alpha} \oplus \mathcal{L}_{-\omega_\beta}$. This gives $\omega_{\tilde{X}} = \mathcal{O}_\xi(-2) \otimes \xi^* \omega_{G_2/B} \otimes \xi^* \mathcal{L}_{\omega_\alpha + \omega_\beta} = \mathcal{O}_\xi(-2) \otimes \xi^* \mathcal{L}_{-\rho}$. Pulling back $\mathcal{O}_X(1)$ along the blow-up of Z or the blow-up of Y , and using the above identifications of normal bundles $\mathcal{O}_{E_1}(E_1)$ and $\mathcal{O}_{E_2}(E_2)$, we get $\pi^* \mathcal{O}_X(1) = \mathcal{O}_{\tilde{X}}(E_1) \otimes \xi^* \mathcal{L}_{\omega_\beta} = \mathcal{O}_{\tilde{X}}(E_2) \otimes \xi^* \mathcal{L}_{\omega_\alpha}$. Hence, there is an isomorphism $\mathcal{O}_\xi(-1) = \pi^* \mathcal{O}_X(-1)$, and $\xi_* \pi^* \mathcal{O}_X(1) = \xi_* \mathcal{O}_\xi(1) = \mathcal{L}_{\omega_\alpha} \oplus \mathcal{L}_{\omega_\beta}$.

Consider the distinguished triangles for the divisors E_1 and E_2 :

$$\begin{aligned} \cdots \rightarrow \mathcal{O}_{\tilde{X}}(-E_1) \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow i_{1*} \mathcal{O}_{E_1} \rightarrow \mathcal{O}_{\tilde{X}}(-E_1)[1] \rightarrow \cdots, \\ \cdots \rightarrow \mathcal{O}_{\tilde{X}}(-E_2) \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow i_{2*} \mathcal{O}_{E_2} \rightarrow \mathcal{O}_{\tilde{X}}(-E_2)[1] \rightarrow \cdots \end{aligned}$$

Proposition 8.2. *There is a distinguished triangle in $\mathcal{D}^b(\tilde{X})$:*

$$(7) \quad \cdots \rightarrow \xi^* \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1) \rightarrow \mathcal{O}_{\tilde{X}} \oplus \xi^* \mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow \mathcal{O}_{\tilde{X}}(E_1) \rightarrow \cdots$$

Proof. By [42] we have $\mathcal{D}^b(\tilde{X}) = \langle \xi^* \mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1), \xi^* \mathcal{D}^b(G_2/B) \rangle$, hence there is a natural morphism $\xi^* \xi_* \mathcal{O}_{\tilde{X}}(E_1) \rightarrow \mathcal{O}_{\tilde{X}}(E_1)$. From the discussion above we know that $\mathcal{O}_{\tilde{X}}(E_1) = \pi^* \mathcal{O}_X(1) \otimes \xi^* \mathcal{L}_{-\omega_\beta} = \mathcal{O}_\xi(1) \otimes \xi^* \mathcal{L}_{-\omega_\beta}$. This gives

$$(8) \quad \xi_* \mathcal{O}_{\tilde{X}}(E_1) = \mathcal{O}_{G_2/B} \oplus \mathcal{L}_{\omega_\alpha - \omega_\beta}.$$

The cone of the natural morphism $\xi^* \xi_* \mathcal{O}_{\tilde{X}}(E_1) \rightarrow \mathcal{O}_{\tilde{X}}(E_1)$ is isomorphic to $\xi^*(?) \otimes \mathcal{O}_\xi(-1)$ for some object $? \in \mathcal{D}^b(G_2/B)$:

$$\cdots \rightarrow \mathcal{O}_{\tilde{X}} \oplus \xi^* \mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow \mathcal{O}_{\tilde{X}}(E_1) \rightarrow \xi^*(?) \otimes \mathcal{O}_\xi(-1) \rightarrow \cdots$$

To compute $?$ in the above triangle we tensor it with $\mathcal{O}_\xi(-1)$ and apply ξ_* ; we obtain an isomorphism $? \otimes \mathcal{L}_{-\rho}[-1] = \xi_*(\mathcal{O}_{\tilde{X}}(E_1) \otimes \mathcal{O}_\xi(-1))$. Tensoring equation (8) with $\mathcal{O}_\xi(-1)$ and applying ξ_* , we obtain $\xi_*(\mathcal{O}_{\tilde{X}}(E_1) \otimes \mathcal{O}_\xi(-1)) = \mathcal{L}_{-\omega_\beta}$. Finally, we get $? = \mathcal{L}_{\omega_\alpha}[1]$, arriving at distinguished triangle (7). \square

The triangle (7) is induced by the short exact sequence

$$(9) \quad 0 \rightarrow \xi^* \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1) \rightarrow \mathcal{O}_{\tilde{X}} \oplus \xi^* \mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow \mathcal{O}_{\tilde{X}}(E_1) \rightarrow 0.$$

Since $\xi^* \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1) = \mathcal{O}_{\tilde{X}}(-E_2)$, the sequence (9) is isomorphic to

$$0 \rightarrow \mathcal{O}_{\tilde{X}}(-E_2) \rightarrow \mathcal{O}_{\tilde{X}} \oplus \xi^* \mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow \mathcal{O}_{\tilde{X}}(E_1) \rightarrow 0.$$

Observe that since $\mathcal{O}_{\tilde{X}}(E_1) = \xi^* \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(1)$, the above extension corresponds to a unique non-trivial extension in the group $\text{Ext}^1(\xi^* \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(1), \xi^* \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1)) = \text{H}^1(G_2/B, \mathcal{L}_\rho \otimes \mathcal{L}_{-\rho}[-1]) = \text{H}^0(G_2/B, \mathcal{O}_{G_2/B}) = \mathbb{C}$. By duality, we obtain:

$$0 \rightarrow \mathcal{O}_{\tilde{X}}(-E_1) \rightarrow \mathcal{O}_{\tilde{X}} \oplus \xi^* \mathcal{L}_{\omega_\beta - \omega_\alpha} \rightarrow \mathcal{O}_{\tilde{X}}(E_2) \rightarrow 0,$$

with $(\mathcal{O}_{\tilde{X}} \oplus \xi^* \mathcal{L}_{\omega_\alpha - \omega_\beta}) = (\mathcal{O}_{\tilde{X}} \oplus \xi^* \mathcal{L}_{\omega_\alpha - \omega_\beta})^\vee \otimes \xi^* \mathcal{L}_{\omega_\alpha - \omega_\beta}$.

Using the isomorphisms $\pi_*\mathcal{O}_{\tilde{X}}(E_1) = \pi_*\mathcal{O}_{\tilde{X}}(E_2) = \pi_*\mathcal{O}_{\tilde{X}}(E_1 + E_2) = \mathcal{O}_X$, we can sum up the results as follows:

Lemma 8.3. *We have the following isomorphisms:*

$$\begin{aligned}\mathcal{O}_{\tilde{X}}(E_1) &= \xi^*\mathcal{L}_{-\omega_\beta} \otimes \pi^*\mathcal{O}_X(1) \text{ and } \mathcal{O}_{\tilde{X}}(E_2) = \xi^*\mathcal{L}_{-\omega_\alpha} \otimes \pi^*\mathcal{O}_X(1); \\ \pi_*\xi^*\mathcal{L}_{-\omega_\beta} &= \mathcal{O}_X(-1) = \pi_*\xi^*\mathcal{L}_{-\omega_\alpha} \text{ and } \pi_*\xi^*\mathcal{L}_{-\rho} = \mathcal{O}_X(-2).\end{aligned}$$

There are also short exact sequences:

$$\begin{aligned}0 \rightarrow \xi^*\mathcal{L}_{\omega_\alpha} \otimes \pi^*\mathcal{O}_X(-1) &\rightarrow \mathcal{O}_{\tilde{X}} \oplus \xi^*\mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow \xi^*\mathcal{L}_{-\omega_\beta} \otimes \pi^*\mathcal{O}_X(1) \rightarrow 0, \\ 0 \rightarrow \xi^*\mathcal{L}_{\omega_\beta} \otimes \pi^*\mathcal{O}_X(-1) &\rightarrow \mathcal{O}_{\tilde{X}} \oplus \xi^*\mathcal{L}_{\omega_\beta - \omega_\alpha} \rightarrow \xi^*\mathcal{L}_{-\omega_\alpha} \otimes \pi^*\mathcal{O}_X(1) \rightarrow 0.\end{aligned}$$

8.5. Vector bundles on X . Pulling back an exceptional vector bundle on G_2/B along $\xi : \tilde{X} \rightarrow G_2/B$ gives an exceptional vector bundle on \tilde{X} . However, we are looking for exceptional vector bundles on X ; to construct these, we start with appropriate exceptional vector bundles on \tilde{X} and modify it across the divisors E_1 and E_2 in a way that the modified bundles be trivial when restricted to the fibers of π_α and π_β . Proposition 7.1 ensures that in this case the modified bundle is pulled back from X .

We start with the tautological vector bundle \mathcal{U}_2 on G_2^{ad} and its pullback $\pi_\beta^*\mathcal{U}_2$ to $E_2 \cong G_2/B$. Recall that $\xi \circ i_2 = \xi \circ i_1 = \text{id}_{G_2/B}$, where the divisors E_1 and E_2 are identified with G_2/B . Thus, $i_1^*\xi^*\pi_\beta^*\mathcal{U}_2 = \pi_\beta^*\mathcal{U}_2$, and we have a short exact sequence, see sequence (6):

$$0 \rightarrow \mathcal{L}_{-\omega_\alpha} \rightarrow i_1^*\xi^*\pi_\beta^*\mathcal{U}_2 \rightarrow \mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow 0,$$

and by adjunction of i_1^* and i_{1*} we obtain a surjective morphism $\xi^*\pi_\beta^*\mathcal{U}_2 \rightarrow i_{1*}\mathcal{L}_{\omega_\alpha - \omega_\beta}$ of coherent sheaves on \tilde{X} .

Definition 8.4. Define $\tilde{\mathcal{U}}$ to be the kernel of the above surjective map:

$$(10) \quad 0 \rightarrow \tilde{\mathcal{U}} \rightarrow \xi^*\pi_\beta^*\mathcal{U}_2 \rightarrow i_{1*}\mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow 0,$$

Proposition 8.5. *Let $\tilde{\mathcal{U}}$ be as above.*

- (1) *The coherent sheaf $\tilde{\mathcal{U}}$ is locally free of rank two on \tilde{X} .*
- (2) *$\tilde{\mathcal{U}}$ is the pullback of a vector bundle \mathbb{U} of rank two on X .*
- (3) *The bundle \mathbb{U} is exceptional.*
- (4) *The bundles $\langle \mathbb{U}, \mathcal{O}_X \rangle$ form an exceptional pair.*
- (5) *We have $\det(\tilde{\mathcal{U}}) = \pi^*\det(\mathbb{U}) = \pi^*\mathcal{O}_X(-1)$.*

Proof. With notation as in Section 7, we have $\mathcal{U}_2 = F_Y$, $\mathcal{L}_{\omega_\alpha - \omega_\beta} = \mathcal{O}_p(1)$ and $\mathcal{L}_{-\omega_\alpha} = \Omega_{E/Y}^1 \otimes \mathcal{O}_p(1)$. In particular $\tilde{\mathcal{U}} = \tilde{F}_Y$ and the results follow from Propositions 7.3, 7.4, 7.7 and 7.8. We have $\mathbb{U} = \mathcal{F}_Y$, and the isomorphism in (5) is obtained by passing to determinants in the short exact sequence (10). Note that we have $\xi_*\tilde{\mathcal{U}} = \xi_*\tilde{F}_Y = \mathcal{L}_{-\omega_\alpha}$. \square

The bundle \mathcal{U}_2 in the above construction is pulled back to G_2/B from the Grassmannian G_2^{ad} via π_β^* . One can also apply the above construction to vector bundles that are pulled back from \mathbb{Q}_5 via the projection π_α . Next, we construct another exceptional bundle $\pi_\alpha^*\mathcal{S}$ on X coming from the spinor bundle \mathcal{S} on \mathbb{Q}_5 . To this end we need an auxiliary fact.

Proposition 8.6. *On G_2/B , we have an exact sequence of vector bundles*

$$(11) \quad 0 \rightarrow \pi_\beta^* \mathcal{U}_2 \rightarrow \pi_\alpha^* \mathcal{S} \rightarrow \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \rightarrow 0.$$

Proof. This was shown in [36, Proposition 3 and Lemma 4]. For convenience of the reader, we give a sketch of its proof, providing details of some cohomological calculations which are paramount to this section.

We first verify that the graded vector space $\text{Hom}_{G_2/B}^\bullet(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}, \pi_\beta^* \mathcal{U}_2) = \text{Hom}_{G_2/B}^\bullet(\pi_\beta^* \mathcal{U}_2^\vee, \pi_\beta^* \mathcal{U}_2 \otimes \mathcal{L}_{\omega_\alpha})$ is isomorphic to $\mathbb{C}[-1]$. To see this, recall that the bundle $\pi_\beta^* \mathcal{U}_2^\vee$ has a two-step filtration by the line bundles $\mathcal{L}_{\omega_\beta - \omega_\alpha}$ and $\mathcal{L}_{\omega_\alpha}$, as can be seen from sequence (6). Similarly, the bundle $\pi_\beta^* \mathcal{U}_2 \otimes \mathcal{L}_{\omega_\alpha}$ has a two-step filtration by the line bundles $\mathcal{L}_{2\omega_\alpha - \omega_\beta}$ and $\mathcal{O}_{G_2/B}$. Applying Theorem 8.1, we immediately see that the pairwise Ext-groups between all line bundles above are trivial, except one which is isomorphic to \mathbb{C} in degree 1. Specifically, the weights of line bundles in question are $\omega_\alpha - \omega_\beta, -\omega_\alpha, 3\omega_\alpha - 2\omega_\beta, \omega_\alpha - \omega_\beta$, and all these weights except the weight $3\omega_\alpha - 2\omega_\beta = -\beta$ are singular, since $\langle \omega_\alpha - \omega_\beta, \beta^\vee \rangle = \langle -\omega_\alpha, \alpha^\vee \rangle = -1$ (recall that $\langle \rho, \gamma^\vee \rangle = 1$ for any simple root γ). Hence, by Theorem 8.1, (1) the line bundles $\mathcal{L}_{\omega_\alpha - \omega_\beta}$ and $\mathcal{L}_{-\omega_\alpha}$ are acyclic. As for the weight $3\omega_\alpha - 2\omega_\beta = -\beta$, after applying the simple reflection s_β to it, we obtain $s_\beta \cdot (-\beta) = 0$, and Theorem 8.1, (3) gives that $H^1(G_2/B, \mathcal{L}_{-\beta}) = \text{Ext}_{G_2/B}^1(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}, \pi_\beta^* \mathcal{U}_2) = \mathbb{C}$.

Associated to a unique non-trivial extension is a short exact sequence:

$$(12) \quad 0 \rightarrow \pi_\beta^* \mathcal{U}_2 \rightarrow ? \rightarrow \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \rightarrow 0.$$

The coherent sheaf in the middle, being an extension of two locally free sheaves, is also locally free. We check that it is the pullback of a vector bundle on \mathbb{Q}_5 . To ensure that, it is sufficient to show that $\pi_{\alpha*} (? \otimes \mathcal{L}_{-\omega_\beta}) = 0$. Indeed, if $? = \pi_\alpha^* \mathcal{E}$ for a vector bundle \mathcal{E} on \mathbb{Q}_5 , then by the projection formula $\pi_{\alpha*} (? \otimes \mathcal{L}_{-\omega_\beta}) = \pi_{\alpha*} (\pi_\alpha^* \mathcal{E} \otimes \mathcal{L}_{-\omega_\beta}) = \mathcal{E} \otimes \pi_{\alpha*} \mathcal{L}_{-\omega_\beta} = 0$, since the line bundle $\mathcal{L}_{-\omega_\beta}$ has the relative degree -1 with respect to the projection π_α . From [42] it follows that this sufficient condition is also necessary that a vector bundle on G_2/B be pulled back from \mathbb{Q}_5 . Using the above filtrations for the bundles $\pi_\beta^* \mathcal{U}_2$ and $\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}$, tensoring (12) with $\mathcal{L}_{-\omega_\beta}$, and finally applying $\pi_{\alpha*}$ to the resulting sequence, one may check that $\pi_{\alpha*} (? \otimes \mathcal{L}_{-\omega_\beta})$ fits into a triangle $\cdots \rightarrow \pi_{\alpha*} (? \otimes \mathcal{L}_{-\omega_\beta}) \rightarrow \mathcal{L}_{-2\omega_\alpha} \rightarrow \mathcal{L}_{-2\omega_\alpha} \rightarrow \cdots$ which forces $\pi_{\alpha*} (? \otimes \mathcal{L}_{-\omega_\beta})$ to be zero.

Thus, there is a vector bundle \mathcal{E} on \mathbb{Q}_5 , such that $\pi_\alpha^* \mathcal{E} = ?$. By [27, Theorem 4.19], the category $\mathcal{D}^b(\mathbb{Q}_5)$ has a full exceptional collection $\langle \mathcal{L}_{-3\omega_\alpha}, \mathcal{L}_{-2\omega_\alpha}, \mathcal{L}_{-\omega_\alpha}, \mathcal{S}, \mathcal{O}_{\mathbb{Q}_5}, \mathcal{L}_{\omega_\alpha} \rangle$. To ensure that \mathcal{E} is isomorphic to the spinor bundle \mathcal{S} , we check first that \mathcal{E} is right orthogonal to line bundles $\mathcal{O}_{\mathbb{Q}_5}$ and $\mathcal{L}_{\omega_\alpha}$ and is left orthogonal to line bundles $\mathcal{L}_{-3\omega_\alpha}, \mathcal{L}_{-2\omega_\alpha}$, and $\mathcal{L}_{-\omega_\alpha}$. This easy cohomological computation which we are now skipping gives that \mathcal{E} is a direct sum of copies of the bundle \mathcal{S} . However, the rank of \mathcal{S} is equal to four which is the same as the rank of \mathcal{E} as is seen from the sequence (12). Hence, \mathcal{E} is isomorphic to \mathcal{S} , and the statement follows. \square

Definition 8.7. Define $\tilde{\mathcal{S}}$ as the kernel of the natural map

$$(13) \quad 0 \rightarrow \tilde{\mathcal{S}} \rightarrow \xi^* \pi_\alpha^* \mathcal{S} \rightarrow i_{2*} (\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \rightarrow 0.$$

Proposition 8.8. *With notation as above:*

- (1) *The coherent sheaf $\tilde{\mathcal{S}}$ is locally free of rank four on \tilde{X} .*

- (2) $\tilde{\mathcal{S}}$ is the pullback of a vector bundle \mathcal{S} of rank four on X .
 (3) We have $\text{Ext}_{\tilde{X}}^i(\mathcal{S}, \mathcal{S}) = 0$ for $i > 0$ and $\text{End}_X(\mathcal{S}) = \mathbb{C} \oplus \mathbb{C}$.
 (4) We have $\det(\tilde{\mathcal{S}}) = \pi^* \det(\mathcal{S}) = \pi^* \mathcal{O}_X(-2) = \mathcal{O}_\xi(-2)$.

Proof. Item (1) is similiar to Proposition 8.5, (i) since E_2 is a smooth divisor in \tilde{X} and $\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}$ is a locally free sheaf on E_2 .

For (2) note that the restriction of $\tilde{\mathcal{S}}$ to E_1 is isomorphic to $\pi_\alpha^* \mathcal{S}$. To compute the restriction of $\tilde{\mathcal{S}}$ to E_2 , apply i_2^* to (13):

$$0 \rightarrow L^1 i_{2*} i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \rightarrow i_2^* \tilde{\mathcal{S}} \rightarrow \pi_\alpha^* \mathcal{S} \rightarrow \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \rightarrow 0.$$

We have $L^1 i_{2*} i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) = \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{L}_{-\omega_\beta + \omega_\alpha} = \pi_\beta^* \mathcal{U}_2$. We deduce that

$$(14) \quad 0 \rightarrow \pi_\beta^* \mathcal{U}_2 \rightarrow i_2^* \tilde{\mathcal{S}} \rightarrow \pi_\beta^* \mathcal{U}_2 \rightarrow 0,$$

hence $i_2^* \tilde{\mathcal{S}}$ is trivial when restricted to fibers of π_β . As the bundle $\pi_\beta^* \mathcal{U}_2$ is exceptional, the above short exact sequence splits. Thus, $i_2^* \tilde{\mathcal{S}} = \pi_\beta^* \mathcal{U}_2 \oplus \pi_\beta^* \mathcal{U}_2$.

To prove (3) we apply $\text{Hom}_{\tilde{X}}(-, \tilde{\mathcal{S}})$ to (13). Since $\xi_* \tilde{\mathcal{S}} = \pi_\beta^* \mathcal{U}_2$, we need to compute the extensions $\text{Ext}_{G_2/B}^i(\pi_\alpha^* \mathcal{S}, \pi_\beta^* \mathcal{U}_2)$ and $\text{Ext}_{\tilde{X}}^i(i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \tilde{\mathcal{S}})$. To compute the first one, apply $\text{Hom}_{G_2/B}(\pi_\alpha^* \mathcal{S}, -)$ to (6). We get $\text{Ext}_{G_2/B}^i(\pi_\alpha^* \mathcal{S}, \mathcal{L}_{-\omega_\alpha}) = H^i(G_2/B, \pi_\alpha^* \mathcal{S}) = 0$ and $\text{Ext}_{G_2/B}^i(\pi_\alpha^* \mathcal{S}, \mathcal{L}_{\omega_\alpha - \omega_\beta}) = 0$; the latter group is trivial since $\pi_{\alpha*} \mathcal{L}_{-\omega_\beta} = 0$.

The group $\text{Ext}_{\tilde{X}}^i(i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \tilde{\mathcal{S}})$ is isomorphic to $\text{Ext}_{G_2/B}^{i-1}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}, i_2^* \tilde{\mathcal{S}} \otimes \mathcal{L}_{\omega_\beta - \omega_\alpha})$; remembering (14) we see that it is isomorphic to $\text{Ext}_{G_2/B}^{i-1}(\pi_\beta^* \mathcal{U}_2^\vee, \pi_\beta^* \mathcal{U}_2^\vee \oplus \pi_\beta^* \mathcal{U}_2^\vee)$. This proves (3).

Finally, item (4) is an easy computation. \square

The top row in the following diagram represents the bundle $\tilde{\mathcal{S}}$ as an extension. The middle column is (13), while the last column is the tensor product of $\xi^*(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha})$ with the short exact sequence

$$0 \rightarrow \mathcal{O}_\xi(-E_2) \rightarrow \mathcal{O}_{\tilde{X}} \rightarrow i_{2*} \mathcal{O}_{E_2} \rightarrow 0,$$

where we use Proposition 8.2 and Lemma 8.3 to identify $\xi^*(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \otimes \mathcal{O}_{\tilde{X}}(-E_2)$ with $\xi^* \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{O}_\xi(-1)$:

$$\begin{array}{ccccccc} & & & 0 & & 0 & \\ & & & \downarrow & & \downarrow & \\ 0 & \longrightarrow & \xi^* \pi_\beta^* \mathcal{U}_2 & \longrightarrow & \tilde{\mathcal{S}} & \longrightarrow & \xi^* \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{O}_\xi(-1) \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \xi^* \pi_\beta^* \mathcal{U}_2 & \longrightarrow & \xi^* \pi_\alpha^* \mathcal{S} & \longrightarrow & \xi^*(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) & \equiv & i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0 \end{array}$$

The first row gives an element in $\text{Ext}_{\tilde{X}}^1(\xi^*\pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{O}_\xi(-1), \xi^*\pi_\beta^*\mathcal{U}_2) = \mathbb{C}$ and is the non-trivial extension. The group $\text{Ext}_{\tilde{X}}^2(i_{2*}(\pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \xi^*\pi_\beta^*\mathcal{U}_2)$ is trivial, and the bundle $\xi^*\pi_\alpha^*\mathcal{S}$ in the center of the above diagram trivializes this extension.

Proposition 8.9. *As \mathbb{C} -algebras, we have $\text{End}_{\tilde{X}}(\tilde{\mathcal{S}}) \simeq \mathbb{C}[t]/t^2$.*

Proof. By [1], the vector bundle $\tilde{\mathcal{S}}$ is uniquely decomposed, up to a reordering, into a direct sum of indecomposable vector bundles. If $\tilde{\mathcal{S}}$ were decomposable, it would be the direct sum of two vector bundles, namely, $\tilde{\mathcal{S}} = \tilde{\mathcal{S}}_1 \oplus \tilde{\mathcal{S}}_2$. Recall that the restriction $i_1^*\tilde{\mathcal{S}}$ of $\tilde{\mathcal{S}}$ to E_1 is isomorphic to $\pi_\alpha^*\mathcal{S}$, hence it is an indecomposable bundle on E_1 ; if the bundle $\tilde{\mathcal{S}}$ was split as above, then applying i_1^* to the direct sum decomposition, one would obtain an isomorphism $i_1^*\tilde{\mathcal{S}} = i_1^*\tilde{\mathcal{S}}_1 \oplus i_1^*\tilde{\mathcal{S}}_2$, with each of the summands being necessarily non-trivial. That would contradict the indecomposability of $i_1^*\tilde{\mathcal{S}}$.

Therefore the bundle $\tilde{\mathcal{S}}$ is indecomposable, hence the algebra $\text{End}_{\tilde{X}}(\tilde{\mathcal{S}})$ is local. By Proposition 8.8, (3), $\dim_{\mathbb{C}} \text{End}_{\tilde{X}}(\tilde{\mathcal{S}}) = 2$, proving the claim. \square

Proposition 8.10. *We have $\text{Hom}_{\tilde{X}}^\bullet(\tilde{\mathcal{S}}, \tilde{\mathcal{U}}) = \text{Hom}_{\tilde{X}}^\bullet(\tilde{\mathcal{U}}, \tilde{\mathcal{S}}) = \mathbb{C}$.*

Proof. Applying $\text{Hom}_{\tilde{X}}(-, \tilde{\mathcal{U}})$ to (13), we need to compute the following groups: $\text{Ext}_{\tilde{X}}^i(\xi^*\pi_\alpha^*\mathcal{S}, \tilde{\mathcal{U}})$ and $\text{Ext}_{\tilde{X}}^i(i_{2*}(\pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \tilde{\mathcal{U}})$. The first computation comes from the fact that

$$\text{Ext}_{\tilde{X}}^i(\xi^*\pi_\alpha^*\mathcal{S}, \tilde{\mathcal{U}}) = \text{Ext}_{G_2/B}^i(\pi_\alpha^*\mathcal{S}, \xi_*\tilde{\mathcal{U}}) = \text{Ext}_{G_2/B}^i(\pi_\alpha^*\mathcal{S}, \mathcal{L}_{-\omega_\alpha}) = H^i(\mathbb{Q}_5, \mathcal{S}) = 0.$$

For the second one, we get an isomorphism

$$\text{Ext}_{\tilde{X}}^i(i_{2*}(\pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \tilde{\mathcal{U}}) = \text{Ext}_{G_2/B}^{i-1}(\pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}, i_2^*\tilde{\mathcal{U}} \otimes \mathcal{L}_{\omega_\beta - \omega_\alpha}).$$

Taking into account (10) we see that the last group is $\text{Ext}_{G_2/B}^{i-1}(\pi_\beta^*\mathcal{U}_2^\vee, \pi_\beta^*\mathcal{U}_2^\vee) = \mathbb{C}$ for $i = 1$. This proves the statement for $\text{Hom}_{\tilde{X}}^\bullet(\tilde{\mathcal{S}}, \tilde{\mathcal{U}})$. For the Hom-groups in the opposite direction, apply $\text{Hom}_{\tilde{X}}(-, \tilde{\mathcal{S}})$ to (10). We get

$$\text{Hom}_{\tilde{X}}^\bullet(i_{1*}\mathcal{L}_{\omega_\alpha - \omega_\beta}, \tilde{\mathcal{S}}) = H^\bullet(G_2/B, \pi_\alpha^*\mathcal{S}) = 0$$

and

$$\text{Hom}_{\tilde{X}}^\bullet(\xi^*\pi_\beta^*\mathcal{U}_2, \tilde{\mathcal{S}}) = \text{Hom}_{\tilde{X}}^\bullet(\pi_\beta^*\mathcal{U}_2, \xi_*\tilde{\mathcal{S}}) = \text{Hom}_{\tilde{X}}^\bullet(\pi_\beta^*\mathcal{U}_2, \pi_\beta^*\mathcal{U}_2) = \mathbb{C}. \quad \square$$

Definition 8.11. Let $\hat{\mathcal{S}} = \text{Cone}(\tilde{\mathcal{S}} \rightarrow \tilde{\mathcal{U}}) = \text{Cone}(\tilde{\mathcal{S}} \otimes \text{Hom}_{\tilde{X}}^\bullet(\tilde{\mathcal{S}}, \tilde{\mathcal{U}}) \rightarrow \tilde{\mathcal{U}})$, where $\text{Hom}_{\tilde{X}}^\bullet(\tilde{\mathcal{S}}, \tilde{\mathcal{U}})$ is identified with \mathbb{C} by Proposition 8.10.

Remark 8.12. By definition, we have $\text{Hom}_{\tilde{X}}^\bullet(\tilde{\mathcal{U}}, \hat{\mathcal{S}}) = 0$.

We have a distinguished triangle

$$(15) \quad \cdots \rightarrow \tilde{\mathcal{S}} \rightarrow \tilde{\mathcal{U}} \rightarrow \hat{\mathcal{S}} \rightarrow \tilde{\mathcal{S}}[1] \rightarrow \cdots$$

The cohomology of this triangle gives $0 \rightarrow \mathcal{H}^{-1}(\hat{\mathcal{S}}) \rightarrow \tilde{\mathcal{S}} \rightarrow \tilde{\mathcal{U}} \rightarrow \mathcal{H}^0(\hat{\mathcal{S}}) \rightarrow 0$.

Proposition 8.13. *We have $\mathcal{H}^{-1}(\hat{\mathcal{S}}) = \xi^*\pi_\beta^*\mathcal{U}_2$ and $\mathcal{H}^0(\hat{\mathcal{S}}) = i_{1*}\mathcal{L}_{-\omega_\alpha}$.*

Proof. By the previous proof, we have $\text{Ext}_{\tilde{X}}^1(i_{2*}(\pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \tilde{\mathcal{U}}) = \mathbb{C}$. Consider the unique non-trivial extension:

$$(16) \quad 0 \rightarrow \tilde{\mathcal{U}} \rightarrow \hat{\mathcal{U}} \rightarrow i_{2*}(\pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \rightarrow 0.$$

We claim that the middle term $\widehat{\mathcal{U}}$ fits into a short exact sequence:

$$(17) \quad 0 \rightarrow \xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\widehat{X}}(-E_1) \rightarrow \xi^*(\mathcal{L}_{-\omega_\alpha} \oplus \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \rightarrow \widehat{\mathcal{U}} \rightarrow 0.$$

Indeed, start with a distinguished triangle

$$(18) \quad \dots \rightarrow \xi^* \xi_* \widehat{\mathcal{U}} \rightarrow \widehat{\mathcal{U}} \rightarrow \xi^*(?) \otimes \mathcal{O}_\xi(-1) \rightarrow \xi^* \xi_* \widehat{\mathcal{U}}[1] \rightarrow \dots$$

To compute $\xi_* \widehat{\mathcal{U}}$ and $?$, apply ξ_* :

$$\dots \rightarrow \xi_* \widetilde{\mathcal{U}} \rightarrow \xi_* \widehat{\mathcal{U}} \rightarrow \pi_\beta^* \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \rightarrow \dots$$

Remembering that $\xi_* \widetilde{\mathcal{U}} = \mathcal{L}_{-\omega_\alpha}$, the above triangle reduces to

$$0 \rightarrow \mathcal{L}_{-\omega_\alpha} \rightarrow \xi_* \widehat{\mathcal{U}} \rightarrow \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \rightarrow 0,$$

which splits since $\text{Ext}_{G_2/B}^1(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}, \mathcal{L}_{-\omega_\alpha}) = H^1(G_2/B, \pi_\beta^* \mathcal{U}_2) = 0$. Thus, $\xi_* \widehat{\mathcal{U}} = \mathcal{L}_{-\omega_\alpha} \oplus \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}$. To compute $?$, tensor (18) with $\mathcal{O}_\xi(-1)$ and apply ξ_* . We have $\xi_*(\xi^* \xi_* \widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = 0$ as $\xi_* \mathcal{O}_\xi(-1) = 0$. Further, $\xi_*(\xi^*(?) \otimes \mathcal{O}_\xi(-2)) = ? \otimes \xi_* \mathcal{O}_\xi(-2) = ? \otimes \mathcal{L}_{-\rho}[-1]$. Thus, $? \otimes \mathcal{L}_{-\rho}[-1] = \xi_*(\widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1))$, and $? = \xi_*(\widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) \otimes \mathcal{L}_\rho[1]$.

To compute $\xi_*(\widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1))$, tensor (16) with $\mathcal{O}_\xi(-1)$:

$$0 \rightarrow \widetilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1) \rightarrow \widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1) \rightarrow i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha - \omega_\beta}) \rightarrow 0.$$

Applying ξ_* and writing $\mathcal{G} = \widetilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)$, we get

$$0 \rightarrow R^0 \xi_*(\mathcal{G}) \rightarrow R^0 \xi_*(\widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) \rightarrow \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\rho} \rightarrow R^1 \xi_*(\mathcal{G}) \rightarrow \dots$$

To compute $\xi_*(\widetilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1))$, tensor (10) with $\mathcal{O}_\xi(-1)$ and apply ξ_* , obtaining $\xi_*(\widetilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = \xi_* i_{1*}(\mathcal{L}_{\omega_\alpha - \omega_\beta - \omega_\alpha})[-1]$ (using again $\xi_* \mathcal{O}_\xi(-1) = 0$ and the identities $i_1^* \mathcal{O}_\xi(-1) = i_1^* \pi^* \mathcal{O}_X(-1) = \mathcal{L}_{-\omega_\alpha}$). Thus, $\xi_*(\widetilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = \mathcal{L}_{-\omega_\beta}[-1]$. We get

$$0 \rightarrow \xi_*(\widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) \rightarrow \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\rho} \rightarrow \mathcal{L}_{-\omega_\beta} \rightarrow 0,$$

which is the sequence (6) tensored with $\mathcal{L}_{-\omega_\alpha}$. In particular we have $\xi_*(\widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = \mathcal{L}_{-2\omega_\alpha}$ and $R^1 \xi_*(\widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = 0$. Returning to $?$ we obtain an isomorphism $? = \xi_*(\widehat{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) \otimes \mathcal{L}_\rho[1] = \mathcal{L}_{\rho - 2\omega_\alpha}[1] = \mathcal{L}_{\omega_\beta - \omega_\alpha}[1]$. Thus, the triangle (18) gives an exact sequence:

$$(19) \quad 0 \rightarrow \xi^* \mathcal{L}_{\omega_\beta - \omega_\alpha} \otimes \mathcal{O}_\xi(-1) \rightarrow \xi^*(\mathcal{L}_{-\omega_\alpha} \oplus \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \rightarrow \widehat{\mathcal{U}} \rightarrow 0.$$

Lemma 8.3 implies that $\xi^* \mathcal{L}_{\omega_\beta - \omega_\alpha} \otimes \mathcal{O}_\xi(-1) = \xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\widehat{X}}(-E_1)$. The morphism $\xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\widehat{X}}(-E_1) \rightarrow \xi^* \mathcal{L}_{-\omega_\alpha}$ in (19) is obtained by tensoring the map $\mathcal{O}_{\widehat{X}}(-E_1) \rightarrow \mathcal{O}_{\widehat{X}}$ with $\xi^* \mathcal{L}_{-\omega_\alpha}$, thus it is injective. The morphism $\xi^* \mathcal{L}_{\omega_\beta - \omega_\alpha} \otimes \mathcal{O}_\xi(-1) \rightarrow \xi^*(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha})$ is also injective since its source is an invertible sheaf and its target is a locally free sheaf (we also compute $\text{Hom}_{\widehat{X}}^\bullet(\xi^* \mathcal{L}_{\omega_\beta - \omega_\alpha} \otimes \mathcal{O}_\xi(-1), \xi^*(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha})) = V(\omega_\beta)$). This proves (17).

Consider the push-out of the extension (13) along the map $\tilde{\mathcal{S}} \rightarrow \tilde{\mathcal{U}}$ in (15). It follows from the above that the push-out extension is isomorphic to $\hat{\mathcal{U}}$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \tilde{\mathcal{S}} & \longrightarrow & \xi^* \pi_\alpha^* \mathcal{S} & \longrightarrow & i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \longrightarrow 0 \\ & & \downarrow & & \downarrow \tilde{f} & & \parallel \\ 0 & \longrightarrow & \tilde{\mathcal{U}} & \longrightarrow & \hat{\mathcal{U}} & \longrightarrow & i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) \longrightarrow 0. \end{array}$$

In the next diagram we pull back (17) along the map $\tilde{f} : \xi^* \pi_\alpha^* \mathcal{S} \rightarrow \hat{\mathcal{U}}$. We also write $\mathcal{A} := \xi^*(\mathcal{L}_{-\omega_\alpha} \oplus \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha})$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\tilde{X}}(-E_1) & \longrightarrow & ? & \longrightarrow & \xi^* \pi_\alpha^* \mathcal{S} \longrightarrow 0 \\ & & \parallel & & \downarrow & & \downarrow \tilde{f} \\ 0 & \longrightarrow & \xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\tilde{X}}(-E_1) & \longrightarrow & \mathcal{A} & \longrightarrow & \hat{\mathcal{U}} \longrightarrow 0. \end{array}$$

We compute $\text{Ext}_{\tilde{X}}^1(\xi^* \pi_\alpha^* \mathcal{S}, \xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\tilde{X}}(-E_1)) = 0$, thus obtaining an isomorphism $? = \xi^* \pi_\alpha^* \mathcal{S} \oplus \xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\tilde{X}}(-E_1)$. We finally see that the cone $\hat{\mathcal{S}}$ is isomorphic to the cone of the morphism

$$(20) \quad \xi^* \pi_\alpha^* \mathcal{S} \oplus \xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\tilde{X}}(-E_1) \rightarrow \xi^*(\mathcal{L}_{-\omega_\alpha} \oplus \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha})$$

in which $\text{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\alpha^* \mathcal{S}, \xi^* \mathcal{L}_{-\omega_\alpha}) = 0$ and $\text{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\alpha^* \mathcal{S}, \xi^* \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}) = \mathbb{C}$. The unique non-trivial element of the latter group gives rise to sequence (11) from Proposition 8.6. The map $\xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\tilde{X}}(-E_1) \rightarrow \xi^* \mathcal{L}_{-\omega_\alpha}$ is obtained by tensoring the embedding map $\mathcal{O}_{\tilde{X}}(-E_1) \rightarrow \mathcal{O}_{\tilde{X}}$ with $\xi^* \mathcal{L}_{-\omega_\alpha}$. Finally, $\text{Hom}_{\tilde{X}}^\bullet(\xi^* \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_{\tilde{X}}(-E_1), \xi^*(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha})) = V(\omega_\beta)$. It follows that the kernel $\mathcal{H}^{-1}(\hat{\mathcal{S}})$ is isomorphic to $\xi^* \pi_\beta^* \mathcal{U}_2$ and that the cokernel of (20) is isomorphic to $\mathcal{H}^0(\hat{\mathcal{S}}) = \xi^* \mathcal{L}_{-\omega_\alpha} \otimes i_{1*} \mathcal{O}_{E_1} = i_{1*} \mathcal{L}_{-\omega_\alpha}$, and Proposition 8.13 follows. \square

Proposition 8.14. *The object $\hat{\mathcal{S}}$ is exceptional.*

Proof. By Proposition 8.13, there is a distinguished triangle:

$$(21) \quad \cdots \rightarrow \xi^* \pi_\beta^* \mathcal{U}_2[1] \rightarrow \hat{\mathcal{S}} \rightarrow i_{1*} \mathcal{L}_{-\omega_\alpha} \rightarrow \xi^* \pi_\beta^* \mathcal{U}_2[2] \rightarrow \cdots$$

The coboundary map in this triangle lies in the group

$$\text{Hom}_{\tilde{X}}(i_{1*} \mathcal{L}_{-\omega_\alpha}, \xi^* \pi_\beta^* \mathcal{U}_2[2]) = \text{Ext}_{\tilde{X}}^2(i_{1*} \mathcal{L}_{-\omega_\alpha}, \xi^* \pi_\beta^* \mathcal{U}_2) = \mathbb{C}$$

and corresponds to a unique non-trivial extension in the latter group. Applying the functor $\text{Hom}_{\tilde{X}}(-, \hat{\mathcal{S}})$ to (21), and using (15) to compute the groups $\text{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\beta^* \mathcal{U}_2, \hat{\mathcal{S}})$ and $\text{Hom}_{\tilde{X}}^\bullet(i_{1*} \mathcal{L}_{-\omega_\alpha}, \hat{\mathcal{S}})$, we arrive at the following:

$$\begin{aligned} \text{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\beta^* \mathcal{U}_2[1], \hat{\mathcal{S}}) &= \text{Hom}_{G_2/B}^\bullet(\pi_\beta^* \mathcal{U}_2[1], \pi_\beta^* \mathcal{U}_2) = \mathbb{C}[-1]; \\ \text{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\beta^* \mathcal{U}_2, \hat{\mathcal{U}}) &= 0; \quad \text{Hom}_{\tilde{X}}^\bullet(i_{1*} \mathcal{L}_{-\omega_\alpha}, \hat{\mathcal{S}}) = 0; \quad \text{Hom}_{\tilde{X}}^\bullet(i_{1*} \mathcal{L}_{-\omega_\alpha}, \hat{\mathcal{U}}) = 0. \end{aligned}$$

We get $\text{Hom}_{\tilde{X}}^\bullet(\hat{\mathcal{S}}, \hat{\mathcal{S}}) = \text{Hom}_{\tilde{X}}(\xi^* \pi_\beta^* \mathcal{U}_2[1], \hat{\mathcal{S}}[1]) = \mathbb{C}$. \square

Lemma 8.15. *We have $\hat{\mathcal{S}} = \pi^* \hat{\mathcal{S}}$ for some $\hat{\mathcal{S}} \in \mathcal{D}^b(X)$.*

Proof. Indeed, $\widehat{\mathbb{S}}$ is the cone of the map $\mathbb{S} \rightarrow \mathbb{U}$. We give a direct proof. By Proposition 7.1 we have to make sure that the restrictions $i_1^*\widehat{\mathcal{S}}$ and $i_2^*\widehat{\mathcal{S}}$ are pulled-back from \mathbb{Q}_5 and G_2^{ad} , respectively. Applying i_2^* to (21), we immediately see that $i_2^*\widehat{\mathcal{S}} \in \pi_\beta^*\mathcal{D}^b(G_2^{\text{ad}})$. Applying i_1^* , we obtain:

$$(22) \quad \cdots \rightarrow \pi_\beta^*\mathcal{U}_2[1] \rightarrow i_1^*\widehat{\mathcal{S}} \rightarrow i_1^*i_{1*}\mathcal{L}_{-\omega_\alpha} \rightarrow \pi_\beta^*\mathcal{U}_2[2] \rightarrow \cdots$$

Applying to the above triangle the cohomology functor \mathcal{H}^0 , we compute $\mathcal{H}^0(i_1^*\widehat{\mathcal{S}}) = \mathcal{L}_{-\omega_\alpha}$, while $\mathcal{H}^{-1}(i_1^*\widehat{\mathcal{S}})$ fits into a short exact sequence:

$$(23) \quad 0 \rightarrow \pi_\beta^*\mathcal{U}_2 \rightarrow \mathcal{H}^{-1}(i_1^*\widehat{\mathcal{S}}) \rightarrow \mathcal{L}_{\omega_\beta-2\omega_\alpha} \rightarrow 0,$$

as $\mathcal{H}^{-1}(i_1^*i_{1*}\mathcal{L}_{-\omega_\alpha}) = L^1i_1^*i_{1*}\mathcal{L}_{-\omega_\alpha} = \mathcal{L}_{\omega_\beta-2\omega_\alpha}$. To see that $\mathcal{H}^{-1}(i_1^*\widehat{\mathcal{S}})$ belongs to $\pi_\alpha^*\mathcal{D}^b(\mathbb{Q}_5)$, it is sufficient to show that $\pi_{\alpha*}(\mathcal{H}^{-1}(i_1^*\widehat{\mathcal{S}}) \otimes \mathcal{L}_{-\omega_\beta}) = 0$. Tensoring (23) with $\mathcal{L}_{-\omega_\beta}$ and applying $\pi_{\alpha*}$ we see that $R^0\pi_{\alpha*}\mathcal{L}_{-2\omega_\alpha} = R^1\pi_{\alpha*}(\pi_\beta^*\mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}) = \mathcal{L}_{-2\omega_\alpha}$, hence $\pi_{\alpha*}(\mathcal{H}^{-1}(i_1^*\widehat{\mathcal{S}}) \otimes \mathcal{L}_{-\omega_\beta}) = 0$. \square

Remark 8.16. Up to tensoring with $\mathcal{L}_{-\omega_\alpha}$, the dual extension to (22) is the push-out of (11) along the map $\pi_\beta^*\mathcal{U}_2 \rightarrow \mathcal{L}_{\omega_\alpha-\omega_\beta}$ from (5):

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_\beta^*\mathcal{U}_2 & \longrightarrow & \pi_\alpha^*\mathcal{S} & \longrightarrow & \pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{L}_{\omega_\alpha-\omega_\beta} & \longrightarrow & \mathcal{G} & \longrightarrow & \pi_\beta^*\mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \longrightarrow 0. \end{array}$$

That is, we have an isomorphism $\mathcal{H}^{-1}(i_1^*\widehat{\mathcal{S}}) = \mathcal{G}^\vee \otimes \mathcal{L}_{-\omega_\alpha}$.

Proposition 8.17. *The triple $\langle \widehat{\mathbb{S}}, \mathbb{U}, \mathcal{O}_X \rangle$ of $\mathcal{D}^b(X)$ is exceptional.*

Proof. We already proved that $\langle \mathbb{U}, \mathcal{O}_X \rangle$ is exceptional. Since $\pi^*\widehat{\mathbb{S}} = \widehat{\mathcal{S}}$ and $\widehat{\mathcal{S}}$ is exceptional, so is $\widehat{\mathbb{S}}$. Furthermore, by Remark 8.12, we have $\text{Hom}_{\widehat{X}}^\bullet(\mathbb{U}, \widehat{\mathbb{S}}) = \text{Hom}_{\widehat{X}}^\bullet(\mathcal{U}, \widehat{\mathcal{S}}) = 0$. Finally from the distinguished triangle (15) and from the vanishings $\text{Hom}_{\widehat{X}}^\bullet(\mathcal{O}_{\widehat{X}}, \mathcal{U}) = 0 = \text{Hom}_{\widehat{X}}^\bullet(\mathcal{O}_{\widehat{X}}, \mathcal{S})$, we get $\text{Hom}_{\widehat{X}}^\bullet(\mathcal{O}_X, \widehat{\mathbb{S}}) = \text{Hom}_{\widehat{X}}^\bullet(\mathcal{O}_{\widehat{X}}, \widehat{\mathcal{S}}) = 0$. \square

8.6. A rectangular Lefschetz exceptional collection on X . Since X is of index 4, Proposition 8.17 suggests to consider the following collection of 12 exceptional objects on X :

$$(24) \quad \left\langle \begin{array}{cccccc} \widehat{\mathbb{S}}(-3), & \mathbb{U}(-3), & \mathcal{O}_X(-3), & \widehat{\mathbb{S}}(-2), & \mathbb{U}(-2), & \mathcal{O}_X(-2), \\ \widehat{\mathbb{S}}(-1), & \mathbb{U}(-1), & \mathcal{O}_X(-1), & \widehat{\mathbb{S}}, & \mathbb{U}, & \mathcal{O}_X \end{array} \right\rangle$$

Each block $\langle \widehat{\mathbb{S}} \otimes \mathcal{O}_X(-i), \mathbb{U} \otimes \mathcal{O}_X(-i), \mathcal{O}_X(-i) \rangle$ for $i \in [1, 3]$ is exceptional.

Lemma 8.18. *We have the following vanishing results.*

- (a) $\text{Hom}_{\widehat{X}}^\bullet(\mathbb{U}, \mathcal{O}_X(-i)) = 0$ for $i \in [1, 3]$.
- (b) $\text{Hom}_{\widehat{X}}^\bullet(\mathbb{S}, \mathcal{O}_X(-i)) = 0$ for $i \in [1, 3]$.
- (c) $\text{Hom}_{\widehat{X}}^\bullet(\mathbb{U}, \mathbb{S} \otimes \mathcal{O}_X(-1)) = 0$.
- (d) $\text{Hom}_{\widehat{X}}^\bullet(\mathbb{S}, \mathbb{S} \otimes \mathcal{O}_X(-1)) = 0$.
- (e) $\text{Hom}_{\widehat{X}}^\bullet(\mathbb{S}, \mathbb{U} \otimes \mathcal{O}_X(-1)) = 0$.
- (f) $\text{Hom}_{\widehat{X}}^\bullet(\mathcal{O}_X, \mathbb{U} \otimes \mathcal{O}_X(-i)) = \text{Hom}_{\widehat{X}}^\bullet(\mathcal{O}_X, \mathbb{S} \otimes \mathcal{O}_X(-i)) = 0$ for $i \in [1, 3]$.

- (g) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{U}, \mathbb{U} \otimes \mathcal{O}_X(-1)) = 0.$
- (h) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{S}, \mathbb{S} \otimes \mathcal{O}_X(-2)) = 0.$
- (i) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{U}, \mathbb{U} \otimes \mathcal{O}_X(-2)) = 0.$
- (j) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{U}, \mathbb{S} \otimes \mathcal{O}_X(-2)) = 0.$
- (k) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{S}, \mathbb{U} \otimes \mathcal{O}_X(-2)) = 0.$
- (l) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{S}, \mathbb{U} \otimes \mathcal{O}_X(-3)) = 0.$
- (m) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{S}, \mathbb{S} \otimes \mathcal{O}_X(-3)) = 0.$
- (n) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{U}, \mathbb{S} \otimes \mathcal{O}_X(-3)) = 0.$
- (o) $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\mathbb{U}, \mathbb{U} \otimes \mathcal{O}_X(-3)) = 0.$

Proof. Since $\pi^* : \mathcal{D}^b(X) \rightarrow \mathcal{D}^b(\tilde{X})$ is fully faithful, the above groups can be computed on \tilde{X} . Recall the isomorphism $\pi^* \mathcal{O}_X(-1) = \mathcal{O}_{\xi}(-1)$.

(a) Applying $\mathrm{Hom}_{\tilde{X}}(-, \mathcal{O}_{\xi}(-i))$ to (10), we need to compute the following groups: $\mathrm{Hom}_{\tilde{X}}(\xi^* \pi_{\beta}^* \mathcal{U}_2, \mathcal{O}_{\xi}(-i))$ and $\mathrm{Hom}_{\tilde{X}}(i_{1*} \mathcal{L}_{\omega_{\alpha} - \omega_{\beta}}, \mathcal{O}_{\xi}(-i))$. We have

$$(25) \quad \begin{aligned} \xi_* \mathcal{O}_{\xi}(-1) &= 0, & \xi_* \mathcal{O}_{\xi}(-2) &= \mathcal{L}_{-\rho}[-1], & \text{and} \\ \xi_* \mathcal{O}_{\xi}(-3) &= (\mathcal{L}_{-\omega_{\alpha} - \rho} \oplus \mathcal{L}_{-\omega_{\beta} - \rho})[-1]. \end{aligned}$$

Using this and (6), we get $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\xi^* \pi_{\beta}^* \mathcal{U}_2, \mathcal{O}_{\xi}(-i)) = 0$ for $i \in [1, 3]$. Moreover

$$\mathrm{Hom}_{\tilde{X}}^{\bullet}(i_{1*} \mathcal{L}_{\omega_{\alpha} - \omega_{\beta}}, \mathcal{O}_{\xi}(-i)) = \mathrm{H}^{\bullet}(\mathbf{Q}_5, \mathcal{L}_{-i\omega_{\alpha}}[-1]) = 0$$

for $i \in [1, 3]$.

(b) Applying $\mathrm{Hom}_{\tilde{X}}(-, \mathcal{O}_{\xi}(-1))$ to (13), we need to compute the following groups: $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\xi^* \pi_{\alpha}^* \mathcal{S}, \mathcal{O}_{\xi}(-i))$ and $\mathrm{Hom}_{\tilde{X}}^{\bullet}(i_{2*}(\pi_{\beta}^* \mathcal{U}_2^{\vee} \otimes \mathcal{L}_{-\omega_{\alpha}}), \mathcal{O}_{\xi}(-i))$.

For $i = 1$, we have $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\xi^* \pi_{\alpha}^* \mathcal{S}, \mathcal{O}_{\xi}(-1)) = 0$. For $i \in [2, 3]$, use (13) and (25). The weights of \mathcal{S}^{\vee} are $0, 2\omega_{\alpha} - \omega_{\beta}, \omega_{\beta} - \omega_{\alpha}, \omega_{\alpha}$. We see that the cohomology of $\pi_{\alpha}^* \mathcal{S}^{\vee} \otimes \mathcal{L}_{-\rho}$ and of $\pi_{\alpha}^* \mathcal{S}^{\vee} \otimes (\mathcal{L}_{-\omega_{\alpha} - \rho} \oplus \mathcal{L}_{-\omega_{\beta} - \rho})$ is trivial. Indeed their respective weights are equal to

$$\begin{aligned} &-\rho, \quad \omega_{\alpha} - 2\omega_{\beta}, \quad -2\omega_{\alpha}, \quad -\omega_{\beta}, \quad \text{and} \\ &-\omega_{\alpha} - \rho, \quad -2\omega_{\beta}, \quad -3\omega_{\alpha}, \quad -\rho - \omega_{\beta} - \rho, \quad \omega_{\alpha} - 3\omega_{\beta}, \quad -\omega_{\alpha} - \rho, \quad -2\omega_{\beta}. \end{aligned}$$

We have $\mathrm{Hom}_{\tilde{X}}^{\bullet}(i_{2*}(\pi_{\beta}^* \mathcal{U}_2^{\vee} \otimes \mathcal{L}_{-\omega_{\alpha}}), \mathcal{O}_{\xi}(-i)) = \mathrm{H}^{\bullet}(G_2^{\mathrm{ad}}, \mathcal{U}_2^{\vee} \otimes \mathcal{L}_{-i\omega_{\beta}}[-1]) = 0$ for $i \in [1, 3]$.

(c) Applying $\mathrm{Hom}_{\tilde{X}}(-, \tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1))$ to (10), we need to compute the Hom-groups $\mathrm{Hom}_{G_2/B}^{\bullet}(\pi_{\beta}^* \mathcal{U}_2, \xi_*(\tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1)))$ and $\mathrm{Hom}_{\tilde{X}}^{\bullet}(i_{1*} \mathcal{L}_{\omega_{\alpha} - \omega_{\beta}}, \tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1))$. Tensoring (13) with $\mathcal{O}_{\xi}(-1)$ and applying ξ_* , we see that $\xi_*(\tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1)) = \pi_{\beta}^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_{\alpha}}[-1]$, and $\mathrm{Hom}_{G_2/B}^{\bullet}(\pi_{\beta}^* \mathcal{U}_2, \pi_{\beta}^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_{\alpha}}[-1]) = 0$. Finally, $\mathrm{Hom}_{\tilde{X}}^{\bullet}(i_{1*} \mathcal{L}_{\omega_{\alpha} - \omega_{\beta}}, \tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1)) = \mathrm{H}^{\bullet}(G_2/B, i_1^* \tilde{\mathcal{S}} \otimes \mathcal{L}_{-\omega_{\alpha}}[-1])$. Since $i_1^* \tilde{\mathcal{S}} = \pi_{\alpha}^* \mathcal{S}$, we conclude by $\mathrm{H}^{\bullet}(G_2/B, \pi_{\alpha}^* \mathcal{S} \otimes \mathcal{L}_{-\omega_{\alpha}}) = 0$.

(d) Applying $\mathrm{Hom}_{\tilde{X}}(-, \tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1))$ to (13), we need to compute $\mathrm{Hom}_{\tilde{X}}^{\bullet}(\xi^* \pi_{\alpha}^* \mathcal{S}, \tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1))$ and $\mathrm{Hom}_{\tilde{X}}^{\bullet}(i_{2*}(\pi_{\beta}^* \mathcal{U}_2^{\vee} \otimes \mathcal{L}_{-\omega_{\alpha}}), \tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1))$. As above, we have $\xi_*(\tilde{\mathcal{S}} \otimes \mathcal{O}_{\xi}(-1)) = \pi_{\beta}^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_{\alpha}}[-1]$. The first group is

$$\mathrm{Hom}_{G_2/B}^{\bullet}(\pi_{\alpha}^* \mathcal{S}, \pi_{\beta}^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_{\alpha}}[-1]) = \mathrm{H}^{\bullet}(G_2/B, \pi_{\alpha}^* \mathcal{S}^{\vee} \otimes \pi_{\beta}^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_{\alpha}}[-1]).$$

It vanishes since the weights of $\pi_{\alpha}^* \mathcal{S}^{\vee} \otimes \pi_{\beta}^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_{\alpha}} = \pi_{\alpha}^* \mathcal{S} \otimes \pi_{\beta}^* \mathcal{U}_2$ are

$$-2\omega_{\alpha}, \quad -\omega_{\beta}, \quad \omega_{\beta} - 3\omega_{\alpha}, \quad -\omega_{\alpha}, \quad -\omega_{\beta}, \quad 2\omega_{\alpha} - 2\omega_{\beta}, \quad -\omega_{\alpha}, \quad \omega_{\alpha} - \omega_{\beta}.$$

The second group is $\mathrm{Hom}_{G_2/B}^\bullet(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}, \pi_\beta^* \mathcal{U}_2^{\oplus 2} \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{L}_{\omega_\beta - \omega_\alpha}[-1])$ and vanishes since $\mathrm{H}^\bullet(G_2/B, \pi_\beta^* \mathcal{U}_2^{\otimes 2}) = 0$.

(e) Apply $\mathrm{Hom}_{\tilde{X}}(-, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1))$ to (13). We need to compute $\mathrm{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\alpha^* \mathcal{S}, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1))$ and $\mathrm{Hom}_{\tilde{X}}^\bullet(i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1))$. Tensoring (10) with $\mathcal{O}_\xi(-1)$ and applying ξ_* we get $\xi_*(\tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = \mathcal{L}_{-\omega_\beta}[-1]$, and

$$\mathrm{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\alpha^* \mathcal{S}, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = \mathrm{H}^{\bullet-1}(G_2/B, \pi_\alpha^* \mathcal{S}^\vee \otimes \mathcal{L}_{-\omega_\beta}) = 0.$$

The second group is isomorphic to $\mathrm{Hom}_{G_2/B}^\bullet(\pi_\beta^* \mathcal{U}_2^\vee, \pi_\beta^* \mathcal{U}_2[-1]) = 0$.

(f) For $i = 0$, the statement follows from Proposition 8.17. By (d) and (e), we have $\xi_*(\tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-1)) = \pi_\beta^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha}[-1]$ and $\xi_*(\tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = \mathcal{L}_{-\omega_\beta}[-1]$, which gives the statement for $i = 1$. Note that $\tilde{\mathcal{U}}^\vee = \tilde{\mathcal{U}} \otimes \pi^* \mathcal{O}_X(1)$ therefore $\mathbb{U}^\vee = \mathbb{U} \otimes \mathcal{O}_X(1)$. By Serre duality, we have $\mathrm{H}^\bullet(X, \mathbb{U} \otimes \mathcal{O}_X(-i)) = \mathrm{H}^{7-\bullet}(X, \mathbb{U} \otimes \mathcal{O}_X(i-3))^\vee$, proving the result for $\mathbb{U} \otimes \mathcal{O}_X(-i)$ and $i = 2, 3$. Tensoring (13) with $\mathcal{O}_\xi(-i)$ for $i = 2, 3$, and applying ξ_* , we get that $\xi_*(\tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-i))$ is isomorphic to the cone of $\pi_\alpha^* \mathcal{S} \otimes \xi_* \mathcal{O}_\xi(-i) \rightarrow \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{L}_{-i\omega_\beta}$. Since $\pi_{\beta*} \mathcal{L}_{-\omega_\alpha} = 0$, we have $\mathrm{H}^\bullet(G_2/B, \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{L}_{-i\omega_\beta}) = 0$. Taking into account the fact that $\xi_* \mathcal{O}_\xi(-i) = \mathcal{S}^{i-2}(\mathcal{L}_{-\omega_\alpha} \oplus \mathcal{L}_{-\omega_\beta}) \otimes \mathcal{L}_{-\rho}[-1]$, we are reduced to showing the vanishing of the cohomology of $\pi_\alpha^* \mathcal{S} \otimes \mathcal{L}_{-\rho}$ and of $\pi_\alpha^* \mathcal{S} \otimes (\mathcal{L}_{-\omega_\alpha} \oplus \mathcal{L}_{-\omega_\beta}) \otimes \mathcal{L}_{-\rho}$. We have $\mathrm{H}^\bullet(G_2/B, \pi_\alpha^* \mathcal{S} \otimes \mathcal{L}_{-\rho}) = \mathrm{H}^\bullet(G_2/B, \pi_\alpha^* \mathcal{S} \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{L}_{-\rho}) = 0$. Remembering that $\pi_{\alpha*} \mathcal{L}_{-2\omega_\beta} = \mathcal{L}_{-3\omega_\alpha}[-1]$, we compute $\mathrm{H}^\bullet(G_2/B, \pi_\alpha^* \mathcal{S} \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{L}_{-\rho}) = \mathrm{H}^\bullet(\mathbb{Q}_5, \mathcal{S} \otimes \mathcal{L}_{-4\omega_\alpha}) = \mathrm{H}^{5-\bullet}(\mathbb{Q}_5, \mathcal{S}^\vee \otimes \mathcal{L}_{-\omega_\alpha})^\vee = \mathrm{H}^\bullet(\mathbb{Q}_5, \mathcal{S}) = 0$, by Serre duality.

(g) Apply $\mathrm{Hom}_{\tilde{X}}(-, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1))$ to (10). We need to compute the group

$$\mathrm{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\beta^* \mathcal{U}_2, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = \mathrm{Hom}_{G_2/B}^{\bullet-1}(\pi_\beta^* \mathcal{U}_2, \mathcal{L}_{-\omega_\beta}) = 0$$

and the group

$$\mathrm{Hom}_{\tilde{X}}^\bullet(i_{1*} \mathcal{L}_{\omega_\alpha - \omega_\beta}, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-1)) = \mathrm{H}^{\bullet-1}(G_2/B, \mathcal{L}_{-\omega_\alpha} \oplus \mathcal{L}_{-2\omega_\alpha}) = 0.$$

(h) Tensoring (13) with $\mathcal{O}_\xi(-2)$ and applying ξ_* , we get

$$0 \rightarrow \pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha - 2\omega_\beta} \rightarrow \mathrm{R}^1 \xi_*(\tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2)) \rightarrow \pi_\alpha^* \mathcal{S} \otimes \mathcal{L}_{-\rho} \rightarrow 0.$$

The extension is trivial as $\mathrm{Ext}_{G_2/B}^1(\pi_\alpha^* \mathcal{S}, \pi_\beta^* \mathcal{U}_2) = 0$. Thus,

$$\xi_*(\tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2)) = (\pi_\beta^* \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \oplus \pi_\alpha^* \mathcal{S} \otimes \mathcal{L}_{-\rho})[-1].$$

Applying $\mathrm{Hom}_{\tilde{X}}(-, \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2))$ to (13), we need to compute $\mathrm{Hom}_{\tilde{X}}^\bullet(i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2))$ and $\mathrm{Hom}_{\tilde{X}}^\bullet(\xi^* \pi_\alpha^* \mathcal{S}, \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2)) = \mathrm{Hom}_{G_2/B}^{\bullet-1}(\pi_\alpha^* \mathcal{S}, \pi_\beta^* \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \oplus \pi_\alpha^* \mathcal{S} \otimes \mathcal{L}_{-\rho})$.

Since $\pi_{\alpha*} \mathcal{L}_{-\omega_\beta} = 0$, we get $\mathrm{Hom}_{G_2/B}^\bullet(\pi_\alpha^* \mathcal{S}, \pi_\alpha^* \mathcal{S} \otimes \mathcal{L}_{-\rho}) = 0$. From (6), we get that $\pi_{\alpha*}(\pi_\beta^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}) = \mathcal{L}_{-2\omega_\alpha}[-1]$, hence

$$\begin{aligned} \mathrm{Hom}_{G_2/B}^\bullet(\pi_\alpha^* \mathcal{S}, \pi_\beta^* \mathcal{U}_2 \otimes \mathcal{L}_{-\rho}) &= \mathrm{Hom}_{G_2/B}^\bullet(\pi_\alpha^* \mathcal{S}, \pi_\beta^* \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta - \omega_\alpha}) \\ &= \mathrm{Hom}_{\mathbb{Q}_5}^{\bullet-1}(\mathcal{S}, \mathcal{L}_{-3\omega_\alpha}) = 0. \end{aligned}$$

Finally, we have $\mathrm{Hom}_{\tilde{X}}^\bullet(i_{2*}(\pi_\beta^* \mathcal{U}_2^\vee \otimes \mathcal{L}_{-\omega_\alpha}), \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2)) = \mathrm{Hom}_{G_2/B}^{\bullet-1}(\pi_\beta^* \mathcal{U}_2^\vee, \pi_\beta^* \mathcal{U}_2^{\oplus 2} \otimes \mathcal{L}_{-\omega_\beta}) = \mathrm{H}^{\bullet-1}(G_2/B, \pi_\beta^*(\mathcal{U}_2 \otimes \mathcal{U}_2)^{\oplus 2} \otimes \mathcal{L}_{-\omega_\beta})$. The latter group is isomorphic to

$$\mathrm{H}^{\bullet-1}(G_2/B, \mathcal{L}_{-2\omega_\beta}) \oplus \mathrm{H}^{\bullet-1}(G_2^{\mathrm{ad}}, \mathcal{S}^2 \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}) = 0$$

as $\omega_{G_2^{\text{ad}}} = \mathcal{L}_{-3\omega_\beta}$ and $S^2\mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}[-1] = \pi_{\beta*}\mathcal{L}_{-4\omega_\alpha}$ that is also acyclic, since $\omega_{Q_5} = \mathcal{L}_{-5\omega_\alpha}$.

(i) Tensoring (10) with $\mathcal{O}_\xi(-2)$ and applying ξ_* , we get

$$0 \rightarrow \mathcal{L}_{-\omega_\alpha - \omega_\beta} \rightarrow R^1 \xi_*(\tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-2)) \rightarrow \pi_\beta^*\mathcal{U} \otimes \mathcal{L}_{-\rho} \rightarrow 0.$$

The extension is trivial since $\text{Ext}_{G_2/B}^1(\pi_\beta^*\mathcal{U} \otimes \mathcal{L}_{-\rho}, \mathcal{L}_{-\rho}) = 0$. Thus, we have the equality $\xi_*(\tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-2)) = (\pi_\beta^*\mathcal{U} \otimes \mathcal{L}_{-\rho} \oplus \mathcal{L}_{-\rho})[-1]$. Applying $\text{Hom}_{\tilde{X}}(-, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-2))$ to (10), we get that the group $\text{Hom}_{\tilde{X}}^\bullet(i_{1*}\mathcal{L}_{\omega_\alpha - \omega_\beta}, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-2))$ equals

$$\text{Hom}_{G_2/B}^{\bullet-1}(\mathcal{L}_{\omega_\alpha - \omega_\beta}, (\mathcal{O}_{G_2/B} \oplus \mathcal{L}_{-\omega_\alpha}) \otimes \mathcal{L}_{-2\omega_\alpha} \otimes \mathcal{L}_{\omega_\alpha - \omega_\beta}) = 0 \quad \text{and}$$

$$\text{Hom}_{\tilde{X}}^\bullet(\xi^*\pi_\beta^*\mathcal{U}_2, \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-2)) = \text{Hom}_{G_2/B}^{\bullet-1}(\pi_\beta^*\mathcal{U}_2, \pi_\beta^*\mathcal{U} \otimes \mathcal{L}_{-\rho} \oplus \mathcal{L}_{-\rho}) = 0.$$

(j) Apply $\text{Hom}_{\tilde{X}}(-, \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2))$ to (10). We need to compute the groups

$$\text{Hom}_{\tilde{X}}^\bullet(\xi^*\pi_\beta^*\mathcal{U}_2, \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2)) = \text{Hom}_{G_2/B}^{\bullet-1}(\pi_\beta^*\mathcal{U}_2, \pi_\beta^*\mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \oplus \pi_\alpha^*\mathcal{S} \otimes \mathcal{L}_{-\rho})$$

and

$$\text{Hom}_{\tilde{X}}^\bullet(i_{1*}\mathcal{L}_{\omega_\alpha - \omega_\beta}, \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-2)) = H^{\bullet-1}(E_1, \pi_\alpha^*\mathcal{S} \otimes \mathcal{L}_{-2\omega_\alpha}).$$

We have $\text{Hom}_{G_2/B}^\bullet(\pi_\beta^*\mathcal{U}_2, \pi_\beta^*\mathcal{U}_2 \otimes \mathcal{L}_{-\rho}) = 0$ as in (i), and $\text{Hom}_{G_2/B}^\bullet(\pi_\beta^*\mathcal{U}_2, \pi_\alpha^*\mathcal{S} \otimes \mathcal{L}_{-\rho}) = 0$ by applying $\text{Hom}_{G_2/B}(-, \pi_\alpha^*\mathcal{S} \otimes \mathcal{L}_{-\rho})$ to (6). Moreover, we have $H^\bullet(G_2/B, \pi_\alpha^*\mathcal{S} \otimes \mathcal{L}_{-2\omega_\alpha}) = 0$.

For (k)-(o), use Serre duality and the previous results:

- $\text{Hom}_{\tilde{X}}^\bullet(\mathbb{S}, \mathbb{U} \otimes \mathcal{O}_X(-2)) = \text{Hom}_{\tilde{X}}^\bullet(\mathbb{U}, \mathbb{S} \otimes \mathcal{O}_X(-2)[7])^\vee = 0$ by (j);
- $\text{Hom}_{\tilde{X}}^\bullet(\mathbb{S}, \mathbb{U} \otimes \mathcal{O}_X(-3)) = \text{Hom}_{\tilde{X}}^\bullet(\mathbb{U}, \mathbb{S} \otimes \mathcal{O}_X(-1)[7])^\vee = 0$ by (c);
- $\text{Hom}_{\tilde{X}}^\bullet(\mathbb{S}, \mathbb{S} \otimes \mathcal{O}_X(-3)) = \text{Hom}_{\tilde{X}}^\bullet(\mathbb{S}, \mathbb{S} \otimes \mathcal{O}_X(-1)[7])^\vee = 0$ by (d);
- $\text{Hom}_{\tilde{X}}^\bullet(\mathbb{U}, \mathbb{S} \otimes \mathcal{O}_X(-3)) = \text{Hom}_{\tilde{X}}^\bullet(\mathbb{S}, \mathbb{U} \otimes \mathcal{O}_X(-1)[7])^\vee = 0$ by (e);
- $\text{Hom}_{\tilde{X}}^\bullet(\mathbb{U}, \mathbb{U} \otimes \mathcal{O}_X(-3)) = \text{Hom}_{\tilde{X}}^\bullet(\mathbb{U}, \mathbb{U} \otimes \mathcal{O}_X(-1)[7])^\vee = 0$ by (g).

This completes the proof. □

Corollary 8.19. *The collection (24) is exceptional.*

Proof. Using Lemma 8.18 and triangle (15), we obtain the following vanishing of Hom-groups:

- (1) $\text{Hom}_{\tilde{X}}^\bullet(\widehat{\mathbb{S}}, \mathcal{O}_X(-i)) = 0$ for $i = 1, 2, 3$, using (a) and (b) of Lemma 8.18.
- (2) $\text{Hom}_{\tilde{X}}^\bullet(\mathcal{O}_X, \widehat{\mathbb{S}}(-i)) = 0$ for $i = 0, 1, 2, 3$, using (f) of Lemma 8.18.
- (3) $\text{Hom}_{\tilde{X}}^\bullet(\widehat{\mathbb{S}}, \mathbb{U}(-i)) = 0$ for $i = 1, 2, 3$, using (c), (g), (i), (k), (l), and (o) of Lemma 8.18.
- (4) $\text{Hom}_{\tilde{X}}^\bullet(\widehat{\mathbb{S}}, \widehat{\mathbb{S}}(-i)) = 0$ for $i = 1, 2, 3$, using (c), (d), (h), (j), (l), and (m) of Lemma 8.18. □

The above statements imply:

Theorem 8.20. *The collection (24) is a rectangular Lefschetz exceptional collection on X whose length is equal to the rank of $K^0(X)$.*

8.7. **Fullness of the exceptional collection in Theorem 8.20.** Denote by \mathcal{A}_0 the admissible subcategory $\langle \widehat{\mathcal{S}}, \mathcal{U}, \mathcal{O}_X \rangle$ of $\mathcal{D}^b(X)$.

Theorem 8.21. *The semiorthogonal sequence $\mathcal{A} = \langle \mathcal{A}_0(-3), \mathcal{A}_0(-2), \mathcal{A}_0(-1), \mathcal{A}_0 \rangle$ is a rectangular Lefschetz semiorthogonal decomposition of $\mathcal{D}^b(X)$ of length 4 with the starting block \mathcal{A}_0 .*

To unburden the notation, in what follows we will suppress the pullback symbol $\xi^*(-) \in \mathcal{D}^b(\widetilde{X})$ at an object $(-) \in \mathcal{D}^b(G_2/B)$, and whenever it doesn't lead to confusion we will also be suppressing the pullback symbols π_α^* and π_β^* at objects in $\mathcal{D}^b(\mathbb{Q}_5)$ and $\mathcal{D}^b(G_2^{\text{ad}})$, respectively. Thus, typically an object like \mathcal{U}_2 means $\xi^*\pi_\beta^*\mathcal{U}_2$, etc., or $\pi_\beta^*\mathcal{U}_2$, etc., depending on the variety an object is being considered.

Proof. By [34, 6.4], the category $\mathcal{D}^b(G_2^{\text{ad}})$ has a full exceptional collection

$$\langle \mathcal{U}_2 \otimes \mathcal{L}_{-2\omega_\beta}, \mathcal{L}_{-2\omega_\beta}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}, \mathcal{L}_{-\omega_\beta}, \mathcal{U}_2, \mathcal{O}_{G_2^{\text{ad}}} \rangle.$$

By [42], there is a semiorthogonal decomposition $\mathcal{D}^b(G_2/B) = \langle \pi_\beta^*\mathcal{D}^b(G_2^{\text{ad}}) \otimes \mathcal{L}_{-\omega_\alpha}, \pi_\beta^*\mathcal{D}^b(G_2^{\text{ad}}) \rangle$, and combining these two statements we obtain a full exceptional collection in $\mathcal{D}^b(G_2/B)$:

$$(26) \quad \begin{aligned} & \mathcal{U}_2 \otimes \mathcal{L}_{-2\omega_\beta}, \mathcal{L}_{-2\omega_\beta}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}, \mathcal{L}_{-\omega_\beta}, \mathcal{U}_2, \mathcal{O}_{\widetilde{X}} \rangle \\ & \langle \mathcal{U}_2 \otimes \mathcal{L}_{-2\omega_\beta - \omega_\alpha}, \mathcal{L}_{-2\omega_\beta - \omega_\alpha}, \mathcal{U}_2 \otimes \mathcal{L}_{-\rho}, \mathcal{L}_{-\rho}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha}, \mathcal{L}_{-\omega_\alpha}, \end{aligned}$$

Remembering that ξ is a \mathbb{P}^1 -bundle and using once again [42], we obtain a semiorthogonal decomposition $\mathcal{D}^b(\widetilde{X}) = \langle \xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1), \xi^*\mathcal{D}^b(G_2/B) \rangle$. Taking the collection (26) in $\mathcal{D}^b(G_2/B)$ and tensoring it with $\mathcal{O}_\xi(-1)$, we obtain a full exceptional collection in $\mathcal{D}^b(\widetilde{X})$. We first perform a series of mutations of the collection thus obtained inside $\mathcal{D}^b(\widetilde{X}) = \langle \xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1), \xi^*\mathcal{D}^b(G_2/B) \rangle$. Throughout, we use Lemma 6.11.

Step 1. Starting from the decomposition (26) and recalling that $\omega_{G_2^{\text{ad}}}^{-1} = \mathcal{L}_{3\omega_\beta}$, we mutate $\mathcal{U}_2 \otimes \mathcal{L}_{-2\omega_\beta}$ to the right inside $\pi_\beta^*\mathcal{D}^b(G_2^{\text{ad}})$, obtaining a full exceptional collection in $\xi^*\mathcal{D}^b(G_2/B) = \xi^*\langle \pi_\beta^*\mathcal{D}^b(G_2^{\text{ad}}) \otimes \mathcal{L}_{-\omega_\alpha}, \pi_\beta^*\mathcal{D}^b(G_2^{\text{ad}}) \rangle$:

$$(27) \quad \begin{aligned} & \mathcal{L}_{-2\omega_\beta}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}, \mathcal{L}_{-\omega_\beta}, \mathcal{U}_2, \mathcal{O}_{\widetilde{X}}, \mathcal{U}_2^* \rangle \\ & \langle \mathcal{L}_{-2\omega_\beta - \omega_\alpha}, \mathcal{U}_2 \otimes \mathcal{L}_{-\rho}, \mathcal{L}_{-\rho}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha}, \mathcal{L}_{-\omega_\alpha}, \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha}, \end{aligned}$$

Step 2. Recall that $\omega_{G_2/B}^{-1} = \mathcal{L}_{2\rho}$ and mutate $\mathcal{L}_{-2\omega_\beta - \omega_\alpha}$ to the right in $\xi^*\mathcal{D}^b(G_2/B)$, we obtain

$$(28) \quad \begin{aligned} & \mathcal{L}_{-2\omega_\beta}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}, \mathcal{L}_{-\omega_\beta}, \mathcal{U}_2, \mathcal{O}_{\widetilde{X}}, \mathcal{U}_2^*, \mathcal{L}_{\omega_\alpha} \rangle \\ & \langle \mathcal{U}_2 \otimes \mathcal{L}_{-\rho}, \mathcal{L}_{-\rho}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha}, \mathcal{L}_{-\omega_\alpha}, \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha}, \end{aligned}$$

Step 3. Consider $\mathcal{D}^b(\widetilde{X}) = \langle \xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1), \xi^*\mathcal{D}^b(G_2/B) \rangle$, and let the full exceptional collection in $\xi^*\mathcal{D}^b(G_2/B)$ be the one from (28), and the collection in $\xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1)$ be the one from (27) tensored with $\mathcal{O}_\xi(-1)$.

Step 4. In $\xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1)$, consider the term $\mathcal{L}_{-2\omega_\beta} \otimes \mathcal{O}_\xi(-1) \in \xi^*\pi_\beta^*\mathcal{D}^b(G_2^{\text{ad}})$ and mutate it to the right inside $\xi^*\pi_\beta^*\mathcal{D}^b(G_2^{\text{ad}}) \otimes \mathcal{O}_\xi(-1) \subset \xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1)$, obtaining $\mathcal{L}_{\omega_\beta} \otimes \mathcal{O}_\xi(-1)$ and a full exceptional collection in $\xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1)$:

$$(29) \quad \begin{aligned} & \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{O}_\xi(-1), \mathcal{O}_\xi(-1), \mathcal{U}_2^* \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{\omega_\beta} \otimes \mathcal{O}_\xi(-1) \\ & \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \\ & \langle \mathcal{L}_{-2\omega_\beta-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \end{aligned}$$

Step 5. In (29), mutate $\mathcal{L}_{\omega_\beta} \otimes \mathcal{O}_\xi(-1)$ to the left inside $\xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1)$, obtaining $\mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1)$ and a full exceptional collection in $\xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1)$:

$$(30) \quad \begin{aligned} & \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{O}_\xi(-1), \mathcal{O}_\xi(-1), \mathcal{U}_2^* \otimes \mathcal{O}_\xi(-1) \\ & \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \\ & \langle \mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-2\omega_\beta-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \end{aligned}$$

Note that $\text{Hom}_{\tilde{X}}^\bullet(\mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-2\omega_\beta-\omega_\alpha} \otimes \mathcal{O}_\xi(-1)) = \text{H}^\bullet(G_2/B, \mathcal{L}_{\omega_\alpha-\omega_\beta}) = 0$ and also $\text{Hom}_{\tilde{X}}^\bullet(\mathcal{L}_{-2\omega_\beta-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1)) = \text{H}^\bullet(G_2/B, \mathcal{L}_{\omega_\beta-\omega_\alpha}) = 0$. Thus, mutating $\mathcal{L}_{-2\omega_\beta-\omega_\alpha} \otimes \mathcal{O}_\xi(-1)$ in (30) to the left through $\mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1)$ amounts to just interchanging these two objects. We obtain

$$(31) \quad \begin{aligned} & \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{O}_\xi(-1), \mathcal{O}_\xi(-1), \mathcal{U}_2^* \otimes \mathcal{O}_\xi(-1) \\ & \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \\ & \langle \mathcal{L}_{-2\omega_\beta-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \end{aligned}$$

Step 6. In (31) we mutate $\mathcal{L}_{-2\omega_\beta-\omega_\alpha} \otimes \mathcal{O}_\xi(-1)$ to the right inside $\xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1)$, obtaining $\mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1)$ and a full exceptional collection in $\xi^*\mathcal{D}^b(G_2/B) \otimes \mathcal{O}_\xi(-1)$:

$$(32) \quad \begin{aligned} & \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{O}_\xi(-1), \mathcal{O}_\xi(-1), \mathcal{U}_2^* \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1) \\ & \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \\ & \langle \mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \end{aligned}$$

Step 7. Finally, recalling that $\omega_{\tilde{X}}^{-1} = \xi^*\mathcal{L}_\rho \otimes \mathcal{O}_\xi(2)$ we mutate $\mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1) \in \mathcal{D}^b(\tilde{X})$ to the right inside the whole of $\mathcal{D}^b(\tilde{X})$, obtaining $\mathcal{L}_{-\omega_\beta-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1) \otimes \xi^*\mathcal{L}_\rho \otimes \mathcal{O}_\xi(2) = \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(1) = \mathcal{O}_{\tilde{X}}(E_2)$ and a full exceptional collection in $\mathcal{D}^b(\tilde{X})$:

$$(33) \quad \begin{aligned} & \mathcal{L}_{-2\omega_\beta}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}, \mathcal{L}_{-\omega_\beta}, \mathcal{U}_2, \mathcal{O}_{\tilde{X}}, \mathcal{U}_2^*, \mathcal{L}_{\omega_\alpha}, \mathcal{O}_{\tilde{X}}(E_2) \\ & \mathcal{U}_2^* \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\rho}, \mathcal{L}_{-\rho}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha}, \mathcal{L}_{-\omega_\alpha}, \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha}, \\ & \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{O}_\xi(-1), \mathcal{O}_\xi(-1), \\ & \langle \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \end{aligned}$$

We now assume that the semiorthogonal sequence \mathcal{A} is not a semiorthogonal decomposition, that is, that there exists a nonzero object $E \in {}^\perp\mathcal{A}$.

Lemma 8.22. *Let \mathcal{B} denote the admissible subcategory $\langle \mathcal{U}_2^*, \mathcal{L}_{\omega_\alpha}, \mathcal{O}_{\tilde{X}}(E_2) \rangle \subset \mathcal{D}^b(\tilde{X})$ of (33). Consider a nonzero object $E \in {}^\perp \mathcal{A}$. Then $\pi^*E \in \mathcal{B}$.*

Proof. We have a semiorthogonal decomposition $\mathcal{D}^b(\tilde{X}) = \langle \mathcal{B}^\perp, \mathcal{B} \rangle$. The statement amounts to showing that π^*E is left orthogonal to \mathcal{B}^\perp , that is to all the objects to the left from $\mathcal{U}_2^*, \mathcal{L}_{\omega_\alpha}, \mathcal{O}_{\tilde{X}}(E_2)$ in (33). We break up the proof into several steps. We first recall the basic short exact sequences that will be used throughout:

$$(34) \quad 0 \rightarrow \mathcal{L}_{-\omega_\alpha} \rightarrow \mathcal{U}_2 \rightarrow \mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow 0.$$

$$(35) \quad 0 \rightarrow \tilde{\mathcal{U}} \rightarrow \mathcal{U}_2 \rightarrow i_{1*} \mathcal{L}_{\omega_\alpha - \omega_\beta} \rightarrow 0,$$

$$(36) \quad 0 \rightarrow \mathcal{U}_2 \rightarrow \mathcal{S} \rightarrow \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \rightarrow 0.$$

$$(37) \quad 0 \rightarrow \tilde{\mathcal{S}} \rightarrow \mathcal{S} \rightarrow i_{2*}(\mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha}) \rightarrow 0.$$

Note that the triangulated subcategory $\pi^* \mathcal{A}_0 = \langle \hat{\mathcal{S}}, \tilde{\mathcal{U}}, \mathcal{O}_{\tilde{X}} \rangle \subset \mathcal{D}^b(\tilde{X})$ is equivalent to a subcategory $\pi^* \tilde{\mathcal{A}}_0 \subset \mathcal{D}^b(\tilde{X})$, where $\tilde{\mathcal{A}}_0 \subset \mathcal{D}^b(X)$ is a subcategory generated by $\langle \tilde{\mathcal{S}}, \mathbb{U}, \mathcal{O}_X \rangle$ (recall Definition 8.11 of the object $\hat{\mathcal{S}}$). Write $\tilde{\mathcal{A}} = \langle \tilde{\mathcal{A}}_0 \otimes \mathcal{O}_X(-3), \tilde{\mathcal{A}}_0 \otimes \mathcal{O}_X(-2), \tilde{\mathcal{A}}_0 \otimes \mathcal{O}_X(-1), \tilde{\mathcal{A}}_0 \rangle \subset \mathcal{D}^b(X)$; we have $\tilde{\mathcal{A}} = \mathcal{A}$ and $E \in {}^\perp \tilde{\mathcal{A}}, \pi^*E \in \pi^{*\perp} \tilde{\mathcal{A}}$. We show that the latter orthogonality condition implies that $\pi^*E \in \mathcal{B}$.

Step 1. We prove the inclusion

$$\pi^*E \in {}^\perp \langle \pi^* \tilde{\mathcal{A}}, \pi^* \tilde{\mathcal{A}} \otimes \mathcal{O}_{\tilde{X}}(E_1), \pi^* \tilde{\mathcal{A}} \otimes \mathcal{O}_{\tilde{X}}(E_2), \pi^* \tilde{\mathcal{A}} \otimes \mathcal{O}_{\tilde{X}}(E_1 + E_2) \rangle.$$

Indeed, isomorphisms $\pi_* \mathcal{O}_{\tilde{X}}(E_1) = \pi_* \mathcal{O}_{\tilde{X}}(E_2) = \pi_* \mathcal{O}_{\tilde{X}}(E_1 + E_2) = \mathcal{O}_X$ imply that $\mathrm{Hom}_{\tilde{X}}^\bullet(\pi^*E, \pi^* \tilde{\mathcal{A}} \otimes \mathcal{O}_{\tilde{X}}(E_1)) = \mathrm{Hom}_{\tilde{X}}^\bullet(\pi^*E, \pi^* \tilde{\mathcal{A}} \otimes \mathcal{O}_{\tilde{X}}(E_2)) = \mathrm{Hom}_{\tilde{X}}^\bullet(\pi^*E, \pi^* \tilde{\mathcal{A}} \otimes \mathcal{O}_{\tilde{X}}(E_1 + E_2)) = \mathrm{Hom}_X^\bullet(E, \tilde{\mathcal{A}}) = 0$.

Step 2. Taking $\mathcal{O}_\xi(-i) = \pi^* \mathcal{O}_X(-i)$ for $i = 0, 1, 2, 3$ and tensoring it with $\mathcal{O}_X(E_1) = \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(1)$ and $\mathcal{O}_X(E_2) = \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(1)$ (resp., with $\mathcal{O}_X(E_1 + E_2) = \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(2)$), we have $\pi^*E \in {}^\perp \langle \mathcal{O}_\xi(-i) \rangle$ for $i = 0, 1, 2, 3$, $\pi^*E \in {}^\perp \langle \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-i), \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-i) \rangle$ for $i = -1, 0, 1, 2$, $\pi^*E \in {}^\perp \langle \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-i) \rangle$ for $i = -2, -1, 0, 1$.

Taking into account that $\pi_* i_{1*} \mathcal{L}_{\omega_\alpha - \omega_\beta} = \pi_* i_{2*}(\mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha}) = 0$ we see from (35) and (37) that $\pi^*E \in {}^\perp \langle \mathcal{U}_2, \mathcal{S} \rangle$. Indeed, applying $\mathrm{Hom}_{\tilde{X}}(\pi^*E, -)$ to the first sequence (35), we obtain an isomorphism $\mathrm{Hom}_{\tilde{X}}(\pi^*E, \tilde{\mathcal{U}}) = \mathrm{Hom}_{\tilde{X}}(\pi^*E, \mathcal{U}_2)$, since $\mathrm{Hom}_{\tilde{X}}(\pi^*E, i_{1*} \mathcal{L}_{\omega_\alpha - \omega_\beta}) = \mathrm{Hom}_{\tilde{X}}(E, \pi_* i_{1*} \mathcal{L}_{\omega_\alpha - \omega_\beta}) = 0$, the last isomorphism being a consequence of $\pi_* i_{1*} \mathcal{L}_{\omega_\alpha - \omega_\beta} = 0$. Now $\mathrm{Hom}_{\tilde{X}}(\pi^*E, \tilde{\mathcal{U}}) = 0$ by the assumption on E : it is supposed to be a non-trivial object of the left orthogonal to \mathcal{A} and $\tilde{\mathcal{U}} \in \pi^* \mathcal{A}$. It follows that $\mathrm{Hom}_{\tilde{X}}(\pi^*E, \mathcal{U}_2) = 0$, as claimed.

Tensoring (35) and (37) with $\mathcal{O}_\xi(-i)$ for $i = 1, 2, 3$, we obtain, respectively

$$(38) \quad 0 \rightarrow \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-i) \rightarrow \mathcal{U}_2 \otimes \mathcal{O}_\xi(-i) \rightarrow i_{1*} \mathcal{L}_{(1-i)\omega_\alpha - \omega_\beta} \rightarrow 0,$$

and

$$(39) \quad 0 \rightarrow \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-i) \rightarrow \mathcal{S} \otimes \mathcal{O}_\xi(-i) \rightarrow i_{2*}(\mathcal{U}_2^* \otimes \mathcal{L}_{-i\omega_\beta - \omega_\alpha}) \rightarrow 0.$$

Taking into account that $\pi_* i_{1*} \mathcal{L}_{(1-i)\omega_\alpha - \omega_\beta} = \pi_* i_{2*} (\mathcal{U}_2^* \otimes \mathcal{L}_{-i\omega_\beta - \omega_\alpha}) = 0$ we obtain $\pi^* E \in {}^\perp \langle \mathcal{U}_2 \otimes \mathcal{O}_\xi(-i), \mathcal{S} \otimes \mathcal{O}_\xi(-i) \rangle$ for $i = 1, 2, 3$. Tensoring (36) with $\mathcal{O}_\xi(-i)$ for $i = 0, 1, 2, 3$ we obtain $\pi^* E \in {}^\perp \langle \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-i) \rangle$ for $i = 0, 1, 2, 3$.

Tensoring (38) with $\mathcal{O}_{\tilde{X}}(E_2) = \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(1)$ and (39) with $\mathcal{O}_{\tilde{X}}(E_1) = \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(1)$ we obtain, respectively:

$$(40) \quad \begin{aligned} 0 \rightarrow \tilde{\mathcal{U}} \otimes \mathcal{O}_\xi(-i) \otimes \mathcal{O}_{\tilde{X}}(E_2) &\rightarrow \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(1-i) \rightarrow \\ &i_{1*} \mathcal{L}_{(1-i)\omega_\alpha - \omega_\beta} \otimes \mathcal{O}_{\tilde{X}}(E_2) \rightarrow 0, \end{aligned}$$

and

$$(41) \quad \begin{aligned} 0 \rightarrow \tilde{\mathcal{S}} \otimes \mathcal{O}_\xi(-i) \otimes \mathcal{O}_{\tilde{X}}(E_1) &\rightarrow \mathcal{S} \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(1-i) \rightarrow \\ &i_{2*} (\mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha}) \otimes \mathcal{O}_{\tilde{X}}(E_1) \rightarrow 0. \end{aligned}$$

Taking into account isomorphisms $i_{1*} \mathcal{L}_{(1-i)\omega_\alpha - \omega_\beta} \otimes \mathcal{O}_{\tilde{X}}(E_2) = i_{1*} \mathcal{L}_{(1-i)\omega_\alpha - \omega_\beta}$ and $i_{2*} (\mathcal{U}_2^* \otimes \mathcal{L}_{-i\omega_\beta - \omega_\alpha}) \otimes \mathcal{O}_{\tilde{X}}(E_1) = i_{2*} (\mathcal{U}_2^* \otimes \mathcal{L}_{-i\omega_\beta - \omega_\alpha})$, we obtain

$$\pi^* E \in {}^\perp \langle (\mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha}) \otimes \mathcal{O}_\xi(1-i), (\mathcal{S} \otimes \mathcal{L}_{-\omega_\beta}) \otimes \mathcal{O}_\xi(1-i) \rangle$$

for $i = 0, 1, 2, 3$. Moreover, tensoring the relative Euler sequence

$$(42) \quad 0 \rightarrow \mathcal{O}_\xi(-1) \rightarrow \mathcal{L}_{-\omega_\beta} \oplus \mathcal{L}_{-\omega_\alpha} \rightarrow \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(1) \rightarrow 0$$

with $\mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1)$, we obtain

$$0 \rightarrow \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-2) \rightarrow (\mathcal{L}_{-\rho} \oplus \mathcal{L}_{-2\omega_\alpha}) \otimes \mathcal{O}_\xi(-1) \rightarrow \mathcal{L}_{-\rho - \omega_\alpha} \rightarrow 0.$$

We have $\pi_*(\mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-2)) = \pi_*(\mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1)) = \mathcal{O}_X(-3)$, while $\pi_*(\mathcal{L}_{-2\omega_\alpha} \otimes \mathcal{O}_\xi(-1))$ can be found from the short exact sequence

$$(43) \quad 0 \rightarrow \mathcal{L}_{-2\omega_\alpha} \rightarrow \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \rightarrow \mathcal{L}_{-\omega_\beta} \rightarrow 0.$$

We deduce that $\pi_* \mathcal{L}_{-2\omega_\alpha} = \text{Cone}(\mathbb{U}(-1) \rightarrow \mathcal{O}_X(-1))$. Thus, $\pi^* E \in {}^\perp \langle \mathcal{L}_{-\rho - \omega_\alpha} \rangle$. This also implies $\pi^* E \in {}^\perp \langle \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-i) \rangle$ for $i = 0, 1$ via tensoring (36) with $\mathcal{L}_{-\omega_\beta}$.

Tensoring (42) with $\mathcal{L}_{\omega_\alpha}$ we obtain:

$$0 \rightarrow \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1) \rightarrow \mathcal{L}_{\omega_\alpha - \omega_\beta} \oplus \mathcal{O}_{\tilde{X}} \rightarrow \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(1) \rightarrow 0,$$

Using that $\mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(1) = \mathcal{O}_{\tilde{X}}(E_1)$, while $\mathcal{L}_{\omega_\alpha - \omega_\beta} \in \langle \mathcal{U}_2, \mathcal{L}_{-\omega_\alpha} \rangle$, we also get that $\pi^* E \in {}^\perp \langle \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1) \rangle$. Tensoring (42) with $\mathcal{U}_2 \otimes \mathcal{O}_\xi(-i)$, $i = 1, 2$, we obtain

$$(44) \quad \begin{aligned} 0 \rightarrow \mathcal{U}_2 \otimes \mathcal{O}_\xi(-i-1) &\rightarrow (\mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \oplus \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}) \otimes \mathcal{O}_\xi(-i) \rightarrow \\ &\mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(1-i) \rightarrow 0 \end{aligned}$$

We have seen that $\pi^* E \in {}^\perp \langle \mathcal{U}_2 \otimes \mathcal{O}_\xi(-i-1) \rangle$ and $\pi^* E \in {}^\perp \langle \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-i) \rangle$ for $i = 1, 2$. For this we refer to the lines after (39) and (41). Tensoring (36) with $\mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-i-1)$ and using again the lines after (41) we obtain that $\pi^* E \in {}^\perp \langle \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-i-1) \rangle$. Hence $\pi^* E \in {}^\perp \langle \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(1-i) \rangle$ for $i = 1, 2$.

We conclude from Steps 1 and 2 that $\pi^* E$ is left orthogonal to the following objects:

$$\begin{aligned}
 (45) \quad & \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta}, \mathcal{L}_{-\omega_\beta}, \mathcal{L}_{-\omega_\alpha}, \mathcal{U}_2, \mathcal{O}_{\tilde{X}} \\
 & \mathcal{O}_\xi(-1), \mathcal{L}_{\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\rho}, \mathcal{L}_{-\rho}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha}, \\
 & \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\beta} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{O}_\xi(-1), \\
 & \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\rho} \otimes \mathcal{O}_\xi(-1), \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1), \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(-1),
 \end{aligned}$$

Step 3. We check that $\pi^*E \in {}^\perp\langle \mathcal{U}_2^* \otimes \mathcal{O}_\xi(-1) \rangle$. This follows immediately from the short exact sequence representing the vector bundle $\tilde{\mathcal{S}}$ as an extension of two vector bundles (cf. the diagram before Proposition 8.9):

$$0 \rightarrow \mathcal{U}_2 \rightarrow \tilde{\mathcal{S}} \rightarrow \mathcal{U}_2^* \otimes \mathcal{O}_\xi(-1) \rightarrow 0.$$

Step 4. We check that $\pi^*E \in {}^\perp\langle \mathcal{L}_{-2\omega_\beta} \rangle$. Tensoring (34) with $\mathcal{L}_{-\rho}$, we get

$$0 \rightarrow \mathcal{L}_{-\rho-\omega_\alpha} \rightarrow \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \rightarrow \mathcal{L}_{-2\omega_\beta} \rightarrow 0.$$

We have $\pi^*E \in {}^\perp\langle \mathcal{L}_{-\rho-\omega_\alpha} \rangle$ (see the line after (43)) and $\pi^*E \in {}^\perp\langle \mathcal{U}_2 \otimes \mathcal{L}_{-\rho} \rangle$ (see the paragraph after (44)). This implies the statement. \square

Lemma 8.22 implies that $\pi^*E \in \mathcal{B}$ belongs to the full subcategory generated by $\langle \mathcal{U}_2^*, \mathcal{L}_{\omega_\alpha}, \mathcal{O}_{\tilde{X}}(E_2) \rangle$. We conclude the proof by showing that an object of the form π^*E belonging to the admissible subcategory $\langle \mathcal{U}_2^*, \mathcal{L}_{\omega_\alpha}, \mathcal{O}_{\tilde{X}}(E_2) \rangle \subset \mathcal{D}^b(\tilde{X})$ is necessarily trivial, i.e. $E = 0$. The first step is to mutate the subcategory $\langle \mathcal{U}_2^*, \mathcal{L}_{\omega_\alpha} \rangle$ to the right through $\langle \mathcal{O}_{\tilde{X}}(E_2) \rangle$. One computes $\mathrm{Hom}_{\tilde{X}}^\bullet(\mathcal{L}_{\omega_\alpha}, \mathcal{O}_{\tilde{X}}(E_2)) = \mathrm{Hom}_{\tilde{X}}^\bullet(\mathcal{L}_{\omega_\alpha} \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(1)) = \mathrm{H}^\bullet(\tilde{X}, \mathcal{L}_{-2\omega_\alpha} \otimes \mathcal{O}_\xi(1)) = \mathrm{H}^\bullet(G_2/B, \mathcal{L}_{-\omega_\alpha} \oplus \mathcal{L}_{-2\omega_\alpha+\omega_\beta}) = k[-1]$, hence the right mutation $\mathbf{R} := \mathbf{R}_{\mathcal{O}_{\tilde{X}}(E_2)}(\mathcal{L}_{\omega_\alpha})$ of $\mathcal{L}_{\omega_\alpha}$ through $\langle \mathcal{O}_{\tilde{X}}(E_2) \rangle$ is given by

$$(46) \quad \cdots \rightarrow \mathcal{L}_{\omega_\alpha}[-1] \rightarrow \mathcal{O}_{\tilde{X}}(E_2) \rightarrow \mathbf{R} \rightarrow \mathcal{L}_{\omega_\alpha} \rightarrow \cdots$$

which is in fact a short exact sequence $0 \rightarrow \mathcal{O}_{\tilde{X}}(E_2) \rightarrow \mathbf{R} \rightarrow \mathcal{L}_{\omega_\alpha} \rightarrow 0$ corresponding to a unique non-trivial extension in $\mathrm{Ext}^1(\mathcal{L}_{\omega_\alpha}, \mathcal{O}_{\tilde{X}}(E_2)) = k$ above. Further, $\mathrm{Hom}_{\tilde{X}}(\mathcal{U}_2^*, \mathcal{O}_{\tilde{X}}(E_2)) = \mathrm{H}^\bullet(\tilde{X}, \mathcal{U}_2 \otimes \mathcal{L}_{-\omega_\alpha} \otimes \mathcal{O}_\xi(1)) = \mathrm{H}^\bullet(G_2/B, \mathcal{U}_2 \oplus \mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\alpha}) = 0$. Thus, the right mutation of \mathcal{U}_2^* through $\mathcal{O}_{\tilde{X}}(E_2)$ is just a transposition, and we arrive at the collection $\langle \mathcal{O}_{\tilde{X}}(E_2), \mathcal{U}_2^*, \mathbf{R} \rangle$.

We have seen that $\mathrm{Hom}_{\tilde{X}}^\bullet(\pi^*E, \mathcal{O}_{\tilde{X}}(E_2)) = \mathrm{Hom}_{\tilde{X}}^\bullet(E, \pi_*\mathcal{O}_{\tilde{X}}(E_2)) = 0$ coming from $\pi_*\mathcal{O}_{\tilde{X}}(E_2) = \mathcal{O}_X$. Thus, the object π^*E fits into an exact triangle

$$\cdots \rightarrow \mathbf{R} \otimes V_1^\bullet \rightarrow \pi^*E \rightarrow \mathcal{U}_2^* \otimes V_2^\bullet \rightarrow \cdots,$$

where V_1^\bullet and V_2^\bullet are graded vector spaces. Restricting that triangle to E_1 , then tensoring it with $\mathcal{L}_{-\omega_\beta}$, and finally applying $\pi_{\alpha*}$, we obtain an exact triangle

$$(47) \quad \cdots \rightarrow \pi_{\alpha*}(i_1^*\mathbf{R} \otimes \mathcal{L}_{-\omega_\beta}) \otimes V_1^\bullet \rightarrow \pi_{\alpha*}(i_1^*\pi^*E \otimes \mathcal{L}_{-\omega_\beta}) \rightarrow \pi_{\alpha*}(\mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\beta}) \otimes V_2^\bullet \rightarrow \cdots$$

The middle term $\pi_{\alpha*}(i_1^*\pi^*E \otimes \mathcal{L}_{-\omega_\beta})$ is equal to 0, while the term $\pi_{\alpha*}(i_1^*\mathbf{R} \otimes \mathcal{L}_{-\omega_\beta}) \otimes V_1^\bullet$ can be read off of the sequence (46):

$$(48) \quad \cdots \rightarrow \pi_{\alpha*}\mathcal{L}_{\omega_\alpha-\omega_\beta}[-1] \rightarrow \pi_{\alpha*}(\mathcal{O}_{E_1} \otimes \mathcal{L}_{-\omega_\beta}) \rightarrow \pi_{\alpha*}(i_1^*\mathbf{R} \otimes \mathcal{L}_{-\omega_\beta}) \rightarrow \pi_{\alpha*}\mathcal{L}_{\omega_\alpha-\omega_\beta} \rightarrow \cdots$$

and since $\pi_{\alpha*}\mathcal{L}_{\omega_\alpha-\omega_\beta} = \pi_{\alpha*}(\mathcal{O}_{E_1} \otimes \mathcal{L}_{-\omega_\beta}) = 0$, we obtain $\pi_{\alpha*}(i_1^*\mathbf{R} \otimes \mathcal{L}_{-\omega_\beta}) = 0$. On the other hand, $\pi_{\alpha*}(\mathcal{U}_2^* \otimes \mathcal{L}_{-\omega_\beta}) = \mathcal{L}_{-\omega_\alpha}$ as can be seen from (34). Using also (47) we conclude that $V_2^\bullet = 0$. Thus, $\pi^*E = \mathbf{R} \otimes V_1^\bullet$. Recall that \mathbf{R} is the middle term in the short exact sequence $0 \rightarrow \mathcal{O}_{\tilde{X}}(E_2) \rightarrow \mathbf{R} \rightarrow \mathcal{L}_{\omega_\alpha} \rightarrow 0$, hence \mathbf{R} is a locally free sheaf having a non-trivial class in $K^0(\tilde{X})$. For consistency of the calculations, let us check that \mathbf{R} belongs to $\pi^*\mathcal{D}^b(X)$. Indeed, restricting \mathbf{R} to E_1 we obtain a short exact sequence $0 \rightarrow \mathcal{O}_{E_1} \rightarrow i_1^*\mathbf{R} \rightarrow \mathcal{L}_{\omega_\alpha} \rightarrow 0$, hence $i_1^*\mathbf{R}$ is pulled back from \mathbf{Q}_5 and has a trivial restriction to the fibers of π_α . Restricting \mathbf{R} to E_2 we obtain a short exact sequence $0 \rightarrow \mathcal{L}_{\omega_\beta-\omega_\alpha} \rightarrow i_2^*\mathbf{R} \rightarrow \mathcal{L}_{\omega_\alpha} \rightarrow 0$, and then $i_2^*\mathbf{R} = \mathcal{U}_2^*$, hence $i_2^*\mathbf{R}$ is pulled back from G_2^{ad} and has a trivial restriction to the fibers of π_β .


Let \mathbb{R} denote the vector bundle on X such that $\pi^*\mathbb{R} = \mathbf{R}$. Now $E = \mathbb{R} \otimes V_1^\bullet \in {}^\perp\mathcal{A}$, where \mathcal{A} is a full subcategory of $\mathcal{D}^b(X)$ generated by a semiorthogonal sequence. We have $\text{Hom}_X^\bullet(E, \mathcal{A}) = \text{Hom}_X^\bullet(\mathbb{R} \otimes V_1^\bullet, \mathcal{A}) = V_1^\bullet \otimes \text{Hom}_X^\bullet(\mathbb{R}, \mathcal{A}) = 0$, where the second isomorphism holds since $E = \mathbb{R} \otimes V_1^\bullet \in \mathcal{D}^b(X)$, and hence V_1^\bullet is a bounded complex of finite-dimensional vector spaces. In the case where $V_1^\bullet \neq 0$ we necessarily obtain that $\text{Hom}_X^\bullet(\mathbb{R}, \mathcal{A}) = 0$, but this would contradict the fact that the terms of \mathcal{A} generate the group $K^0(X) \cong \mathbb{Z}^{12}$. Thus $V_1^\bullet = 0$, and $E = 0$.

Therefore, ${}^\perp\mathcal{A} = 0$, and the sequence \mathcal{A} is a semiorthogonal decomposition, in other words the collection (24) is full. \square


9. APPENDIX

In this appendix, we give the beginning (these are infinite graphs) of the quantum Hasse diagrams in case (1), $n = 3$, case (2), case (3) $n = 3 = m$ and case (5). Each column corresponds to a degree starting from the left with degree zero. The black vertices correspond to the classes σ'_u and where there are two classes of the same degree, the classes are σ'_u and $\sigma'_{u'}$ and the top class is always σ'_u while the bottom one is $\sigma'_{u'}$. We use the same convention for blue vertices which correspond to τ_v classes. Black and blue edges are from the Hasse diagram of Y and Z . Grey edges are from the Hasse diagram of Y to the one of Z while dotted edges correspond to quantum multiplication.


9.1. **Case (1), $n = 3$.** We give the quantum hasse diagram in case (1) for $n = 3$.




9.2. **Case (2).** We give the quantum hasse diagram in case (2).



9.3. **Case (3)**, $n = 3 = m$. We give the quantum hasse diagram in case (3) for $n = 3 = m$.



9.4. **Case (5)**. We give the quantum hasse diagram in case (5).



REFERENCES

[1] M. Atiyah. On the Krull-Schmidt theorem with application to sheaves. *Bull. Soc. Math. France*, 84:307–317, 1956.
 [2] K. Behrend and B. Fantechi. The intrinsic normal cone. *Invent. Math.*, 128(1):45–88, 1997.
 [3] Michel Brion and Baohua Fu. Minimal rational curves on wonderful group compactifications. *J. Éc. polytech. Math.*, 2:153–170, 2015.
 [4] Anders S. Buch, Pierre-Emmanuel Chaput, Leonardo C. Mihalcea, and Nicolas Perrin. Finiteness of cominuscule quantum K -theory. *Ann. Sci. Éc. Norm. Supér. (4)*, 46(3):477–494 (2013), 2013.

- [5] Anders S. Buch, Pierre-Emmanuel Chaput, Leonardo C. Mihalcea, and Nicolas Perrin. Rational connectedness implies finiteness of quantum K -theory. *Asian J. Math.*, 20(1):117–122, 2016.
- [6] Anders Skovsted Buch. Quantum cohomology of Grassmannians. *Compositio Math.*, 137(2):227–235, 2003.
- [7] Anders Skovsted Buch, Andrew Kresch, and Harry Tamvakis. Gromov-Witten invariants on Grassmannians. *J. Amer. Math. Soc.*, 16(4):901–915 (electronic), 2003.
- [8] Anders Skovsted Buch, Andrew Kresch, and Harry Tamvakis. Quantum Pieri rules for isotropic Grassmannians. *Invent. Math.*, 178(2):345–405, 2009.
- [9] P.-E. Chaput, L. Manivel, and N. Perrin. Quantum cohomology of minuscule homogeneous spaces. II. Hidden symmetries. *Int. Math. Res. Not. IMRN*, (22):Art. ID rnm107, 29, 2007.
- [10] P. E. Chaput, L. Manivel, and N. Perrin. Quantum cohomology of minuscule homogeneous spaces. *Transform. Groups*, 13(1):47–89, 2008.
- [11] P.-E. Chaput, L. Manivel, and N. Perrin. Affine symmetries of the equivariant quantum cohomology ring of rational homogeneous spaces. *Math. Res. Lett.*, 16(1):7–21, 2009.
- [12] P. E. Chaput, L. Manivel, and N. Perrin. Quantum cohomology of minuscule homogeneous spaces III. Semi-simplicity and consequences. *Canad. J. Math.*, 62(6):1246–1263, 2010.
- [13] P. E. Chaput and N. Perrin. On the quantum cohomology of adjoint varieties. *Proc. Lond. Math. Soc. (3)*, 103(2):294–330, 2011.
- [14] P.-E. Chaput and N. Perrin. Rationality of some Gromov-Witten varieties and application to quantum K -theory. *Commun. Contemp. Math.*, 13(1):67–90, 2011.
- [15] C. Chevalley. Sur les décompositions cellulaires des espaces G/B . In *Algebraic groups and their generalizations: classical methods (University Park, PA, 1991)*, volume 56 of *Proc. Sympos. Pure Math.*, pages 1–23. Amer. Math. Soc., Providence, RI, 1994. With a foreword by Armand Borel.
- [16] J. A. Cruz Morales, A. Kuznetsov, A. Mellit, N. Perrin, and M. Smirnov. On quantum cohomology of grassmannians of isotropic lines, unfoldings of A_n -singularities, and lefschetz exceptional collections. *To appear in Annales de l'Institut Fourier*, preprint *arXiv:1705.01819*, 2015.
- [17] Michel Demazure. A very simple proof of Bott's theorem. *Invent. Math.*, 33(3):271–272, 1976.
- [18] Anton Fonarev. On the bounded derived category of $IGr(3, 7)$. preprint: *arXiv:1804.06946v2*, 2018.
- [19] W. Fulton and R. Pandharipande. Notes on stable maps and quantum cohomology. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 45–96. Amer. Math. Soc., Providence, RI, 1997.
- [20] William Fulton and Piotr Pragacz. *Schubert varieties and degeneracy loci*, volume 1689 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1998. Appendix J by the authors in collaboration with I. Ciocan-Fontanine.
- [21] S. Galkin, A. Mellit, and M. Smirnov. Dubrovin's conjecture for $IG(2, 6)$. *Int. Math. Res. Not. IMRN*, (18):8847–8859, 2015.
- [22] J. Gandini. Sanya lectures on embeddings of spherical homogeneous spaces. *To appear in Acta Math. Sinica.*, 2017.
- [23] Tom Graber. Enumerative geometry of hyperelliptic plane curves. *J. Algebraic Geom.*, 10(4):725–755, 2001.
- [24] Daniel Huybrechts. *Fourier-Mukai Transforms in Algebraic Geometry*. Oxford Mathematical Monographs. Oxford University Press, 2006.
- [25] V. A. Iskovskikh and Yu. G. Prokhorov. Fano varieties. In *Algebraic geometry, V*, volume 47 of *Encyclopaedia Math. Sci.*, pages 1–247. Springer, Berlin, 1999.
- [26] J. C. Jantzen. *Representations of algebraic groups*, volume 131 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1987.
- [27] M.M. Kapranov. On the derived categories of coherent sheaves on some homogeneous spaces. *Invent. Math.*, 92:479–508, 1988.
- [28] B. Kim and R. Pandharipande. The connectedness of the moduli space of maps to homogeneous spaces. In *Symplectic geometry and mirror symmetry (Seoul, 2000)*, pages 187–201. World Sci. Publ., River Edge, NJ, 2001.

- [29] Friedrich Knop. The Luna-Vust theory of spherical embeddings. In *Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989)*, pages 225–249. Manoj Prakashan, Madras, 1991.
- [30] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1996.
- [31] Andrew Kresch and Harry Tamvakis. Quantum cohomology of the Lagrangian Grassmannian. *J. Algebraic Geom.*, 12(4):777–810, 2003.
- [32] Andrew Kresch and Harry Tamvakis. Quantum cohomology of orthogonal Grassmannians. *Compos. Math.*, 140(2):482–500, 2004.
- [33] S. Kumar. *Kac-Moody groups, their flag varieties and representation theory*, volume 204 of *Progress in Mathematics*. Birkhäuser Boston, 2002.
- [34] A. G. Kuznetsov. Hyperplane sections and derived categories. *Izv. Ross. Akad. Nauk Ser. Mat.*, 70(3):23–128, 2006.
- [35] Alexander Kuznetsov. Derived categories of cubic fourfolds. In *Cohomological and geometric approaches to rationality problems*, volume 282 of *Progr. Math.*, pages 219–243. Birkhäuser Boston, Inc., Boston, MA, 2010.
- [36] Alexander Kuznetsov. Derived equivalence of Ito-Miura-Okawa-Ueda Calabi-Yau 3-folds. *JMSJ*, 70(3):1007–1013, 2018.
- [37] Alexander Kuznetsov and Smirnov Maxim. On residual categories for grassmannians. *Proc. London Math. Soc.*, 120:617–641, 2020.
- [38] Y.-P. Lee. Quantum K -theory. I. Foundations. *Duke Math. J.*, 121(3):389–424, 2004.
- [39] Changzheng Li, Leonardo C. Mihalcea, and Ryan M. Shiffler. Conjecture \mathcal{O} holds for the odd symplectic grassmannian. *preprint arXiv:1706.00744*, 2017.
- [40] Ion Alexandru Mihai. Odd symplectic flag manifolds. *Transform. Groups*, 12(3):573–599, 2007.
- [41] Leonardo C. Mihalcea and Ryan M. Shiffler. Equivariant quantum cohomology of the odd symplectic grassmannian. *to appear in Math. Zeitschrift*, *preprint arXiv:1706.00385*, 2017.
- [42] D. O. Orlov. Projective bundles, monoidal transformations, and derived categories of coherent sheaves. *Izv. Ross. Akad. Nauk Ser. Mat.*, 56(4):852–862, 1992.
- [43] Giorgio Ottaviani. Spinor bundles on quadrics. *Trans. Amer. Math. Soc.*, 307(1):301–316, 1988.
- [44] B. Pasquier. *Variétés horosphériques de Fano*. PhD thesis, Université Grenoble 1, available at <http://tel.archives-ouvertes.fr/tel-00111912>, 2006.
- [45] Boris Pasquier. On some smooth projective two-orbit varieties with Picard number 1. *Math. Ann.*, 344(4):963–987, 2009.
- [46] Boris Pasquier and Nicolas Perrin. Local rigidity of quasi-regular varieties. *Math. Z.*, 265(3):589–600, 2010.
- [47] C. Pech. *Cohomologie quantique des grassmanniennes symplectiques impaires*. PhD thesis, Université Grenoble 1, available at <http://tel.archives-ouvertes.fr/tel-00650211>, 2011.
- [48] Clélia Pech. Quantum cohomology of the odd symplectic Grassmannian of lines. *J. Algebra*, 375:188–215, 2013.
- [49] Nicolas Perrin. Courbes rationnelles sur les variétés homogènes. *Ann. Inst. Fourier (Grenoble)*, 52(1):105–132, 2002.
- [50] Nicolas Perrin. On the geometry of spherical varieties. *Transform. Groups*, 19(1):171–223, 2014.
- [51] Nicolas Perrin. Semisimple quantum cohomology of some Fano varieties. *preprint arXiv:1405.5914*, 2014.
- [52] Nicolas Perrin. Sanya lectures: geometry of spherical varieties. *Acta Math. Sin. (Engl. Ser.)*, 34(3):371–416, 2018.
- [53] Piotr Pragacz and Jan Ratajski. A Pieri-type theorem for Lagrangian and odd orthogonal Grassmannians. *J. Reine Angew. Math.*, 476:143–189, 1996.
- [54] Bernd Siebert and Gang Tian. On quantum cohomology rings of Fano manifolds and a formula of Vafa and Intriligator. *Asian J. Math.*, 1(4):679–695, 1997.
- [55] Jesper Funch Thomsen. Irreducibility of $\overline{M}_{0,n}(G/P, \beta)$. *Internat. J. Math.*, 9(3):367–376, 1998.

DEPARTMENT OF SCIENCES, PONTIFICIA UNIVERSIDAD CATÓLICA DEL PERÚ, SAN MIGUEL,
LIMA 32, LIMA, PERÚ

E-mail address: rgonzalesv@pucp.edu.pe

SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF KENT,
CANTERBURY CT2 7NF, UNITED KINGDOM

E-mail address: c.m.a.pech@kent.ac.uk

LABORATOIRE DE MATHÉMATIQUES DE VERSAILLES, UVSQ, CNRS, UNIVERSITÉ PARIS-
SACLAY, 78035 VERSAILLES, FRANCE

E-mail address: nicolas.perrin@uvsq.fr

INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS, B. KARETNYI PER., 19, 127994,
GSP-4, MOSCOW, RUSSIA

E-mail address: alexander.samokhin@gmail.com