



# Kent Academic Repository

**Romanov, Michael N, Talbot, Richard T., Wilson, Peter W. and Sharp, Peter J. (2003) Erratum to "Genetic Control of Incubation Behavior in the Domestic Hen". Poultry Science, 82 (10). p. 1659. ISSN 0032-5791.**

## Downloaded from

<https://kar.kent.ac.uk/45705/> The University of Kent's Academic Repository KAR

## The version of record is available from

<https://doi.org/10.1093/ps/82.10.1659>

## This document version

Publisher pdf

## DOI for this version

## Licence for this version

UNSPECIFIED

## Additional information

## Versions of research works

### Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

### Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in **Title of Journal**, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

### Enquiries

If you have questions about this document contact [ResearchSupport@kent.ac.uk](mailto:ResearchSupport@kent.ac.uk). Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

# ASSOCIATION NOTES

## Erratum to “Genetic Control of Incubation Behavior in the Domestic Hen”

M. N. Romanov, R. T. Talbot, P. W. Wilson, and P. J. Sharp

*Division of Integrative Biology, Roslin Institute (Edinburgh), Roslin, Midlothian EH25 9PS, Scotland, United Kingdom*

2002 Poultry Science 81:928–931

Table 1 is being rerun to correct the table footnote.

**TABLE 1. Observed and predicted incidence of incubation behavior in White Leghorns (WL), Bantams (B), and their crosses assuming major sex-linked or permissive autosomal genes control of incubation behavior and that their expression results in 78.6% incubation behavior in the environmental conditions to which all groups were exposed**

Prediction tested	Group no.	Mating	Hens incubating/total (n)	Percentage incubating	
				Predicted	Observed
None	1	♂ B × ♀ B	22/28	...	78.6
None	2	♂ WL × ♀ WL	0/28	...	0
Major sex-linkage	3	♂ WL × ♀ B	45/73	0	61.6***
	4	♂ B × ♀ WL	21/37	78.6	56.8*
	5	♂ F <sub>1</sub> (♂ WL × ♀ B) × ♀ WL	6/103	39.3	5.8***
Major permissive autosomal	3	♂ WL × ♀ B	45/73	78.6	61.6***
	4	♂ B × ♀ WL	21/37	78.6	56.8*
	5	♂ F <sub>1</sub> (♂ WL × ♀ B) × ♀ WL	6/103	39.3	5.8***

\* $P < 0.05$ , \*\*\* $P < 0.001$  for the difference between predicted and observed.