



Kent Academic Repository

Simkin, Andrew J., Lopez-Calcano, Patricia E., Davey, Philip A., Headland, Lauren R., Lawson, Tracy, Timm, Stefan, Bauwe, Hermann and Raines, Christine A. (2017) *Simultaneous stimulation of sedoheptulose 1,7-bisphosphatase, fructose 1,6-bisphosphate aldolase and the photorespiratory glycine decarboxylase-1 protein increases CO₂ assimilation, vegetative biomass and seed yield in Arabidopsis*. *Plant Biotechnology Journal*, 15 (7). pp. 805-816. ISSN 1467-7644.

Downloaded from

<https://kar.kent.ac.uk/93875/> The University of Kent's Academic Repository KAR

The version of record is available from

<https://doi.org/10.1111/pbi.12676>

This document version

Publisher pdf

DOI for this version

Licence for this version

CC BY (Attribution)

Additional information

Versions of research works

Versions of Record

If this version is the version of record, it is the same as the published version available on the publisher's web site. Cite as the published version.

Author Accepted Manuscripts

If this document is identified as the Author Accepted Manuscript it is the version after peer review but before type setting, copy editing or publisher branding. Cite as Surname, Initial. (Year) 'Title of article'. To be published in *Title of Journal*, Volume and issue numbers [peer-reviewed accepted version]. Available at: DOI or URL (Accessed: date).

Enquiries

If you have questions about this document contact ResearchSupport@kent.ac.uk. Please include the URL of the record in KAR. If you believe that your, or a third party's rights have been compromised through this document please see our [Take Down policy](https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies) (available from <https://www.kent.ac.uk/guides/kar-the-kent-academic-repository#policies>).

Simultaneous stimulation of sedoheptulose 1,7-bisphosphatase, fructose 1,6-bisphosphate aldolase and the photorespiratory glycine decarboxylase-H protein increases CO₂ assimilation, vegetative biomass and seed yield in *Arabidopsis*

Andrew J. Simkin¹, Patricia E. Lopez-Calcagno¹, Philip A. Davey¹, Lauren R. Headland¹, Tracy Lawson¹, Stefan Timm², Hermann Bauwe² and Christine A. Raines^{1*}

¹School of Biological Sciences, University of Essex, Colchester, UK

²Plant Physiology Department, University of Rostock, Rostock, Germany

Received 25 August 2016;

revised 25 November 2016;

accepted 30 November 2016.

*Correspondence (Tel 01206 873310; fax 01206 872592; email rainc@essex.ac.uk)

Keywords: chlorophyll fluorescence imaging, FBP aldolase, glycine decarboxylase-H protein, photosynthesis, SBPase, transgenic.

Summary

In this article, we have altered the levels of three different enzymes involved in the Calvin–Benson cycle and photorespiratory pathway. We have generated transgenic *Arabidopsis* plants with altered combinations of sedoheptulose 1,7-bisphosphatase (SBPase), fructose 1,6-bisphosphate aldolase (FBPA) and the glycine decarboxylase-H protein (GDC-H) gene identified as targets to improve photosynthesis based on previous studies. Here, we show that increasing the levels of the three corresponding proteins, either independently or in combination, significantly increases the quantum efficiency of PSII. Furthermore, photosynthetic measurements demonstrated an increase in the maximum efficiency of CO₂ fixation in lines over-expressing SBPase and FBPA. Moreover, the co-expression of GDC-H with SBPase and FBPA resulted in a cumulative positive impact on leaf area and biomass. Finally, further analysis of transgenic lines revealed a cumulative increase of seed yield in SFH lines grown in high light. These results demonstrate the potential of multigene stacking for improving the productivity of food and energy crops.

Introduction

The accumulated photosynthate produced over the season determines the yield of a crop, but improvements in photosynthesis have not been used in traditional breeding approaches to identify high-yielding varieties. The reasons for this are twofold, (i) methodologies to make accurate field measurements have only been available in the last 10–20 years, and also, (ii) there is a lack of evidence to determine whether there is a correlation between the rate of photosynthesis on a leaf area basis and final yield of the crop (Evans, 2013; Fischer *et al.*, 1998; Gifford and Evans, 1981). There is now an urgent need to increase crop productivity and yields to meet the nutritional demands of a growing world population, and there is growing evidence that this may be achieved through improvement of photosynthetic energy conversion to biomass (von Caemmerer and Evans, 2010; Ding *et al.*, 2016; Lefebvre *et al.*, 2005; Long *et al.*, 2006, 2015; Simkin *et al.*, 2015). Evidence from a combination of theoretical studies and transgenic approaches has provided compelling evidence that manipulation of the Calvin–Benson (CB) cycle can improve energy conversion efficiency and lead to an increase in yield potential (Long *et al.*, 2006; Poolman *et al.*, 2000; Raines, 2003, 2006, 2011; Zhu *et al.*, 2007, 2010).

Previous studies have demonstrated that even small reductions in individual CB cycle enzymes such as sedoheptulose 1,7-bisphosphatase (SBPase) and fructose 1,6-bisphosphate aldolase (FBPA) negatively impact on carbon assimilation and growth,

indicating that these enzymes exercise significant control over photosynthetic efficiency (Ding *et al.*, 2016; Haake *et al.*, 1998, 1999; Harrison *et al.*, 1998, 2001; Lawson *et al.*, 2006; Raines, 2003; Raines and Paul, 2006; Raines *et al.*, 1999). Furthermore, the disruption of the chloroplastic fructose-1,6-bisphosphatase (FBPase) gene was also shown to impact negatively on carbon fixation (Kossmann *et al.*, 1994; Rojas-González *et al.*, 2015; Sahrawy *et al.*, 2004). These results strongly suggested that improvements in photosynthetic carbon fixation may be achieved by increasing the activity of individual CB cycle enzymes. Evidence supporting this hypothesis came from transgenic tobacco plants over-expressing SBPase (Lefebvre *et al.*, 2005), the cyanobacterial bifunctional SBPase/FBPase (Miyagawa *et al.*, 2001) or FBPA (Uematsu *et al.*, 2012). These single manipulations resulted in increase in photosynthetic carbon assimilation, enhanced growth and an increase in biomass. More recently, Simkin *et al.* (2015) demonstrated that the combined over-expression of SBPase and FBPA in tobacco resulted in a cumulative increase in biomass and that these increases could be further enhanced by the over-expression of the cyanobacterial inorganic carbon transporter B (ictB), proposed to be involved in CO₂ transport, although its function was not established in these plants (Simkin *et al.*, 2015). These results demonstrate the potential for the manipulation of photosynthesis, using multigene stacking approaches, to increase biomass yield (Simkin *et al.*, 2015).

The efficiency of CO₂ fixation by the CB cycle is compromised by the oxygenase activity of ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) which directly competes with CO₂

fixation at the active site, resulting in the formation of 2-phosphoglycolate (2PG) and subsequently significant energy costs and CO₂ losses in the photorespiratory pathway, resulting in significant losses in yield (Bowes *et al.*, 1971; Tolbert, 1997; Walker *et al.*, 2015, 2016). Therefore, a major target to improve photosynthesis has been to reduce photorespiration, either through protein engineering to improve Rubisco catalysis or by limiting the flux through this pathway, none of which have as yet yielded positive results due to both the complexity of the Rubisco catalytic and assembly processes (Cai *et al.*, 2014; Carmo-Silva *et al.*, 2015; Lin *et al.*, 2014a; Orr *et al.*, 2016; Sharwood *et al.*, 2016; Whitney *et al.*, 2011). More ambitious approaches to this problem are now being taken, including the introduction of cyanobacterial or algal CO₂-concentrating mechanisms, novel synthetic metabolic pathways and the introduction of the C4 pathway into C3 crops (Betti *et al.*, 2016; Lin *et al.*, 2014b; McGrath and Long, 2014; Meyer *et al.*, 2016; Montgomery *et al.*, 2016). However, to date the only successful approach to limiting photorespiration which has resulted in an improvement in photosynthesis has been through the introduction of alternative routes to metabolize 2PG and return CO₂ for use in the CB cycle (Dalal *et al.*, 2015; Kebeish *et al.*, 2007; Maier *et al.*, 2012; Nolke *et al.*, 2014; Peterhänsel *et al.*, 2013; Xin *et al.*, 2015). Reductions in the flux through the photorespiratory cycle by targeted knock-down of GDC-P in potato and GDC-H in rice have been shown to lead to reductions in photosynthesis and growth rates (Engel *et al.*, 2007; Heineke *et al.*, 2001; Lin *et al.*, 2016). The opposite approach, namely over-expression of the glycine decarboxylase (GDC)-H protein (GDC-H) and glycine decarboxylase (GDC)-L protein (GDC-L) in *Arabidopsis thaliana* (*Arabidopsis*), resulted in an improvement of photosynthesis and increased vegetative biomass when compared to wild-type plants (Timm *et al.*, 2012, 2015, 2016). Although the underlying mechanism responsible for this effect has not been fully elucidated, these authors proposed that stimulation of the CB cycle is brought about by the increase in GDC activity, resulting in a reduction in the steady-state levels of photorespiratory metabolites that may negatively impact on the function of the CB cycle (e.g. 2PG, glycolate, glyoxylate or glycine (Anderson, 1971; Kelly and Latzko, 1976; Eisenhut *et al.*, 2007; Lu *et al.*, 2014; Timm *et al.*, 2015, 2016)).

In the light of the results from Timm *et al.* (2012, 2015), the aim of this study was to explore the possibility that the simultaneous increase in the activity of enzymes of both the CB cycle and the photorespiratory pathway could lead to a cumulative positive impact on photosynthetic carbon assimilation and yield. To test this, we have taken a proof-of-concept approach using the model plant *Arabidopsis* in which we have over expressed SBPase, FBPA and GDC-H either alone or in combination. We have shown that the simultaneous manipulation of multiple targets can lead to a cumulative impact on biomass yield under both low- and high-light growing conditions. Interestingly, we have also shown that manipulation of the photorespiratory pathway alone resulted in an increase in vegetation biomass but not seed yield. In contrast, manipulation of both the CB cycle and photorespiratory pathway increased both biomass and seed yield.

Results

Production and selection of *Arabidopsis* transformants

The full-length *Arabidopsis* SBPase (*At3g55800*) and FBPA cDNA (*At4g38970*) were used to generate three over-expression

constructs PTS1-SB, PTS1-FB and PTS1-SBFB in vector pGWPTS1 (Figure S1). These transgenes were under the control of the photosynthetic tissue-specific (PTS) *rbcS2B* (1150 bp; *At5g38420*) promoter. These constructs were transformed into *Arabidopsis* using the floral dip method (Clough and Bent, 1998), and the resulting transgenic plants were selected on kanamycin/hygromycin-containing medium. T2 plants expressing the integrated transgenes were screened by immunoblotting and allowed to self-fertilize to generate seeds for T3 plants. Following initial characterisation of primary independent lines generated, 3–4 independent lines overexpressing either SBPase (S3, S8, S12) or FBPA (F6, F9, F11) and SBPase and FBPA together (SF4, SF6, SF7, SF12) were selected for further study.

Further analysis was carried out on T3 plants grown at 130 μmol/m²/s in an 8-h/16-h light/dark cycle and total extractable SBPase and FBPA activity determined in extracts from newly fully expanded leaves. The results are represented as a percentage (%) of total activity for SBPase (6.7 μmol/m²/s) and FBPA aldolase (22 μmol/m²/s) determined in wild type (WT). This analysis showed that these plants had increased levels of SBPase (137%–185%) and FBPA (146%–180%) activity (Figure 1) compared to WT and nontransformed azygous (A) controls (azygous-control plants used in this study were recovered from the segregating population and verified by PCR). Interestingly, a small increase in endogenous FBPA activity (125%–136%) was also observed in SBPase over-expressing lines (Figure 1a), but no significant increase in SBPase activity was observed in lines over-expressing FBPA.

Plants over-expressing SBPase (S), FBPA (F) and the GDC-H protein (H) were generated by crossing two SBPase + FBPA (SF) lines (SF6 and SF12) with two *Flaveria pringlei* GDC-H protein (Kopriva and Bauwe, 1995) over-expressing lines (*FpHL17* and *FpHL18*) originally generated by Timm *et al.* (2012) under the control of the leaf-specific and light-regulated *Solanum tuberosum* ST-LS1 promoter (Stockhaus *et al.*, 1989). Four independent lines (SFH4, SFH20, SFH23 and SFH31) over-expressing SBPase, FBPA and GDC-H (SFH) were identified by PCR and SBPase and FBPA enzyme activities. SBPase and FBPA protein levels were found to be similar to those observed in SF lines (Figure 1b). No significant difference in SBPase or FBPA activities was observed in lines over-expressing GDC-H alone compared to WT/A controls (C). The full set of assays showing the variation between plants for both SBPase and FBPA activities can be seen in Figure S2.

In addition to total extractable enzyme activity, immunoblot analysis of the T3 progenies of S, F, SF, H and SFH lines was carried out using WT/A as controls (C). This analysis identified a number of plants over-expressing SBPase or FBPA and others with increased levels of both SBPase and FBPA (Figures 1a,b and S3). Interestingly, the over-expression of SBPase in *Arabidopsis* led to an increase in endogenous FBPA protein levels (Figure 1a) in agreement with the observed increase in enzyme activity. The original H lines and the newly generated SFH plants were shown to accumulate GDC-H when compared to both nontransformed control plants and other transgenic lines (Figure 1a,b). Given the change in FBPA protein levels in the SBPase over-expressing line, we used immunoblot analysis to determine whether there were any changes in other CB cycle enzymes. No detectable changes in the levels of transketolase (TK), phosphoribulokinase (PRK), fructose-1,6-bisphosphatase (FBPase), Rubisco or the ADP glucose pyrophosphorylase (ssAGPase) small protein were observed when compared to levels in C plants (Figure 2).




Figure 1 Molecular and biochemical analysis of the transgenic plants over-expressing SBPase (S), FBPA (F) and GDC-H (H). SBPase and FBPA enzyme activity (SBPase, FBPA) and immunoblot blot analysis (SBPase, FBPA, GDC-H) of protein extracts from two independent leaves of (a) S, F and H lines and (b) SF and SFH lines used in this study compared to non-transformed control (C). Enzyme assays represent data from 12 to 24 independent plants per group compared to 12–16 C plants. The results are represented as a percentage (%) of total activity for SBPase ($6.7 \mu\text{mol}/\text{m}^2/\text{s}$) and FBPA aldolase ($22 \mu\text{mol}/\text{m}^2/\text{s}$) determined in wild type (WT). Enzyme activities per plant can be seen in Figure S2. Columns represent mean values, and standard errors are displayed. Lines that are significantly different to C are indicated ($*P < 0.05$).

Chlorophyll fluorescence imaging reveals increased photosynthetic efficiency in young over-expressing seedlings

To explore the impact of increased levels of SBPase, FBPA and the GDC-H protein on photosynthesis, plants were grown at $130 \mu\text{mol}/\text{m}^2/\text{s}$ in an 8-h/16-h light/dark cycle and the quantum efficiency of PSII photochemistry (F_q'/F_m') analysed using chlorophyll a fluorescence imaging (Baker, 2008; Murchie and Lawson, 2013). Plants over-expressing SBPase and FBPA, either independently or in combination (including with GDC-H), had a significantly higher F_q'/F_m' at an irradiance of $200 \mu\text{mol}/\text{m}^2/\text{s}$ when compared to C plants (Figure 3a,b). Plants over-expressing GDC-H alone showed a small increase in the average levels of F_q'/F_m' compared to C ($P = 0.061$). When measurements were made at a higher light level ($600 \mu\text{mol}/\text{m}^2/\text{s}$), all lines analysed, with the exception of SFH, showed a significant increase in F_q'/F_m' compared to C plants (Figure S4a). From images taken as part of the chlorophyll fluorescence analysis, leaf area was determined and shown to be significantly larger for all transgenic lines compared with WT and azygous (A) controls (Figure 3c). Interestingly, SFH plants showed the greatest leaf area in all experiments. No significant differences in leaf area were observed

between WT and A. From this point on, C plants used were the combined data from WT and A plants.

Photosynthetic CO_2 assimilation rates are increased in mature plants grown in low light

To explore the impact of changes in the levels of enzymes in both the CB cycle and photorespiratory pathway, CO_2 assimilation rates were determined as a function of light intensity (Figure 4a, b). From these light response curves, the maximum light-saturated rate of photosynthesis (A_{sat}) was shown to be significantly higher in all transgenic plants when compared to C plants (Figure 4c). Small differences in CO_2 assimilation rates (A) were also observed in the S, F, SF and SFH plants even at light intensities as low as $150 \mu\text{mol}/\text{m}^2/\text{s}$, which is close to that of the growth conditions (Figure S5).

We also determined A as a function of internal CO_2 concentration (C_i) in the same plants (Figure 4d,e). In all transgenic plants, except those over-expressing GDC-H alone, A was significantly greater at C_i concentrations above $400 \mu\text{mol}/\text{mol}$ than in C plants (Figure 4d,e). Although A in SFH plants was higher than in the control plants at $400 \mu\text{mol}/\text{mol}$, it was lower than that observed in the S, F or SF plants. The maximum rate of CO_2 assimilation (A_{max}) was significantly higher in lines S, F, SF




Figure 2 Molecular and biochemical analysis of the transgenic plants over-expressing SBPase (S), FBPA (F) and GDC-H (H). Immunoblot blot analysis of protein extracts from two independent leaves of (a) S, F and H lines and (b) SF and SFH lines used in this study compared to C. Transketolase (TK), phosphoribulokinase (PRK), fructose-1,6-bisphosphase (FBPase) the small subunit of ADP glucose pyrophosphorylase (ssAGPase) and Rubisco.




Figure 3 Photosynthetic capacity and leaf area in transgenic seedlings determined using chlorophyll fluorescence imaging. C and transgenic plants were grown in controlled environment conditions with a light intensity $130 \mu\text{mol}/\text{m}^2/\text{s}$, 8-h light/16-h dark cycle for 15 days and chlorophyll fluorescence imaging used to determine F_q'/F_m' (maximum PSII operating efficiency) values of the whole plant at (a, b) $200 \mu\text{mol}/\text{m}^2/\text{s}$ and (c) leaf area at time of analysis. Azygous controls (A) recovered from a segregating population. Lines over-expressing SBPase (S), FBPA (F), GDC-H protein (H), SBPase and FBPA (SF) and SBPase, FBPA and GDC-H (SFH) are represented. The data were obtained using six individual plants from two (H), three (S, F, SF) or four (SFH) independent transgenic lines (18–24 plants total) compared to 12 C. Columns represent mean values, and standard errors are displayed. Significant differences between lines ($P < 0.05$) are represented as capital letters indicating whether each specific line is significantly different from another (i.e. SBPase lines (S) are significantly bigger than wild type (WT) and azygous lines (A)). Numbers indicate % increases over WT.

and SFH compared to C; however, no significant differences were observed between these lines (Figure 4f). Interestingly, the H plants show no increase in A_{max} when compared to C plants. Further analysis of the A/C_i curves using the equations published by von Caemmerer and Farquhar (1981) illustrated that the maximum rate of carboxylation by Rubisco (V_{cmax} ; Figure S4b) in lines S, SF and SFH was significantly greater than in C, and V_{cmax} in these lines was also significantly greater than in lines expressing GDC-H alone. Maximum electron transport rates (J_{max} ; Figure S4c) were also elevated in lines S, F, SF and SFH compared to C and were also shown to be significantly elevated compared to H.

To further assess the effect of the manipulation of the CB cycle and/or the GDC-H protein, the rates of photosynthetic carbon

assimilation and electron transport were determined in mature plants as a function of light intensity at 2% $[\text{O}_2]$ to eliminate photorespiration (Figure 5a). Electron transport rates through PSII in H and SFH over-expression plants were significantly greater than in the C and SF plants at light levels above $300 \mu\text{mol}/\text{m}^2/\text{s}$ (Figure 5b). A_{sat} was also significantly higher, 12%–19%, in all lines compared to C although no significant differences were observed between the different transgenic lines (Figure 5c).

Increased SBPase and FBPA activity and over-expression of the glycine decarboxylase-H protein stimulates growth in low light

The growth of the different transgenic and C plants was determined using image analysis of total leaf area over a period of 38 days from planting (Figure 6a), which showed all transgenic lines had a significantly greater leaf area than C, as early as 16 days after planting (Figure 6b). Furthermore, plants over-expressing all three transgenes (SFH) were shown to have a significantly larger leaf area when compared to all other transgenic lines including G and SF, indicating a cumulative advantage from combining these transgenes at this stage in development. This growth trend continued through to 15 days postplanting (Figure S6a). By 20 days after planting (Figure S6b), plants over-expressing the glycine decarboxylase-H protein (H) were shown to be significantly bigger than S, F and SF at the same time point, and triple over-expressing lines SFH remained significantly bigger than all other lines studied (Figure 6b).

Plants were allowed to continue growing until harvest at 38 days (Figure S7). At this stage of development, no significant difference in leaf area or dry weight could be observed between S, F, H or SF lines when compared to each other (Figure 6c). However, all lines attained a significantly larger leaf area and dry weight when compared to C. Notably, at this stage, the triple over-expressing lines SFH were significantly larger with a higher dry weight (+70%) than all other transgenic and C plants. Furthermore, lines SF and SFH both showed a significant increase in leaf number after 38 days (Figure S8).

Increased SBPase and FBPA activity and expression of the glycine decarboxylase-H protein impacts on the carbohydrate profile of selected lines

To determine how the over-expression of these key proteins impacts on downstream processes, leaf tissue was harvested and starch and sugar content were evaluated. No significant difference in starch levels were detected at the end of the day in any of the transgenic lines compared to C (Figure 7). Interestingly, slightly higher starch levels were detected 1 h before sunrise (dark) in transgenic lines F, H and SFH compared to C. All transgenic lines were shown to have consistently higher levels of sucrose, with these levels being significantly higher than C in F and SF lines. SF lines were also shown to have a significantly higher amount of glucose (Figure 7) compared to C, although other lines were consistently elevated but not significantly so.

Impact of light intensity on biomass and seed yield

A subset of plants was allowed to seed in either low or high light, and final vegetative biomass and seed yield determined per plant. In low-light grown plants, the final vegetative biomass was higher in all of the transgenic lines compared to C; however, no significant differences were observed between the different transgenic lines (Figure 8a). Furthermore, seed yield was




Figure 4 Photosynthetic responses of C and transgenic plants. (a, b) Photosynthesis carbon fixation rates were determined as a function of increasing light intensity. (c) A_{sat} determined from light response curves. (d, e) Photosynthetic carbon fixation rates were determined as a function of increasing CO_2 concentrations (A/C_i) at saturating-light levels ($1000 \mu\text{mol}/\text{m}^2/\text{s}$). (f) A_{max} determined from A/C_i response curves. C and transgenic plants were grown in controlled environment conditions with a light intensity $130 \mu\text{mol}/\text{m}^2/\text{s}$, 8-h light/16-h dark cycle for 4 weeks. Lines over-expressing SBPase (S), FBPA (F), GDC-H protein (H), SBPase and FBPA (SF) and SBPase, FBPA and GDC-H (SFH) are represented. Columns represent mean values, and standard errors are displayed. Significant differences between lines ($P < 0.05$) are represented as capital letters indicating whether each specific line is significantly different from another. Results are based on 4–7 plants per line. (i.e. SBPase lines (S) are significantly different to controls (C)).

increased by 35%–53% in transgenic lines S, SF and SFH (Figure 8b). Interestingly, no increase in seed yield was observed in the H plants.

We next compared the impact of growth of plants in high light to explore further the potential positive impact of these transgenic manipulations on growth. In high-light grown plants, an increase in vegetative biomass from 14% to 51% was observed (Figures 8c and S9). Notably, the H and SFH plant produced significantly more vegetative biomass than the S, F, SF or C plants. Furthermore, seed yield in high-light-grown plants was increased by 39%–62% in transgenic lines S, F, SF and SFH, when compared to C (Figure 8d). Although the highest increase in seed yield was observed in lines SFH in high light, no increase in seed yield was observed in the H plants in high-light-grown plants. The seed yield for individual plants can be seen in Figure S10.

Discussion

In this study, we have shown that simultaneously increasing the levels of two enzymes of the CB cycle, SBPase and FBPA, and the H protein of the glycine decarboxylase enzyme of the photorespiratory pathway in the same plant, resulted in a substantial and significant increase in both vegetative biomass and seed yield of *Arabidopsis* grown in controlled environment conditions. An increase in both biomass and yield was also observed in plants

overexpressing SBPase or FBPA alone or in combination. However, although overexpression of GDC-H alone resulted in an increase in vegetative biomass, no increase in seed yield was evident in these plants, grown in either low- or high-light conditions. The reasons for this differential effect on seed yield have not yet been elucidated but may be due to changes in carbon status brought about by altered source/sink allocation which is supported by changes to starch and sucrose levels at the end of the night period in some of these lines. Higher levels of sucrose (and fructose, maltose) have also been observed in GDC-L over-expressors (Timm *et al.*, 2015), and the over-expression of GDC-L enhances the metabolic capacity of photorespiration and is believed to alter the carbon flow through the TCA cycle (Timm *et al.*, 2015).

It was shown in earlier studies that over-expression of FBPA or SBPase alone in tobacco results in a stimulation of photosynthesis and biomass, with the greatest effect being seen in plants grown under elevated CO_2 (Lefebvre *et al.*, 2005; Rosenthal *et al.*, 2011; Uematsu *et al.*, 2012). Furthermore, when FBPA was over-expressed in combination with SBPase in tobacco, this led to a cumulative increase in biomass in plants grown in ambient CO_2 under greenhouse conditions (Simkin *et al.*, 2015). Interestingly, in this current study, we have shown that in *Arabidopsis* that the over-expression of FBPA alone, under current atmospheric CO_2 levels, results in a stimulation of photosynthesis and increase in




Figure 5 Photosynthetic responses of the transgenic plants at 2% $[\text{O}_2]$ (a) and (b) chloroplast electron transport rates in transgenic plants at 2% $[\text{O}_2]$. (c) Mean values of A_{sat} determined from light response curves. C and transgenic plants were grown in controlled environment conditions with a light intensity $130 \mu\text{mol}/\text{m}^2/\text{s}$, 8-h light/16-h dark cycle for 4 weeks. Lines over-expressing the GDC-H protein (H), SBPase and FBPA (SF) and SBPase, FBPA and GDC-H (SFH) are represented. Columns represent mean values, and standard errors are displayed. Significant differences between lines ($P < 0.05$) are represented as capital letters. Results are based on 5–6 plants per line compared to six controls.

biomass on a similar level to that observed by over-expression of SBPase alone. However, contrary to the results obtained in tobacco, the co-expression of SBPase and FBPA in Arabidopsis did not lead to a further significant increase in either leaf area or biomass when compared to plants independently expressing SBPase (resulting in higher endogenous FBPA activity) or FBPA. This lack of differential effect of the co-overexpression of SBPase and FBPA in this study can likely be explained by the fact that over-expression of SBPase in Arabidopsis also led to a small but significant increase in endogenous FBPA protein levels and activity

(25%–36%). Given that no increase in SBPase was present in the FBPA plants, this would suggest that in Arabidopsis, the stimulation in the SBPase, FBPA and the SF over-expression lines may be due to increased FBPA activity. This is in contrast to tobacco where over-expression of SBPase alone led to an increase in biomass and no increases in endogenous FBPA activity, highlighting the differences between species (Lefebvre *et al.*, 2005; Rosenthal *et al.*, 2011; Simkin *et al.*, 2015).

Detailed analysis of a range of photosynthesis parameters revealed a similar increase in A_{sat} at low $[\text{O}_2]$ for all of the transgenic lines studied. The most significant increase was observed in SF lines which showed a 44% increase over control plants, with the lowest increase of 19% being observed in the H plants. An evaluation of the electron transport rates at low $[\text{O}_2]$ in a subset of these plants showed that lines over-expressing GDC-H (both H and SFH) displayed higher photosynthetic electron transport rates compared to C and plants over-expressing SBPase and FBPA (SF). These results are in keeping with the previous study by Timm *et al.* (2012). All of the transgenic lines analysed here showed an increase in photosynthesis under high light and ambient CO_2 conditions. However, under high light and saturated levels of CO_2 the rate of assimilation in the H plants was similar to C, and this is in contrast to all other transgenic lines. This observation is in keeping with the proposal that over-expression of the H protein stimulates the flow of carbon through the photorespiratory pathway, thereby reducing steady-state levels of inhibitory photorespiratory metabolites, which in turn stimulates flux through the CB cycle. Whilst this hypothesis is supported by metabolite data and the observation that growth of GDC-H plants is not stimulated when these plants are grown in elevated CO_2 conditions (Timm *et al.*, 2012), the exact mechanism of such feedback from photorespiration to the CB cycle is not yet known. The effect of these manipulations on photosynthesis was also determined at the growth light intensity where small differences in A are observed even at light levels as low as $150 \mu\text{mol}/\text{m}^2/\text{s}$. This together with the increased leaf area observed at early stages in development provides evidence that the small differences in photosynthesis lead to an increase in leaf area. The cumulative impact of this over time results in increased biomass and yield.

Conclusion

In this proof-of-concept study in Arabidopsis, we have demonstrated that the simultaneous over-expression of two CB cycle enzymes leads to an increase in photosynthesis and an increase in overall biomass and seed yield. We also show that when the transgenic SF lines were crossed with GDC-H over-expressing plants (Timm *et al.*, 2012), the combined effects of these three transgenes (SFH) resulted in a cumulative impact on biomass (+71%) which was significantly higher than H (+50%) and SF (+41%) under low light. Importantly, the work here also allowed a parallel comparative analysis between the different manipulations under different environmental conditions.

Although it is still necessary to address the importance of these manipulations in crop species and also under field conditions, this study provides additional evidence that multigene manipulation of photosynthesis and photorespiration can form an important tool to improve crop yield. These results also provide new information indicating that it will be necessary to tailor the targets for manipulation for different crops and for either biomass or seed yield.




Figure 6 Growth analysis of C and transgenic lines grown in low light. (a) Plants were grown at 130 $\mu\text{mol}/\text{m}^2/\text{s}$ light intensity in short days (8 h/16 h days) for 15 days. (b) Plant growth rate evaluated over the first 38 days. Significant differences $^*(P < 0.05)$, $^{**}(P < 0.01)$, $^{***}(P < 0.001)$ are indicated. (c) Final dry weight (g) after 38 days of development and statistical differences between lines. % increases over C are indicated within the columns. Lines over-expressing SBPase (S), FBPA (F), GDC-H protein (H), SBPase and FBPA (SF) and SBPase, FBPA and GDC-H (SFH) are represented. Columns represent mean values, and standard errors are displayed. Significant differences between lines ($P < 0.03$) are represented as capital letters indicating whether each specific line is significantly different from another. Results are representative of 9–12 plants from two (H), three (S, F, SF) or four (SFH) independent lines (C plants including wild type and azygous lines segregated from primary transformants).

Materials and methods

Generation of pGW photosynthetic tissue-specific destination vector pGWPTS1

pGWB1 (Nakagawa *et al.*, 2007: AB289764) was cut with HindIII at 37 °C. Following purification, digested vectors were treated

with alkaline phosphatase (BioLabs, UK) for 60 min at 37 °C. The *rbcS2B* (1150 bp; *At5 g38420*) promoter was amplified with primers Pr_ *rbcS2B*_F_HindIII (5' CACCaagcttATgACATCATAgCAAgCAAggACACg'3) and Pr_ *rbcS2B*_R_HindIII (5' CTGAGAAagcttTACTTCTTCTgTTgTTTCTCTTCTTC'3). The amplicon was




Figure 7 Leaf starch and sugar content at end of 8-h light period (light grey) and end of 16-h dark period (dark grey). Results are mean values based on 12–18 individual plants from two (H), three (S, F, SF) or four (SFH) independent transgenic lines. Columns represent mean values, and standard errors are displayed. Lines over-expressing SBPase (S), FBPA (F), GDC-H protein (H), SBPase and FBPA (SF) and SBPase, FBPA and GDC-H (SFH) are represented. Significant differences between C and over-expressing lines ($*P < 0.01$) are represented.

digested with HindIII and cloned into the corresponding site of pGWB1 to make vector pGWPTS1 (Figure S1a).

Constructs were generated using Gateway cloning technology and vector pGWPTS1. All transgenes were under the control of the *rbcS2B* (1150 bp; *At5 g38420*) promoter. Full details of PTS1-

SB, PTS1-FB and PTS1-SBFB construct assembly can be seen in the supplementary data. Construct maps are shown in Figure S1b–d.

Generation of transgenic plants

The recombinant plasmids PTS1-SB, PTS1-FB and PTS1-SBFB were introduced into wild-type *Arabidopsis* by floral dipping (Clough and Bent, 1998) using *Agrobacterium tumefaciens* GV3101. Positive transformants were regenerated on MS medium containing kanamycin (50 mg/L) and hygromycin (20 mg/L). Kanamycin/hygromycin-resistant primary transformants (T1 generation) with established root systems were transferred to soil and allowed to self-fertilize. Plants over-expressing SBPase, FBPA and the GDC-H protein were generated by floral inoculation of two SBPase + FBPA lines (SF6 and SF12) with the pollen from two GDC-H protein over-expressing lines (*FpH17* and *FpH18*) provided by Timm *et al.* (2012). Lines *FpH17* and 18 were originally generated by floral dipping and over-expressing the *Flaveria pringlei* GDC-H protein (Kopriva and Bauwe, 1995) under the control of the leaf-specific and light-regulated *Solanum tuberosum* *ST-LS1* promoter (Stockhaus *et al.*, 1989). Following initial characterization of generated lines, three lines for SBPase (S3, S8, S12), FBPA (F6, F9, F11) and SF (SF6, SF7, SF12) were selected for further study from all lines generated.


Plant growth conditions

Wild-type T2 *Arabidopsis* plants resulting from self-fertilization of transgenic plants were germinated in sterile agar medium containing Murashige and Skoog salts (plus kanamycin 50 mg/L for the transformants) and grown to seed in soil (Levington F2, Fisons, Ipswich, UK). Lines of interest were identified by immunoblot and qPCR. For experimental study, T3 progeny seeds from selected lines were germinated on soil in controlled environment chambers at an irradiance of 130 $\mu\text{mol photons/m}^2/\text{s}$, 22 °C, relative humidity of 60%, in an 8-h/16-h square-wave photoperiod. Plants were sown randomly, and trays rotated daily. Four leaf discs (0.6 cm diameter) from two individual leaves, for the analysis of SBPase and FBPA activities, were taken and immediately plunged into liquid N_2 , and stored at -80 °C. Leaf areas were calculated using standard photography and ImageJ software (imagej.nih.gov/ij). Wild-type plants and null segregants (azygous) used in this study were initially evaluated independently. However, once it was determined that no significant difference were observed between these two groups (see figures and supplementary figures), wild-type plants and null segregants were combined (null segregants from the transgenic lines verified by PCR for nonintegration of the transgene) and used as a combined 'control' group (C).

Protein extraction and immunoblotting

Leaf discs sampled as described above were ground in liquid nitrogen. Total protein was extracted in extraction buffer (50 mM 4-(2-hydroxyethyl)piperazine-1-ethanesulphonic acid (HEPES) pH 8.2, 5 mM MgCl_2 , 1 mM ethylenediaminetetraacetic acid tetrasodium salt (EDTA), 10% glycerol, 0.1% Triton X-100, 2 mM benzamidine, 2 mM aminocaproic acid, 0.5 mM phenylmethanesulphonyl fluoride (PMSF) and 10 mM DTT), and the insoluble material was removed by centrifugation at 14 000 g for 10 min (4 °C) and protein quantification determined (Harrison *et al.*, 1998). Samples were loaded on an equal protein basis, separated using 12% (w/v) SDS-PAGE, transferred to polyvinylidene difluoride membrane and probed using antibodies raised against SBPase, FBPA and the GDC-H protein (Timm *et al.*, 2012).

Figure 8 GDC-H and GDC-H with SBPase and FBPA overexpression in *Arabidopsis* differentially impact biomass and seed yield. (a, c) Dry weight and (b, d) seed weight were determined at seed harvest. C and transgenic plants were grown in controlled environment conditions at either 130 $\mu\text{mol}/\text{m}^2/\text{s}$, 8-h light/16-h dark cycle (a, b.) or 390 $\mu\text{mol}/\text{m}^2/\text{s}$, 8-h light/16-h dark cycle (c, d.). Lines over-expressing SBPase (S), FBPA (F), GDC-H protein (H), SBPase and FBPA (SF) and SBPase, FBPA and GDC-H (SFH) are represented. The data were obtained using 10–17 individual plants from two (H), three (S, F, SF) or four (SFH) independent transgenic lines (2 H lines. See Timm *et al.*, 2012) compared to 12–13 C. Columns represent mean values and standard errors are displayed. Significant differences between lines ($P < 0.05$) are represented as capital letters indicating whether each specific line is significantly different from another. Numbers indicate % increases over C. Seed weights per plant and full statistical evaluation between groups can be seen in Figure S10.



Proteins were detected using horseradish peroxidase conjugated to the secondary antibody and ECL chemiluminescence detection reagent (Amersham, Buckinghamshire, UK). SBPase antibodies are previously characterized in Lefebvre *et al.* (2005), and FBPA antibodies were raised against a peptide from a conserved region of the protein [C]-ASIGLENTANRQAYR-amide, Cambridge Research Biochemicals, Cleveland, UK (Simkin *et al.*, 2015). In addition to the aforementioned antibodies, samples were probed using antibodies raised against the phosphoribulokinase (AS09464), ssAGPase (AS111739), purchased from Agrisera (via Newmarket Scientific, UK) and FBPase (see Lefebvre *et al.*, 2005), transketolase (Henkes *et al.*, 2001) and Rubisco (Foyer *et al.*, 1993).

Determination of SBPase activity by phosphate release

SBPase activity was determined by phosphate release. Immediately after photosynthesis measurement, leaf discs were isolated from the same leaves and frozen in liquid nitrogen. For analysis, leaf discs were ground to a fine powder in liquid nitrogen in extraction buffer (50 mM HEPES, pH 8.2; 5 mM MgCl_2 ; 1 mM EDTA; 1 mM EGTA; 10% glycerol; 0.1% Triton X-100; 2 mM benzamide; 2 mM aminocaproic acid; 0.5 mM phenylmethylsulphonyl fluoride; 10 mM dithiothreitol), and the resulting solution was centrifuged 1 min at 14 000 g , 4 $^{\circ}\text{C}$. The resulting supernatant was desalted through an NAP-10 column (Amersham) and eluted, aliquoted and stored in liquid nitrogen. For the assay, the reaction was started by adding 20 μL of extract to 80 μL of assay buffer (50 mM Tris, pH 8.2; 15 mM MgCl_2 ; 1.5 mM EDTA; 10 mM dithiothreitol; 2 mM SBP) and incubated at 25 $^{\circ}\text{C}$ for 30 min as described previously (Simkin *et al.*, 2015). The reaction was stopped by the addition of 50 μL of 1 M perchloric acid and centrifuged for 10 min at 14 000 g , 4 $^{\circ}\text{C}$. Samples (30 μL) and standards (30 μL , 0.125–4 nmol PO_4^{3-}) in triplicate were incubated 30 min at room temperature following the addition of 300 μL of Biomol Green (Affiniti Research Products, Exeter, UK), and the A_{620} was measured using a microplate reader (VERSAmax, Molecular Devices, Sunnyvale, CA).

Determination of FBPA activity

Desalted protein extracts, as described above, were evaluated for FBPA activity as described previously (Haake *et al.*, 1998).

Chlorophyll fluorescence imaging

Measurements were performed on 2-week-old *Arabidopsis* seedlings that had been grown in a controlled environment chamber providing 130 $\mu\text{mol}/\text{m}^2/\text{s}$ PPFD and ambient CO_2 . Chlorophyll fluorescence parameters were obtained using a chlorophyll fluorescence (CF) imaging system (Technologica, Colchester, UK; Barbagallo *et al.*, 2003; Baker and Rosenqvist, 2004). The operating efficiency of photosystem two (PSII) photochemistry, F_q'/F_m' , was calculated from the measurements of steady-state fluorescence in the light (F') and maximum fluorescence in the light (F_m') since $F_q'/F_m' = (F_m' - F')/F_m'$. Images of F' were taken when fluorescence was stable at 130 $\mu\text{mol}/\text{m}^2/\text{s}$ PPFD, whilst images of maximum fluorescence were obtained after a saturating 600 ms pulse of 6200 $\mu\text{mol}/\text{m}^2/\text{s}$ PPFD (Baker *et al.*, 2001; Oxborough and Baker, 1997). Parallel measurements of plants grown in high light (390 $\mu\text{mol}/\text{m}^2/\text{s}$ PPFD and ambient CO_2) were also performed on 2-week-old *Arabidopsis* (Supporting Information).

Gas exchange measurements

The response of net photosynthesis (A) to intracellular CO_2 (C_i) was measured using a portable gas exchange system (CIRAS-1, PP Systems Ltd, Ayrshire, UK). Leaves were illuminated with an integral red-blue LED light source (PP systems Ltd) attached to the gas exchange system, and light levels were maintained at saturating photosynthetic photon flux density (PPFD) of 1000 $\mu\text{mol}/\text{m}^2/\text{s}$ for the duration of the A/C_i response curve. Measurements of A were made at ambient CO_2 concentration (C_a) at 400 $\mu\text{mol}/\text{mol}$, before C_a was decreased to 300, 200, 150, 100 and 50 $\mu\text{mol}/\text{mol}$ before returning to the initial value and increased to 500, 600, 700, 800, 900, 1000, 1100 and 1200 $\mu\text{mol}/\text{mol}$. Leaf temperature and vapour pressure deficit

(VPD) were maintained at 22 °C and 1 ± 0.2 kPa, respectively. The maximum rates of Rubisco- ($V_{C_{max}}$) and the maximum rate of electron transport for RuBP regeneration (J_{max}) were determined and standardized to a leaf temperature of 25 °C based on equations from Bernacchi *et al.* (2001) and McMurtrie and Wang (1993), respectively.

Photosynthetic light response curves

A/Q response curves were measured using a CIRAS-1 portable gas exchange system (PP Systems (CIRAS-1, PP Systems Ltd). Cuvette conditions were maintained at a leaf temperature of 22 °C, relative humidity of 50%–60% and ambient growth CO₂ concentration (400 mmol/mol for plants grown in ambient conditions). Leaves were initially stabilized at saturating irradiance 1000 $\mu\text{mol}/\text{m}^2/\text{s}$, after which *A* and *g_s* were measured at the following PPFD levels: 0, 50, 100, 150, 200, 250, 300, 350, 400, 500, 600, 800 and 1000 $\mu\text{mol}/\text{m}^2/\text{s}$. Measurements were recorded after *A* reached a new steady state (1–2 min) and before stomatal conductance (*g_s*) changed to the new light levels. A/Q analyses were performed at 21% and 2% O₂.

Determination of sucrose and starch

Carbohydrates and starch were extracted from 20 mg leaf tissue, and samples were collected at two time points, 1 h before dawn (15 h into the dark period) and 1 h before sunset (7 h into the light period). Four leaf discs collected from two different leaves were ground in liquid nitrogen, and 20 mg/FW of tissue was incubated in 80% (v/v) ethanol for 20 min at 80 °C and then repeated three times with ethanol 80% (v/v) at 80 °C. The resulting solid pellet and pooled ethanol samples were freeze-dried. Suc was measured from the extracts in ethanol using an enzyme-based protocol (Stitt *et al.*, 1989), and the starch contents were estimated from the ethanol-insoluble pellet according to Stitt *et al.* (1978), with the exception that the samples were boiled for 1 h and not autoclaved.

Statistical analysis

All statistical analyses were performed by comparing ANOVA, using Sys-stat, University of Essex, UK. The differences between means were tested using the *post hoc* Tukey test (SPSS, Chicago, IL).

Acknowledgements

We thank Dr Lorna McAusland and Mr Jack Matthews for help with gas exchange, Dr Stuart Fisk for help with enzyme assays and James E Fox for help with pigment analysis. A.J.S, L.R.H and P.E.L were supported by the British Biological Sciences Research Council (BBSRC, grant: BB/J004138/1 awarded to C.A.R and T.L).

Author contributions

C.A.R. conceived this project, provided the funding and led the supervision of this research with input from T.L. A.J.S generated transgenic plants and performed molecular, biochemical and plant phenotypic analysis. L.R.H and P.E.L contributed to the generation and analysis of the transgenic plants. A.J.S and P.A.D carried out data analysis on their respective contributions. S.T and H.B generated and provided glycine decarboxylase over-expressing lines used for crosses. The manuscript was drafted by A.J.S and finalised by C.A.R. All authors reviewed and commented on the final manuscript.

Conflict of interest

The authors declare no conflict of interest.

References

- Anderson, L.E. (1971) Chloroplast and cytoplasmic enzymes. 2. Pea leaf triose phosphate isomerases. *Biochim. Biophys. Acta* **235**, 237–244.
- Baker, N.R. (2008) Chlorophyll fluorescence: a probe of photosynthesis in vivo. *Annu. Rev. Plant Biol.* **59**, 89–113.
- Baker, N.R. and Rosenqvist, E. (2004) Applications of chlorophyll fluorescence can improve crop production strategies: an examination of future possibilities. *J. Exp. Bot.* **55**, 1607–1621.
- Baker, N.R., Oxborough, K., Lawson, T. and Morison, J.I. (2001) High resolution imaging of photosynthetic activities of tissues, cells and chloroplasts in leaves. *J. Exp. Bot.* **52**, 615–621.
- Barbagallo, R.P., Oxborough, K., Pallett, K.E. and Baker, N.R. (2003) Rapid, non-invasive screening for perturbations of metabolism and plant growth using chlorophyll fluorescence imaging. *Plant Physiol.* **132**, 485–493.
- Bernacchi, C.J., Singaas, E.L., Pimentel, C., Portis, A.R. Jr and Long, S.P. (2001) Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant, Cell Environ.* **24**, 253–260.
- Betti, M., Bauwe, H., Busch, F.A., Fernie, A.R., Keech, O., Levey, M., Ort, D.R. *et al.* (2016) Manipulating photorespiration to increase plant productivity: recent advances and perspectives for crop improvement. *J. Exp. Bot.* **67**, 2977–2988.
- Bowes, G., Ogren, W.L. and Hageman, R.H. (1971) Phosphoglycolate production catalysed by ribulose diphosphate carboxylase. *Biochem. Biophys. Res. Commun.* **45**, 716–722.
- von Caemmerer, S. and Evans, J.R. (2010) Enhancing C₃ Photosynthesis. *Plant Physiol.* **154**, 589–592.
- von Caemmerer, S. and Farquhar, G.D. (1981) Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta*, **153**, 376–387.
- Cai, Z., Liu, G., Zhang, J. and Li, Y. (2014) Development of an activity-directed selection system enabled significant improvement of the carboxylation efficiency of Rubisco. *Protein Cell*, **5**, 552–562.
- Carmo-Silva, E., Scales, J.C., Madgwick, P.J. and Parry, M.A. (2015) Optimizing Rubisco and its regulation for greater resource use efficiency. *Plant, Cell Environ.* **38**, 1817–1832.
- Clough, S.J. and Bent, A.F. (1998) Floral dip: a simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**, 735–743.
- Dalal, J., Lopez, H., Vasani, N.B., Hu, Z., Swift, J.E., Yalamanchili, R., Dvora, M. *et al.* (2015) A photorespiratory bypass increases plant growth and seed yield in biofuel crop *Camelina sativa*. *Biotechnol. Biofuels* **8**, 175.
- Ding, F., Wang, M., Zhang, S. and Ai, X. (2016) Changes in SBPase activity influence photosynthetic capacity, growth, and tolerance to chilling stress in transgenic tomato plants. *Sci. Rep.* **6**, 32741.
- Eisenhut, M., Bauwe, H. and Hagemann, M. (2007) Glycine accumulation is toxic for the cyanobacterium *Synechocystis* sp. strain PCC 6803, but can be compensated by supplementation with magnesium ions. *FEMS Microbiol. Lett.* **277**, 232–237.
- Engel, N., van den Daele, K., Kolkusaoglu, U., Morgenthal, K., Weckwerth, W., Pärnik, T., Keerberg, O. *et al.* (2007) Deletion of glycine decarboxylase in *Arabidopsis* is lethal under nonphotorespiratory conditions. *Plant Physiol.* **144**, 1328–1335.
- Evans, J.R. (2013) Improving photosynthesis. *Plant Physiol.* **162**, 1780–1793.
- Fischer, R.A., Rees, D., Sayre, K.D., Lu, Z.M., Condon, A.G. and Saavedra, A.L. (1998) Wheat yield progress associated with higher stomatal conductance and photosynthetic rate, and cooler canopies. *Crop Sci.* **38**, 1467–1475.
- Foyer, F.H., Nurmi, A., Dulieu, H. and Parry, M.A.J. (1993) Analysis of two rubisco-deficient tobacco mutants, H7 and Sp25; evidence for the production of rubisco large subunits in the Sp25 mutant that form clusters and are inactive. *J. Exp. Bot.* **44**, 1445–1452.
- Gifford, R.M. and Evans, L. (1981) Photosynthesis, carbon partitioning, and yield. *Annu. Rev. Plant Physiol.* **32**, 485–509.

- Haake, V., Zrenner, R., Sonnewald, U. and Stitt, M. (1998) A moderate decrease of plastid aldolase activity inhibits photosynthesis, alters the levels of sugars and starch, and inhibits growth of potato plants. *Plant J.* **14**, 147–157.
- Haake, V., Geiger, M., Walch-Liu, P., Engels, C., Zrenner, R. and Stitt, M. (1999) Changes in aldolase activity in wild-type potato plants are important for acclimation to growth irradiance and carbon dioxide concentration, because plastid aldolase exerts control over the ambient rate of photosynthesis across a range of growth conditions. *Plant J.* **17**, 479–489.
- Harrison, E.P., Willingham, N.M., Lloyd, J.C. and Raines, C.A. (1998) Reduced sedoheptulose-1,7-bisphosphatase levels in transgenic tobacco lead to decreased photosynthetic capacity and altered carbohydrate accumulation. *Planta*, **204**, 27–36.
- Harrison, E.P., Ölçer, H., Lloyd, J.C., Long, S.P. and Raines, C.A. (2001) Cell and molecular biology, biochemistry and molecular physiology—small decreases in SBPase cause a linear decline in the apparent RuBP regeneration rate, but do not affect Rubisco carboxylation. *J. Exp. Bot.* **52**, 1779–1784.
- Heineke, D., Bykova, N., Gardeström, P. and Bauwe, H. (2001) Metabolic response of potato plants to an antisense reduction of the P-protein of glycine decarboxylase. *Planta*, **212**, 880–887.
- Henkes, S., Sonnewald, U., Badur, R., Flachmann, R. and Stitt, M. (2001) A small decrease of plastid transketolase activity in antisense tobacco transformants has dramatic effects on photosynthesis and phenylpropanoid metabolism. *Plant Cell*, **13**, 535–551.
- Kebeish, R., Niessen, M., Thiruveedhi, K., Bari, R., Hirsch, H.J., Rosenkranz, R., Stähler, N. *et al.* (2007) Chloroplastic photorespiratory bypass increases photosynthesis and biomass production in *Arabidopsis thaliana*. *Nat. Biotechnol.* **25**, 593–599.
- Kelly, G.J. and Latzko, E. (1976) Inhibition of spinach-leaf phosphofructokinase by 2-phosphoglycolate. *FEBS Lett.* **68**, 55–58.
- Kopriva, S. and Bauwe, H. (1995) H-protein of glycine decarboxylase is encoded by multigene families in *Flaveria pringlei* and *F. cronquistii* (Asteraceae). *Mol. Gen. Genet.* **248**, 111–116.
- Kossmann, J., Sonnewald, U. and Willmitzer, L. (1994) Reduction of the chloroplastic fructose-1,6-bisphosphatase in transgenic potato plants impairs photosynthesis and plant growth. *Plant J.* **6**, 637–650.
- Lawson, T., Bryant, B., Lefebvre, S., Lloyd, J.C. and Raines, C.A. (2006) Decreased SBPase activity alters growth and development in transgenic tobacco plants. *Plant, Cell Environ.* **29**, 48–58.
- Lefebvre, S., Lawson, T., Zakhleniuk, O.V., Lloyd, J.C. and Raines, C.A. (2005) Increased sedoheptulose-1,7-bisphosphatase activity in transgenic tobacco plants stimulates photosynthesis and growth from an early stage in development. *Plant Physiol.* **138**, 451–460.
- Lin, M.T., Occhialini, A., Andralojc, P.J., Parry, M.A. and Hanson, M.R. (2014a) A faster Rubisco with potential to increase photosynthesis in crops. *Nature*, **513**, 547–550.
- Lin, M.T., Occhialini, A., Andralojc, P.J., Devonshire, J., Hines, K.M., Parry, M.A. and Hanson, M.R. (2014b) β -Carboxysomal proteins assemble into highly organized structures in Nicotiana chloroplasts. *Plant J.* **79**, 1–12.
- Lin, H., Karki, S., Coe, R.A., Bagha, S., Khoshravesh, R., Balahadia, C.P., Ver Sagun, J. *et al.* (2016) Targeted knockdown of GDCH in rice leads to a photorespiratory-deficient phenotype useful as a building block for C4 rice. *Plant Cell Physiol.* **57**, 919–932.
- Long, S.P., Zhu, X.G., Naidu, S.L. and Ort, D.R. (2006) Can improvement in photosynthesis increase crop yields? *Plant, Cell Environ.* **29**, 315–330.
- Long, S.P., Marshall-Colon, A. and Zhu, X.G. (2015) Meeting the global food demand of the future by engineering crop photosynthesis and yield potential. *Cell*, **161**, 56–66.
- Lu, Y., Li, Y., Yang, Q., Zhang, Z., Chen, Y., Zhang, S. and Peng, X.X. (2014) Suppression of glycolate oxidase causes glyoxylate accumulation that inhibits photosynthesis through deactivating Rubisco in rice. *Plant Physiol.* **150**, 463–476.
- Maier, A., Fahnenstich, H., von Caemmerer, S., Engqvist, M.K., Weber, A.P., Flügge, U.I. and Maurino, V.G. (2012) Transgenic introduction of a glycolate oxidative cycle into a thaliana chloroplasts leads to growth improvement. *Front Plant Sci.* **3**, 38.
- McGrath, J.M. and Long, S.P. (2014) Can the cyanobacterial carbon-concentrating mechanism increase photosynthesis in crop species? A theoretical analysis *Plant Physiol.* **164**, 2247–2261.
- McMurtrie, R.E. and Wang, Y.P. (1993) Mathematical models of the photosynthetic response of tree stands to rising CO₂ concentrations and temperature. *Plant, Cell Environ.* **16**, 1–13.
- Meyer, M.T., McCormick, A.J. and Griffiths, H. (2016) Will an algal CO₂-concentrating mechanism work in higher plants? *Curr. Opin. Plant Biol.* **31**, 181–188.
- Miyagawa, Y., Tamoi, M. and Shigeoka, S. (2001) Over-expression of a cyanobacterial fructose-1,6/sedoheptulose-1,7-bisphosphatase in tobacco enhances photosynthesis and growth. *Nat. Biotechnol.* **19**, 965–969.
- Montgomery, B.L., Lechno-Yossef, S. and Kerfeld, C.A. (2016) Interrelated modules in cyanobacterial photosynthesis: the carbon-concentrating mechanism, photorespiration, and light perception. *J. Exp. Bot.* **67**, 2931–2940.
- Murchie, E.H. and Lawson, T. (2013) Chlorophyll fluorescence analysis: guide to good practice and understanding some new applications. *J. Exp. Bot.* **64**, 3983–3998.
- Nakagawa, T., Kurose, T., Hino, T., Tanaka, K., Kawamukai, M., Niwa, Y., Toyooka, K. *et al.* (2007) Development of series of gateway binary vectors, pGWBs, for realizing efficient construction of fusion genes for plant transformation. *J. Biosci. Bioeng.* **104**, 34–41.
- Nolke, G., Houdelet, M., Kreuzaler, F., Peterhänsel, C. and Schillberg, S. (2014) The expression of a recombinant glycolate dehydrogenase polyprotein in potato (*Solanum tuberosum*) plastids strongly enhances photosynthesis and tuber yield. *Plant Biotechnol. J.* **12**, 734–742.
- Orr, D.J., Alcantara, A., Kapralov, M.V., Andralojc, P.J., Carmo-Silva, E. and Parry, M.A. (2016) Surveying Rubisco diversity and temperature response to improve crop photosynthetic efficiency. *Plant Physiol.* **172**, 707–717. doi:10.1104/pp.16.00750
- Oxborough, K. and Baker, N.R. (1997) An instrument capable of imaging chlorophyll a fluorescence from intact leaves at very low irradiance and at cellular and subcellular levels. *Plant, Cell Environ.* **20**, 1473–1483.
- Peterhänsel, C., Krause, K., Braun, H.P., Espie, G.S., Fernie, A.R., Hanson, D.T., Keech, O. *et al.* (2013) Engineering photorespiration: current state and future possibilities. *Plant Biol.* **15**, 754–758.
- Poolman, M.G., Fell, D.A. and Thomas, S. (2000) Modelling photosynthesis and its control. *J. Exp. Bot.* **51**, 319–328.
- Raines, C.A. (2003) The Calvin cycle revisited. *Photosynth. Res.* **75**, 1–10.
- Raines, C.A. (2006) Transgenic approaches to manipulate the environmental responses of the C₃ carbon fixation cycle. *Plant, Cell Environ.* **29**, 331–339.
- Raines, C.A. (2011) Increasing photosynthetic carbon assimilation in C₃ plants to improve crop yield: current and future strategies. *Plant Physiol.* **155**, 36–42.
- Raines, C.A. and Paul, M.J. (2006) Products of leaf primary carbon metabolism modulate the developmental programme determining plant morphology. *J. Exp. Bot.* **57**, 1857–1862.
- Raines, C.A., Lloyd, J.C. and Dyer, T.A. (1999) New insights into the structure and function of sedoheptulose-1, 7-bisphosphatase; an important but neglected Calvin cycle enzyme. *J. Exp. Bot.* **50**, 1–8.
- Rojas-González, J.A., Soto-Súarez, M., García-Díaz, Á., Romero-Puertas, M.C., Sandalio, L.M., Mérida, Á., Thormählen, I. *et al.* (2015) Disruption of both chloroplastic and cytosolic FBPase genes results in a dwarf phenotype and important starch and metabolite changes in *Arabidopsis thaliana*. *J. Exp. Bot.* **66**, 2673–2689.
- Rosenthal, D.M., Locke, A.M., Khozaei, M., Raines, C.A., Long, S.P. and Ort, D.R. (2011) Over-expressing the C₃ photosynthesis cycle enzyme Sedoheptulose-1-7 Bisphosphatase improves photosynthetic carbon gain and yield under fully open air CO₂ fumigation (FACE). *BMC Plant Biol.* **11**, 123.
- Sahrawy, M., Avila, C., Chueca, A., Canovas, F.M. and Lopez-Gorge, J. (2004) Increased sucrose level and altered nitrogen metabolism in *Arabidopsis thaliana* transgenic plants expressing antisense chloroplastic fructose-1,6-bisphosphatase. *J. Exp. Bot.* **55**, 2495–2503.

- Sharwood, R.E., Ghannoum, O. and Whitney, S.M. (2016) Prospects for improving CO₂ fixation in C3-crops through understanding C4-Rubisco biogenesis and catalytic diversity. *Curr. Opin. Plant Biol.* **31**, 135–342.
- Simkin, A.J., McAusland, L., Headland, L.R., Lawson, T. and Raines, C.A. (2015) Multigene manipulation of photosynthetic carbon assimilation increases CO₂ fixation and biomass yield. *J. Exp. Bot.* **66**, 4075–4090.
- Stitt, M., Bulpin, P.V. and ap Rees, T.. (1978) Pathway of starch breakdown in photosynthetic tissues of *Pisum sativum*. *Biochim. Biophys. Acta* **544**, 200–214.
- Stitt, M., Lilley, R.M., Gerhardt, R. and Heldt, H.W. (1989) Metabolite levels in specific cells and subcellular compartments of plant tissues. *Methods Enzymol.* **174**, 518–552.
- Stockhaus, J., Schell, J. and Willmitzer, L. (1989) Correlation of the expression of the nuclear photosynthetic gene ST-LS1 with the presence of chloroplast. *EMBO J.* **8**, 2445–2451.
- Timm, S., Florian, A., Arrivault, S., Stitt, M., Fernie, A.R. and Bauwe, H. (2012) Glycine decarboxylase controls photosynthesis and plant growth. *FEBS Lett.* **586**, 3692–3697.
- Timm, S., Wittmiß, M., Gamlien, S., Ewald, R., Florian, A., Frank, M., Wirtz, M. et al. (2015) Mitochondrial dihydrolipoyl dehydrogenase activity shapes photosynthesis and photorespiration of *Arabidopsis thaliana*. *Plant Cell* **27**, 1968–1984.
- Timm, S., Florian, A., Fernie, A.R. and Bauwe, H. (2016) The regulatory interplay between photorespiration and photosynthesis. *J. Exp. Bot.* **67**, 2923–2929.
- Tolbert, N.E. (1997) The C2 oxidative photosynthetic carbon cycle. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **48**, 1–25.
- Uematsu, K., Suzuki, N., Iwamae, T., Inui, M. and Yukawa, H. (2012) Increased fructose 1,6-bisphosphate aldolase in plastids enhances growth and photosynthesis of tobacco plants. *J. Exp. Bot.* **63**, 3001–3009.
- Walker, B.J., Van Locke, A., Bernacchi, C.J. and Ort, D.R. (2015) The costs of photorespiration to food production now and in the future. *Annu. Rev. Plant Biol.* **67**, 107–129.
- Walker, B.J., South, P.F. and Ort, D.R. (2016) Physiological evidence for plasticity in glycolate/glycerate transport during photorespiration. *Photosynth. Res.* **129**, 93–103.
- Whitney, S.M., Houtz, R.L. and Alonso, H. (2011) Advancing our understanding and capacity to engineer nature's CO₂-sequestering enzyme, Rubisco. *Plant Physiol.* **155**, 27–35.
- Xin, C.P., Tholen, D., Devloo, V. and Zhu, X.G. (2015) The benefits of photorespiratory bypasses: how can they work? *Plant Physiol.* **167**, 574–585.
- Zhu, X.G., de Sturler, E. and Long, S.P. (2007) Optimizing the distribution of resources between enzymes of carbon metabolism can dramatically increase photosynthetic rate: a numerical simulation using an evolutionary algorithm. *Plant Physiol.* **145**, 513–526.
- Zhu, X.G., Long, S.P. and Ort, D.R. (2010) Improving photosynthetic efficiency for greater yield. *Annu. Rev. Plant Biol.* **61**, 235–261.

Supporting information

Additional Supporting Information may be found online in the supporting information tab for this article:

Figure S1 Schematic representation of the (a) vector pGWPTS1, (b) *A. thaliana* SBPase (PTS1SB) over-expression construct, and the (c) *A. thaliana* FBPA (PTS1-FB) over-expression construct, (d) shows the structure of a dual construct for the expression of both SBPase and FBPA (PTS1-SBFB).

Figure S2 (a) Complete data set for SBPase enzyme assays in plants analysed. (b) Complete data set for FBPA aldolase enzyme assays in plants analysed.

Figure S3 Molecular and biochemical analysis of the transgenic plants overexpressing SBPase (S), FBPA (F) or both (SF).

Figure S4 (a) The operating efficiency of PSII photochemistry of C and transgenic plants at 600 µmol/m²/s light. Capacity determined using chlorophyll fluorescence imaging. (b) the maximum carboxylation activity of Rubisco and (c) *J*_{max} were derived from *A/C_i* response curves (Figure 4).

Figure S5 Photosynthesis carbon fixation rates determined as a function of light intensity in developing leaves.

Figure S6. Complete data set for all transgenic lines evaluated. (a) leaf area at 15 days, (b) Leaf area at 20 days (c) Leaf area at 25 days.

Figure S7 Growth analysis of the transgenic and control plants grown in low light.

Figure S8 Leaf number in control and transgenic lines.

Figure S9 Complete data set for leaf area of all transgenic lines evaluated at high light (390 µmol/m²/s).

Figure S10 Complete data set for seed yield (g) from all transgenic lines evaluated in (a) low light (130 µmol/m²/s) and (b) high light (390 µmol/m²/s).